

## Role of $\alpha\beta$ and $\gamma\delta$ T Cells in the Host Response to *Salmonella* Infection as Demonstrated in T-Cell-Receptor-Deficient Mice of Defined *Ity* Genotypes

BENNETT C. WEINTRAUB,<sup>1</sup>† LARS ECKMANN,<sup>2</sup> SHARON OKAMOTO,<sup>2</sup> MARC HENSE,<sup>1</sup>‡  
STEPHEN M. HEDRICK,<sup>1</sup> AND JOSHUA FIERER<sup>2,3\*</sup>

Department of Biology, Cancer Center,<sup>1</sup> and Department of Medicine,<sup>2</sup> University of California, San Diego, La Jolla, California 92093, and Infectious Diseases Section, Department of Veterans Affairs Medical Center, San Diego, California 92161<sup>3</sup>

Received 8 November 1996/Returned for modification 13 December 1996/Accepted 29 March 1997

*Salmonella* spp. are facultative intracellular bacteria which enter the body through the intestinal tract. We studied the roles of T cells expressing either the  $\alpha$  and  $\beta$  chains or the  $\gamma$  and  $\delta$  chains of the T-cell receptor ( $\alpha\beta$  T cells or  $\gamma\delta$  T cells, respectively) in the host defense against *Salmonella* using mice genetically deficient in either  $\alpha\beta$  T cells,  $\gamma\delta$  T cells, or both T-cell subsets. These mutant strains of mice were infected orally or intraperitoneally with *Salmonella dublin*, and the progression of the disease was monitored by determining bacterial numbers in the feces, gut wall, Peyer's patches, mesenteric lymph nodes, spleen, and liver. Since susceptibility to *Salmonella* infection in mice is strongly affected by the alleles at the *Ity* locus, T-cell-mutant mice with either the *Ity*-sensitive or *Ity*-resistant phenotype were tested for resistance to *S. dublin* infection. We found that even though large numbers of intraepithelial and mucosal  $\alpha\beta$  and  $\gamma\delta$  T cells populate the normal intestine, they have no role in controlling the invasion of *S. dublin* into the intestine or the subsequent bacterial replication in the Peyer's patches or gut wall. Furthermore, systemic infections were equally severe for the first 6 days in normal,  $\alpha\beta$  T-cell-deficient, and  $\gamma\delta$  T-cell-deficient mice, and  $\alpha\beta$  but not  $\gamma\delta$  T cells were required for clearance of *S. dublin*, regardless of the *Ity* phenotype. However, mice that lacked both T-cell subsets had higher bacterial counts in their livers 15 to 18 days after infection than did  $\alpha\beta$  T-cell-deficient mice, suggesting that  $\gamma\delta$  T cells can contribute to acquired immunity to *S. dublin*.

Nontyphoid *Salmonella* spp. including *S. typhimurium*, *S. dublin*, *S. choleraesuis*, and *S. enteritidis* cause a spectrum of diseases in humans ranging from self-limited gastrointestinal infections to systemic infections with high mortality (38). *S. dublin*, which is host adapted to bovines, can cause severe systemic disease in humans, and the incidence of *S. dublin* infections has increased more than 10-fold over the last 2 decades (10). Like *S. typhimurium*, *S. dublin* efficiently infects mice, which are commonly used as a model to study the pathogenesis of the infection and the host response to *Salmonella* (16).

Multiple bacterial and host factors determine the outcome of *Salmonella* infections. An important determinant of the innate host response to *Salmonella* is the *Nramp1* gene, which is located at the *Ity/Bcg/Lsh* locus on mouse chromosome 1 (36). *Nramp1* encodes a protein that belongs to a family of integral membrane proteins with 10 highly conserved transmembrane domains (42). *Nramp1* genes which have a point mutation at nucleotide position 596 resulting in a glycine to aspartate substitution are susceptible not only to infection with nontyphoid *Salmonella* (e.g., they have the *Ity*<sup>s</sup> phenotype) but also to *Mycobacterium bovis* (BCG) and *Leishmania donovani*

(41). Macrophages from *Ity*<sup>s</sup> mice cannot efficiently kill *Salmonella* or *L. donovani* (5, 25), and in vivo *Salmonella* grow more rapidly in the livers and spleens of susceptible mice (1). *Ity*<sup>s</sup> mice are >10<sup>4</sup>-fold more susceptible than *Ity*<sup>r</sup> mice to intraperitoneal (i.p.) infection with *S. dublin* (6).

*Salmonella* enter the body through the small intestine and colon. Following penetration of the epithelial barrier, *Salmonella* are taken up by phagocytic cells in Peyer's patches (PP) and the lamina propria. Bacteria then spread to the mesenteric lymph nodes (MLN) and from there to systemic sites, most importantly the spleen and liver (3). In experimental infections, i.p. administration can be used to mimic the systemic phase of the infection, bypassing the entry of the bacteria through epithelial cells in the gut. Oral *Salmonella* infection differs in some aspects from i.p. infection, since it results in a different kinetics of bacterial counts early after infection. Furthermore, the invading bacteria depend on a set of invasion-related genes after oral infection that are not required for systemic virulence (39).

During the early, innate host response against *Salmonella*, high levels of gamma interferon (IFN- $\gamma$ ) and tumor necrosis factor alpha (TNF- $\alpha$ ) are produced (6), and in vivo neutralization of either IFN- $\gamma$  or TNF- $\alpha$  with antibodies severely compromises the host's ability to control the infection (28, 29, 34, 40). It is not clear whether T cells, especially those in the gut, i.e., intraepithelial lymphocytes (IELs) and lamina propria T cells, are important for production of these critical cytokines during the early phase of infection or if they influence the immune response to *Salmonella* through other mechanisms. IELs are particularly interesting in this regard, since they are potentially the first T cells to interact with *Salmonella* during gut invasion. IELs are in a position to play a role early in

\* Corresponding author. Mailing address: Infectious Diseases Section (111F), Department of Veterans Affairs Medical Center, 3350 La Jolla Village Dr., San Diego, CA 92161. Phone: (619) 552-7446. Fax: (619) 552-4398. E-mail: jfierer@ucsd.edu.

† Present address: Howard Hughes Medical Institute, Stanford University, Stanford, CA 94305-5428.

‡ Present address: Department of Cell Biology and Immunology, National Research Center for Biotechnology, 38124 Braunschweig, Germany.




FIG. 1. Typing mice for *Ity* genotype by using PCR. DNA from tails was amplified by PCR by using a common downstream primer and either Primer S containing the nucleotide associated with *Ity<sup>s</sup>* genotype or Primer R containing the nucleotide associated with the *Ity<sup>r</sup>* genotype. Aliquots of the PCRs were run on a 1% agarose-ethidium bromide gel. DNA from BALB/c, BALB/c.D2 *Ity<sup>r</sup>* congenics, DBA/2, and C57BL/6 mice (A) and DNA from TCR $\delta$  knockout mice that we bred (B) are shown. TCR $\delta^{-/-}$  94 mice did not yield a product and could not be typed in this round of PCR analysis. TCR $\delta^{-/-}$  87, 89, and 92 mice are *Ity<sup>r</sup>* and the rest are *Ity<sup>s</sup>*, since the presence of both bands indicates that mice are heterozygous at the *Ity* locus (TCR $\delta^{-/-}$  85, 90, 91, 93, 95, and 96), resulting in an *Ity<sup>r</sup>* phenotype.

infection because IELs have some properties of activated T cells, such as large size and expression of the T-cell activation marker CD69 (24), and because some IEL subsets produce keratinocyte growth factor (2), which may be important for wound healing and protection of epithelial surfaces. IELs in mice are composed of similar percentages of  $\alpha\beta$  and  $\gamma\delta$  T cells (37), so that either  $\alpha\beta$  or  $\gamma\delta$  IELs could affect invasion or growth of *Salmonella*.

The clearance of bacteria from tissues in the later stages of *Salmonella* infection is dependent on T cells, as athymic nude mice or T-cell-depleted mice survive the initial stages but fail to resolve the infection (17, 26, 27, 33, 35). Resolution of infection is potentially mediated by  $\alpha\beta$ ,  $\gamma\delta$ , or both subsets of T cells. Several studies have addressed the relative role of  $\alpha\beta$  and  $\gamma\delta$  T cells in *Salmonella* infection. Humans infected with *Salmonella typhi* have an increased percentage of circulating  $\gamma\delta$  T cells (15). In mice, Emoto and colleagues observed an influx of  $\gamma\delta$  T cells into the peritoneal cavity after i.p. injection of *S. choleraesuis* (7) and suggested that this influx was greater in *Ity<sup>r</sup>* mice than *Ity<sup>s</sup>* mice (8), but they did not establish in that report whether  $\gamma\delta$  T cells protect against *Salmonella* infection. Mixer et al. used antibodies to deplete *Ity<sup>s</sup>* mice of  $\alpha\beta$  T cells or  $\gamma\delta$  T cells and found a protective role for both T-cell subsets in *S. enteritidis* infection (30). In contrast, Emoto et al. found that mice genetically deficient for  $\gamma\delta$  T cells controlled *Salmonella* infection better than wild-type controls (9), but these authors did not control for the *Ity* genotype of the mice.

Because of these seemingly contradictory results, we studied the role of  $\alpha\beta$  and  $\gamma\delta$  T cells in the resistance to *S. dublin* using mutant mice which were genetically deficient for  $\alpha\beta$  T cells,  $\gamma\delta$  T cells, or both T-cell subsets and which were segregated based on their *Ity* genotype. We found that  $\alpha\beta$  T cells, but not  $\gamma\delta$  T cells, are required for resolution of *Salmonella* infection in both *Ity<sup>r</sup>* and *Ity<sup>s</sup>* mice. Neither  $\alpha\beta$  nor  $\gamma\delta$  T cells were important in limiting bacterial entry into the intestine or in slowing bacterial growth outside the intestine during the first 6 days after oral infection.

#### MATERIALS AND METHODS

**Mice.** Mutant and control mice were purchased from Jackson Laboratories (Bar Harbor, Me.). TCR $\alpha^{-/-}$  mice were typed by Southern blotting as previously described (32). TCR $\delta^{-/-}$  mice were typed by PCR as previously described (18). For *Ity* typing, tail DNA was amplified by PCR to yield a 210-bp fragment by using 10 pmol of a common 3' primer (5'-ACAGCCCGGACAGGTGGG-3') and 10 pmol of either a 5' primer specific for the *Ity<sup>s</sup>* genotype (5'-ACGCATC CCGCTGTGGGA-3') or a 5' primer specific for the *Ity<sup>r</sup>* genotype (5'-ACGCA TCCCGCTGTGGG-3'). Reaction mixtures containing 1.5 mM MgCl<sub>2</sub> were heated to 94°C for 3 min; this was followed by 27 cycles of 20 s at 94°C, 20 s at 60°C, 20 s at 72°C, and a final extension at 72°C for 7 minutes. *Ity<sup>r</sup>* TCR $\alpha^{-/-}$  and TCR $\delta^{-/-}$  mice were bred to *Ity<sup>r</sup>* 129 strain mice, and F1 mice were bred to each other or to the parental mutant strain to obtain *Ity<sup>r</sup>* mutant mice. Mice used for experiments were >6 weeks old. Mice were housed under specific pathogen-free conditions in microisolator cages with sterile food and water.

**Bacterial strains.** The following strains of *S. dublin* were used: Lane, a patient isolate; Lane pSD6, a kanamycin-resistant derivative; and LD842, a plasmid-cured derivative of Lane (4). Lane pSD6 has Tn5 inserted in the virulence plasmid outside the *spv* regulon, and it is a fully virulent *S. dublin* strain (4). After overnight culture in tryptic soy broth (Difco, Detroit, Mich.), bacteria were washed and resuspended in 0.1 M sodium bicarbonate for oral infection or 0.9% NaCl for i.p. injection. Mice were fasted overnight before oral infection (16) and were inoculated with the indicated number of *S. dublin* in a 200- $\mu$ l volume for feeding or a 100- $\mu$ l volume for i.p. injection.

**Salmonella colony counts.** After the mice were sacrificed, various organs were removed. The distal portion of the small intestine was removed and flushed of contents. The three most distal PP of the small intestine were removed as previously described (16). A 3-cm-long piece of the remaining small intestine lacking PP was used in some experiments to determine bacterial numbers in the gut wall. Feces and pieces of liver were weighed before grinding. Each organ was ground in a homogenizer (Tri-R Instruments, Rockville Center, N.Y.) and then plated at various dilutions in saline on either tryptic soy agar (Difco) or Hektoen Enteric agar (Becton Dickinson, Cockeysville, Md.) with kanamycin (20  $\mu$ g/ml). CFU were counted after incubation at 37°C overnight. Statistical analysis was performed with StatView 4.0 (Abacus Concepts, Berkeley, Calif.) by using unpaired Student's *t* tests.

**IFN- $\gamma$  ELISA.** Blood was collected from the tail vein of mice immediately prior to sacrificing. Levels of IFN- $\gamma$  in the serum were determined by enzyme-linked immunosorbent assay (ELISA) as described previously (6).

#### RESULTS

**Typing of mutant mice for *Ity* genotype.** Alleles of the *Nramp1* gene at the *Ity* locus are important for determining resistance and susceptibility to *Salmonella* infection in mice (41). Since the mice used for gene targeting were of mixed




FIG. 2. Oral infection of *Ity*<sup>-/-</sup>  $\alpha\beta$  T-cell-deficient mice and wild-type littermates. Mice were fed  $5.6 \times 10^7$  CFU of *S. dublin* Lane pSD6 and sacrificed on the indicated days after infection, and CFU were determined. Circles represent counts from individual mice and lines represent the geometric means of the mice in the groups. (○), TCR $\alpha^{-/-}$  mice; (●), wild-type littermates. Two mutant mice (†, top left graph) died during the experiment on day 12 and day 28, and *Salmonella* counts were not determined for these mice. No wild-type mice died during the experiment.

genetic background, i.e., 129 mice are *Ity*<sup>-/-</sup> and C57BL/6 (B6) mice are *Ity*<sup>+/+</sup>, we developed a PCR-based assay to determine the *Ity* genotype of the mutant mice. Since *Taq* polymerase cannot extend a primer that does not anneal to the final nucleotide, two sense PCR primers were designed which have a 3' nucleotide complementary to either the wild-type or mutant base in the *Nramp1* gene. A common antisense primer complementary to a region 210 bp downstream in the *Nramp1* gene was used to generate PCR products. By using these two primer pairs the mutant and wild-type *Nramp1* alleles could be accurately distinguished, as determined by appropriate amplification of the product associated with the *Ity*<sup>-/-</sup> but not the *Ity*<sup>+/+</sup> allele in stock B6 mice and amplification of the *Ity*<sup>+/+</sup> product but not the *Ity*<sup>-/-</sup> product in stock 129 mice. Since *Nramp2* has significant sequence homology with *Nramp1* in this region (1), it was important to establish the specificity of the primers under the conditions we used to amplify the DNA. Figure 1A shows that the appropriate primer amplified an *Nramp1* gene from each of four strains of known *Ity* phenotypes. Since the BALB/c.D2 congenic mice have the same *Nramp2* allele but different alleles of *Nramp1*, the fact that the two primers differentiate between these congenic strains supports the conclusion that the primers are specific for *Nramp1*. An example of PCR amplification of tail DNA from TCR $\delta^{-/-}$  mice is shown in Fig. 1B.

Typing of the stock TCR $\alpha^{-/-}$ , TCR $\beta^{-/-}$ , and TCR $\delta^{-/-}$  mice showed that they were all *Ity*<sup>+/+</sup>. In order to generate *Ity*<sup>-/-</sup> mutant mice, *Ity*<sup>-/-</sup> mice were back crossed to 129 mice, and the

resultant F1 mice were bred to each other or to the parental mice.

**$\alpha\beta$  T cells but not  $\gamma\delta$  T cells are required to resolve *S. dublin* infection in *Ity*<sup>-/-</sup> mice.** To examine the role of  $\alpha\beta$  T cells in controlling *Salmonella* infection, *Ity*<sup>-/-</sup>  $\alpha\beta$  T-cell-deficient mice and wild-type *Ity*<sup>+/+</sup> littermates were orally infected with *S. dublin*, and the severity of infection was monitored by determining bacterial numbers in the gut wall, PP, MLN, spleen, and liver (Fig. 2). Wild-type mice had begun to resolve the infection in the liver and spleen by day 15 p.i. and between day 15 and 30 in the MLN, whereas the bacterial counts in  $\alpha\beta$  T-cell-deficient mice continued to increase throughout the infection, and some T-cell-deficient mice died of the infection. The results of PP cultures are hard to interpret since those structures were very small in the  $\alpha\beta$  T-cell-deficient mice, making them hard to find and remove in their entirety. This result demonstrates that  $\alpha\beta$  T cells are necessary to resolve *Salmonella* infection. However, no difference was observed between mutant and control mice during the first 6 days of infection, indicating that  $\alpha\beta$  T cells are important only later in infection.

In contrast to the requirement for  $\alpha\beta$  T cells, comparison of *Ity*<sup>-/-</sup>  $\gamma\delta$  T-cell-deficient mice with *Ity*<sup>+/+</sup> wild-type littermates showed that  $\gamma\delta$  T cells are not necessary for resolving *S. dublin* infection. Throughout the 38-day experiment, no significant difference between  $\gamma\delta$  T-cell-deficient and wild-type mice was observed in the number of bacteria in any organ cultured (Fig. 3), including the intestinal wall, the contents of the terminal




FIG. 3. Oral infection of *Ity*<sup>-/-</sup>  $\gamma\delta$  T-cell-deficient mice and wild-type littermates. Mice were fed  $3.3 \times 10^7$  CFU of *S. dublin* Lane pSD6. Methods and symbols are as described in the legend for Fig. 2.  $\circ$ , *TCR\delta*<sup>-/-</sup> mice;  $\bullet$ , wild-type littermates. Two mice died during the course of the experiment, a wild-type mouse on day 10 p.i. ( $\dagger$ ) and a mutant mouse on day 20 p.i. ( $\ddagger$ ).

ileum, and the feces (data not shown). (This experiment was repeated once with the same result.)

**Role of  $\alpha\beta$  and  $\gamma\delta$  T cells in *Ity*<sup>-/-</sup> mice infected with *S. dublin*.** We then repeated the experiments using T-cell-deficient mice that were *Ity*<sup>-/-</sup>, including *TCR\gamma*<sup>-/-</sup>  $\times$  *TCR\beta*<sup>-/-</sup> mice. Since *S. dublin* Lane kills *Ity*<sup>-/-</sup> mice 6 to 8 days after oral infection, we could only study earlier time points in this experiment. The infection was equally severe in all mice, regardless of their T-cell status, on days 2 and 4 after infection. We also found no significant difference in CFU in the livers, spleens, MLN, PP, intestinal walls of the ileum, and the feces (data not shown). By day 4 the geometric means of CFU were  $5 \times 10^4$  to  $5 \times 10^5$  in the spleen and  $8 \times 10^4$  to  $2 \times 10^5$  per g of liver in wild-type and all mutant mice.

Since *Salmonella* can be detected very rapidly in the distal ileum after feeding (within 15 to 20 min) and  $\gamma\delta$  T cells in the gut could potentially limit the initial invasion of *Salmonella* into the gut wall, we sacrificed another group of mice relatively early (4 to 9 h) after oral infection to find evidence for increased intestinal invasion in the  $\gamma\delta$  T-cell-deficient mice. However, no significant difference was found in the bacterial counts in the gut wall or PP between  $\gamma\delta$  T-cell-deficient mice and wild-type littermates regardless of the *Ity* phenotype (Fig. 4). We also cultured the contents of the terminal ileum and the feces and found no differences in CFU per gram between the groups (data not shown). These results show that  $\gamma\delta$  IELs and other intestinal  $\gamma\delta$  T cells do not affect the invasion of *S. dublin* into the gut wall or PP after oral infection.

Even though there was no detectable difference in the ability of  $\gamma\delta$  T-cell-deficient mice and wild-type *Ity*<sup>-/-</sup> mice to control *Salmonella* infection (Fig. 3), nor any difference for the first 4 days in *Ity*<sup>-/-</sup> mice, it was possible that  $\gamma\delta$  T cells had a function in controlling *Salmonella* infection later in infection that was not apparent because  $\alpha\beta$  T cells compensated for their absence. To test this possibility, we compared *Ity*<sup>-/-</sup> *TCR\beta*<sup>-/-</sup> mice with *Ity*<sup>-/-</sup> *TCR\beta*<sup>-/-</sup>  $\times$  *TCR\delta*<sup>-/-</sup> mice. Since the doubly deficient mice were available only with the *Ity*<sup>-/-</sup> phenotype, we infected the mice with *S. dublin* LD842, an isogenic plasmid-cured strain that has a 50% lethal dose ( $LD_{50}$ ) of  $10^4$  in *Ity*<sup>-/-</sup> mice after i.p. infection (11). The i.p. route was chosen for these experiments to reduce mouse-to-mouse variability, and this experiment focused on the later stages of infection, which are similar for oral and i.p. infection. No significant difference among *TCR\beta*<sup>-/-</sup>  $\times$  *TCR\delta*<sup>-/-</sup>, *TCR\beta*<sup>-/-</sup>, and wild-type mice in the bacterial counts in spleens or livers was detected on day 10 p.i. in two independent experiments (data not shown). Furthermore, as expected, in each of four independent experiments, mice lacking  $\alpha\beta$  T cells or all T cells had significantly more *S. dublin* in their spleens and livers than wild-type mice after 16 to 25 days of infection (Fig. 5). However, mice with no  $\alpha\beta$  or  $\gamma\delta$  T cells had, on average, about 10-fold more *Salmonella* in their spleens and livers than  $\alpha\beta$  T-cell-deficient mice (Fig. 5). Although the two groups overlapped, the difference between the groups was significant ( $P < 0.01$ ). In addition, 2 of 24 *TCR\beta*<sup>-/-</sup>  $\times$  *TCR\delta*<sup>-/-</sup> mice died during the experiment, whereas none of the 22 *TCR\beta*<sup>-/-</sup> mice died. *Ity*<sup>-/-</sup> mice lacking






FIG. 4. *Salmonella* in the gut wall and PP early after oral infection. Mice were infected orally with *S. dublin* LD842 (2.6 × 10<sup>7</sup> CFU for *Ity<sup>R</sup>* mice and 1.3 × 10<sup>7</sup> CFU for *Ity<sup>S</sup>* mice). To determine bacterial counts in the gut wall, 3 cm of the distal end of the ileum was removed from the animal, all PP were removed from this piece, and the section was rinsed thoroughly with saline. In some experiments, gentamicin (50 μg/ml) was included in the saline rinse to kill extracellular *Salmonella* which had not invaded intestinal epithelial cells, but no differences in CFU were observed between samples rinsed with saline alone and those rinsed with saline containing gentamicin. To determine the number of bacteria in PP, the three most distal PP in the ileum were removed and rinsed thoroughly in saline. Data are individual results from 4 to 6 mice per group. ●, wild-type mice; ○, TCRδ<sup>-/-</sup> mice. The bars represent geometric means for each experimental group. Differences between mutant and wild-type mice were not significant ( $P \geq 0.4$  by Student's *t* test for all groups).

γδ T cells cleared *S. dublin* LD842 infection as well as wild-type *Ity<sup>S</sup>* mice (data not shown), demonstrating that the effect of γδ T cells was revealed because of the absence of αβ T cells, not because *Ity<sup>S</sup>* mice were used or because *S. dublin* LD842 was the infectious agent. These results suggest that γδ T cells play a role in the resolution of *Salmonella* infections, although this effect is small compared to the role of αβ T cells and is detectable only in the absence of αβ T cells.

**IFN-γ production in T-cell-deficient mice.** Since production of IFN-γ is critical for clearance of *Salmonella* (34), we tested the ability of TCRβ<sup>-/-</sup> and TCRβ<sup>-/-</sup> × TCRδ<sup>-/-</sup> mice to produce IFN-γ. αβ T-cell-deficient mice from the experiment shown in Fig. 2 produced levels of IFN-γ that were comparable to those in wild-type mice in response to *S. dublin* infection (Table 1). In both strains of mice, serum IFN-γ levels correlated with bacterial load, as was shown by Eckmann et al. (6). Furthermore, sera from both αβ T-cell-deficient and αβ and γδ T-cell-deficient mice contained high levels of IFN-γ following *S. dublin* LD842 infection (Table 1). These data show not

only that IFN-γ is produced by cells other than T cells in response to infection, but also that IFN-γ production is not sufficient to induce clearance of *Salmonella* infection in the absence of αβ T cells.

## DISCUSSION

In order to define the role of αβ and γδ T cells in the host defense against *Salmonella*, we used mice lacking αβ T cells, γδ T cells, and both αβ and γδ T cells. Several aspects of the immune response to *Salmonella* have been revealed in this study. First, the critical role of αβ T cells was demonstrated, since αβ T-cell-deficient mice were not able to resolve an oral infection with *S. dublin* regardless of their *Ity* phenotype. Second, γδ T cells do not contribute a unique function to the




FIG. 5. i.p. infection of *Ity<sup>S</sup>* αβ T-cell-deficient, αβ and γδ T-cell-deficient, and wild-type control mice. Mice were injected i.p. with 3.5 × 10<sup>3</sup> to 4.0 × 10<sup>3</sup> CFU *S. dublin* LD842 in four independent experiments. TCRβ<sup>-/-</sup> mice were from a B6 background while TCRβ<sup>-/-</sup> × TCRδ<sup>-/-</sup> mice were from a mixed B6 and 129 background. Each point represents CFU from a single mouse from one of the four experiments. Two mice, both TCRβ<sup>-/-</sup> × TCRδ<sup>-/-</sup>, died during the experiments on day 15 and day 20 p.i. and are not included in the figure. Differences between the mean log CFU in both the livers and spleens of TCRβ<sup>-/-</sup> mice compared to those of TCRβ<sup>-/-</sup> × TCRδ<sup>-/-</sup> mice 16 to 25 days after infection were significant at  $P < 0.01$  by Student's *t* test.

TABLE 1. Serum IFN- $\gamma$  levels in *S. dublin*-infected T-cell-subset-deficient mice<sup>a</sup>

<i>Ity</i> phenotype <sup>b</sup>	Mice	Days p.i.	CFU/spleen	CFU/g of liver	Serum IFN- $\gamma$ (pg/ml)
R	TCR $\alpha^{-/-}$	6	$2.5 \times 10^3$	$1.0 \times 10^4$	$600 \pm 113$
		15	$7.2 \times 10^2$	$4.6 \times 10^3$	$308 \pm 128$
	Wild-type	6	$3.4 \times 10^3$	$1.2 \times 10^4$	$965 \pm 430$
		15	$<10^1$	$1.5 \times 10^1$	$<300$
S	TCR $\beta^{-/-}$	20	$4.2 \times 10^5$	$4.9 \times 10^4$	$645 \pm 151$
		20	$2.5 \times 10^7$	$1.5 \times 10^7$	$560 \pm 282$
	Wild-type	20	$2.0 \times 10^3$	$1.7 \times 10^1$	$<250$
		20	$2.0 \times 10^3$	$1.7 \times 10^1$	$<250$

<sup>a</sup> *Ity*<sup>r</sup> mice were infected with *S. dublin* Lane as described in the legend to Fig. 2, and *Ity*<sup>s</sup> mice were infected with *S. dublin* LD842 as described in the legend for Fig. 5. CFU are the geometric means of 4 to 5 mice. Serum IFN- $\gamma$  levels from four mice in each group were determined by ELISA, and values are means  $\pm$  standard errors.

<sup>b</sup> R, resistant, S, sensitive.

immunity to *Salmonella* that cannot be compensated by  $\alpha\beta$  T cells. Thus, in mice with  $\alpha\beta$  T cells, we could not demonstrate a role for  $\gamma\delta$  T cells either early in infection when innate immunity predominates or later during the acquired immune response. However, our data suggest that  $\gamma\delta$  T cells can play a role in the later stages of *Salmonella* infection, although this effect can be detected only in the absence of  $\alpha\beta$  T cells. Taken together, this demonstrates that  $\alpha\beta$  T cells can fully compensate for the lack of  $\gamma\delta$  T cells, but the converse is true only to a very limited extent. Third, we show that neither  $\alpha\beta$  nor  $\gamma\delta$  T cells, in the intestine or in other sites, contribute to limiting bacterial entry into the intestine or bacterial growth during the early phase of *Salmonella* infection. This is further supported by the finding that *Ity*<sup>r</sup> Rag2<sup>-/-</sup> mice, which lack all B and T cells, did not show increased severity of disease during the first 6 days of infection with *S. dublin* (data not shown). Similarly, a recent study from Guilloteau and colleagues, who infected SCID mice, which also lack all B and T cells, i.p. with *S. dublin* came to the same conclusion (14).

Two recent reports addressed the role of  $\alpha\beta$  and  $\gamma\delta$  T cells in *Salmonella* infection. One concludes that both  $\alpha\beta$  and  $\gamma\delta$  T cells are protective against *Salmonella*, while the other concludes that  $\gamma\delta$  T cells increase susceptibility to *Salmonella* infection. Emoto et al. found that  $\gamma\delta$  T-cell-deficient mice are more resistant to *S. choleraesuis* infection than wild-type mice (9). However, Emoto's report contains no indication that their mice were categorized on the basis of *Ity* phenotype. Since the mice they used could have been heterozygous at the *Ity* locus, and since their groups contained small numbers of mice, variable distribution of the *Ity* phenotypes between the wild-type and  $\gamma\delta$  T-cell-deficient mice could account for their results. Other possible explanations for the difference between Emoto et al.'s results and ours, such as differences in response to *S. choleraesuis* and *S. dublin*, cannot be ruled out.

The other report, by Mixer and colleagues (30), showed that *Ity*<sup>s</sup> mice depleted of either  $\alpha\beta$  or  $\gamma\delta$  T cells by the injection of anti-T-cell-receptor antibodies are more susceptible to oral infection with *S. enteritidis*, as measured by lower LD<sub>50</sub> values determined 14 days after infection. Since we assessed resistance by measuring bacterial counts in various organs and did not determine LD<sub>50</sub>, Mixer et al.'s data could be reconciled with our findings if T-cell-deficient mice die with fewer numbers of *Salmonella* in the liver and spleen than wild-type mice. We tested this possibility and found that TCR $\beta^{-/-}$ , TCR $\delta^{-/-}$ , and wild-type mice with an *Ity*<sup>s</sup> phenotype all die with approx-

imately  $3 \times 10^7$  CFU per spleen (data not shown). Therefore, we do not favor the hypothesis that T-cell-deficient mice have a different cause of death than wild-type mice. We suspect that some of the differences in LD<sub>50</sub> in Mixer's experiments were due to a secondary effect of antibody-mediated depletion of large numbers of cells, rather than being directly due to lack of  $\alpha\beta$  or  $\gamma\delta$  T cells, but there may also be differences between the host responses to *S. enteritidis* and *S. dublin*.

Studies similar to ours using *Listeria monocytogenes*, BCG, and *Mycobacterium tuberculosis* reveal some similarities but also some interesting differences among the immune responses to different facultative intracellular bacteria. Mombaerts and colleagues (31) showed that  $\alpha\beta$  T-cell-deficient and  $\gamma\delta$  T-cell-deficient mice can resolve *Listeria* infection but that mice without any T cells (Rag-1<sup>-/-</sup> or TCR $\beta^{-/-}$   $\times$  TCR $\delta^{-/-}$ ) could not resolve infection. This shows that either  $\alpha\beta$  or  $\gamma\delta$  T cells are sufficient to clear *Listeria* infection, whereas our data show that  $\alpha\beta$  but not  $\gamma\delta$  T cells are sufficient to clear *Salmonella*. Another difference between the immune responses to *Listeria* and *Salmonella* is that the  $\alpha\beta$  T-cell response to *Listeria* primarily involves CD8<sup>+</sup> cytotoxic T cells (22), whereas CD4<sup>+</sup> helper T cells are more important for the host response to *Salmonella* (33).

The importance of  $\alpha\beta$  and  $\gamma\delta$  T cells was also studied in response to infection with BCG and *M. tuberculosis* (19, 23). In BCG infection, as in *Salmonella* infection,  $\alpha\beta$  T cells but not  $\gamma\delta$  T cells play a critical role in controlling infection. Studies with major histocompatibility class I- or class II-deficient mice show that CD4<sup>+</sup>  $\alpha\beta$  T cells are more important than CD8<sup>+</sup>  $\alpha\beta$  T cells in the clearance of BCG (21). Resistance to *M. tuberculosis*, a mycobacterium closely related to, but more virulent than BCG, is dependent upon both CD4<sup>+</sup> and CD8<sup>+</sup>  $\alpha\beta$  T cells (12, 13).  $\gamma\delta$  T-cell-deficient mice infected with *M. tuberculosis* have slightly higher CFU than wild-type mice, implying that  $\gamma\delta$  T cells are more important for clearance of *M. tuberculosis* than BCG. Together with the *Salmonella* and *Listeria* results, these findings suggest a correlation between the importance of  $\gamma\delta$  T cells and the relative importance of CD8<sup>+</sup> versus CD4<sup>+</sup>  $\alpha\beta$  T cells. One explanation for these data is that  $\gamma\delta$  T cells are able to functionally compensate for CD8<sup>+</sup>  $\alpha\beta$  T cells but not CD4<sup>+</sup>  $\alpha\beta$  T cells.

The mechanism of action of  $\alpha\beta$  and  $\gamma\delta$  T cells in host defense against *Salmonella* is poorly understood. Bacteria are eliminated in the late stages of infection mainly by activated macrophages, which are stimulated by a combination of bacterial signals, e.g., lipopolysaccharide, and T-cell-mediated signals. We show here that T-cell-deficient mice produce large amounts of IFN- $\gamma$  after *Salmonella* infection, indicating that T cells are not an important source of this cytokine during *Salmonella* infection. Presumably, IFN- $\gamma$  is produced by NK cells in *Salmonella*-infected mice, as has been proposed for early *Listeria* infection (20). However, despite the continued production of IFN- $\gamma$ , mutant mice infected with *Salmonella* had a progressive infection and eventually died. Macrophages in T-cell-deficient mice may not be able to be activated by IFN- $\gamma$  (or other T-cell cytokines), or T-cell surface ligands may be required to fully activate macrophages. Alternatively, the contribution of T cells to the immune response against *Salmonella* may also involve functions that are independent of macrophages.

#### ACKNOWLEDGMENTS

We thank Martin F. Kagnoff for helpful advice and discussion. We thank Antonis Vassiloyanakopoulos, Tomoko Yamamoto, Brian Hel-mich, and Jennifer R. Smith for excellent technical assistance, and David Schwarz for critically reading this manuscript.

This work was supported in part by National Institutes of Health grant P07DK35108 and a grant from the Research Service of the Department of Veterans Affairs. L. Eckmann is a recipient of a Career Development Award of the Crohn's and Colitis Foundation of America. M. Hense was supported by a fellowship from the German Academic Exchange Service (DAAD; HSPH).

## REFERENCES

- Benjamin, W. H., Jr., P. Hall, S. J. Roberts, and D. E. Briles. 1990. The primary effect of the *Ity* locus is on the rate of growth of *Salmonella typhimurium* that are relatively protected from killing. *J. Immunol.* **144**:3143–3151.
- Boismenu, R., and W. L. Havran. 1994. Modulation of epithelial cell growth by intraepithelial gamma delta T cells. *Science* **266**:1253–1255.
- Carter, P. B., and F. M. Collins. 1974. The route of enteric infection in normal mice. *J. Exp. Med.* **139**:1189–1203.
- Chikami, G. K., J. Fierer, and D. G. Guiney. 1985. Plasmid-mediated virulence in *Salmonella dublin* demonstrated by use of a *Tn5-orit* construct. *Infect. Immun.* **50**:420–424.
- Crocker, P. R., J. M. Blackwell, and D. J. Bradley. 1984. Expression of the natural resistance gene *Lsh* in resident liver macrophages. *Infect. Immun.* **43**:1033–1040.
- Eckmann, L., J. Fierer, and M. F. Kagnoff. 1996. Genetically resistant (*Ity*<sup>r</sup>) and susceptible (*Ity*<sup>s</sup>) congenic mouse strains show similar cytokine responses following infection with *Salmonella dublin*. *J. Immunol.* **156**:2894–2900.
- Emoto, M., H. Danbara, and Y. Yoshikai. 1992. Induction of gamma/delta T cells in murine salmonellosis by an avirulent but not by a virulent strain of *Salmonella choleraesuis*. *J. Exp. Med.* **176**:363–372.
- Emoto, M., T. Naito, R. Nakamura, and Y. Yoshikai. 1993. Different appearance of gamma delta T cells during salmonellosis between *Ity*<sup>r</sup> and *Ity*<sup>s</sup> mice. *J. Immunol.* **150**:3411–3420.
- Emoto, M., H. Nishimura, T. Sakai, K. Hiromatsu, H. Gomi, S. Itohara, and Y. Yoshikai. 1995. Mice deficient in  $\gamma\delta$  T cells are resistant to lethal infection with *Salmonella choleraesuis*. *Infect. Immun.* **63**:3736–3738.
- Fang, F. C., and J. Fierer. 1991. Human infection with *Salmonella dublin*. *Medicine (Baltimore)* **70**:198–207.
- Fierer, J., G. Chikami, L. Hatlen, E. J. Heffernan, and D. Guiney. 1988. Active immunization with LD842, a plasmid-cured strain of *Salmonella dublin*, protects mice against group D and group B *Salmonella* infection. *J. Infect. Dis.* **158**:460–463.
- Flynn, J. L., M. M. Goldstein, K. J. Triebold, and B. R. Bloom. 1993. Major histocompatibility complex class I-restricted T cells are necessary for protection against *M. tuberculosis* in mice. *Infect. Agents Dis.* **2**:259–262.
- Flynn, J. L., M. M. Goldstein, K. J. Triebold, B. Koller, and B. R. Bloom. 1992. Major histocompatibility complex class I-restricted T cells are required for resistance to *Mycobacterium tuberculosis* infection. *Proc. Natl. Acad. Sci. USA* **89**:12013–12017.
- Guilloteau, L. A., A. J. Lax, S. MacIntyre, and T. S. Wallis. 1996. The *Salmonella dublin* virulence plasmid does not modulate early T-cell responses in mice. *Infect. Immun.* **64**:222–229.
- Hara, T., Y. Mizuno, K. Takaki, H. Takada, H. Akeda, T. Aoki, M. Nagata, K. Ueda, G. Matsuzaki, Y. Yoshikai, and K. Nomoto. 1992. Predominant activation and expansion of V gamma 9-bearing gamma delta T cells in vivo as well as in vitro in *Salmonella* infection. *J. Clin. Invest.* **90**:204–210.
- Heffernan, E. J., J. Fierer, G. Chikami, and D. Guiney. 1987. Natural history of oral *Salmonella dublin* infection in BALB/c mice: effect of an 80-kilobase-pair plasmid on virulence. *J. Infect. Dis.* **155**:1254–1259.
- Hormaeche, C. E., P. Mastroeni, A. Arena, J. Uddin, and H. S. Joysey. 1990. T cells do not mediate the initial suppression of a *Salmonella* infection in the RES. *Immunology* **70**:247–250.
- Itohara, S., P. Mombaerts, J. Lafaille, J. Iacomini, A. Nelson, A. R. Clarke, M. L. Hooper, A. Farr, and S. Tonegawa. 1993. T cell receptor delta gene mutant mice: independent generation of alpha beta T cells and programmed rearrangements of gamma delta TCR genes. *Cell* **72**:337–348.
- Ladel, C. H., C. Blum, A. Dreher, K. Reifenberg, and S. H. Kaufmann. 1995. Protective role of gamma/delta T cells and alpha/beta T cells in tuberculosis. *Eur. J. Immunol.* **25**:2877–2881.
- Ladel, C. H., C. Blum, and S. H. E. Kaufmann. 1996. Control of natural killer cell-mediated innate resistance against the intracellular pathogen *Listeria monocytogenes* by  $\gamma\delta$  T lymphocytes. *Infect. Immun.* **64**:1744–1749.
- Ladel, C. H., S. Daugelat, and S. H. Kaufmann. 1995. Immune response to *Mycobacterium bovis* bacille Calmette Guerin infection in major histocompatibility complex class I- and II-deficient knock-out mice: contribution of CD4 and CD8 T cells to acquired resistance. *Eur. J. Immunol.* **25**:377–384.
- Ladel, C. H., I. E. Flesch, J. Arnoldi, and S. H. Kaufmann. 1994. Studies with MHC-deficient knock-out mice reveal impact of both MHC I- and MHC II-dependent T cell responses on *Listeria monocytogenes* infection. *J. Immunol.* **153**:3116–3122.
- Ladel, C. H., J. Hess, S. Daugelat, P. Mombaerts, S. Tonegawa, and S. H. Kaufmann. 1995. Contribution of alpha/beta and gamma/delta T lymphocytes to immunity against *Mycobacterium bovis* bacillus Calmette Guerin: studies with T cell receptor-deficient mutant mice. *Eur. J. Immunol.* **25**:838–846.
- Lin, T., G. Matsuzaki, M. Umesue, K. Omoto, H. Yoshida, M. Harada, C. Singaram, K. Hiromatsu, and K. Nomoto. 1995. Development of TCR-gamma delta CD4-CD8+ alpha alpha but not TCR-alpha beta CD4-CD8+ alpha alpha I-IEL is resistant to cyclosporin A. *J. Immunol.* **155**:4224–4230.
- Lissner, C. R., R. N. Swanson, and A. D. O'Brien. 1983. Genetic control of the innate resistance of mice to *Salmonella typhimurium*: expression of the *Ity* gene in peritoneal and splenic macrophages isolated *in vitro*. *J. Immunol.* **131**:3006–3013.
- Maier, T., and H. C. Oels. 1972. Role of the macrophage in natural resistance to salmonellosis in mice. *Infect. Immun.* **6**:438–443.
- Maskell, D. J., C. E. Hormaeche, K. A. Harrington, H. S. Joysey, and F. Y. Liew. 1987. The initial suppression of bacterial growth in a salmonella infection is mediated by a localized rather than a systemic response. *Microb. Pathog.* **2**:295–305.
- Mastroeni, P., J. N. Skepper, and C. E. Hormaeche. 1995. Effect of anti-tumor necrosis factor alpha antibodies on histopathology of primary *Salmonella* infections. *Infect. Immun.* **63**:3674–3682.
- Mastroeni, P., B. Villarreal-Ramos, and C. E. Hormaeche. 1992. Role of T cells, TNF alpha and IFN gamma in recall of immunity to oral challenge with virulent salmonellae in mice vaccinated with live attenuated *aro*<sup>-</sup> salmonella vaccines. *Microb. Pathog.* **13**:477–491.
- Mixter, P. F., V. Camerini, B. J. Stone, V. L. Miller, and M. Kronenberg. 1994. Mouse T lymphocytes that express a  $\gamma\delta$  T-cell antigen receptor contribute to resistance to *Salmonella* infection in vivo. *Infect. Immun.* **62**:4618–4621.
- Mombaerts, P., J. Arnoldi, F. Russ, S. Tonegawa, and S. H. E. Kaufmann. 1993. Different roles of alpha beta and gamma delta T cells in immunity against an intracellular bacterial pathogen. *Nature* **365**:53–56.
- Mombaerts, P., A. R. Clarke, M. A. Rudnicki, J. Iacomini, S. Itohara, J. J. Lafaille, L. Wang, Y. Ichikawa, R. Jaenisch, M. L. Hooper et al. 1992. Mutations in T-cell antigen receptor genes alpha and beta block thymocyte development at different stages. *Nature* **360**:225–231.
- Nauciel, C. 1990. Role of CD4<sup>+</sup> T cells and T-independent mechanisms in acquired resistance to *Salmonella typhimurium* infection. *J. Immunol.* **145**:1265–1269.
- Nauciel, C., and F. Espinasse-Maes. 1992. Role of gamma interferon and tumor necrosis factor alpha in resistance to *Salmonella typhimurium* infection. *Infect. Immun.* **60**:450–454.
- O'Brien, A. D., and E. S. Metcalf. 1982. Control of early *Salmonella typhimurium* growth in innately *Salmonella*-resistant mice does not require functional T lymphocytes. *J. Immunol.* **129**:1349–1351.
- Plant, J., and A. A. Glynn. 1979. Locating *Salmonella* resistance gene on mouse chromosome 1. *Clin. Exp. Immunol.* **37**:1–6.
- Raulet, D. H. 1989. The structure, function, and molecular genetics of the gamma/delta T cell receptor. *Annu. Rev. Immunol.* **7**:175–207.
- Saphra, I., and J. W. Winter. 1957. Clinical manifestations of salmonellosis in man. An evaluation of 7779 human infections identified at the New York Salmonella Center. *N. Engl. J. Med.* **256**:1128–1134.
- Stein, M. A., S. D. Mills, and B. B. Finlay. 1994. *Salmonella*: now you see it, now you don't. *Bioessays* **16**:537–538.
- Tite, J. P., G. Dougan, and S. N. Chatfield. 1991. The involvement of tumor necrosis factor in immunity to *Salmonella* infection. *J. Immunol.* **147**:3161–3164.
- Vidal, S., M. L. Tremblay, G. Govoni, S. Gauthier, G. Sebastiani, D. Malo, E. Skamene, M. Olivier, S. Jothy, and P. Gros. 1995. The *Ity/Lsh/Bcg* locus: natural resistance to infection with intracellular parasites is abrogated by disruption of the *Nramp1* gene. *J. Exp. Med.* **182**:655–666.
- Vidal, S. M., D. Malo, K. Vogan, E. Skamene, and P. Gros. 1993. Natural resistance to infection with intracellular parasites: isolation of a candidate for *Bcg*. *Cell* **73**:469–485.