

Fluid management in diabetic ketoacidosis

C D Inward, T L Chambers

Have we got it right yet?

Young people with insulin dependent diabetes mellitus are three times more likely to die in childhood than the general population.¹ Despite advances in management over the past 20 years, the incidence of mortality associated with diabetic ketoacidosis (DKA) remains unchanged. Cerebral oedema is the predominant cause of this mortality; young children are particularly at risk, with an incidence of 0.7–1% of episodes of DKA.^{2,3} The mortality appears to be greatest among patients at first presentation,^{1,3,4} if there has been a long history of symptoms prior to admission,³ and during the first 24 hours of treatment.⁴ In a recently published retrospective multicentre analysis of children with DKA, low p_{CO_2} levels and high serum sodium concentration at presentation were identified as particular risk factors for the development of cerebral oedema, together with bicarbonate therapy.⁵ However, in the accompanying editorial, Dunger and Edge point out that this may simply be revealing an association between severe DKA and dehydration and the risk of cerebral oedema.⁶ The pathogenesis of cerebral oedema remains poorly understood but there may be many contributing factors.⁷

The aim of management of DKA is to restore metabolic homeostasis while minimising the risks of complications including hypoglycaemia, hypokalaemia, cardiac failure, and in children the development of cerebral oedema. How best to achieve this remains contentious, with particular controversy centred on optimal fluid management. The most appropriate volume, type, and rate of fluid to be given have all been the subject of debate. A survey in 1994 of UK paediatricians found a threefold variation in the amount of fluid recommended within the first 12 hours.⁸ Since then national guidelines have been developed by the British Society for Paediatric Endocrinology (BSPE). However, it remains uncertain whether the recommended strategy, using a volume calculated by “maintenance plus deficit”, is the safest and most appropriate way to manage rehydration in children with DKA. We suggest that childhood DKA

may be considered and treated as a variant of renal electrolyte disorder with hypertonic dehydration.

FLUID AND ELECTROLYTE LOSSES

The fluid and electrolyte losses of DKA are predominantly caused by hyperglycaemia with resultant glycosuria and osmotic diuresis. In addition, the kidney has a low threshold for ketoacids, which are excreted into the urine with an accompanying cation, further exacerbating the electrolyte loss. Relative acute dehydration with a larger sodium loss will result primarily in intravascular dehydration, but a more prolonged development will result in a greater degree of intracellular dehydration, with relative maintenance of intravascular volume, particularly if the patient remains polydipsic.⁹ There are thus wide variations in fluid, acid–base, and electrolyte deficiencies, during both development and treatment of DKA, yet few protocols acknowledge this.

There is clear evidence that volume depletion triggers the release of counter regulatory hormones including catecholamines, growth hormone, and cortisol as well as renin, aldosterone, and arginine vasopressin.^{10,11} These hormones, whose combined actions are directed towards preserving intravascular volume, also cause insulin resistance. An important effect of fluid administration is therefore reducing the stimulus for the secretion of the counter regulatory hormones. Indeed it has been shown that plasma glucose starts to fall following fluid administration prior to giving insulin, although insulin is required for correction of the metabolic acidosis.¹²

RISK FACTORS FOR CEREBRAL OEDEMA

The importance of prompt restoration of the circulating volume for patients presenting with DKA is clear. However, the observation that symptomatic cerebral oedema often occurs several hours after treatment has begun, usually at a time when plasma glucose has started to fall and metabolic acidosis to improve, has led to concern that the treatment itself may contribute to the development of cerebral oedema. There have been few

case controlled studies of cerebral oedema complicating DKA and there is conflicting evidence as to the role of iatrogenic factors in fatal cases.^{6,13} However, excessive rates of fluid administration, particularly early in resuscitation,¹⁴ and failure of plasma sodium to rise as plasma glucose declines, have been identified as risk factors.

Studies using computed tomography suggest that subclinical cerebral swelling may be a common occurrence in children presenting with DKA,^{15,16} although this has been disputed.¹⁷ If cerebral swelling is a relatively common event it is not clear why only a minority progress to herniation. In addition, fatal herniation has occurred in the absence of any fluid therapy. Therefore, although clearly not the only cause, it seems reasonable to conclude that fluid management may be a factor contributing to the development of overt cerebral oedema. On the basis of the available evidence it would seem prudent to avoid rapid rates of fluid infusion and to maintain serum sodium concentrations in order to avoid the potential danger of excess free water administration which may exacerbate the development of symptomatic brain swelling.

Failure to follow guidelines for the management of DKA has been well documented.¹⁸ Of particular concern is the evidence that clinicians have been consistently shown to overestimate the degree of dehydration in this condition.^{19,20} Nevertheless, the assumption underlying most DKA protocols is that rehydration is desirable and achievable in 24 hours, using a volume calculated by “maintenance plus deficit”.

THE NEED FOR CHANGE?

From a renal perspective the term “maintenance fluid” is a misnomer. True maintenance fluid volume in children is the sum of the insensible losses plus the obligate urine production and is probably only 40% of recommended amounts.²¹ Water requirements are related to energy expenditure. The recommended “maintenance” volumes are based on calculations of insensible losses (estimated as 50 ml/100 cal/day) plus 66.7 ml/100 cal/day to replace urine output.²² This allows for production of isotonic urine. As normally functioning kidneys are capable of fourfold concentration of the glomerular filtrate, “maintenance fluids” will provide sufficient volume for both insensible and obligate losses as well as the extra needed for rehydration. A 5% dehydrated child with normally functioning kidneys can be rehydrated using “maintenance” fluid only—albeit slowly.

In non-diabetic dehydration it is customary to assess whether the dehydration is hypertonic, hypotonic, or isotonic

and whether there is simple contraction of the extracellular compartment or whether compensatory mechanisms are operating.

It is accepted that safe therapy of hypertonic dehydration should be both prolonged and cautious to avoid the development of fits and cerebral oedema during rehydration.²³⁻²⁴

The cells of the central nervous system protect cell volume under hyperosmolar conditions by producing intracellular osmotically active molecules, "idiogenic osmoles".²⁵ Once formed these molecules dissipate slowly. Rapid reduction of plasma osmolality by the administration of free water under these conditions will create a gradient, causing movement of water into brain cells and thus brain swelling. These considerations probably apply equally to hypernatraemic dehydration and diabetic ketoacidosis.

We suggest that the treatment of dehydration of severe DKA should be based on the general principles used to treat dehydration combined with renal failure, taking into account the additional relatively brief¹⁹ problem of an osmotic diuresis. Halting the decompensation process is the initial therapeutic goal, followed by provision of sufficient fluid to allow the establishment of renal and hormonal homeostasis over a period of time comparable to that taken for the problem to develop. There is no evidence that the use of modest fluid regimens is harmful.²⁶⁻²⁷ Harris and colleagues¹⁹⁻²⁸ have shown that in 149 episodes of childhood DKA, rehydration over 48 hours was associated with "no deaths or near death episodes", and Johnston reported that in adults a regimen of 48-72 ml/kg/day was safe.²⁹

A PROPOSAL

Twenty or so years ago there was a revolution in the management of DKA with the introduction of continuous intravenous insulin infusion. Perhaps it is now time for another review, not only of the fluid management, but also of all other aspects. There are several interested parties—endocrinologists, nephrologists, general paediatricians, and specialists in emergency medicine. We wonder whether a summit meeting should be convened to consider this; perhaps to commission research (the NHS and the corporate spirit of British paediatrics are ideal for such multicentre studies) and propose a second revolution in DKA management. Who will step forward?

ACKNOWLEDGEMENTS

We would like to thank Dr Fiona Cowan, Dr Mary McGraw, and Mrs Rachel Owen for their invaluable help in the preparation of the manuscript.

Arch Dis Child 2002;**86**:443-445

Authors' affiliations

C D Inward, T L Chambers, Bristol Royal Hospital for Children, Bristol, UK

Correspondence to: Dr C D Inward, Bristol Royal Hospital for Children, Upper Maudlin Street, Bristol BS2 8BJ, UK

REFERENCES

- Edge JA, Ford-Adams ME, Dunger DB. Causes of death in children with insulin dependent diabetes 1990-96. *Arch Dis Child* 1999;**81**:318-23.
- Edge JA, Hawkins MM, Winter DL, et al. Incidence presentation management and outcome of cerebral oedema associated with diabetic ketoacidosis (DKA) in Great Britain. *Arch Dis Child* 1999;**80**(suppl 1):A11.
- Bello FA, Sotos JF. Cerebral oedema in diabetic ketoacidosis in children. *Lancet* 1990;**336**:64.
- Rosenbloom AL. Intracerebral crises during treatment of diabetic ketoacidosis. *Diabetes Care* 1990;**13**:22-33.
- Glaser N, Barnett P, McCaslin I, et al. Risk factors for cerebral edema in children with diabetic ketoacidosis. *N Engl J Med* 2001;**344**:264-9.
- Dunger DB, Edge JA. Predicting cerebral oedema during diabetic ketoacidosis. *N Engl J Med* 2001;**344**:302-3.
- Edge JA. Cerebral oedema during treatment of diabetic ketoacidosis: are we any nearer finding a cause? *Diabetes Metab Res Rev* 2000;**16**:316-24.
- Edge JA, Dunger DB. Variations in the management of diabetic ketoacidosis in children. *Diabet Med* 1994;**11**:984-6.
- Adrogué HJ, Wilson H, Boyd AE, et al. Plasma acid-base patterns in diabetic ketoacidosis. *N Engl J Med* 1982;**307**:1603-10.
- Tulassay T, Rascher W, Korner A, et al. Atrial natriuretic peptide and other vasoactive hormones during treatment of severe diabetic ketoacidosis in children. *J Pediatr* 1987;**111**:329-34.
- Luzi L, Barrett EJ, Groop LC, et al. Metabolic effects of low-dose insulin therapy on glucose metabolism in diabetic ketoacidosis. *Diabetes* 1988;**37**:1470-7.
- Owen OE, Licht JH, Sapir DG. Renal function and effects of partial rehydration during diabetic ketoacidosis. *Diabetes* 1981;**30**:510-18.
- Mahoney CP, Vleck BW, DelAguila M. Risk factors for developing brain herniation during diabetic ketoacidosis. *Pediatr Neurol* 1999;**21**:721-7.
- Rosenbloom AL, Schatz DA, Krischer JP, et al. Therapeutic controversy: prevention and treatment of diabetes in children. *J Clin Endocrinol Metab* 2000;**85**:494-522.
- Krane EJ, Rockoff MA, Wallman JK, et al. Subclinical brain swelling in children during treatment of diabetic ketoacidosis. *N Engl J Med* 1985;**312**:1147-51.
- Hoffman WH, Steinhart CM, ElGammal T, et al. Cranial CT in children and adolescents with diabetic ketoacidosis. *Am J Neuroradiol* 1988;**9**:733-9.
- Smedman L, Escobar R, Hesser U, et al. Subclinical cerebral oedema does not occur regularly during treatment for diabetic ketoacidosis. *Acta Paediatr* 1997;**86**:1172-6.
- Singh RK, Perros P, Frier BM. Hospital management of diabetic ketoacidosis: are clinical guidelines implemented effectively? *Diabet Med* 1997;**14**:482-6.
- Harris GD, Fiordalisi I, Harris WL, et al. Minimizing the risk of brain herniation during treatment of diabetic ketoacidemia: a retrospective and prospective study. *J Pediatr* 1990;**117**:22-31.
- Grove LM, Nobel-Jamieson CM, Barnes ND, et al. Assessment of dehydration, fluid balance and insulin requirements in diabetic ketoacidosis. *Proc Br Paediatr Assoc Annu Meet* 1995;**67**:26.

- Molteni KH. Initial management of hypernatraemic dehydration in the breastfed infant. *Clin Pediatr* 1994;**33**:731-40.
- Holliday MA, Segar WE. The maintenance need for water in parenteral fluid therapy. *Pediatrics* 1957;**19**:823-32.
- Holliday M. The evolution of therapy for dehydration: should deficit therapy still be taught? *Pediatrics* 1996;**98**:171-7.
- Oddie S, Richmond S, Coulter M. Hypernatraemic dehydration and breast feeding: a population study. *Arch Dis Child* 2001;**85**:318-20.
- Lee JH, Arcinue E, Ross BD. Organic osmolytes in the brain of an infant with hypernatremia. *N Engl J Med* 1994;**331**:439-42.
- Lebovitz HA. Diabetic ketoacidosis. *Lancet* 1995;**345**:767-71.
- Adrogué H, Barrero J, Eknoyan G. Salutary effects of modest fluid replacement in the treatment of adults with ketoacidosis. *JAMA* 1989;**262**:2108-13.
- Harris GD, Fiordalisi I. Physiologic management of diabetic ketoacidemia. A 5-year prospective pediatric experience in 231 episodes. *Arch Pediatr Adolesc Med* 1994;**148**:1046-52.
- Johnston C. Fluid replacement in diabetic ketoacidosis. *BMJ* 1992;**305**:522.

COMMENTARY

The treatment of the child with diabetic ketoacidosis (DKA) is unfortunately not based on very much evidence, but rather on anecdote and precedent. Fluid management has developed over several decades, with changes occurring gradually in an attempt to avoid complications (initially hypokalaemia, more recently cerebral oedema), rather than a logical approach to the particular fluid and electrolyte deficits. This has resulted in some anomalies in the management of DKA, and Inward and Chambers have now opened the debate about whether a revolution in fluid and electrolyte management is necessary.

DKA is associated with severe fluid losses. Much of the information about the specific fluid and electrolyte deficits in DKA was obtained in the 1930s from work on two young men with diabetes whose fluid and electrolyte balance was studied in great detail before, while, and after they were allowed experimentally to become ill with DKA.¹ The reported fluid deficit appeared to be approximately half from the intracellular and half from the extracellular compartments.¹⁻³ This work formed the basis for most of the DKA management protocols in existence.^{3,4} For obvious reasons, these studies have never been repeated in children, and yet body composition and fluid requirements are very different from those in adults.

Moreover, as Inward and Chambers point out, factors such as the duration of the episode of DKA will produce wide variations in fluid and electrolyte deficiencies. The water deficit is made up of varying combinations from the osmotic