

# Comparing tobacco use among incoming recruits and military personnel on active duty in the United States

Michael C Chisick, Forrest R Poindexter, Andrew K York

## Abstract

**Objective**—To compare the tobacco use profile of recruits with that of military personnel on active duty to determine whether the military environment in some way induces service members to initiate tobacco use.

**Design and setting**—Cross-sectional survey of United States armed forces active duty and recruit personnel in 1994–95.

**Subjects**—2711 military recruits and 4603 military personnel on active duty.

**Main outcome measures**—Comparative cigarette smoking and smokeless tobacco use prevalence between recruits and personnel on active duty controlling for age, sex, and race. Impact of demographic factors on the odds of smoking or using smokeless tobacco.

**Results**—Increases in tobacco use in American military personnel occurred exclusively in men. The highest tobacco use resided with white men on active duty (43% cigarette smoking; 24% smokeless tobacco use) and represents a doubling of tobacco use seen among white male recruits. Among non-white men, tobacco use increased 2–4 times between recruits and personnel on active duty.

**Conclusions**—Efforts to reduce tobacco use by American military personnel on active duty should focus more on discouraging the initiation of tobacco use.

(*Tobacco Control* 1998;7:236–240)

Keywords: tobacco use, military personnel, United States

## Introduction

Historically, tobacco use by American military personnel on active duty has greatly exceeded that found in the civilian population. Roughly two decades ago, the first Department of Defense (DoD) survey of health-related behaviours reported an overall cigarette smoking rate of 51% for military personnel on active duty. According to the most recent DoD survey (1995), the overall cigarette smoking rate for service members on active duty had declined to 33%. This decline has been attributed to an increased emphasis on smoking cessation and prevention in the military.<sup>1</sup>

Comparing the cigarette smoking rate among service members on active duty to that among the civilian population requires standardising the data to control for sociodemographic differences between the two

populations.<sup>2</sup> When this is done, overall cigarette smoking prevalence in the military population (33%) appears to compare favourably with the civilian population (31%).<sup>1</sup> However, overall population comparisons can obscure important differences between subgroups within those populations. For example, results from the 1995 DoD survey also show that, depending on branch of service, 33–45% of 18–25 year old military personnel on active duty smoke cigarettes, compared with 36% of similarly aged civilians.<sup>1</sup>

One limitation of the DoD surveys is that their samples did not include military recruits, that is, individuals newly ascended into the military from civilian life. It would be unwise to assume that military recruits possess traits similar to those found in civilian tobacco surveys. Military recruits are a self-selected subgroup that differ from the general civilian population in many ways. For one, they are generally healthier, having met physical fitness standards for entry into military service. In addition, the services prefer to recruit individuals with at least a high-school education, yet few college-bound prospects enter the military for enlisted service.

Given the potentially different demographic profile of military recruits and the general civilian population, it is unknown how the tobacco use profile of military recruits compares with that of service members on active duty. Knowing the comparative tobacco use profile of military recruits versus personnel on active duty is important for determining whether the existing tobacco use profile of personnel on active duty is merely a reflection of the tobacco use behaviours that recruits bring with them when they enter the military, or whether events subsequent to enlistment play a role. Such knowledge will give some indication of whether exposure to the military environment in some way induces service members to begin using tobacco. The military may well be contributing to its tobacco use problem by its co-existing, contradictory policies on tobacco. Despite DoD-wide smoking cessation programmes, smoking bans in the workplace, and bans on tobacco use during recruit basic training, tobacco products are still readily available in military stores at a substantial price discount compared with the civilian community. Knowing the comparative tobacco use profile of military recruits versus personnel on active duty will also help determine whether current

US Army Center for Health Promotion and Preventive Medicine, Aberdeen Proving Ground, Maryland, USA  
M C Chisick

Bolling Air Force Base, Washington, DC  
F R Poindexter

Naval Dental Research Institute-Detachment, Bethesda, Maryland  
A K York

Correspondence to:  
LTC Mike Chisick, 2422 Bibury Lane, Apartment 203, Baltimore, Maryland 21244–1993, USA.  
LTC\_Mike\_Chisick@chppm-ccmail.apgea.army.mil

Table 1 Unweighted distribution of recruit (R) and active duty (AD) samples, by age, sex, and race

Age (years)	Whites		Blacks		Others	
	R	AD	R	AD	R	AD
<b>Men</b>						
18-19	701	157	198	35	118	18
20-24	566	2128	190	502	116	286
25-29	56	459	17	127	15	74
30-34	10	71	4	26	6	22
Total	1333	2815	409	690	255	400
<b>Women</b>						
18-19	221	24	60	8	48	2
20-24	216	311	67	138	34	46
25-29	36	87	13	42	2	15
30-34	8	15	6	7	3	3
Total	481	437	146	195	87	66

military tobacco control policies and programmes, which emphasise tobacco cessation, are adequate or whether a more expanded strategy, including tobacco interdiction, is indicated.

Only one previous study has explored the comparative tobacco use profiles of military recruits and personnel on active duty. This 1988 survey of male navy personnel<sup>3</sup> has several serious design limitations. First, the authors did not adequately describe their sampling strategy. They merely described the source and size of their recruit and active duty samples, but not how either was selected. Nor did the authors mention non-response. It is unclear whether either the recruit or active duty samples were random and representative of the populations from which they were drawn or whether they were convenience samples. Second, the authors chose an inappropriate comparison sample by including officers and by including enlisted personnel beyond the age range of recruits in its active duty sample. Third, the age and race composition of the comparison active duty sample was not standardised to that of the recruit sample so that overall smoking prevalence rates between the two groups could be directly compared. These limitations could introduce serious bias into the findings.

Our study overcomes these limitations and expands the scope of the research to include smokeless tobacco (SLT), service members from the other armed services beyond the navy, and women.

## Methods

The data for this study come from the 1994 Tri-Service Comprehensive Oral Health Survey, a 30-site, cross-sectional survey of American army, navy, marine, and air force active duty and recruit personnel.<sup>4</sup> We drew the recruit sample (n = 2711) at recruit in-processing centres using stratified, systematic random sampling. Personnel on active duty (n = 15 915) were selected using stratified, multistage random sampling and were surveyed at military dental clinics. Women and minorities were oversampled. All of the recruit sample and 82% of the active duty sample (n = 13 050) participated in the survey.

All participants, whether recruits or personnel on active duty, completed a self-administered, electronic questionnaire<sup>5</sup> that included queries on cigarette smoking and SLT use. For both groups, the questions were worded identically and were administered in an identical format. We defined a cigarette smoker as someone who answered "yes" to the question: "Do you smoke cigarettes now?". We defined a SLT user as someone who answered "yes" to either of the following questions: "Do you use snuff now?" and "Do you use chewing tobacco now?". Data were collected on the recruit sample between February and July 1994 during in-processing, not during basic training. In-processing occurs during the first three days following the recruit's arrival at a basic training centre. Data were collected on the active duty sample between April 1994 and January 1995.

To make the two groups comparable in demographic composition, we removed from the active duty sample all officers, all personnel over 34 years of age, and all personnel with greater than five years of military service. This created a final active duty sample of 4603. A comparison of the two unweighted samples stratified by age, sex, and race is presented in table 1.

We did two types of analysis. First, we compared tobacco use between active duty and recruit military personnel after stratifying the weighted samples by age, sex, and race. We weighted the data to reflect the respective total populations of 18-34 year old recruits and personnel on active duty—101 072 and 711 534, respectively. We calculated point estimates and 95% confidence intervals (CIs). For all

Table 2 Proportion of 18-34 year old recruits (R) and military personnel on active duty (AD) who smoke, by age, sex, and race

Age (years)	Whites				Blacks				Others			
	R		AD		R		AD		R		AD	
	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI
<b>Men</b>												
18-19	20.3	17.3-23.3	42.0	34.3-49.7	4.5	1.6-7.4	25.9	11.4-40.4	9.3	4.1-14.5	28.6	7.7-49.5
20-24	21.4	18.0-24.8	36.8	34.8-38.8	7.0	3.4-10.6	15.9	7.9-23.9	14.8	8.3-21.3	24.5	19.5-29.5
25-29	25.8	14.4-37.2	32.3	28.0-36.6	0	0	15.0	8.8-21.2	9.8	-5.2-24.8	25.0	15.1-34.9
30-34	*	*	43.5	32-55.0	*	*	23.0	6.9-39.1	*	*	21.4	4.3-38.5
Total	21.1	18.9-23.3	42.8	41.0-44.6	5.4	3.2-7.6	20.4	17.4-23.4	11.5	7.6-15.4	26.4	22.1-30.7
<b>Women</b>												
18-19	28.7	22.7-34.7	15.6	31.3-30.1	3.3	-1.2-7.8	*	*	28.8	16.0-41.6	*	*
20-24	24.6	18.9-30.3	30.8	25.7-35.9	13.2	5.1-21.3	6.8	2.6-11.0	13.4	2.0-24.8	15.4	5.0-25.8
25-29	26.8	12.3-41.3	38.5	28.3-48.7	*	*	17.9	6.3-29.5	*	*	41.8	16.8-66.8
30-34	*	*	12.2	-4.4-28.8	*	*	*	*	*	*	*	*
Total	26.3	22.4-30.2	24.0	20.0-28.0	9.9	5.1-14.7	6.8	3.3-10.3	26.2	17.0-35.4	36.0	24.4-47.6

\*Insufficient sample size for stable estimate.  
CI = confidence intervals.

Table 3 Proportion of 18–34 year old recruits (R) and military personnel on active duty (AD) who use smokeless tobacco, by age, sex, and race

Age (years)	Whites				Blacks				Others			
	R		AD		R		AD		R		AD	
	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI	(%)	95% CI
<b>Men</b>												
18–19	11.7	9.3–14.1	23.0	16.4–29.6	0	0	0.7	–2.1–3.5	2.3	–0.4–5.0	9.4	–4.1–22.9
20–24	10.6	8.1–13.1	22.9	21.1–24.7	0	0	1.6	0.5–2.7	2.7	–0.2–5.6	8.3	5.1–11.5
25–29	3.4	–1.3–8.1	19.6	16.0–23.2	8.9	–4.6–22.4	3.2	0.1–6.3	0	0	8.1	1.9–14.3
30–34	*	*	19.8	10.5–29.1	*	*	4.4	–3.5–12.3	*	*	0	0
Total	10.9	9.2–12.6	24.4	22.8–26.0	0.5	–0.2–1.2	1.3	0.5–2.1	2.2	0.4–4.0	8.6	5.9–11.3
<b>Women</b>												
18–19	0.9	–0.3–2.1	0	0	0	0	*	*	0	0	*	*
20–24	0	0	0.9	–0.1–1.9	0	0	0.8	–0.6–2.2	0	0	0	0
25–29	2.6	–2.6–7.8	0.9	–1.1–2.9	*	*	0	0	*	*	0	0
30–34	*	*	6.4	–6.0–18.8	*	*	*	*	*	*	*	*
Total	0.6	–0.1–1.3	0.6	–0.1–1.3	2.4	–0.1–4.9	0.4	–0.5	0	0	0	0

\*Insufficient sample size for stable estimate.  
CI = confidence intervals.

stratified analyses, estimates for cells containing fewer than 15 observations were excluded because the size of these subsamples was too small to provide stable estimates. To compare tobacco use across all age levels for each sex and race subgroup between the two populations, we standardised the demographic profile of the population on active duty to that of the recruit population.<sup>2</sup>

For our second type of analysis, we conducted backwards, stepwise logistic regression analyses on the unweighted samples, using the regression strategy advocated by Hosmer and Lemshov.<sup>6</sup> We performed all analyses using Stata statistical software.

## Results

Results of our post-stratification analyses for cigarette smoking and SLT use are presented in tables 2 and 3. The highest levels of smoking (42.8%) and SLT use (24.4%) among 18–34 year old enlisted military personnel reside with white men on active duty, and represent a doubling of tobacco use found among white male recruits. Between recruits and personnel on active duty, smoking prevalence doubled for non-white/non-black men from 11.5% to 26.4%, quadrupled for black men from 5.4% to 20.4%, but remained unchanged for women of all races at roughly 25% for whites and non-whites/non-blacks and 10% for blacks.

Table 4 Proportion of 18–34 year old recruits and military personnel on active duty who smoke cigarettes or use smokeless tobacco, by branch of service

Service	Smoking						Smokeless tobacco (men only)					
	Recruits			Active duty			Recruits			Active duty		
	n	(%)	95% CI	n	(%)	95% CI	n	(%)	95% CI	n	(%)	95% CI
Army	535	16.0	12.9–19.1	2002	34.2	32.1–36.3	373	6.5	4.0–9.0	1764	19.8	18.0–21.6
Air force	751	6.3	4.6–8.0	1261	25.9	23.5–28.3	563	4.5	2.8–7.2	1008	12.5	10.5–15.5
Marines	716	19.0	16.1–21.9	243	39.8	33.6–46.0	648	10.4	6.5–14.3	234	21.0	17.9–24.1
Navy	709	27.4	24.3–30.7	1097	33.9	31.1–36.7	413	9.4	6.6–12.2	899	13.2	11.0–15.4

Table 5 Statistically significant adjusted odds ratios for smoking by 18–34 year old recruits and military personnel on active duty

	Recruits				Active duty			
	Men		Women		Men		Women	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<b>Race</b>								
White	1	NA	1	NA	1	NA	1	NA
Black	0.23	0.14–0.36	0.28	0.15–0.54	0.33	0.26–0.41	0.24	0.14–0.40
Hispanic	NA	NA	NA	NA	0.48	0.35–0.65	NA	NA
Asians	NA	NA	NA	NA	1.41	0.84–2.38	NA	NA
Others	0.56	0.37–0.84	0.70	0.39–1.26	0.82	0.47–1.43	0.55	0.29–1.04
<b>Service</b>								
Army	1	NA	1	NA	1	NA	1	NA
Air force	0.28	0.18–0.46	0.28	0.14–0.55	0.58	0.48–0.70	0.61	0.39–0.96
Marines	1.16	0.81–1.66	0.62	0.28–1.36	0.69	0.50–0.94	0.53	0.33–0.86
Navy	1.69	1.16–2.46	1.65	1.04–2.64	0.67	0.56–0.81	0.53	0.33–0.86
<b>Education years</b>								
<12	1	NA	1	NA	1	NA	1*	NA
12	0.56	0.25–1.18	0.43	0.10–1.76	1.03	0.56–1.88	1	NA
13–15	0.33	0.15–0.72	0.37	0.09–1.52	0.63	0.34–1.16	0.85	0.57–1.27
≥16	0.20	0.07–0.57	0.27	0.05–1.50	0.43	0.21–0.89	0.68	0.32–1.48
<b>Years of service</b>								
1	NA	NA	NA	NA	1	NA	1	NA
2	NA	NA	NA	NA	1.3	1.0–1.6	0.94	0.56–1.56
3	NA	NA	NA	NA	1.5	1.2–2.0	1.03	0.59–1.80
4	NA	NA	NA	NA	1.6	1.1–2.1	0.66	0.35–1.26
5	NA	NA	NA	NA	1.6	1.1–2.2	0.75	0.38–1.47

\*The reference group for women on active duty was 12 years of education or less.  
OR = odds ratios; CI = confidence intervals; NA = not applicable.

Table 6 Statistically significant adjusted odds ratios for smokeless tobacco use by 18–34 year old recruits and men on active duty

	Recruits		Active duty	
	OR	95% CI	OR	95% CI
<b>Race</b>				
White	1	NA	1	NA
Black	0.02	0.00–0.15	0.06	0.04–0.11
Hispanic	NA	NA	0.19	0.11–0.32
Asian	NA	NA	0.33	0.13–0.83
Others	0.20	0.08–0.47	0.78	0.40–1.52
<b>Service</b>				
Army	1	NA	1	NA
Air force	0.50	0.28–0.91	0.36	0.28–0.45
Marines	1.92	1.13–3.26	1.04	0.74–1.47
Navy	1.22	0.70–2.12	0.53	0.42–0.67
<b>Home region</b>				
New England	1	NA	NA	NA
Midwest	3.09	1.3–7.2	NA	NA
Southwest	3.17	1.2–8.6	NA	NA
West coast	2.97	1.1–7.8	NA	NA
Region 2	0.88	0.33–2.30	NA	NA
Region 4	1.89	0.83–4.31	NA	NA
Region 6	2.40	0.64–8.97	NA	NA
Region 8	1.17	0.23–6.08	NA	NA
<b>Rank</b>				
E-1	NA	NA	1	NA
E-2	NA	NA	0.57	0.27–1.24
E-3	NA	NA	0.51	0.25–1.03
E-4	NA	NA	0.41	0.20–0.82
E-5	NA	NA	0.52	0.25–1.08

OR = odds ratios; CI = confidence intervals; NA = not applicable.

SLT use quadrupled from 2.2% to 8.6% between recruit non-white/non-black men but remained unchanged, at near zero, for black men and for women of all races.

Tobacco use varied across branch of service (table 4). Navy personnel showed smaller gains in tobacco consumption between recruits and active duty than other military personnel.

Regression results for smoking (table 5) show that race is the only explanatory variable common to all four smoking models. Race is the only explanatory variable to stay in the smoking model for women on active duty.

Branch of service is retained as an explanatory variable in three of the four smoking models. For men, although navy recruits enter the service *more* likely to smoke than army recruits, and marine recruits are as likely as army recruits to smoke, sailors on active duty and marines are *less* likely to smoke than soldiers on active duty. In contrast, although both male air force recruits and personnel on active duty are less likely to smoke than their army counterparts, relatively more air force personnel on active duty smoke than air force recruits.

Among women, navy recruits are more likely, marine recruits are as likely, and air force recruits are less likely to smoke than army recruits. However, there is no difference in smoking prevalence across branch of service for personnel on active duty.

Where significant, higher education is associated with decreased smoking prevalence. Years of service on active duty is associated with increased smoking prevalence.

Table 6 shows SLT regression findings for men only because too few women used SLT to allow building SLT models for women. For both recruits and personnel on active duty, whites are much more likely to use SLT than other racial groups. Recruits from the midwest, southwest, and west coast are more likely to

use SLT than recruits from elsewhere in the United States. Comparing non-army with army personnel, relatively fewer service members on active duty use SLT than recruits.

## Discussion

The chief limitation of this study is its cross-sectional design. As such, causal conclusions cannot be drawn from the associations uncovered in the results. Nonetheless, cross-sectional studies can suggest causal hypothesis that can be tested with other study designs.

Overall, results from this study suggest, but do not prove, that exposure to the military leads to dramatic increases in tobacco use by young enlisted men, but not women. Further, increases in tobacco use are more likely to be found in the army than in the other services and among whites than among minorities.

The striking differential in smoking prevalence between genders in personnel on active duty is particularly noteworthy because no such differential exists in military recruits or in civilian society. Achieving gender parity in smoking prevalence in American civilians, largely the result of reducing smoking prevalence in men, took more than 30 years of government-supported public education and marketing.<sup>7</sup> Military service somehow appears to undo these gains in as few as five years.

However, when inspecting the data more closely across age groups, we do not see an increase in tobacco use with increasing age. This suggests that there may be strong secular changes in smoking prevalence in military personnel on active duty that cannot be dissected out in this cross-sectional study.

These findings present a challenge and an opportunity. The challenge is to find ways to curb the initiation of tobacco use in male enlisted personnel. The opportunity is to explore and understand what causes the differential appeal of tobacco consumption across gender, race, and branch of service among enlisted service members so that effective tobacco control strategies might be devised. Also, the issues explored in this research need to be studied in officer personnel as well. Curbing tobacco use among military personnel should be a high priority to DoD for societal, economic, and readiness reasons.

From the societal perspective, tobacco use is the single most preventable cause of premature death in the United States, accounting for roughly 20% of all deaths annually.<sup>8</sup> Many of these deaths follow costly, chronic, debilitating diseases, such as cardiovascular disease, respiratory illnesses, and various cancers that have been linked to tobacco use.<sup>9</sup> Because fewer than 4% of Americans die before age 35, smoking is *not* a major determinant of death between 17 and 35 years of age.<sup>10</sup> Thus, the military medical system is unlikely to be burdened with many of the long-term cost consequences of service members' tobacco use. Nonetheless, these costs ultimately will be borne by society, including both the civilian and the Department of Veterans Affairs health-care systems. Currently, 1.6 million men and

women serve in the American armed forces with many of them re-entering civilian life after a short tour of duty. DoD has an obligation to the society from which it draws its members to return service members to civilian life as healthy and productive as possible.

Regarding economic self-interest, even though the military medical system may evade many of the long-term costs of tobacco use by service members, it will not avoid the short-term costs.

Smokers, on average, account for 18% more annual medical expenditures than non-smokers.<sup>10</sup> Tobacco use has been linked to adverse pregnancy outcomes and to periodontal disease.<sup>9-11</sup> These are excess costs that the military medical system can ill afford.

Tobacco consumption has an adverse impact on military readiness. Tobacco users have higher absenteeism, impaired perceptual and motor skills, and poorer endurance than non-tobacco users.<sup>10-14</sup>

DoD anti-smoking efforts already include a smoking ban in DoD buildings, a ban on access to tobacco during military recruit training, an increase in tobacco prices in military commissaries, as well as tobacco education and cessation programmes. Most of these efforts are aimed at punishing or changing the behaviour of current smokers. Because of the strong addictive qualities of tobacco, this approach can only have a limited effect on curbing tobacco use. A recent review of 188 randomised controlled trials on smoking cessation reveals that only 2% of all smokers successfully quit.<sup>15</sup>

More emphasis needs to be placed on convincing service members not to initiate tobacco use. This strategy will require a strong

marketing approach that attempts to change favourable cultural attitudes toward tobacco within the military.

This paper was presented in a poster session at the annual scientific session of the Society of Behavioral Medicine in San Francisco, 16–19 April 1997. The views of the authors do not purport to reflect the views of the Department of the Army, the Department of the Navy, the Department of the Air Force, or the Department of Defense.

- 1 US Department of Defense. *Survey of health related behaviors among military personnel*. Report No RTI/6019/06-FR. Research Triangle Park, North Carolina: Research Triangle Institute, 1995.
- 2 Fletcher RH, Fletcher SW, Wagner EH. *Clinical epidemiology: the essentials*. 2nd ed. Baltimore, Maryland: Williams and Wilkins, 1988.
- 3 Cronan TA, Conway TL. Is the Navy attracting or creating smokers? *Mil Med* 1988;153:175–7.
- 4 Chisick MC, Arthur JS, York AK, et al. Designing a standardized oral health survey for the tri-services. *Mil Med* 1994;159:179–86.
- 5 Poindexter FR, York AK, Chisick MC, et al. A computer program for epidemiological data collection. *Toward an electronic patient record '96. Proceedings of the twelfth international symposium on the creation of an electronic health record system and global conference on patient cards*. 1996;1:276–79.
- 6 Hosmer DW, Lemshow S. *Applied logistic regression*. New York: John Wiley, 1989.
- 7 Geronimus AT, Neidert LJ, Bound J. Age patterns of smoking in US black and white women of childbearing age. *Am J Public Health* 1993;83:1258–64.
- 8 Nelson DE, Kirdendall RS, Lawton RL, et al. Surveillance for smoking-attributable mortality and years of potential life lost, by state—United States, 1990. *MMWR* 1994; 43:1–8.
- 9 Fielding JE. Smoking: health effects and control. *N Engl J Med* 1985;313:491–8.
- 10 Hodgson TA. Cigarette smoking and lifetime medical expenditures. *Milbank Q* 1992;70:81–125.
- 11 Mandel I. Smoke signals: an alert for oral disease. *J Am Dent Assoc* 1994;125:872–8.
- 12 Cooper KH, Gey GO, Bottenberg RA. Effects of cigarette smoking on endurance performance. *JAMA* 1968; 203:123–6.
- 13 Bahrke MS, Baur TS, Poland DF, et al. Tobacco use and performance on the US Army physical fitness test. *Mil Med* 1988;153:229–35.
- 14 Jensen RG. The effect of cigarette smoking on Army physical readiness test performance of enlisted Army Medical Department personnel. *Mil Med* 1986;151:83–5.
- 15 Law M, Tang JL. An analysis of the effectiveness of interventions intended to help people stop smoking. *Arch Intern Med* 1995;155:1933–41.