

REVIEW

The links between axin and carcinogenesis

S Salahshor, J R Woodgett

J Clin Pathol 2005;58:225–236. doi: 10.1136/jcp.2003.009506

The products of the two mammalian Axin genes (Axin1 and its homologue Axin2) are essential for the degradation of β catenin, a component of Wnt signalling that is frequently dysregulated in cancer cells. Axin is a multidomain scaffold protein that has many functions in biological signalling pathways. Overexpression of axin results in axis duplication in mouse embryos. Wnt signalling activity determines dorsal–ventral axis formation in vertebrates, implicating axin as a negative regulator of this signalling pathway. In addition, Wnts modulate pattern formation and the morphogenesis of most organs by influencing and controlling cell proliferation, motility, and fate. Defects in different components of the Wnt signalling pathway promote tumorigenesis and tumour progression. Recent biochemical studies of axins indicate that these molecules are the primary limiting components of this pathway. This review explores the intriguing connections between defects in axin function and human diseases.

Axin was originally identified as the product of the mouse gene called “fused” or fu (renamed Axin), and has since been shown to play a crucial role in controlling axis formation during embryonic development. Axin overexpression in frog embryos inhibits dorsal axis formation. Furthermore, mutation of the mouse gene “fused” was found to cause axis duplication in homozygous mouse embryos.¹ Wnt signalling activity determines dorsal–ventral duplication in vertebrates and these results suggested that Axin somehow negatively regulates this signalling pathway. The subsequent demonstration of the effect of Axin on β catenin concentrations, together with its biallelic inactivation in some human hepatocellular carcinomas (HCCs), indicated that Axin is a tumour suppressor gene.² To date, axin has been implicated in at least three different signalling pathways: the stress activated protein kinase (SAPK), transforming growth factor β (TGF β), and Wnt signalling pathways (fig 1). The multimeric nature of axin complexes suggests that axin might play an important role in other cell signalling systems, in addition to the coordination of these signals.

“The effect of Axin on β catenin concentrations, together with its biallelic inactivation in some human hepatocellular carcinomas, indicated that Axin is a tumour suppressor gene”

WNT SIGNALLING

The Wnt signalling pathway regulates cellular proliferation, differentiation, and motility and is essential for development and morphogenesis.^{3–5} Alterations in protein phosphorylation status are central to the regulation of Wnt signalling. Several components of the Wnt signalling pathway—including axin, adenomatous polyposis coli (APC), glycogen synthase kinase 3 (GSK3; both GSK β and GSK α), and β catenin—are phosphoproteins that are regulated through phosphorylation. In unstimulated cells, GSK3 α and GSK3 β phosphorylate cytoplasmic β catenin, which creates a recognition motif for β transducin repeat containing protein, an E3 ubiquitin ligase.⁶ The ubiquitinated β catenin is rapidly degraded by the 26S proteasome, ensuring that cytoplasmic concentrations of β catenin are very low. Secreted Wnt ligands act on the cell surface receptor Frizzled and activate Dishevelled (DVL) through a poorly understood mechanism. Activated DVL binds and inhibits the phosphorylation of β catenin by GSK3 β/α , blocking β catenin degradation (fig 2), so that β catenin accumulates and translocates to the nucleus, where it interacts with the T cell specific factor (TCF)/lymphoid enhancer binding factor 1 (LEF-1) transcription factor and induces the transcription of target genes such as c-jun, c-myc, and cyclin D1.^{7–11} Several in vitro and in vivo studies suggest that axins serve as scaffold proteins that bind directly to many proteins involved in the Wnt signalling pathway, and promote the phosphorylation of β catenin by driving the formation of a complex with APC and GSK3.^{12–14}

The β catenin molecule is crucial in this pathway and shows abnormal expression and localisation in a wide variety of human cancers.¹⁵ In recent years, so called “canonical” Wnt signalling has received considerable attention because many components of this signalling cascade have been shown to play a role in tumorigenesis.¹ The concentration of β catenin in cells is tightly regulated by the APC–axin–GSK3 destruction complex.¹⁶ APC, another component

Abbreviations: aa, amino acids; AD, Alzheimer’s disease; APC, adenomatous polyposis coli; CKI/II, casein kinase I/II; DIX, Dishevelled and axin binding domain; DVL, Dishevelled; Frat, frequently rearranged in activated T cells; GSK3, glycogen synthase kinase 3; HB, hepatoblastoma; HCC, hepatocellular cancer; JNK, Jun N-terminal kinase; LEF-1, lymphoid enhancer binding factor 1; MB, medulloblastoma; MEKK1, mitogen activated protein/extracellular regulated kinase kinase kinase 1; MID, MEKK binding domain; PP2A, protein phosphatase 2A; RGS, regulator of G protein signalling; SAPK, stress activated protein kinase; SCC, oesophageal squamous cell carcinoma; TCF, T cell specific factor; TGF β R, transforming growth factor β receptor

See end of article for authors’ affiliations

Correspondence to: Dr S Salahshor, Ontario Cancer Institute, Division of Experimental Therapeutics, 610 University Ave, M5G 2M9, Toronto, ON, Canada; Salahsho@Uhnres.Utoronto.Ca

Accepted for publication 26 October 2004




Figure 1 At least three different signalling pathways are regulated by axin. (A) In the absence of Wnt ligands, axin stimulates β catenin phosphorylation and subsequent protease mediated degradation limits its transcriptional activity. (B) In the presence of transforming growth factor β (TGF β) signals, axin stimulates Smad phosphorylation by TGF β receptors (TGF β receptors I and II). The activated Smads then translocate to the nucleus and activate transcription of downstream target genes. (C) In cells subjected to stress, axin binds to mitogen activated protein/extracellular regulated kinase kinase kinase 1 (MEKK1) and stimulates stress activated protein kinase (SAPK)/JNK (Jun N-terminal kinase) mediated apoptosis. GSK3, glycogen synthase kinase 3.

of Wnt signalling, is mutated in most (~ 70%) human colorectal cancers and is also linked to familial adenomatous polyposis.¹⁷ Inappropriate activation of Wnt signalling by mutation of different components of this pathway has been seen in a large number of other human cancers including colon carcinoma,¹⁸⁻¹⁹ medulloblastoma,²⁰⁻²¹ melanoma,²² hepatocellular carcinoma,²³ and ovarian and uterine cancer.²⁴⁻²⁶ Although both β catenin and Wnt are positive regulators of

the pathway and have been identified as protooncogenes, APC and axin are considered as negative effectors of the pathway and function as tumour suppressors.²⁷ This model asserts that because axin-APC-GSK3 mediated degradation of β catenin is inhibited by either Wnt signals or mutation of components of the Wnt signalling pathway, the concentration of β catenin in cells is raised, leading to increased cell proliferation and cancer.




Figure 2 The role of axin in Wnt signalling. The Wnt signalling pathway plays an important role in the regulation of cellular proliferation, differentiation, motility, and morphogenesis. Axin serves as a scaffold protein that binds many of the proteins involved in this pathway. In the absence of Wnt ligands, the β catenin destruction complex, which is composed of adenomatous polyposis coli (APC), glycogen synthase kinase 3 (GSK3), and axin, is formed and leads to the phosphorylation and degradation of β catenin. In the presence of Wnt ligands, the formation of this complex is inhibited and the now stabilised β catenin is translocated into the nucleus and activates the transcription of downstream target genes. Axin is also implicated in shuttling β catenin out of the nucleus. However, β catenin can move in and out of the nucleus independently of axin. It is unclear whether axin participates in the regulation of all of the different pools of β catenin in the cell. Deregulation of many components of the Wnt pathway has been found in human cancer. DVL, Dishevelled; LEF, lymphoid enhancer binding factor; P, organic phosphate; TCF, T cell specific factor.

TGF β SIGNALLING

Similar to the Wnt signalling pathway, TGF β signalling regulates several cellular functions, often in concert with Wnt signalling, including proliferation, differentiation, migration, and apoptosis.²⁸ TGF β also plays an important role in carcinogenesis. Upon stimulation, TGF β receptors phosphorylate Smads—the TGF β effector proteins (fig 1B).²⁹ The phosphorylated Smads then translocate into the nucleus, where they regulate the transcription of target genes. Although TGF β can often appear to have different effects in different cell types, its primary effect on colonic epithelial cells is to reduce proliferation and induce differentiation.³⁰ Mutations of the TGF β receptors are often found in cancer cells with defects in mismatch repair systems.³¹ Axin has been shown to regulate the TGF β signalling pathway by acting as an adaptor for Smad3, one of the TGF β effectors (fig 1B).³² Both axin1 and axin2 physically interact with Smad3 through interaction with a domain that is located between the β catenin and DVL binding domains within the C-terminal region. The binding of DVL to axin has been shown to inhibit axin mediated downregulation of β catenin. Although both Smad3 and DVL interact with the same region of axin, biochemical studies have shown that the interaction of DVL with axin does not compete with Smad3 binding to axin. Colocalisation of axin with Smad2 and Smad3 in the cytoplasm has been observed. Upon receptor activation, Smad3 bound to axin is efficiently phosphorylated by TGF β receptor 1 and dissociates from the axin complex. Thus, axin may facilitate the phosphorylation and transcriptional activity of Smad3.³² TGF β is one of the anti-oncogenic factors that inhibit nuclear β catenin signalling in the Wnt pathway. Smad4, another effector of TGF β , has been shown to facilitate Wnt signalling through interaction with β catenin and TCF/LEF.³³ Although axin is a negative regulator of the Wnt signalling pathway, it facilitates TGF β signalling. These results indicate that axin has a dual function in signal transduction—acting as a negative regulator of the Wnt signalling pathway and as a positive regulator of the TGF β signalling pathway. Deregulation of both pathways is frequently detected in human cancer cells.

SAPK/JNK SIGNALLING

The SAPK/JNK (stress activated protein kinase/Jun N-terminal kinase) signal transduction pathway is activated in cells in response to stress. It is also involved in many normal physiological processes including tissue morphogenesis, cell proliferation, cell survival, and cell death.³⁴ Depending on the cell type or the context of activation of other signalling pathways, SAPK/JNK can help mediate cell survival or apoptosis.^{35–36} Overexpression of axin1 in cells stimulates SAPK/JNK and can induce apoptosis through the activation of MEKK (mitogen activated protein/extracellular regulated kinase kinase).^{37–38} A MEKK1 interacting domain (MID) on axin is flanked by the APC and GSK3 binding sites (fig 3A), and the C-terminal region of axin, which contains an oligomerisation domain, is necessary for SAPK activation (fig 3A). The effect of axin on the induction of apoptosis is mediated by SAPK/JNK activation and the destabilisation of β catenin.³⁹ The importance of the axin–MEKK interaction and subsequent SAPK/JNK activation has been shown by the Axin1 mutations found in cancer cells that are resistant to apoptosis.^{2–39–41} Induced expression of Axin in transgenic mice leads to massive cell death in different organs as a consequence of the increased capacity to degrade β catenin.⁴² The introduction of wild-type Axin1 into hepatocellular and colorectal cancer cells induces apoptosis in a SAPK/JNK dependent manner. GSK3 competes with a MEKK binding site on axin1 and thereby inhibits SAPK/JNK induced apoptosis. This is consistent with observations that

GSK3 can prevent cell death under certain circumstances.⁴³ Thus, axin can bind to either GSK3 or MEKK1 and form distinct complexes with opposing effects on apoptosis. Although axin can be “switched” to activate the SAPK/JNK pathway instead of Wnt signalling, it is still capable of downregulating β catenin.⁴⁴ Axin domains essential for Wnt signalling activation are distinct from those domains required for SAPK/JNK activation.

“Depending on the cell type or the context of activation of other signalling pathways, SAPK/JNK can help mediate cell survival or apoptosis”

MEKK4, which is related to MEKK1, was recently identified as another axin binding protein that mediates SAPK/JNK activation.⁴⁵ Interestingly, MEKK1 and MEKK4 compete for axin binding, even though the binding domain for MEKK4 is separate from the MEKK1 binding domain.⁴⁶ Axin dimerisation and oligomerisation appear to be necessary for the activation of the SAPK/JNK pathway.³⁷ Other components of Wnt signalling such as GSK3 and CKIe (casein kinase 1e) also compete for binding to axin, and thereby inhibit SAPK/JNK signalling, whereas the binding of APC and β catenin to axin does not interfere with axin mediated SAPK/JNK activation.⁴⁴

AXIN GENE STRUCTURE, EXPRESSION, AND SUBCELLULAR LOCALISATION

Axin1 (also simply called Axin), which encodes isoforms a and b, and Axin2 (also called Axil or Conductin) have 45% identity at the nucleotide level and the proteins they encode appear to be functionally similar. However, whereas Axin1 is expressed ubiquitously during mouse embryogenesis, Axin2 is expressed in a restricted pattern.¹ Axin1 is the constitutively expressed component of the β catenin degradation complex and is essential for the maintenance of low Wnt signalling activity in the basal state. In contrast, axin2 is upregulated in response to increased β catenin concentrations and thus serves to limit the duration and intensity of the Wnt signal.^{47–48} Axin is downregulated in a Wnt dependent manner and is dephosphorylated after Wnt stimulation, which leads to axin1 destabilisation over time. Cells that receive Wnt ligand signals have low concentrations of axin. Biochemical studies show that the intracellular concentrations of axin are approximately 1000 times lower than those of other destruction complex components, suggesting that axin is the limiting factor in this pathway.⁴⁹

Axin 1

Axin1 was first identified as the product of the mouse fused locus. Its human homologue was mapped to chromosome 16p13.3 and shows 87% similarity to the mouse protein.¹ Axin1 “isoform a” encodes a 862 amino acid (aa) polypeptide (GenBank NP_003493), whereas “isoform b” is a shorter form of axin lacking 36 aa within the N-terminal domain encoded by exon 8 (GenBank NP_851393) (fig 3A). The function of the polypeptide region encoded by exon 8 is as yet unknown (fig 4A). This splice form is conserved between different species, suggesting a conserved role. The polypeptide encoded by exon 8 is located between the β catenin binding and Dishevelled and axin binding domain (DIX) domains (fig 3A). The region contains a predicted CKI phosphorylation site and is also close to the axin oligomerisation site, where axin binds to itself. It has been suggested that axin dimerisation is necessary for its stability and function in cells. The spliced exon is also located between two potential axin nuclear export signals.⁵⁰




Figure 3 Genomic structure of Axin. (A) Axin1 is composed of 10 exons (encoding isoform a). Exon 8 is spliced out in isoform b. (B) Axin2 also consists of 10 exons that encode 843 amino acids (aa) (isoform a) or a 778 aa polypeptide (isoform b). Similar to Axin1, the binding partners of Axin2 are APC (aa 81–200), GSK3 (aa 372–413), β catenin (aa 414–476), and DVL (aa 761–843). So far, all mutations found in Axin2 are located in exon 7. APC, adenomatous polyposis coli; CK1, casein kinase I; DIX, Dishevelled and axin binding domain; DVL, Dishevelled; GSK3, glycogen synthase kinase 3; MEKK1, mitogen activated protein/extracellular regulated kinase kinase 1; NES, nuclear export signal; NLS, nuclear localisation signal; PP2A, protein phosphatase.

Axin2

A homologue of axin, axin2 was identified by virtue of its interaction with β catenin.^{16 51 52} Mutation of the Axin2 gene leads to an increase in β catenin concentrations in colorectal cancers with defective mismatch repair systems.⁵³ Molecular studies revealed that Axin2 contains 10 exons spanning more than 2.5 kb.⁵⁴ Similar to Axin1, Axin2 encodes two isoforms (a and b). In isoform b, exon 6 is spliced out, leading to a transcript with 65 fewer aa (S Salahshor, unpublished data, 2004). Fluorescence in situ hybridisation analysis assigned it to human chromosome 17q24, a region that shows frequent loss of heterozygosity in breast cancer, neuroblastoma, and other tumours.⁵⁴ Similar to Axin1, Axin2 has binding domains for APC, GSK3, and β catenin (fig 3B).¹⁶

Subcellular localisation

Based on the known function and proposed activities of axin, both axins are expected to be located in the cytoplasm of cells where they promote β catenin phosphorylation and degradation. Confocal microscopy of whole mount crypts from patients with colorectal cancer shows that staining for axin1 is diffuse within the nucleus and along lateral cell membranes where β catenin is located, and also in the cytoplasm

where both GSK3 and the cytoplasmic pool of β catenin are found. Cell lysate fractionation of two human epithelial cell lines—HCT116 (colorectal cancer cells with β catenin mutation) and HEK293 (embryonic kidney cells)—reflects the axin localisation pattern seen in tissue. Although axin1 was found mainly in the nucleus of HCT116 cells, in HEK293 cells it was localised in both the nuclear and cytoplasmic compartments.⁵⁵ Staining for axin1 is seen in the cytoplasm of normal epithelial cells, whereas it is located in the nucleus in adenocarcinomas and tumours (fig 4B). The subcellular localisation of axin1 appears to be cell type dependent (fig 5). In contrast, axin2 shows strong nuclear staining in normal tissue. Nuclear localisation of axin2 could also be detected in polyps and carcinomas with some cytoplasmic translocation.⁵⁵ Despite the similarity between these two molecules, it appears that they are not found within the same compartment of the cell, and thus may have distinct functions. Axin1 can shuttle between the cytoplasm and the nucleus and act as a β catenin chaperone. Two nuclear export (between aa 532 and 667) and three nuclear import signals have been identified on axin1 (fig 3A). Axin1 translocation to the nucleus and its interaction with β catenin is required for axin induced cytoplasmic shifting of β catenin.^{50 56}




Figure 4 Axin expression pattern and subcellular localisation in tumour cell lines and tumour tissue. (A) Western blot analysis of protein extracted from MCF12A and SW480 cells confirmed the existence of two axin1 isoforms (a and b). (B) The subcellular localisation of axin1 was examined in a sporadic colorectal cancer case and axin1 was found to have a predominantly nuclear localisation.




Figure 5 Subcellular localisation of axin1 in MCF12A cells. Based on the known function of axin1, it is expected to be mainly localised in the cytoplasm. Confocal sections of MCF12A cells stained with (A and D) anti-axin1 and (B and E) anti- β catenin antibodies (C and F merged) show both nuclear and cytoplasmic axin1 staining, with a pronounced nuclear localisation, whereas β catenin is located mainly at the cell membrane (scale bars, 20 μ m). The following antibodies were used: polyclonal anti-axin1 (1/500 dilution; Zymed, South San Francisco, California, USA), monoclonal anti- β catenin antibody (1/500 dilution; Signal Transduction Laboratories; Lexington, Kentucky, USA), secondary monoclonal and polyclonal antibodies (1/1000 dilution; Molecular Probes, Eugene, Oregon, USA). Other studies show a more cytoplasmic localisation for axin1. However, the subcellular localisation of axin1 appears to be cell type dependent.

AXIN BINDING PARTNERS

Both axins contain several domains that mediate direct binding to low density lipoprotein related protein receptor, the Frizzled coreceptor,⁵⁷⁻⁵⁹ APC (which appears to assist axin in recruiting β catenin to the axin complex),¹³ GSK3 β/α ,⁶⁰ β catenin.⁶¹⁻⁶² DVL (DVL-1, DVL-2, and DVL-3),⁶³ MEKK1, CKI,⁶⁴ protein phosphatase 2A (PP2A),⁶⁵ frequently rearranged in advanced T cell lymphomas (Frat1),⁶⁶ and a homodimerisation domain (figs 3 and 6).⁶⁷

APC

The region of axin involved in APC binding shows significant homology to members of the regulators of G protein signalling (RGS) family.¹⁻¹² The region of APC that interacts with axin consists of a conserved sequence of approximately 20 aa containing a Ser-Ala-Met-Pro (SAMP) motif.¹⁶ The APC gene contains three SAMP repeat sequences, all of which are located after its mutation cluster region, where most truncating mutations in cancer are found. These mutations eliminate all axin binding sites on APC, but preserve some of the β catenin binding domains, indicating the important role that the interaction between axin and APC plays in APC tumour suppressor activity.²²⁻⁶⁸ Thus, the tumorigenic potential of mutated APC correlates with loss of binding to axin1/axin2, rather than loss of β catenin binding. APC binding to axin via its RGS domain is required for the efficient downregulation of β catenin by APC.²²⁻⁵¹ However, overexpression of axin in colorectal cancer cell lines bearing a mutated form of APC that lacks an axin binding domain can

still promote β catenin downregulation.¹⁶⁻⁶¹⁻⁶⁹ Similarly, axin mutants lacking the APC binding domain are still capable of β catenin downregulation when overexpressed.⁵¹⁻⁶² These results suggest that APC might be only a cofactor for axin1/axin2. However, multiple binding sites on APC for β catenin (10 binding sites) and axin1/axin2 (three binding sites) may be important for the efficient assembly of the β catenin degradation machinery in different cell types or various *in vivo* conditions. These results further indicate that the concentration of axin rather than its interaction with other components of Wnt signalling might control its activity in cells, and that axin is the rate limiting factor in these reactions. In contrast, in SW480 colorectal cancer cell lines, which carry a truncated form of APC, β catenin is stabilised. Expression of ectopic APC leads to β catenin degradation.²²⁻⁷⁰ Endogenous axin1 and axin2 are present in this cell line, but appear not be capable of downregulating β catenin in the absence of wild-type APC.

“Direct binding of APC to β catenin is not essential for APC to degrade β catenin, whereas the binding of axin to β catenin is necessary for it to degrade β catenin”

However, overexpression of exogenous axin2 can induce β catenin degradation in SW480 colon carcinoma cells. Overexpression of an axin2 mutant that lacks the β catenin binding domain does not induce β catenin degradation, whereas an axin2 mutant lacking the RGS domain (APC

Table 1 Sequence variants of the Axin1 gene found in human cancer

Tissue type	Exon/nt or aa	Position (aa)	Domain(s) affected	Ref
CRC	1	S215L	APC-MEKK1?	94
CRC	1	R103M	APC	95
CRC	1	L101P	APC	95
CRC	3	R353H	GSK3	95
CRC	1	T122A	APC	95
CRC	4	E411D	GSK3	95
CRC	4	M418I	GSK3	95
CRC	4	R395C	GSK3	95
CRC	5	G584S	β Catenin	95
CRC	2	Y305X	MEKK1	96
CRC	1	K203M	APC	96
CRC	2	N307K	MEKK1	96
CRC	4	H394N	GSK3	96
CRC	10	P848L	Axin-DVL	96
CRC	10	E852G	Axin-DVL	96
CRC	1	S215L	APC-MEKK1?	94
MB	1-5 (aa 89-528)	Del	APC-GSK3- β catenin	21
MB	1-5 (aa 111-571)	Del	APC-GSK3- β catenin	21
MB	1-5 (aa 59-532)	Del	APC-GSK3- β catenin	21
MB	9-10	Del	Axin-DVL	21
MB	8-10	Del	Axin-DVL	21
MB	6-10	Del	PP2A-axin-DIX	21
MB	7-10	Del	PP2A-axin-DIX	21
MB	1	P255S	APC	21
MB	1	P255S	MEKK1	20
MB	1	S263C	MEKK1	20
HCC	1	Stop	APC-DVL-axin	2
HCC	4	Stop	GSK3-DVL-axin	2
HCC	4	5 bp del	β Catenin	2
HCC	5	13 bp del	β Catenin	2
HCC	3	P345L	MEKK1	97
HCC	1	L106R	APC	97
HCC	2 (1076 del 1 bp)	Frameshift	MEKK1?	97
HCC	5	Ins QVHH	β Catenin-PP2A?	97
HCC	5	G425S	β Catenin	97
HB	10	R842Q	Axin-DIX	97
HB	6	G651S	PP2A?	97
HB	2	T95M	APC	98
EM	4	A397V	GSK3	99
SCC	4	G419A	GSK3	100
OEA	4	L416X	GSK3	101
ACC	7	P661L	PP2A	102
ACC	5	G433E	β Catenin	102
ACC	10	E842K	Axin-DVL	102
ACC	10	F824K	Axin-DVL	102
ACC	6	S628Y	PP2A?	102

Mutations found in the axin1 coding sequence are scattered throughout the whole gene. There are some differences between the alterations found in different forms of tumours. Large deletions are more common in medulloblastomas, whereas missense mutations occur more frequently in hepatocellular carcinomas. Silent mutations and polymorphisms reported in axin1 are not included in this list. Protein and mRNA sequences are based on Swissprot and GenBank accession numbers O15169 and AF009674, respectively.

aa, amino acids; ACC, adenoid cystic carcinoma; APC, adenomatous polyposis coli; bp, base pair; CRC, colorectal cancer; del, deletion; DVL, Dishevelled; EM, endometrial cancer; GSK3, glycogen synthase kinase 3; HB, hepatoblastoma; HCC, hepatocellular cancer; MB, medulloblastomas; MEKK1, mitogen activated protein/extracellular regulated kinase kinase kinase 1; nt, nucleotide; OEA, ovarian endometrioid adenocarcinoma; PP2A, protein phosphatase 2A; SCC, oesophageal squamous cell carcinoma.

CKI/II

CKI has recently been found to be an activator of Wnt signalling because microinjection of CKI into xenopus induces a secondary axis.⁸⁷ The overexpression of CKI mimics Wnt signalling by stabilising β catenin and thereby inducing the expression of β catenin target genes. CKI has been detected in the axin-GSK3 complex and appears to phosphorylate APC and β catenin (at Ser45) in an axin dependent manner.^{64 79 88} CKI also competes for MEKK1 binding and can attenuate axin-SAPK/JNK signalling.⁸⁹ In contrast, although CKII also interacts with axin, it does not have an inhibitory effect on SAPK/JNK signalling.⁸⁹ CKII interacts and phosphorylates DVL, which is thought to inhibit the function of axin, thereby stimulating Wnt signal transduction.⁹⁰

PP2A

PP2A is a broad specificity serine/threonine protein phosphatase that comprises a catalytic "C" subunit and various

"A" and "B" regulatory subunits that confer selective binding to proteins and scaffolds within cells. PP2A physically interacts with axin and APC in the axin-GSK3- β catenin destruction complex and decreases the phosphorylation of axin, which suggests a positive role for PP2A in Wnt signal transduction.¹⁴ Phosphorylation of APC is also important for β catenin binding and its subsequent degradation. PP2A has been implicated as a positive effector for Wnt mediated signalling.⁹¹ However, the phosphatase has other, confounding effects, probably because of its action on both positive and negative effectors of the pathway, such that its role is clearly complex.^{14 65 92 93}

AXIN VARIANTS IN CANCER

Alterations in both axin1 and axin2 have been detected in several different tumours (tables 1 and 2). Mutations are found in most axin domains including the APC (RGS) and β catenin binding domains. Axin sequence variants have also

Table 2 Axin2 variants in human diseases

Tissue type	Exon/nt (bp)	Position (aa)	Domain affected	Ref
CRC	7	L688X	?	53
CRC	7	E706X	?	53
OEA	7	G665X	?	101
CRC	7	E706Ter	?	103
EM	7	S658C	?	99
CRC/ Oligodentia	7	R656X	?	104
CRC/ Oligodentia	7 (1995 ins G)	?706X	?	104
HCC	7	R659W	?	97
HCC	7 (2102; del 12 bp)	del TTPR	?	97

Approximately 20% of mismatch repair deficient colorectal cancers harbour mutations in axin2. Most mutations are located around a G₇ repeat in exon 7 (nt 637–714), which results in premature protein truncation.

aa, amino acids; APC, adenomatous polyposis coli; bp, base pair; CRC, colorectal cancer; del, deletion; EM, endometrial cancer; GSK3, glycogen synthase kinase 3; HB, hepatoblastoma; HCC, hepatocellular cancer; ins, insertion; nt, nucleotide; OEA, ovarian endometrioid adenocarcinoma.

been found in colon, ovarian, endometrioid, adenocarcinoma, and HCC cell lines (table 3). Biochemical and functional studies have shown that these mutations interfere with the binding of GSK3 and that they also alter the interaction between axin and two upstream activators of TCF dependent transcription, Frat1 and DVL.⁹⁴ Several studies have investigated the role of axin in several types of tumour.

Medulloblastoma

Medulloblastoma (MB) is the most common malignant brain tumour in children.¹⁰⁵ Most MBs are sporadic; however, patients with germline mutations in APC or PTCH (patched) genes carry a higher risk of this disease.¹⁰⁶ In addition, mutations in β catenin, another component in the Wnt pathway, have been detected in MB tumours.^{107–109} Recently, several deletions and single somatic point mutations in Axin1 have been found in sporadic MB cases.^{20 21 110} These aberrations include somatic point mutations, deletions, and loss of heterozygosity. The C-terminal deletion in these cases corresponded to loss of exons 6–10, where the DIX and oligomerisation domains are located. Oligomerisation of Axin1 is crucial for the inhibition of TCF transcription,⁶⁷ and C-terminal deleted axin1 fails to downregulate β catenin and cell growth in the SW480 colon cancer cell line.⁶³ Another type of inframe deletion that removes exons 1–5, the APC binding domain, has also been found in MB. The truncated protein lacks binding sites for APC, GSK3, and β catenin (table 1). This mutant is also incapable of downregulating β catenin. In addition, some truncated forms of axin1 may act as dominant negatives and inactivate endogenous axin1.¹¹¹

Table 3 Axin1 variants identified in cancer cell lines

Cell line	Type	Exon	Position (aa)	Domain(s) affected	Ref
MDAH-2774	OEA	5	V555I	?	101
SNU475	HCC	1–2	Del	APC	2
SNU423	HCC	3	13 bp del	GSK3	2
Alexander	HCC	4	Del	GSK3	2
HCT-8	CRC	4	L396M	GSK3	94

aa, amino acids; APC, adenomatous polyposis coli; bp, base pair; CRC, colorectal cancer; del, deletion; GSK3, glycogen synthase kinase 3; HCC, hepatocellular cancer; OEA, ovarian endometrioid adenocarcinoma.

Colorectal cancer and familial tooth agenesis

Many components of the Wnt signalling system are mutated in colorectal cancer. Germline loss of function mutations in the APC gene are associated with an inherited form of colorectal cancer—familial adenomatous polyposis—with 90–95% penetrance. Somatic APC mutations are also found in most sporadic colorectal cancers.¹¹² Alterations in other components of Wnt signalling, including β catenin, TCF, axin1, and axin2, found in colorectal cancer indicate the important role that this pathway plays in the aetiology of this disease.¹¹³ Most Axin1 mutations in colorectal cancer occur between exon 1 and 5, where the APC, GSK3, and β catenin binding domains are located (table 1). Mutations in axin2 have been found in approximately 20% of mismatch repair deficient colorectal tumours (table 2).^{6 53 94 101 103} In most cases, one base deletion or insertion occurs in the mononucleotide repeat sequences located in exon 7, leading to a frame shift and premature protein truncation.⁵³ These mutations lead to elimination of the DIX domain, where DVL binds and negatively regulates axin activity. This domain is also essential for homo-oligomerisation of axin. The mutant form of axin2 appears to be more stable than the wild-type protein. Transfection of normal fibroblasts with axin2 mutants led to the accumulation of β catenin in the nuclei.

“Alterations in other components of Wnt signalling, including β catenin, TCF, axin1, and axin2, found in colorectal cancer indicate the important role that this pathway plays in the aetiology of this disease”

Immunohistochemical staining of β catenin in tumours with defects in axin2 also showed nuclear accumulation of β catenin in the cells. These results suggest a dominant negative effect of axin2 mutant proteins in cells. High levels of Axin2 gene expression have been detected in most human colon cancer cell lines (SW480, SW620, LoVo, SW620, Caco2, HT29, HCT116, T84, HCT15, and Alab), in addition to colorectal tumours. Axin2 appears to be a transcriptional target of the Wnt signalling pathway. Upregulation of Axin2 by Wnt– β catenin appears to constitute a negative feedback loop that acts to restrain or desensitise Wnt signalling. Axin2 gene expression increases in response to raised concentrations of β catenin, thereby modulating the duration and activity of the Wnt signal.^{47 48} In addition to colorectal cancer, several Axin2 mutations have been found in patients with oligodentia. Linkage analysis of a Finnish family with colorectal cancer in association with oligodentia revealed a high logarithm of the odds score in chromosome 17 between markers D17S949 and D17S1352, where more than 80 known or predicted genes (including Axin2) are located. Direct mutational analysis of the coding sequence of Axin2 identified a base pair transition in exon 7, leading to premature termination of translation of Axin2, whereas family members with normal dentition showed no sign of Axin2 mutation or neoplasia. Axin2 is expressed in developing dental tissues, and germline Axin2 mutations found in patients with oligodentia provided the first evidence for its function.¹⁰⁴ In a form of familial colorectal cancer, Gardner syndrome, the occurrence of odontomas and supernumerary teeth in association with familial adenomatous polyposis has been reported.^{114–117} These results indicate that overactivation of Wnt signalling as a result of mutation of a component of the β catenin destruction complex, in addition to carcinogenesis, may lead to the failure of tooth development (tables 1–3).

Hepatocellular carcinomas

HCC and hepatoblastoma (HB) are primary liver cancers that occur predominantly in adults and children, respectively.

Alteration of β catenin occurs in approximately 20% of HCCs and 40–89% of HBs.^{23 118–121} A fraction of HCCs (~ 10%) with wild-type β catenin have mutations in Axin1 or Axin2 instead.^{2 122} These data suggest that alterations in the Wnt pathway are involved in the pathogenesis of a large proportion of both HBs and HCCs. The axin1 mutations identified in the previous reports include truncation mutations as a result of either small deletion/insertion or nonsense mutations, although Axin1 mutations found in liver cancer show a different spectrum of alterations. Most of the mutations found in these cases were predominantly missense. These Axin mutations apparently only affect one allele, so that their oncogenic effect may result from the dominant negative activity of the mutant Axin, which causes stabilisation of free β catenin.¹⁶ In addition, detection of loss of heterozygosity at the Axin1 locus suggests that axin1 functions as a tumour suppressor. However, in a subset of tumours mutations in Axin1 or Axin2 were found in addition to mutations in β catenin.⁹⁷ The expression of wild-type Axin1 in HCC or colorectal cancer cells with APC, β catenin, or Axin1 mutations led to the induction of apoptosis.² Even though axin1 and axin2 are both able to downregulate β catenin when overexpressed, and show similar biochemical characteristics, axin2 does not compensate for axin1 mutations in HCCs. Therefore, axin1 and axin2 expression is either cell type dependent or, more likely, these two proteins are not functionally equivalent.

Ovarian endometrial adenocarcinomas

Ovarian cancer is one of the most frequent gynaecological malignancies in women.¹²³ A subtype of ovarian cancer, endometriosis, often shows defects in the components of the Wnt signalling pathway. Although mutations in APC are rare in this type of tumour, mutation of β catenin is found in 16–24% of cases.^{24 124 125} Recently, mutations in axin1 and axin2 have been detected in a subset of ovarian endometrial adenocarcinomas, where activation of TCF dependent transcription could be detected in a tumour with an Axin2 frameshift mutation (tables 1 and 2). As noted in cell lines, the mutated form of axin2 appears to be more stable than the wild-type axin2 protein.¹⁰¹

Oesophageal squamous cell carcinoma

Oesophageal cancer accounts for 7% of all gastrointestinal malignancies, but in some regions of Asia its incidence may be as high as 170/100 000 of the population. About half of the oesophageal cancers diagnosed are squamous cell carcinomas (SCCs). Several studies suggest a role for defects in Wnt signalling in the pathogenesis of SCC. Deletion of chromosome 17 (where Axin2 is located) has been found in 45% of SCCs.¹²⁶ Immunohistochemical studies of axin1 have shown cytoplasmic localisation of the protein in normal stratified squamous epithelium of the oesophagus, whereas a mixed pattern of expression was detected in tumours. Protein expression analysis of seven oesophageal SSC cell lines revealed differential axin expression. Four cell lines (TE1, TE15, TT, and TTn) showed high expression, whereas another three cell lines (TE2, TE8 and TE13) showed very low axin1 expression. However, northern blot analysis of the same cells showed similar amounts of RNA in all cases. Several mutations and polymorphisms have been reported in SCC tumours and cell lines (tables 1 and 3), and a correlation between reduced axin1 expression and tumour progression has been suggested in oesophageal SCC.^{100 127}

Roles of axin/GSK3 in Alzheimer's disease

In addition to tumorigenesis, defects in the Wnt signalling pathway have also been postulated to contribute to the pathogenesis of Alzheimer's disease (AD).^{128–130} The pathology of this form of dementia involves the deposition of amyloid

plaques, which are made up of A β peptides derived from the abnormal cleavage of amyloid precursor protein, and neurofibrillary tangles, which are composed of hyperphosphorylated forms of the microtubule associated protein, tau. These two lesions are associated with neuronal cell death. GSK3 is one of several protein kinases that phosphorylates tau at sites observed in the neurofibrillary tangles.^{131–133} Axin negatively affects tau phosphorylation by GSK3.¹³⁴ Tau that has been phosphorylated by GSK3 has a decreased capacity to bind and stabilise microtubules.¹³⁵ Altered expression of PP2A, another protein implicated in Wnt signalling, has also been associated with AD pathology and appears to play a major role in regulating tau phosphorylation in the adult brain.^{136–139} These results have led to the hypothesis that deregulation of GSK3 might be involved in the pathogenesis of AD.^{140 141} Considering the central role that the scaffolding protein axin plays in regulating GSK3 mediated phosphorylation of β catenin, and the fact that axin negatively regulates GSK3 mediated tau phosphorylation, it has been suggested that deregulation of the phosphorylation status of GSK3 may contribute to tau pathology in AD. A direct role for Wnt signalling in AD pathology remains to be demonstrated. However, it can be speculated that events leading to a decrease in GSK3–axin complex formation, and ultimately axin concentrations, may allow the aberrant phosphorylation of tau by GSK3 at primed sites that are increased in the AD brain, and thus contribute to tau dysfunction and pathology.¹⁴²

AXIN: A POTENTIAL TARGET FOR CANCER TREATMENT

Given the clear link between defects in Wnt signalling components and several human cancers, compounds that either positively or negatively regulate these molecules according to their role in the pathway will provide valuable reagents for the development of anticancer drugs. Because axin positively regulates the SAPK/JNK (apoptosis) pathway and negatively regulates the Wnt (survival) pathway, and is the limiting factor in these systems, a search for axin antagonists or agonists might lead to the discovery of compounds that have potential for the treatment of cancer. To ascertain whether the Axin mutations found in cancers were biologically important, normal Axin1 genes were inserted into the cancer cells and their effects on cell growth monitored. In the case of HCC cell lines, the introduction of Axin appeared to induce programmed cell death, suggesting a therapeutic potential. Although this opens up the possibility of using Axin as a cargo gene in gene therapy, small molecule approaches are also being pursued.

“A search for axin antagonists or agonists might lead to the discovery of compounds that have potential for the treatment of cancer”

The high frequency of mutations in the Wnt pathway in specific tumour types suggests that the activated transcriptional output of these pathways efficiently misregulates multiple growth regulatory functions. Because axin can degrade β catenin when overexpressed in cells that lack functional APC it is a good candidate for cancer treatment. One drawback is that although the overexpression of axin1 strongly promotes the downregulation of wild-type β catenin in colon cancer cells, mutant, oncogenic β catenin is unaffected.

Thus, Axin treatments would be selective for APC associated cancers. More studies of the mechanistic details of these molecules are required to elucidate the precise role and effect of axin stimulation in cancer treatment.

Take home messages

- Axins (both Axin1 and its homologue Axin2) are tumour suppressor genes
- Deregulation of axin has been found in different forms of cancer
- Wnt signals modulate axin concentrations in the cell
- Axin is a rate limiting component of different signalling pathways
- The Wnt, stress activated protein kinase/Jun N-terminal kinase, and transforming growth factor β signalling pathways are regulated by axin
- Upregulation of axin leads to the phosphorylation and degradation of β catenin
- The overexpression of axin promotes apoptosis

CONCLUDING REMARKS

A large body of evidence indicates that the regulation of β catenin stability and concentration in cells by the axin-APC-GSK3 complex is crucial for both embryogenesis and carcinogenesis. Axins act as negative regulators for Wnt signalling, and positive regulators for SAPK/JNK signalling. Because axin itself is sensitive to Wnt signals, axin concentrations in cells fluctuate, which probably impacts upon its other functions. Despite being discovered in 1997, there is much to be learned about the physiological functions and roles of axin. In common with several other molecules in the Wnt signalling pathways, axin binds to different components and regulates opposite functions in the cell. The question is how axin operates to maintain the specificity of these different signalling pathways. Downregulation of axin can be just as catastrophic as too much activity. Axin2 overexpression in cells leads to apoptosis, yet upregulation of axin2 is found in most colorectal cancers. Deregulation of axin2 can be detected in adenomas where cells do not show β catenin nuclear accumulation, which indicates that axin2 might be an early marker of tumour initiation. Furthermore, axin2 is mostly located in the nucleus, and has access to the "active" form of its target molecule, β catenin. It remains to be determined how the basal concentrations of axins are controlled and how their nuclear and cytoplasmic localisation is accomplished. Currently, our understanding of axins suggests that these proteins play key assembly functions for promoting efficient protein-protein interaction and stand at the crossroads of several signalling paths—helping to integrate and coordinate the plethora of continuously changing signals.

ACKNOWLEDGEMENTS

Supported by grants from the Canadian Institutes of Health Research and the National Institute of Cancer/Canadian Cancer Society.

Authors' affiliations

S Salahshor, J R Woodgett, Ontario Cancer Institute, Division of Experimental Therapeutics, Toronto, Ontario, Canada, M5G 2M9

APPENDIX 1 ONLINE LINKS AND DATABASES

- http://kinase.uhnres.utoronto.ca/Sima/axins_pathway/ (Axin page)
- <http://www.stanford.edu/~rnusse/wntwindow.html> (Wnt page)
- <http://us.expasy.org/prosite/> (Prosite database)

REFERENCES

- 1 Zeng L, Fagotto F, Zhang T, *et al.* The mouse Fused locus encodes axin, an inhibitor of the Wnt signalling pathway that regulates embryonic axis formation. *Cell* 1997;**90**:181–92.
- 2 Satoh S, Daigo Y, Furukawa Y, *et al.* AXIN1 mutations in hepatocellular carcinomas, and growth suppression in cancer cells by virus-mediated transfer of AXIN1. *Nat Genet* 2000;**24**:245–50.
- 3 Akiyama T. Wnt/ β -catenin signaling. *Cytokine Growth Factor Rev* 2000;**11**:273–82.
- 4 Wodarz A, Nusse R. Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 1998;**14**:59–88.
- 5 Biens M, Clevers H. Linking colorectal cancer to Wnt signaling. *Cell* 2000;**103**:311–20.
- 6 Bellamy CO, Malcomson RD, Harrison DJ, *et al.* Cell death in health and disease: the biology and regulation of apoptosis. *Semin Cancer Biol* 1995;**6**:3–16.
- 7 Molenaar M, van de Wetering M, Oosterwegel M, *et al.* XTCF-3 transcription factor mediates β -catenin-induced axis formation in *Xenopus* embryos. *Cell* 1996;**86**:391–9.
- 8 Behrens J, von Kries JP, Kuhl M, *et al.* Functional interaction of β -catenin with the transcription factor LEF-1. *Nature* 1996;**382**:638–42.
- 9 He TC, Sparks AB, Rago C, *et al.* Identification of c-MYC as a target of the APC pathway. *Science* 1998;**281**:1509–12.
- 10 Tetsu O, McCormick F. β -Catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature* 1999;**398**:422–6.
- 11 Shutman M, Zhurinsky J, Simcha I, *et al.* The cyclin D1 gene is a target of the β -catenin/LEF-1 pathway. *Proc Natl Acad Sci U S A* 1999;**96**:5522–7.
- 12 Ikeda S, Kishida S, Yamamoto H, *et al.* Axin, a negative regulator of the Wnt signaling pathway, forms a complex with GSK-3 β and β -catenin and promotes GSK-3 β -dependent phosphorylation of β -catenin. *EMBO J* 1998;**17**:1371–84.
- 13 Kishida S, Yamamoto H, Ikeda S, *et al.* Axin, a negative regulator of the wnt signaling pathway, directly interacts with adenomatous polyposis coli and regulates the stabilization of β -catenin. *J Biol Chem* 1998;**273**:10823–6.
- 14 Ikeda S, Kishida M, Matsuura Y, *et al.* GSK-3 β -dependent phosphorylation of adenomatous polyposis coli gene product can be modulated by β -catenin and protein phosphatase 2A complexed with axin. *Oncogene* 2000;**19**:537–45.
- 15 Morin PJ. β -Catenin signaling and cancer. *Bioessays* 1999;**21**:1021–30.
- 16 Behrens J, Jerchow BA, Wurtele M, *et al.* Functional interaction of an axin homolog, conductin, with β -catenin, APC, and GSK3 β . *Science* 1998;**280**:596–9.
- 17 Kinzler KW, Nilbert MC, Vogelstein B, *et al.* Identification of a gene located at chromosome 5q21 that is mutated in colorectal cancers. *Science* 1991;**251**:1366–70.
- 18 Morin PJ, Sparks AB, Korinek V, *et al.* Activation of β -catenin-Tcf signaling in colon cancer by mutations in β -catenin or APC. *Science* 1997;**275**:1787–90.
- 19 Ilyas M, Tomlinson IP, Novelli MR, *et al.* Clinico-pathological features and p53 expression in left-sided sporadic colorectal cancers with and without microsatellite instability. *J Pathol* 1996;**179**:370–5.
- 20 Baeza N, Masuoka J, Kleihues P, *et al.* AXIN1 mutations but not deletions in cerebellar medulloblastomas. *Oncogene* 2003;**22**:632–6.
- 21 Dahmen RP, Koch A, Denkhau D, *et al.* Deletions of AXIN1, a component of the WNT/wingless pathway, in sporadic medulloblastomas. *Cancer Res* 2001;**61**:7039–43.
- 22 Rubinfeld B, Robbins P, El-Gamil M, *et al.* Stabilization of β -catenin by genetic defects in melanoma cell lines. *Science* 1997;**275**:1790–2.
- 23 Miyoshi Y, Iwao K, Nagasawa Y, *et al.* Activation of the β -catenin gene in primary hepatocellular carcinomas by somatic alterations involving exon 3. *Cancer Res* 1998;**58**:2524–7.
- 24 Palacios J, Gamallo C. Mutations in the β -catenin gene (CTNBN1) in endometrioid ovarian carcinomas. *Cancer Res* 1998;**58**:1344–7.
- 25 Fukuchi T, Sakamoto M, Tsuda H, *et al.* β -Catenin mutation in carcinoma of the uterine endometrium. *Cancer Res* 1998;**58**:3526–8.
- 26 Polakis P. Wnt signaling and cancer. *Genes Dev* 2000;**14**:1837–51.
- 27 Willert K, Shibamoto S, Nusse R. Wnt-induced dephosphorylation of axin releases β -catenin from the axin complex. *Genes Dev* 1999;**13**:1768–73.
- 28 Massague J. TGF- β signal transduction. *Annu Rev Biochem* 1998;**67**:753–91.
- 29 Heldin CH, Miyazono K, ten Dijke P. TGF- β signalling from cell membrane to nucleus through SMAD proteins. *Nature* 1997;**390**:465–71.
- 30 Chakrabarty S, Fan D, Varani J. Modulation of differentiation and proliferation in human colon carcinoma cells by transforming growth factor β 1 and β 2. *Int J Cancer* 1990;**46**:493–9.
- 31 Jass JR, Do KA, Simms LA, *et al.* Morphology of sporadic colorectal cancer with DNA replication errors. *Gut* 1998;**42**:673–9.
- 32 Furuhashi M, Yagi K, Yamamoto H, *et al.* Axin facilitates Smad3 activation in the transforming growth factor β signaling pathway. *Mol Cell Biol* 2001;**21**:5132–41.
- 33 Nishita M, Hashimoto MK, Ogata S, *et al.* Interaction between Wnt and TGF- β signalling pathways during formation of Spemann's organizer. *Nature* 2000;**403**:781–5.
- 34 Ip YT, Davis RJ. Signal transduction by the c-Jun N-terminal kinase (JNK)—from inflammation to development. *Curr Opin Cell Biol* 1998;**10**:205–19.
- 35 Leppa S, Bohmann D. Diverse functions of JNK signaling and c-Jun in stress response and apoptosis. *Oncogene* 1999;**18**:6158–62.
- 36 Schaeffer HJ, Weber MJ. Mitogen-activated protein kinases: specific messages from ubiquitous messengers. *Mol Cell Biol* 1999;**19**:2435–44.

- 37 **Zhang Y**, Neo SY, Wang X, *et al.* Axin forms a complex with MEKK1 and activates c-Jun NH(2)-terminal kinase/stress-activated protein kinase through domains distinct from Wnt signaling. *J Biol Chem* 1999;**274**:35247–54.
- 38 **Zhang Y**, Neo SY, Han J, *et al.* Dimerization choices control the ability of axin and dishevelled to activate c-Jun N-terminal kinase/stress-activated protein kinase. *J Biol Chem* 2000;**275**:25008–14.
- 39 **Neo SY**, Zhang Y, Yaw LP, *et al.* Axin-induced apoptosis depends on the extent of its JNK activation and its ability to down-regulate beta-catenin levels. *Biochem Biophys Res Commun* 2000;**272**:144–50.
- 40 **Clevers H**. Axin and hepatocellular carcinomas. *Nat Genet* 2000;**24**:206–8.
- 41 **Aaltonen LA**. Hereditary intestinal cancer. *Semin Cancer Biol* 2000;**10**:289–98.
- 42 **Hsu W**, Shakya R, Costantini F. Impaired mammary gland and lymphoid development caused by inducible expression of Axin in transgenic mice. *J Cell Biol* 2001;**155**:1055–64.
- 43 **Hoeflich KP**, Luo J, Rubie EA, *et al.* Requirement for glycogen synthase kinase-3beta in cell survival and NF-kappaB activation. *Nature* 2000;**406**:86–90.
- 44 **Zhang Y**, Qiu WJ, Liu DX, *et al.* Differential molecular assemblies underlie the dual function of Axin in modulating the WNT and JNK pathways. *J Biol Chem* 2001;**276**:32152–9.
- 45 **Luo W**, Lin SC. Axin: a master scaffold for multiple signaling pathways. *Neurosignals* 2004;**13**:99–113.
- 46 **Luo W**, Ng WW, Jin LH, *et al.* Axin utilizes distinct regions for competitive MEKK1 and MEKK4 binding and JNK activation. *J Biol Chem* 2003;**278**:37451–8.
- 47 **Lustig B**, Jerchow B, Sachs M, *et al.* Negative feedback loop of Wnt signaling through upregulation of conductin/axin2 in colorectal and liver tumors. *Mol Cell Biol* 2002;**22**:1184–93.
- 48 **Yan D**, Wiesmann M, Rohan M, *et al.* Elevated expression of axin2 and hnk1 mRNA provides evidence that Wnt/beta-catenin signaling is activated in human colon tumors. *Proc Natl Acad Sci U S A* 2001;**98**:14973–8.
- 49 **Lee E**, Salic A, Kruger R, *et al.* The roles of APC and axin derived from experimental and theoretical analysis of the Wnt pathway. *PLoS Biol* 2003;**E10**.
- 50 **Cong F**, Varmus H. Nuclear-cytoplasmic shuttling of Axin regulates subcellular localization of beta-catenin. *Proc Natl Acad Sci U S A* 2004;**101**:2882–7.
- 51 **Yamamoto H**, Kishida S, Uochi T, *et al.* Axil, a member of the Axin family, interacts with both glycogen synthase kinase 3beta and beta-catenin and inhibits axis formation of *Xenopus* embryos. *Mol Cell Biol* 1998;**18**:2867–75.
- 52 **Mai M**, Qian C, Yokomizo A, *et al.* Cloning of the human homolog of conductin (AXIN2), a gene mapping to chromosome 17q23–q24. *Genomics* 1999;**55**:341–4.
- 53 **Liu W**, Dong X, Mai M, *et al.* Mutations in AXIN2 cause colorectal cancer with defective mismatch repair by activating beta-catenin/TCF signalling. *Nat Genet* 2000;**26**:146–7.
- 54 **Dong X**, Seelan RS, Qian C, *et al.* Genomic structure, chromosome mapping and expression analysis of the human AXIN2 gene. *Cytogenet Cell Genet* 2001;**93**:26–8.
- 55 **Anderson CB**, Neufeld KL, White RL. Subcellular distribution of Wnt pathway proteins in normal and neoplastic colon. *Proc Natl Acad Sci U S A* 2002;**99**:8683–8.
- 56 **Wiechens N**, Heinle K, Englmeier L, *et al.* Nucleo-cytoplasmic shuttling of Axin, a negative regulator of the Wnt-beta-catenin pathway. *J Biol Chem* 2004;**279**:5263–7.
- 57 **Wehrli M**, Dougan ST, Caldwell K, *et al.* Arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* 2000;**407**:527–30.
- 58 **Mao J**, Wang J, Liu B, *et al.* Low-density lipoprotein receptor-related protein-5 binds to Axin and regulates the canonical Wnt signaling pathway. *Mol Cell Biol* 2001;**21**:801–9.
- 59 **Tolwinski NS**, Wehrli M, Rives A, *et al.* Wg/Wnt signal can be transmitted through arrow/LRP5,6 and Axin independently of Zw3/Gsk3beta activity. *Dev Cell* 2003;**4**:407–18.
- 60 **Li L**, Yuan H, Weaver CD, *et al.* Axin and Frat1 interact with dvl and GSK, bridging Dvl to GSK in Wnt-mediated regulation of LEF-1. *EMBO J* 1999;**18**:4233–40.
- 61 **Hart MJ**, de los Santos R, Albert IN, *et al.* Downregulation of beta-catenin by human Axin and its association with the APC tumor suppressor, beta-catenin and GSK3 beta. *Curr Biol* 1998;**8**:573–81.
- 62 **Nakamura T**, Hamada F, Ishidate T, *et al.* Axin, an inhibitor of the Wnt signalling pathway, interacts with beta-catenin, GSK-3beta and APC and reduces the beta-catenin level. *Genes Cells* 1998;**3**:395–403.
- 63 **Kishida S**, Yamamoto H, Hino S, *et al.* DIX domains of Dvl and axin are necessary for protein interactions and their ability to regulate beta-catenin stability. *Mol Cell Biol* 1999;**19**:4414–22.
- 64 **Rubinfeld B**, Tice DA, Polakis P. Axin-dependent phosphorylation of the adenomatous polyposis coli protein mediated by casein kinase I epsilon. *J Biol Chem* 2001;**276**:39037–45.
- 65 **Hsu W**, Zeng L, Costantini F. Identification of a domain of axin that binds to the serine/threonine protein phosphatase 2A and a self-binding domain. *J Biol Chem* 1999;**274**:3439–45.
- 66 **Farr GH 3rd**, Ferkey DM, Yost C, *et al.* Interaction among GSK-3, GBP, axin, and APC in *Xenopus* axis specification. *J Cell Biol* 2000;**148**:691–702.
- 67 **Sakanaka C**, Williams LT. Functional domains of axin. Importance of the C terminus as an oligomerization domain. *J Biol Chem* 1999;**274**:14090–3.
- 68 **Nakamura Y**, Nishisho I, Kinzler KW, *et al.* Mutations of the APC (adenomatous polyposis coli) gene in FAP (familial polyposis coli) patients and in sporadic colorectal tumors. *Tohoku J Exp Med* 1992;**168**:141–7.
- 69 **Salic A**, Lee E, Mayer L, *et al.* Control of beta-catenin stability: reconstitution of the cytoplasmic steps of the wnt pathway in *Xenopus* egg extracts. *Mol Cell* 2000;**5**:523–32.
- 70 **Polakis P**. The adenomatous polyposis coli (APC) tumor suppressor. *Biochim Biophys Acta* 1997;**1332**:F127–47.
- 71 **von Kries JP**, Winbeck G, Asbrand C, *et al.* Hot spots in beta-catenin for interactions with LEF-1, conductin and APC. *Nat Struct Biol* 2000;**7**:800–7.
- 72 **Matsuzawa SI**, Reed JC. Siah-1, SIP, and Ebi collaborate in a novel pathway for beta-catenin degradation linked to p53 responses. *Mol Cell* 2001;**7**:915–26.
- 73 **Liu J**, Stevens J, Rote CA, *et al.* Siah-1 mediates a novel beta-catenin degradation pathway linking p53 to the adenomatous polyposis coli protein. *Mol Cell* 2001;**7**:927–36.
- 74 **Levina E**, Oren M, Ben-Ze'ev A. Downregulation of beta-catenin by p53 involves changes in the rate of beta-catenin phosphorylation and Axin dynamics. *Oncogene* 2004;**23**:4444–53.
- 75 **Woodgett JR**. Judging a protein by more than its name: GSK-3. *Sci STKE* 2001;**100**:RE12.
- 76 **Doble BW**, Woodgett JR. GSK-3: tricks of the trade for a multi-tasking kinase. *J Cell Sci* 2003;**116**:1175–86.
- 77 **Hedgepeth CM**, Deardorff MA, Rankin K, *et al.* Regulation of glycogen synthase kinase 3beta and downstream Wnt signaling by axin. *Mol Cell Biol* 1999;**19**:7147–57.
- 78 **Hinoi T**, Yamamoto H, Kishida M, *et al.* Complex formation of adenomatous polyposis coli gene product and axin facilitates glycogen synthase kinase-3 beta-dependent phosphorylation of beta-catenin and down-regulates beta-catenin. *J Biol Chem* 2000;**275**:34399–406.
- 79 **Amit S**, Hatzubai A, Birman Y, *et al.* Axin-mediated CKI phosphorylation of beta-catenin at Ser 45: a molecular switch for the Wnt pathway. *Genes Dev* 2002;**16**:1066–76.
- 80 **Yamamoto H**, Kishida S, Kishida M, *et al.* Phosphorylation of axin, a Wnt signaling negative regulator, by glycogen synthase kinase-3beta regulates its stability. *J Biol Chem* 1999;**274**:10681–4.
- 81 **Jho E**, Lomvardas S, Costantini F. A GSK3beta phosphorylation site in axin modulates interaction with beta-catenin and Tcf-mediated gene expression. *Biochem Biophys Res Commun* 1999;**266**:28–35.
- 82 **Tolwinski NS**, Wieschaus E. Rethinking WNT signaling. *Trends Genet* 2004;**20**:177–81.
- 83 **Julius MA**, Schelbert B, Hsu W, *et al.* Domains of axin and dishevelled required for interaction and function in wnt signaling. *Biochem Biophys Res Commun* 2000;**276**:1162–9.
- 84 **Kishida M**, Hino Si, Michiue T, *et al.* Synergistic activation of the Wnt signaling pathway by Dvl and casein kinase I epsilon. *J Biol Chem* 2001;**276**:33147–55.
- 85 **Capelluto DG**, Kutateladze TG, Habas R, *et al.* The DIX domain targets dishevelled to actin stress fibres and vesicular membranes. *Nature* 2002;**419**:726–9.
- 86 **Cliffe A**, Hamada F, Bienz M. A role of Dishevelled in relocating Axin to the plasma membrane during wingless signaling. *Curr Biol* 2003;**13**:960–6.
- 87 **Peters JM**, McKay RM, McKay JP, *et al.* Casein kinase I transduces Wnt signals. *Nature* 1999;**401**:345–50.
- 88 **Sakanaka C**, Leong P, Xu L, *et al.* Casein kinase I epsilon in the wnt pathway: regulation of beta-catenin function. *Proc Natl Acad Sci U S A* 1999;**96**:12548–52.
- 89 **Zhang Y**, Qiu WJ, Chan SC, *et al.* Casein kinase I and casein kinase II differentially regulate axin function in Wnt and JNK pathways. *J Biol Chem* 2002;**277**:17706–12.
- 90 **Willert K**, Brink M, Wodarz A, *et al.* Casein kinase 2 associates with and phosphorylates dishevelled. *EMBO J* 1997;**16**:3089–96.
- 91 **Ratcliffe MJ**, Itoh K, Sokol SY. A positive role for the PP2A catalytic subunit in Wnt signal transduction. *J Biol Chem* 2000;**275**:35680–3.
- 92 **Seeling JM**, Miller JR, Gil R, *et al.* Regulation of beta-catenin signaling by the B56 subunit of protein phosphatase 2A. *Science* 1999;**283**:2089–91.
- 93 **Li X**, Yost HJ, Virshup DM, *et al.* Protein phosphatase 2A and its B56 regulatory subunit inhibit Wnt signaling in *Xenopus*. *EMBO J* 2001;**20**:4122–31.
- 94 **Webster MT**, Rozycka M, Sara E, *et al.* Sequence variants of the axin gene in breast, colon, and other cancers: an analysis of mutations that interfere with GSK3 binding. *Genes Chromosomes Cancer* 2000;**28**:443–53.
- 95 **Shimizu Y**, Ikeda S, Fujimori M, *et al.* Frequent alterations in the Wnt signaling pathway in colorectal cancer with microsatellite instability. *Genes Chromosomes Cancer* 2002;**33**:73–81.
- 96 **Jin LH**, Shao QJ, Luo W, *et al.* Detection of point mutations of the Axin1 gene in colorectal cancers. *Int J Cancer* 2003;**107**:696–9.
- 97 **Taniguchi K**, Roberts LR, Aderca IN, *et al.* Mutational spectrum of beta-catenin, AXIN1, and AXIN2 in hepatocellular carcinomas and hepatoblastomas. *Oncogene* 2002;**21**:4863–71.
- 98 **Miao J**, Kusafuka T, Udatsu Y, *et al.* Sequence variants of the Axin gene in hepatoblastoma. *Hepatology* 2003;**25**:174–179.
- 99 **Moreno-Bueno G**, Hardisson D, Sanchez C, *et al.* Abnormalities of the APC/beta-catenin pathway in endometrial cancer. *Oncogene* 2002;**21**:7981–90.
- 100 **Nakajima M**, Fukuchi M, Miyazaki T, *et al.* Reduced expression of Axin correlates with tumour progression of oesophageal squamous cell carcinoma. *Br J Cancer* 2003;**88**:1734–9.
- 101 **Wu R**, Zhai Y, Fearon ER, *et al.* Diverse mechanisms of beta-catenin deregulation in ovarian endometrioid adenocarcinomas. *Cancer Res* 2001;**61**:8247–55.
- 102 **Daa T**, Kashima K, Kaku N, *et al.* Mutations in components of the Wnt signaling pathway in adenoid cystic carcinoma. *Mod Pathol* 2004;**17**:1475–82.

- 103 **Domingo E**, Espin E, Armengol M, *et al.* Activated BRAF targets proximal colon tumors with mismatch repair deficiency and MLH1 inactivation. *Genes Chromosomes Cancer* 2004;**39**:138–42.
- 104 **Lammi L**, Arte S, Somer M, *et al.* Mutations in AXIN2 cause familial tooth agenesis and predispose to colorectal cancer. *Am J Hum Genet* 2004;**74**:1043–50.
- 105 **Stevens MC**, Cameron AH, Muir KR, *et al.* Descriptive epidemiology of primary central nervous system tumours in children: a population-based study. *Clin Oncol (R Coll Radiol)* 1991;**3**:323–9.
- 106 **Hamilton SR**, Liu B, Parsons RE, *et al.* The molecular basis of Turcot's syndrome. *N Engl J Med* 1995;**332**:839–47.
- 107 **Zurawel RH**, Chiappa SA, Allen C, *et al.* Sporadic medulloblastomas contain oncogenic beta-catenin mutations. *Cancer Res* 1998;**58**:896–9.
- 108 **Eberhart CG**, Tihan T, Burger PC. Nuclear localization and mutation of beta-catenin in medulloblastomas. *J Neuropathol Exp Neurol* 2000;**59**:333–7.
- 109 **Huang H**, Mahler-Araujo BM, *et al.* APC mutations in sporadic medulloblastomas. *Am J Pathol* 2000;**156**:433–7.
- 110 **Yokota N**, Nishizawa S, Ohta S, *et al.* Role of Wnt pathway in medulloblastoma oncogenesis. *Int J Cancer* 2002;**101**:198–201.
- 111 **Fagotto F**, Jho E, Zeng L, *et al.* Domains of axin involved in protein-protein interactions, Wnt pathway inhibition, and intracellular localization. *J Cell Biol* 1999;**145**:741–56.
- 112 **Kinzler KW**, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;**87**:159–70.
- 113 **Lustig B**, Behrens J. The Wnt signaling pathway and its role in tumor development. *J Cancer Res Clin Oncol* 2003;**129**:199–221.
- 114 **Gardner E**. Follow-up study of a family group exhibiting dominant inheritance for a syndrome including intestinal polyps, osteomas, fibromas and epidermal cysts. *Am J Hum Genet* 1962;**14**:376–90.
- 115 **Ida M**, Nakamura T, Utsunomiya J. Osteomatous changes and tooth abnormalities found in the jaw of patients with adenomatosis coli. *Oral Surg Oral Med Oral Pathol* 1981;**52**:2–11.
- 116 **Carl W**, Sullivan MA. Dental abnormalities and bone lesions associated with familial adenomatous polyposis: report of cases. *J Am Dent Assoc* 1989;**119**:137–9.
- 117 **Aggarwal VR**, Sloan P, Horner K, *et al.* Dento-osseous changes as diagnostic markers in familial adenomatous polyposis families. *Oral Dis* 2003;**9**:29–33.
- 118 **de La Coste A**, Romagnolo B, Billuart P, *et al.* Somatic mutations of the beta-catenin gene are frequent in mouse and human hepatocellular carcinomas. *Proc Natl Acad Sci U S A* 1998;**95**:8847–51.
- 119 **Kondo Y**, Kanai Y, Sakamoto M, *et al.* Beta-catenin accumulation and mutation of exon 3 of the beta-catenin gene in hepatocellular carcinoma. *Jpn J Cancer Res* 1999;**90**:1301–9.
- 120 **Ihara A**, Koizumi H, Hashizume R, *et al.* Expression of epithelial cadherin and alpha- and beta-catenins in nontumoral livers and hepatocellular carcinomas. *Hepatology* 1996;**23**:1441–7.
- 121 **Huang H**, Fujii H, Sankila A, *et al.* Beta-catenin mutations are frequent in human hepatocellular carcinomas associated with hepatitis C virus infection. *Am J Pathol* 1999;**155**:1795–801.
- 122 **Laurent-Puig P**, Legoix P, Bluteau O, *et al.* Genetic alterations associated with hepatocellular carcinomas define distinct pathways of hepatocarcinogenesis. *Gastroenterology* 2001;**120**:1763–73.
- 123 **Greenlee RT**, Hill-Harmon MB, Murray T, *et al.* Cancer statistics, 2001. *CA Cancer J Clin* 2001;**51**:15–36.
- 124 **Saegusa M**, Okayasu I. Frequent nuclear beta-catenin accumulation and associated mutations in endometrioid-type endometrial and ovarian carcinomas with squamous differentiation. *J Pathol* 2001;**194**:59–67.
- 125 **Wright K**, Wilson P, Morland S, *et al.* Beta-catenin mutation and expression analysis in ovarian cancer: exon 3 mutations and nuclear translocation in 16% of endometrioid tumours. *Int J Cancer* 1999;**82**:625–9.
- 126 **Wagata T**, Ishizaki K, Imamura M, *et al.* Deletion of 17p and amplification of the int-2 gene in esophageal carcinomas. *Cancer Res* 1991;**51**:2113–17.
- 127 **de Castro J**, Gamallo C, Palacios J, *et al.* Beta-catenin expression pattern in primary oesophageal squamous cell carcinoma. Relationship with clinicopathologic features and clinical outcome. *Virchows Arch* 2000;**437**:599–604.
- 128 **Kang DE**, Soriano S, Xia X, *et al.* Presenilin couples the paired phosphorylation of beta-catenin independent of axin: implications for beta-catenin activation in tumorigenesis. *Cell* 2002;**110**:751–62.
- 129 **De Ferrari GV**, Chacon MA, Barria MI, *et al.* Activation of Wnt signaling rescues neurodegeneration and behavioral impairments induced by beta-amyloid fibrils. *Mol Psychiatry* 2003;**8**:195–208.
- 130 **De Strooper B**, Woodgett J. Alzheimer's disease: mental plaque removal. *Nature* 2003;**423**:392–3.
- 131 **Hanger DP**, Hughes K, Woodgett JR, *et al.* Glycogen synthase kinase-3 induces Alzheimer's disease-like phosphorylation of tau: generation of paired helical filament epitopes and neuronal localisation of the kinase. *Neurosci Lett* 1992;**147**:58–62.
- 132 **Ishiguro K**, Shiratsuchi A, Sato S, *et al.* Glycogen synthase kinase 3 beta is identical to tau protein kinase I generating several epitopes of paired helical filaments. *FEBS Lett* 1993;**325**:167–72.
- 133 **Lovestone S**, Reynolds CH, Latimer D, *et al.* Alzheimer's disease-like phosphorylation of the microtubule-associated protein tau by glycogen synthase kinase-3 in transfected mammalian cells. *Curr Biol* 1994;**4**:1077–86.
- 134 **Stoothoff WH**, Bailey CD, Mi K, *et al.* Axin negatively affects tau phosphorylation by glycogen synthase kinase 3beta. *J Neurochem* 2002;**83**:904–13.
- 135 **Spittaels K**, Van den Haute C, Van Dorpe J, *et al.* Glycogen synthase kinase-3beta phosphorylates protein tau and rescues the axonopathy in the central nervous system of human four-repeat tau transgenic mice. *J Biol Chem* 2000;**275**:41340–9.
- 136 **Sontag E**, Luangpirom A, Hladik C, *et al.* Altered expression levels of the protein phosphatase 2A A/BalphaC enzyme are associated with Alzheimer disease pathology. *J Neuropathol Exp Neurol* 2004;**63**:287–301.
- 137 **Planel E**, Miyasaka T, Launey T, *et al.* Alterations in glucose metabolism induce hypothermia leading to tau hyperphosphorylation through differential inhibition of kinase and phosphatase activities: implications for Alzheimer's disease. *J Neurosci* 2004;**24**:2401–11.
- 138 **Kins S**, Kurosinski P, Nitsch RM, *et al.* Activation of the ERK and JNK signaling pathways caused by neuron-specific inhibition of PP2A in transgenic mice. *Am J Pathol* 2003;**163**:833–43.
- 139 **Sun L**, Liu SY, Zhou XW, *et al.* Inhibition of protein phosphatase 2A- and protein phosphatase 1-induced tau hyperphosphorylation and impairment of spatial memory retention in rats. *Neuroscience* 2003;**118**:1175–82.
- 140 **Sperber BR**, Leight S, Goedert M, *et al.* Glycogen synthase kinase-3 beta phosphorylates tau protein at multiple sites in intact cells. *Neurosci Lett* 1995;**197**:149–53.
- 141 **Flaherty DB**, Soria JP, Tomasiewicz HG, *et al.* Phosphorylation of human tau protein by microtubule-associated kinases: GSK3beta and cdk5 are key participants. *J Neurosci Res* 2000;**62**:463–72.
- 142 **Goedert M**, Strittmatter WJ, Roses AD. Alzheimer's disease. Risky apolipoprotein in brain. *Nature* 1994;**372**:45–6.