

## ORIGINAL ARTICLE

## Pilot study for the development of a new campylobacter selective medium at 37°C using aztreonam

G D Thomas

*J Clin Pathol* 2005;58:413–416. doi: 10.1136/jcp.2004.019042**Aims:** To overcome contamination and temperature inhibition by isolating campylobacter at 37°C.**Methods:** The  $\beta$  lactam antibiotic aztreonam was included in a selective medium because of its inhibitory activity against Gram negative organisms but not against *Campylobacter jejuni*. Vancomycin and amphotericin were added to inhibit Gram positive bacteria and yeasts.**Results:** The aztreonam amphotericin vancomycin (AAV) experimental campylobacter selective medium showed growth microaerobically at 37°C of *C jejuni*, *C coli*, *C lari*, *C hyointestinalis*, *C fetus subsp. fetus*, and *C jejuni subsp. doylei* after 24 to 48 hours of incubation. Six campylobacter NCTC strains demonstrated a minimum inhibitory concentration (MIC)  $\geq$  256 mg/litre for vancomycin and aztreonam, whereas *C upsaliensis* and two "campylobacter-like" strains now reclassified under genus helicobacter—*H cinaedi* and *H fennelliae*—had a MIC of 4 mg/litre for vancomycin and aztreonam. In the pilot study (150 samples), AAV medium (37°C) had a higher sensitivity for isolating campylobacters: 14 were isolated on AAV compared with 10 on modified CDA (43°C) over three days, and nine were isolated on AAV medium compared with five on modified CDA (43°C) after 24 hours of incubation. Contamination rates remained low.**Conclusion:** The medium was devised in a pilot study performed between 1990 and 1993; however, this is the first report of AAV medium used as a selective medium capable of growing six campylobacters of pathogenic importance at 37°C. Further studies are indicated.Correspondence to:  
G D Thomas,  
lpi@pi37.comAccepted for publication  
15 October 2004

Campylobacter isolation requires a simple medium that uses antibiotics as selective agents. These antibiotics vary from the "cocktail" of vancomycin, polymixin B, and trimethoprim found in Skirrow's medium,<sup>1</sup> to a single antibiotic (cefazolin) in CDA,<sup>2</sup> and cefoperazone in modified CDA<sup>3</sup>; in addition, less inhibitory agents have also been advocated, such as teicoplanin in CAT medium.<sup>4</sup> However, *Campylobacter jejuni* and *C coli* remain the two main pathogens responsible for campylobacter enteritis, and early diagnosis by culture and isolation can enhance clinical investigation. The recovery of campylobacters remains dependant on the selective media and techniques used, and in their letter to the editor, Bolton *et al* raised the question: "what are we missing?"<sup>5</sup> A large number of comparative studies and evaluations of conventional methods have preceded and followed that question,<sup>6–8</sup> with McClurg and colleagues<sup>9</sup> repeating the question 15 years later; fundamentally, studies have described the temperature of 43°C as prohibitive for the growth of some *C lari* species,<sup>6</sup> and the selective pressure of the antibiotics cefalothin, colistin, and polymixin B, normally incorporated into most conventional media, as a reason for the low recovery of some strains of *C jejuni*, *C coli*, *C hyointestinalis*, and non-thermophiles.<sup>10–11</sup> More recent studies have found that filtration techniques are effective but time consuming,<sup>4</sup> and that molecular methods of detecting campylobacters using DNA probes are not ideal for routine clinical use.<sup>12</sup>

"Studies have described the temperature of 43°C as prohibitive for the growth of some *Campylobacter lari* subspecies"

\* NCTC strain *C jejuni subsp. doylei* became contaminated during the study

A preliminary study evaluating conventional methods<sup>6</sup> was inconclusive, so this present pilot study sought to establish an isolation selective medium that would support the growth of most common enteric campylobacters. The study evaluates six different antibiotics not previously used in campylobacter selective medium, with the hope that one antibiotic would prove active in microaerobic conditions, suppress Gram negative flora, and yet be non-inhibitory to *Campylobacter* spp at the incubation temperature of 37°C. Aztreonam promised to fulfilled these requirements. Furthermore, vancomycin was chosen because of its inhibitory effect on Gram positive contaminating flora, and was assessed for its antagonism to aztreonam; amphotericin was also included in the selective combination for its activity against yeasts. This pilot study for AAV medium used several challenge methods and AAV was compared with modified CDA medium to verify its subsequent suitability as a selective medium for the isolation of *Campylobacter* spp.

## METHODS AND MATERIALS

## Selective media, cultures, and incubation

Modified CDA: colombia blood agar (Oxoid CM 351; Oxoid, Basingstoke, UK) plus 7% (wt/vol) defibrinated horse blood. AAV was developed as follows: campylobacter charcoal base (Oxoid CM 908), 10 mg/litre aztreonam (Squib, Moreton, UK), dissolved in H<sub>2</sub>O (refrigerated for 24 hours before use), 2 mg/litre amphotericin (Squib), and 10 mg/litre vancomycin (hydrochloride) (Lilly, Basingstoke, UK). NCTC strains of campylobacter obtained from NCTC were as follows: *C jejuni* NCTC 11168, *C coli* NCTC 11366, *C lari* NCTC 11352, *C hyointestinalis*, *C jejuni subsp. doylei*\*, *C fetus subsp. fetus*, and *C upsaliensis* NCTC 11926. In addition, two "campylobacter-like" strains now reclassified under genus helicobacter were

**Abbreviations:** AAV, aztreonam amphotericin vancomycin; MIC, minimum inhibitory concentration

**Table 1** Suitability study showing susceptibility of *Campylobacter* spp to the six antibiotics tested

Antibiotic disc	Campylobacter strains				
	<i>C fetus</i>	<i>C jejuni</i>	<i>C hyointestinalis</i>	<i>C coli</i>	<i>C lari</i>
Imipenem (10 mg)	S	S	S	S	S
Amikacin (10 mg)	S	S	S	S	S
Tobramycin (10 mg)	S	S	S	S	S
Azlocillin (75 mg)	S	S	S	S	S
Piperacillin (75 mg)	S	S	S	S	R
Aztreonam (30 mg)	R	R	R	R	R
Aztreonam (30 mg)	<i>C upsaliensis</i>	<i>H cinaedi</i>	<i>H fennelliae</i>		
	S	S	S		

R, resistant; S, sensitive.

tested: *H fennelliae* NCTC 11612 and *H cinaedi* NCTC 11611. A reservoir of campylobacter strains and routine isolates from a preliminary study was preserved for suitability studies. Microaerobic incubation, identification, and biotyping procedures were adapted from Bolton *et al.*,<sup>6</sup> and efficacy, minimum inhibitory concentration (MIC), and challenge procedures by multipoint inoculation were adapted from Karmali *et al.*<sup>7</sup>

**Suitability studies: untried antibiotics and antagonism**

The suitability of the untried antibiotics was assessed by a disc diffusion method, in which six commercially prepared discs (Mast, Bootle, UK) chosen for their inhibitory action against Gram negative organisms were placed on to each lawned control strain. The discs contained the following antibiotics: tobramycin (10 mg), azlocillin (75 mg), imipenem (10 mg), amikacin (10 mg), piperacillin (75 mg), and aztreonam (30 mg), and zones of inhibition were recorded. Any possible antagonism when in combination with vancomycin was established by placing the commercially prepared discs of aztreonam (30 mg) and vancomycin (30 mg) 15 mm apart on non-selective blood plates, lawned with Oxford staphylococcus NCTC 6571 and *Escherichia coli* NCTC 10418 as controls, and inhibition zones were noted for radial symmetry.

**Evaluation of prototype AAV medium**

The suitability of the selective prototype medium was assessed by comparing it with modified CDA; 454 routine isolates—of which 385 were coliforms and 69 Gram positives—were spot inoculated in 0.3 µl volumes by a multipoint inoculation technique (Mast) on to modified CDA and AAV medium, growth selected, and saved before identification and MIC determination of aztreonam and vancomycin.

One hundred and fifty hospital faecal samples were routinely cultured and all plates incubated microaerobically for three days with AAV medium at 37°C and modified CDA at 43°C. Procedures were standardised with *C jejuni* NCTC 11168 and plates read on successive days for three days. Suspect campylobacter growth was confirmed by oxidase, Gram film, motility, and characteristic morphology, with isolates on AAV medium saved for identification and MIC determination.

**Table 3** Numbers of campylobacter strains isolated over three days of incubation on modified CDA (43°C) and AAV (37°C) medium (150 stool samples)

Medium	Day 1	Day 2	Day 3	Total (%)
	No of isolates	No of isolates	No of isolates	
Modified CDA	5	4	1	10 (7%)
AAV	9	4	1	14 (9%)

AAV, aztreonam amphotericin vancomycin.

**Evaluation of prototype: growth efficacy, cephalothin, and thermotolerance**

Methods used a comparison of AAV medium with modified CDA, with serial 10 fold dilutions of NCTC campylobacter strains to 10<sup>-10</sup> (vol/vol) spot inoculated by the multipoint method on to modified CDA and AAV media. All plates were incubated microaerophilically at 37°C for 24 hours, and results were recorded as the lowest dilution to show visible growth. Campylobacters not isolated by modified CDA, but which grew on AAV at 37°C, were re-challenged for susceptibility (inoculum diluted to 10<sup>-10</sup> by plate-out) by plating on to non-selective charcoal agar at 37°C at 43°C using a cefradine disc of 30 mg, and incubated under microaerobic conditions for 48 hours.

**MIC, biotyping, and identification**

Doubling dilutions of the antibiotics vancomycin and aztreonam were prepared in a charcoal agar base in the range 0.2–256 mg/litre, and preserved NCTC campylobacter strains, clinical strains, and routine contaminants that grew on AAV medium (37°C) were recultured and spot inoculated on to each set of prepared plates. Growth was recorded by Mastscan and MIC values read as the lowest concentration giving visible growth. The coliforms were identified using the API E kit (bioMérieux, Marcy-L’Etoile, France), and a simplified biotyping scheme adapted from Penner<sup>13</sup> was used for the identification of campylobacters.

**Table 2** Challenge studies showing growth of routine isolates (using multipoint) tested against modified CDA (43°C) and AAV (37°C) medium

Medium	Total no of organisms tested	Total no of coliforms	Total no of Gram positives	Coliforms resistant on multipoint set	Gram positives resistant on multipoint set	Total resistant organisms
Modified CDA	454	385	69	16 (3%)	2 (0.5%)	18 (4%)
AAV	454	385	69	22 (5%)	0	22 (5%)

AAV, aztreonam amphotericin vancomycin.

**Table 4** Numbers of plates contaminated over three days of incubation on modified CDA (43°C) and AAV medium (37°C) (150 stool samples)

	Day 1		Day 2		Day 3	
	Modified CDA	AAV	Modified CDA	AAV	Modified CDA	AAV
±	–	–	1	7	2	4
+	4	5	6	9	3	8
++	10	3	2	2	4	–
+++	7	–	1	–	–	3
TL	–	–	–	10	1	6
Total	21 (14%)	8 (5%)	10 (7%)	28 (19%)	10 (7%)	21 (14%)

AAV, aztreonam amphotericin vancomycin; TL, thin layer; ±, very scanty growth; +, scanty growth; ++, moderate growth; +++, heavy growth.

## RESULTS

### Suitability studies

The aztreonam disc (30 mg) showed no inhibitory action against *C jejuni* NCTC 11168, *C coli* NCTC 11366, *C lari* NCTC 11352, *C jejuni subsp. doylei*, *C hyointestinalis*, *C fetus subsp. fetus* (table 1), although it did not inhibit *C upsaliensis* NCTC 11926, *H fennelliae* NCTC 11612, and *H cinaedi* NCTC 11611. A combination of aztreonam and vancomycin showed no antagonism and supported the growth of all campylobacter NCTC strains with the exception of *C upsaliensis* NCTC 11926, *H fennelliae* NCTC 11612, and *H cinaedi* NCTC 11611; however a “wild-type” clinical strain of *C upsaliensis* grew on AAV medium after 48 hours of incubation (table 1).

### Challenge studies: isolation and contamination

No significant difference was shown between the selectivity of both media against Gram negative bacteria: 22 coliforms grew on AAV medium compared with 16 on modified CDA; however, the growth of all Gram positive bacteria tested was inhibited by AAV medium (table 2). From 150 routine patient faecal samples, 14 (9%) campylobacters were isolated on AAV medium at 37°C, nine of which grew after 24 hours of incubation, compared with 10 (7%) campylobacters on modified CDA (43°C), five of which grew after 24 hours of incubation (table 3). After one day of incubation, contamination on AAV medium (37°C) remained minimal, with 5% of plates demonstrating mainly coliform contaminants—for example, *E coli*, *Enterobacter cloacae*, *Citrobacter freundii*, and a rare colony of *Acinetobacter* spp, compared with 14% contamination on modified CDA (43°C). If contamination was assessed separately for each day, a reverse trend was seen on the second and third days when 19% and 14% of AAV plates (37°C), respectively, showed contamination by facultative aerobic organisms (such as *Lactobacilli* spp, *Enterococcus* spp, and yeasts) compared with 7% of modified CDA plates (43°C) on both days (table 4).

### Growth efficacy, biotyping, and MIC values

AAV medium proved slightly more successful in supporting the growth of campylobacter NCTC strains, but *C upsaliensis* NCTC 11926, *H fennelliae* NCTC 11612, and *H cinaedi* NCTC 11611 did not grow on this medium. Of the 14 campylobacters isolated, 10 were identified as *C jejuni*; two as *C coli*, one of which was isolated on both AAV (37°C) and modified CDA (43°C) on the third day of incubation (table 3), and the second of which was isolated only on AAV (37°C); the two remaining campylobacters that grew were identified as *C jejuni subsp. doylei*. All campylobacter NCTC strains showed a MIC  $\geq$  256 mg/litre to aztreonam and vancomycin, with the exception of *C upsaliensis* NCTC 11926, *H fennelliae* NCTC 11612, and *H cinaedi* NCTC 11611, each with a MIC of 4 mg/litre to aztreonam and vancomycin (table 5). The MIC values for campylobacter strains from earlier work and the pilot study were within the range of 128–256 mg/litre, with one exception being provided by a *C jejuni* isolate from a child, which had a MIC value of 128 mg/litre. The eight coliforms that failed to grow when re-challenged on AAV medium gave MIC values of 2–128 mg/litre for aztreonam and a MIC value  $\geq$  256 mg/litre for vancomycin; in addition, the four *Lactobacilli* spp, with MIC values of 32–64 mg/litre for vancomycin and MIC values of 128–256 mg/litre for aztreonam, were sensitive when re-challenged. An *Enterococcus* spp re-grew on AAV medium giving a MIC value of  $\geq$ 256 mg/litre for both aztreonam and vancomycin.

## DISCUSSION

No one medium fulfils all the criteria for the isolation of campylobacter; this phrase remains the topic for a discussion for campylobacter isolation. In this study, aztreonam proved to be non-inhibitory for six campylobacter NCTC reference strains and, importantly, vancomycin was found to be non-antagonistic to aztreonam. Therefore, it was initially thought that campylobacters might be inherently resistant to aztreonam, because antibiotic susceptibility studies by Rittenbury<sup>14</sup> had shown that the  $\beta$  lactam antibiotic aztreonam bound poorly to the penicillin binding protein sites of microaerophilic and aerobic bacteria. However, this theory was quickly disproved with the sensitivity of *C upsaliensis* NCTC 11926 to aztreonam. AAV medium supported the growth of six NCTC campylobacter reference strains and grew a “wild-type” clinical strain of *C upsaliensis*, and some complimentary antibiotic activity was exhibited between the antibiotics aztreonam and vancomycin. This last fact would enhance selective pressure and provide an explanation for the equal activity of AAV medium to modified CDA in inhibiting the growth of coliforms and Gram positive contaminants.

In a small comparative study (150 samples), AAV (37°C) was more successful in isolating campylobacters on the first day of incubation: nine campylobacters were isolated on AAV (37°C) medium compared with five on modified CDA (43°C). Medium efficacy provides one explanation in that AAV medium showed more ability to support the growth of

**Table 5** Identification and minimum inhibitory concentration (MIC) values of campylobacter reference strains

Campylobacter reference strains	Identification	MIC vancomycin	MIC aztreonam
<i>C jejuni</i> * NCTC 11168	<i>C jejuni</i>	>256	>256
<i>C lari</i> NCTC 11352	<i>C lari</i>	>256	>256
<i>C coli</i> NCTC 11366	<i>C coli</i>	>256	>256
<i>C fetus subsp. fetus</i>	<i>C fetus</i>	>256	>256
<i>H (C) fennelliae</i> NCTC 11612	<i>H (C) fennelliae</i>	>256	4
<i>H (C) cinaedi</i> NCTC 11611	<i>H (C) cinaedi</i>	>256	4
<i>C upsaliensis</i> NCTC 11926	<i>C upsaliensis</i>	>256	4
<i>C hyointestinalis</i>	<i>C hyointestinalis</i>	>256	>256

\*NCTC strain *C jejuni subsp. doylei* became contaminated during the study.

### Take home messages

- AAV (aztreonam amphotericin vancomycin) campylobacter selective medium supported the growth of six campylobacter strains, although it did not support the growth of the *C upsaliensis* NCTC strain
- In the pilot study (150 samples), AAV medium improved rates of isolation, particularly on the first day, which could be clinically relevant
- Contamination rates were low
- Further studies are indicated

*Campylobacter* spp, but temperature and cephalothin tolerance studies provided no clear indication of why four more campylobacters were isolated on AAV medium at 37°C. All four strains grew equally well at 37°C and 43°C on non-selective charcoal medium and all were resistant to cephalothin. Equally, each isolate gave MIC values  $\geq 256$  mg/litre for vancomycin and aztreonam, but one isolate was identified as *C coli*, which is generally more fastidious than *C jejuni*. It would have been useful to have estimated MIC values for cefoperazone, but time limits did not permit this. AAV medium (37°C) showed good selective pressure to faecal flora, with 5% of plates contaminated after the first day of incubation compared with 14% on modified CDA (43°C); however, a change in the preparation of the AAV plates (the addition of 1 ml of 7% horse blood) affected 30–40 plates and coincided with an increase number of contaminants growing on the second and third day of incubation. This reverse trend was temporary, and contaminants were identified as resistant *Lactobacilli* spp and diphtheroids, which had MIC values  $> 256$  mg/litre for aztreonam and vancomycin. Because these MIC values differed from the other *Lactobacilli* spp isolated in the study, these organisms may have derived from the reagents and not from the faeces. However, it was shown that blood protein probably bound to the deoxycholate component of the charcoal basal agar, reducing the inhibitory effect of deoxycholate over contaminants resistant to aztreonam and vancomycin. Once this procedure was discontinued, and 7% horse blood omitted from the preparation of the AAV plates, rates of contamination returned to the same proportions seen on the first day of incubation; however, the overall results were slightly distorted because of this additional procedure.

“This early confirmation of culture and isolation could remove some of the delays seen at present and provide the clinician with an option for rapid treatment”

Of the eight coliform contaminants sensitive when re-challenged on AAV, records showed that one *E coli* with a MIC of 128 mg/litre was derived from a patient receiving multiple antibiotic treatment; thus, acquired antibiotic

resistance<sup>15</sup> is one possible explanation for the initial growth on AAV medium. With the exception of one *C jejuni* isolate with a MIC value of 128 mg/litre for aztreonam, campylobacter NCTC strains, reserve strains, and those isolated from the study (50 strains) all had MIC values  $\geq 256$  mg/litre for vancomycin and aztreonam. The *C upsaliensis* NCTC 11926 with a MIC value of 4 mg/litre was the only campylobacter sensitive to aztreonam, and genetic modification<sup>15</sup> may provide an explanation of why there was growth of the wild-type clinical strain in the suitability studies.

AAV medium at 37°C improved rates of isolation on the first day of incubation; this early confirmation of culture and isolation could remove some of the delays seen at present and provide the clinician with an option for rapid treatment. Use of the medium enabled the easy recognition of campylobacter growth, with the culture presenting as a silver grey swarming colony, and facilitated the isolation of *C coli*, *C fetus subsp. fetus*, *C lari*, *C hyointestinalis*, and *C jejuni subsp. doylei* without the conventional methods of filtration or the need of incubation at 43°C. The pilot study was small (150 samples) and aztreonam was slightly denatured by acidity under in vitro conditions; the buffering of the medium at 37°C, the type of sample, or the clinical history of the patient may be the responsible factor. Therefore, further studies are needed.

G D Thomas, previously of Sheffield Children's Hospital (UK)

### REFERENCES

- 1 **Skirrow MB**. Campylobacter enteritis. A new disease. *BMJ* 1977;**25**:10–12.
- 2 **Bolton FJ**, Hutchinson DN, Coates G. Blood-free selective medium for isolation of *Campylobacter jejuni* from faeces. *J Clin Microbiol* 1984;**19**:169–71.
- 3 **Hutchinson DN**, Bolton FJ. An improved blood free selective medium for the isolation of *Campylobacter jejuni* from faecal specimens. *J Clin Pathol* 1984;**37**:956–7.
- 4 **Aspinall ST**, Waring DR, Hayward PG, *et al*. A comparison of a new selective medium (CAT) with membrane filter for the isolation of thermophilic campylobacter including *C. upsaliensis*. *J Appl Bacteriol* 1996;**80**:645–50.
- 5 **Bolton FJ**, Hutchinson DN, Parker G. Isolation of campylobacter; what are we missing? *J Clin Pathol* 1987;**40**:702–3.
- 6 **Bolton FJ**, Hutchinson DN, Parker G. Reassessment of selective agars and filtration technique for isolation of campylobacter species from faeces. *Eur J Clin Microbiol Infect Dis* 1988;**7**:155–60.
- 7 **Karmali MA**, Simor AE, Roscoe M, *et al*. Evaluation of a blood-free, charcoal-based selective medium for the isolation of campylobacter organisms from faeces. *J Clin Microbiol* 1986;**23**:456–59.
- 8 **Engberg J**, On SLW, Harrington CS, *et al*. Prevalence of *Campylobacter*, *Archobacter*, *Helicobacter* and *Sutterella* spp. in human faecal samples as estimated by a re-evaluation of isolation methods for campylobacters. *J Clin Microbiol* 2000;**38**:286–91.
- 9 **McClurg KR**, McClurg RB, Moore JE. Efficient isolation of campylobacters from stools: what are we missing? *J Clin Pathol* 2002;**55**:239–40.
- 10 **Corry JE**, Post DE, Colin P, *et al*. Culture media for the isolation of campylobacters. *Int J Food Microbiol* 1995;**26**:43–76.
- 11 **Fennell CL**, Totton PR, Quinn TC, *et al*. Characterisation of campylobacter-like organisms from homosexual men. *J Infect Dis* 1984;**149**:58–66.
- 12 **Maher M**, Finnigan C, Collins E, *et al*. Evaluation of culture methods and a DNA probe-based PCR assay for detection of campylobacter species in clinical specimens of faeces. *J Clin Microbiol* 2003;**41**:2980–6.
- 13 **Penner JL**. The genus campylobacter: a decade of progress. *Clin Microbiol Rev* 1988;**1**:157–72.
- 14 **Rittenbury MS**. “How and why aztreonam works.” *Surg Gynecol Obstet Supp* 1990;**171**:19–23.
- 15 **Lau SKP**, Woo PCY, Leung KW, *et al*. Emergence of cotrimoxazole and quinolone resistant campylobacter infections in bone marrow transplant recipient. *Eur J Clin Infect Dis* 2002;**21**:286–91.