



Published in final edited form as:

J Mol Biol. 2006 December 8; 364(4): 667–677.

***Thermus thermophilus* bacteriophage ϕ YS40 genome and proteomic characterization of virions**

Tatyana Naryshkina^{1,#}, Jing Liu^{2,#}, Laurence Florens², Selene K. Swanson², Andrey R. Pavlov³, Nadejda V. Pavlova³, Ross Inman⁴, Sergei A. Kozyavkin³, Michael Washburn², Arcady Mushegian^{2,5}, and Konstantin Severinov^{1,6,7,*}

¹From the Waksman Institute for Microbiology, Kansas City, MO 64110

²From the Stowers Institute for Medical Research, Kansas City, MO 64110

³From the Fidelity Systems, Inc., Gaithersburg, MD 20879

⁴From the Institute for Molecular Virology, University of Wisconsin, Madison, WI, 53706

⁵From the Department of Microbiology, Kansas University Medical Center, Kansas City KS 66160

⁶From the Department of Molecular Biology and Biochemistry, Rutgers, the State University of New Jersey, Piscataway, NJ, 08854

⁷From the Institute of Molecular Genetics, Russian Academy of Sciences, Moscow, 123182 Russia

Abstract

We determined the sequence of the 152,372-bp genome of ϕ YS40, a lytic tailed bacteriophage of *Thermus thermophilus*. The genome contains 170 putative open reading frames and three tRNA genes. Functions for 25% of ϕ YS40 gene products were predicted on the basis of similarity to proteins of known function from diverse phages and bacteria. ϕ YS40 encodes a cluster of proteins involved in nucleotide salvage, such as flavin-dependent thymidylate synthase, thymidylate kinase, ribonucleotide reductase, and deoxycytidylate deaminase, and in DNA replication, such as DNA primase, helicase, type A DNA polymerase, and predicted terminal protein involved in initiation of DNA synthesis. The structural genes of ϕ YS40, most of which have no similarity to sequences in public databases, were identified by mass-spectrometric analysis of purified virions. Various ϕ YS40 proteins have different phylogenetic neighbors, including Myovirus, Podovirus, and Siphovirus gene products, bacterial genes, and in one case, a dUTPase from a eukaryotic virus. ϕ YS40 has apparently arisen through multiple acts of recombination between different phage genomes as well as through acquisition of bacterial genes.

Keywords

Thermus thermophilus; bacteriophage; genome; virion; proteomics; bioinformatics; DNA polymerase

Introduction

In the last decade, the genomes of several hundred phages have been completely sequenced (282 complete dsDNA phage genomes in the Genome Division of GenBank as of July 2006). While bacterial hosts of these phages are phylogenetically diverse, only ten of those completely sequenced phages are known to infect thermophilic microorganisms. Most of 'thermophilic'

#These authors contributed equally to this work

*Corresponding author Waksman Institute for Microbiology, 190 Frelinghuysen Road, Piscataway, NJ, 08854 Phone: (732) 445-6095, FAX: (732) 445-5735, E-mail: severik@waksman.rutgers.edu

phages were isolated from a small number of archaeal species¹⁻³. Sequence analysis revealed that archaeophages encode mostly uncharacterized proteins with no similarities to sequences in public databases, though more detailed examination revealed a limited number of recognizable ATPases, nucleotide salvage enzymes, and putative transcription factors⁴. As of the time of this writing, the only sequenced genome of a phage from a thermophilic eubacterium is RM 378 that infects *Rhodothermus marinus*⁵.

During their development in a bacterial host, phages are known to regulate host macromolecular synthesis by modifying host transcription and translation machinery and making it serve the needs of the virus. Proteins from thermophilic bacteria are particularly amenable to structural studies of large complexes involved in DNA replication, DNA transcription, and RNA translation. Thus, structural and functional analysis of thermophilic phage-encoded regulators and their complexes with RNA polymerases, ribosomes, and other components of thermophilic bacteria can provide insights into molecular mechanisms of regulation of transcription, translation, and other cellular processes. With these ideas in mind, we determined the genomic sequence of ϕ YS40, a large myophage hosted by the thermophilic bacterium *Thermus thermophilus* (temperature range from 56 to 78°C)⁶. Here, we present the results of a preliminary study of the ϕ YS40 genome and the proteome of ϕ YS40 virions.

Results

Overview of the ϕ YS40 genome

The sequence of the ϕ YS40 genome was determined using the fimer technology and assembled into a single 152,372 bp contig using the phredPhrap package (see Materials and Methods). The G + C content of the ϕ YS40 genome is 32.59%, which is significantly lower than that of its host (69.4%). Though the GC-content of ϕ YS40 is close to values typical of the low-GC Gram-positive bacteria, there is no specific evolutionary affinity between sequences of ϕ YS40 and these bacteria, and the GC-content of the phage may instead reflect specific aspects of phage molecular biology, for example distinct mutational bias of its DNA polymerase. ϕ YS40 DNA appears to be unmodified as it is susceptible to digestion with all common methylation-sensitive restriction endonucleases tested (data not shown).

A total of 170 ORFs were predicted in the ϕ YS40 genome (Table 1, Fig. 1). The intergenic regions were screened for additional genes by searching GenBank, GenPept, and the database of unfinished microbial genomes at NCBI, but no additional conserved ORFs were found. The predicted ϕ YS40 ORFs are between 43 and 1744 codons in length. As with most other phages, the genome of ϕ YS40 is tightly packed: coding sequences occupy 95% of the ϕ YS40 genome. There are 46 cases of overlaps (from 1 to 40 bases long) between neighboring ORFs. The longest non-coding region (390 bp) lies between ORF138 and ORF139. Most of the 170 predicted ORFs start at the AUG codon, 22 ORFs use GUG codon, and three use UUG. At the ends of ϕ YS40 genes, there are 90 TAA stop codons, 66 TGA codons, and 16 TAG codons.

Two-thirds of the ϕ YS40 genes (114 genes) are transcribed in one direction, designated as leftward in the genome map (Fig. 1), and 56 genes are transcribed in the rightward direction. The G+C content is approximately the same for both sets of ORFs. Taking a set of genes transcribed in the same direction and having no more than three consecutive intruders (*i.e.*, genes transcribed in a different direction) as a cluster, we find four gene clusters in the ϕ YS40 genome. The ORF1-ORF36 and ORF62-ORF146 clusters are transcribed in the leftward direction, and ORF37-ORF61 and ORF147-ORF170 clusters are transcribed in the rightwards direction (Fig. 1). The probability of obtaining each of the four clusters by chance, calculated using equation 2 from Durand and Sankoff⁷ is less than 0.1, indicating that at least part of the clustering may be due to evolutionary or functional constraints.

tRNA genes

Using the tRNA scan-SE program, we identified three tRNA genes in the ϕ YS40 genome. The tRNA1 gene overlaps with ORF61, whereas the tRNA2 and tRNA3 genes are both located between ORF139 and ORF140. Other large tailed dsDNA bacteriophages, such as coliphage T4⁸, vibriophage KVP40⁹, and phage phiKZ of *P. aeruginosa*¹⁰ also encode several tRNAs.

The ϕ YS40 tRNA1 and tRNA3 recognize ACA (threonine) and AGA (arginine) codons, respectively. These codons, while overrepresented in the ϕ YS40 genome, are the rarest threonine and arginine codons in *T. thermophilus* genes. tRNA2 has a CAU anticodon, which would correspond to methionine codon AUG if C34 in the wobble position is unmodified. In homologous tRNAs from a number of bacteria and bacteriophages, the corresponding cytidine is converted to lysidine, which results in the AUA (Ile) decoding¹¹⁻¹³. Determinants for tRNA^{Ile} identity are thought to consist of anticodon loop bases A37 and A38, the discriminator base A73, and conserved base pairs in the D-arm (U12-A23), the anticodon arm (C29-G41), and the acceptor arm (C4-G69)¹⁴. All these characteristics are present in ϕ YS40 tRNA2, which therefore may decode the isoleucine codon AUA, another rare *T. thermophilus* codon that is much more frequent in ϕ YS40 ORFs. Thus, ϕ YS40-encoded tRNAs may ensure efficient decoding of codons that are overrepresented in the phage genome relative to its host.

Sequence analysis of predicted ϕ YS40 proteins

Analysis of intrinsic features of protein sequences indicates that seven ϕ YS40 ORFs encode proteins with putative transmembrane domains (from one to three) and four ϕ YS40 proteins are predicted to have coiled-coil regions. Only one protein, gp107, is predicted to be strongly non-globular, and only one protein, gp35, contains an N-terminal secretion signal peptide. All deduced amino acid sequences were compared to proteins in the non-redundant database at NCBI using the PSI-BLAST program with a slightly relaxed cutoff for profile inclusion (-h parameter). The comparison showed that ~25% of ϕ YS40 proteins display sequence similarity to proteins of known function (Table 1).

ϕ YS40 proteins involved in nucleotide metabolism—Like other large phage genomes, ϕ YS40 encodes a number of enzymes involved in nucleotide metabolism. They are gp8, a homolog of mammalian/viral UTPase (EC 3.6.1.23); gp9, related to a predicted flavin-dependent thymidylate synthase (EC 2.1.1.148); GMP reductase gp17 (EC 1.7.1.7); thymidine kinase gp24 (EC 2.7.1.21); deoxycytidylate deaminase gp38 (EC 3.5.4.12); dNMP kinase gp60 (EC 2.7.4.-); and the catalytic α subunit of ribonucleotide reductase encoded by two adjoining ORFs, gp41 and gp42 (EC 1.17.4.1). Except for dUTPase gp8, all these gene products show stronger sequence similarity to prokaryotic or phage enzymes than to their eukaryotic or archaeal counterparts. The best database match and closest phylogenetic neighbor for dUTPase gp8 is dUTPase from *Lymantria dispar* nucleopolyhedrosis virus. Gene exchange between phages and bacteria has been suggested to account for odd gene phylogenies that are sometimes observed in the components of bacterial replication and transcription machinery¹⁵. Our observation indicates that eukaryotic viruses, and perhaps their hosts, may also be involved in such exchange.

ϕ YS40 proteins involved in DNA replication and recombination— ϕ YS40 encodes most of the proteins required for replisome formation, namely gp14, a replication initiation helicase DnaB; gp23, a bacterial DnaG-family DNA primase; gp26, a RecB family exonuclease; gp33, a type A DNA polymerase, and gp27, a DEAD box helicase. Another predicted DEAD-box helicase is encoded by gp79. Based on the fact that gp79 is a part of the ϕ YS40 virion, we suspect that it is involved in viral DNA packaging. ϕ YS40 also encodes two recombination proteins, gp12, a RecA/RadA recombinase, and gp114, an ssDNA-

annealing protein of the ERF family. There are no gene products with detectable sequence similarity to known ssDNA-binding proteins¹⁶ or DNA ligases.

The product of gene 65 is of particular interest for understanding the replication mechanism of ϕ YS40. It shows a striking sequence similarity to a portion of the terminal protein (TP) of *B. subtilis* phage ϕ 29. The Ser²³² residue of the TP protein forms a phosphoester bond with the 5'-terminal dAMP of the phage genome, and is essential for protein-primed replication of linear dsDNA genome of ϕ 29¹⁷⁻¹⁹. This serine is conserved in ϕ YS40 gp65 (Fig. 2). Thus, it is likely that gp65 primes the replication of ϕ YS40 genomic DNA. It should be noted that the ends of the ϕ YS40 genome as presented in Fig. 1 are arbitrary, since no defined ends were revealed during genome sequencing and assembly, indicating that the ϕ YS40 genome may be circularly permuted or may have direct terminal repeats. This matter requires further investigation.

Properties of the ϕ YS40 DNA polymerase—The ϕ YS40 gp33 is a type A DNA polymerase, which contains a conserved nucleotidyltransferase domain and a 3'-5' exonuclease domain, but lacks the 5'→3' exonuclease domain. Since gp33 is the first known example of a type A DNA polymerase from a thermophilic phage, we expressed recombinant gp33 in *E. coli* and studied its properties *in vitro*. At 60-65 °C, recombinant gp33 exhibited moderate polymerization activity and very strong 3'→5' exonuclease activity toward both single-stranded DNA and double-stranded DNA substrates, even in the presence of 1 mM dNTP. As a result, at pH > 8.0 and low salt concentrations, the enzyme mostly hydrolyzed the primer. The increase of salt concentration partially inhibited the exonucleolytic activity and allowed primer elongation, until further increase inhibited the polymerase activity as well. The decay of primer-template substrate by gp33 exonuclease was abolished when primers were protected with thiolate modification, but the interference of the exonucleolytic activity during elongation resulted in poor DNA yield.

Gp33 was moderately thermostable. Both polymerase and exonuclease functions were lost after a 3-min incubation at 85 °C. At 75 °C, the polymerase activity decreased faster than the exonuclease activity; as a result, the enzyme produced shorter elongation products after heating. Similarly low thermostability has been reported for type B DNA polymerase from the *Rhodothermus marinus* phage (a half-life of 2 min at 90 °C⁵). These observations indicate that both processivity of ϕ YS40 DNA polymerase and its stability at elevated temperatures must be conferred by its interactions with other components of the replicative complex, in marked contrast with other DNA polymerases of bacteria and archaea, such as *Taq* or *Pfu*, which are processive and thermostable in the absence of cofactors.

Protein composition of ϕ YS40 virions—To identify ϕ YS40 structural proteins, ϕ YS40 virions were purified by double sedimentation in CsCl gradients. The results of SDS-PAGE analysis of purified ϕ YS40 virions are shown in Fig. 3. The two major protein components of the virion were identified by mass spectrometry as gp73 and gp19 (Fig. 3). These proteins may correspond to major head and tail proteins, but their function could not have been predicted by sequence comparison because of lack of database homologs.

Three independent ϕ YS40 lysates of increasing titer (from 2×10^7 to 2×10^9 pfu/ml) were also directly examined by multidimensional protein identification technology, MudPIT²⁰ a shotgun proteomics approach where proteolytic peptides of a protein complex under study (in our case, phage virions) are generated, loaded onto triphasic microcapillary columns, eluted over several chromatography steps and analyzed directly by tandem mass spectrometry. Peptides matching 33 ϕ YS40 proteins were detected in one or more of these samples. There were also 79 host proteins, all of which decreased in abundance when the lysates of higher titer were used as a starting material for CsCl purification (Supplementary Table A). In contrast,

the NSAF (Normalized Spectral Abundance Factor, see Materials and Methods) values for ϕ YS40 proteins increased with the titer of phage in the starting sample (Supplementary Table B). Gp73 and gp19 were detected at the highest levels in all three analyses in agreement with these being major structural proteins. With the exception of gp52 (annotated as a UDP-3-O-[3-hydroxymyristoyl] glucosamine N-acyltransferase), gp69 (tail sheath protein), gp79 (DEAD-Box helicase), gp150 (putative baseplate assembly protein), and gp152 (fibrin neck whisker), most ϕ YS40 virion proteins identified in this analysis are novel proteins without any detectable database homologs. Interestingly, all multiply detected ϕ YS40 virion proteins are the products of adjacent co-transcribed genes, except for ORF19 (Fig. 4C). In particular, a group of 13 proteins detected at high levels are the products of genes at the end of the largest cluster of ϕ YS40 genes (ORF62-ORF146, above) that therefore may correspond to the late gene cluster.

DISCUSSION

Bacteriophages may be the most abundant living entities on Earth. It has been proposed that the origin of dsDNA bacteriophages is as ancient as DNA replication itself and that the analysis of the currently known bacteriophages may provide clues to early evolution of cellular and viral genomes¹⁵.

Here, we report a preliminary analysis of *Thermus thermophilus* bacteriophage ϕ YS40 genome. The analysis shows that ϕ YS40 does not easily fit into previously established groups of dsDNA bacterial viruses and may represent a distinct branch of the *Myoviridae* family. A substantial fraction of ϕ YS40 genes codes for predicted proteins to which no function can be assigned; however, 25% of the ϕ YS40-encoded proteins show detectable homology to their counterparts in a broad phylogenetic range of microorganisms, and some proteins are homologous to proteins found in other dsDNA bacteriophages infecting diverse hosts, such as *Staphylococcus*, *Rhodothermus marinus*, and *Vibrio parahaemolyticus*. In agreement with morphological data, predicted tail genes are mostly *Myoviridae*-related. Most of other ϕ YS40 genes that have database homologs are, however, closer to either podoviral or siphoviral gene products: for instance, gp26 (RecB family exonuclease) and gp60 (dNMP kinase) are most closely related to homologs from a podovirus SIO1 and a λ -like siphovirus phi-BT1, respectively. Yet other genes are phylogenetically close to bacterial genes, and, in one case, to a homolog from a eukaryotic baculovirus. ϕ YS40 has apparently arisen through multiple acts of recombination between different groups of phages and perhaps even their hosts.

Molecular adaptations to thermophily in various species are of great interest. Comparative studies of the genomes of thermophilic, hyperthermophilic, and mesophilic prokaryotes have suggested several attributes of thermostability at the levels of amino acid sequence, properties of folded proteins, and gene content. The proposed sequence level predictors of thermostability, such as large charged-versus-polar (CvP) amino acid ratio or (E + K)/(Q + H) ratio, are not conclusive in the case of ϕ YS40, and genes that are indicative of the host ability to survive at extreme temperatures²¹ are missing from the ϕ YS40 genome. Moreover, only seven ϕ YS40 gene products have closest phylogenetic neighbors in thermophilic microorganisms.

In its genome size, ϕ YS40 is similar to T4, an *E. coli* phage that is known to rely on host RNA polymerase for expression of its genes. During its development, T4 sequentially modifies host RNA polymerase to shut off transcription of host genes and to ensure correct expression of several classes of its own genes (reviewed in Ref. 22). Like T4, ϕ YS40 does not encode its own RNA polymerase and therefore has to rely on the host enzyme for transcription of its DNA. The early genes of ϕ YS40 should therefore be transcribed by the *T. thermophilus* RNA polymerase holoenzyme, most likely containing general initiation factor σ^A . Preliminary analysis reveals the presence of sequences with strong similarities to bacterial housekeeping

sigma promoters in front of many ϕ YS40 genes, but no such sequences are found in front of genes coding for ϕ YS40 structural proteins (A. Sevostyanova, M. Gelfand and KS, unpublished observations). Structural genes, which should be expressed late in infection, must be therefore transcribed by a modified form of host RNA polymerase. Further biochemical studies may reveal ϕ YS40 proteins that are required for these modifications.

MATERIALS AND METHODS

Cell growth and phage infection

The bacterial strain *Thermus thermophilus* HB8 and ϕ YS40 were generously provided by Dr. Tairo Oshima, Tokyo University of Pharmacy & Life Science. The cells and phage were grown overnight in the Tth medium (0.8% polypeptone, 0.4% yeast extract, 0.2% NaCl, and 0.35 M CaCl₂ and 0.4 M MgSO₄) at 65°C with vigorous agitation.

To isolate individual ϕ YS40 plaques, 1 ml of overnight HB8 culture (**OD600~1.6**) was centrifuged and resuspended in 100 μ l of the Tth medium and combined with 5 μ l dilutions of ϕ YS40 stock, incubated for 15 min at 65°C, plated in soft Tth agar (0.7 %), and incubated overnight at 65°C. An individual plaque was picked up and subjected to two more rounds of plaque purification, before making a phage lysate stock solution. To this end, a single plaque was resuspended in a small volume of the Tth medium and mixed with 0.1 ml of overnight HB8 culture. The mixture was incubated for 15 minutes at 65 °C to allow phage absorption, 5 ml of fresh Tth medium was added and the culture was incubated on a rotary shaker at 65 °C until complete lysis occurred (usually overnight). Cell debris was removed from the lysate by centrifugation at 12,000g for 15 minutes. The resultant phage stock (6×10^9 pfu/ml) was saturated with chloroform and stored at 4 °C. The ϕ YS40 stock was used to prepare larger amounts of phage lysate using a scale-up of the procedure described above.

Purification of ϕ YS40 virions

DNase I and RNase A (each to a final concentration of 1 μ g/ml) were added to ϕ YS40 lysed *T. thermophilus* culture followed by a 30-min incubation at 30°C. Solid NaCl was added a final concentration of 1 M and dissolved by swirling. The lysed culture was left on ice for 1 h and centrifuged at 11,000 g for 10 min at 4°C. To precipitate ϕ YS40, PEG 8000 was added to the supernatant to the final concentration of 10% (w/v) followed by a 1-h incubation on ice. Precipitated ϕ YS40 particles were recovered by centrifugation at 11,000g for 10 min at 4 °C. The phage pellet was resuspended in 2 ml of SM buffer (NaCl, MgSO₄, Tris-HCl, pH7.5, 2% gelatin). The PEG 8000 and cell debris were extracted from the phage suspension by adding an equal volume of chloroform and centrifuged at 3,000g for 15 min at 4°C. 0.5 g of solid CsCl per milliliter of bacteriophage suspension was added to the aqueous phase, which contained the bacteriophage particles, and dissolved by gentle mixing. CsCl step gradients (three steps with 1.45, 1.50, and 1.70 g/l density) were performed in Beckman SW41 polypropylene centrifuge tubes at 22,000 rpm for 2 hrs at 4 °C and at 38,000 rpm for 24 hrs at 4 °C (Beckman SW50.1 rotor, Beckman Coulter, Fullerton, CA). Purified bacteriophage suspension was dialyzed twice at room temperature for 1 h against a 1000-fold volume of 10 mM NaCl, 50 mM Tris-HCl pH 8.0, 10 mM MgCl₂.

Extraction of phage DNA

EDTA (to a final concentration of 20 mM), proteinase K (to a final concentration of 50 μ g/ml), SDS (to a final concentration of 0.5%) were added to bacteriophage solution and incubated at 56°C for 1 h. An equal volume of phenol was added to chilled bacteriophage suspension, mixed, and centrifuged at 3000 g for 5 min at room temperature. The aqueous phase was extracted with a 50:50 mixture of equilibrated phenol and chloroform, and equal volume of chloroform. DNA was precipitated with ethanol.

Genome sequencing

Initial sequence data were obtained using mini shotgun library of phage DNA. Several rounds of sequencing reactions were performed directly on phage DNA using ThermoFidelase and Fimer technology^{23, 24}. Trace assembly was done with phredPhrap package (<http://www.phrap.com/>)²⁵. The final round of sequencing resulted in one pseudocircular contig with a no- errors quality level.

Sequence analysis

ORFs of ϕ YS40 were predicted using the GeneMark server (http://opal.biology.gatech.edu/GeneMark/heuristic_hmm2.cgi, Ref. 26). The PSI-BLAST program²⁷ was used to detect the homologs of ϕ YS40 genes in the DNA and protein databases, with profile inclusion cutoff *E*-value in PSI-BLAST (-h parameter) set at 0.02. Both options for low-complexity filtering (-F parameter) and composition-based statistics (-t parameter) were sometime adjusted for better detection in sequence similarities. Phylogenetic analysis was performed using the programs in the PHYLIP package.²⁸

tRNA genes were searched by using the tRNAscan-SE program.²⁹ Searches for the presence of the transmembrane helices and coiled coil regions were done with the aid of the SEALS package.³⁰

MudPIT

Three independent virion lysates were prepared by double sedimentation in CsCl gradients and had phage titers of 2×10^7 pfu/ml, 4.2×10^8 pfu/ml and 2×10^9 pfu/ml. These lysates were treated with for 30 minutes at 37°C with 0.1U of benzonase (Sigma, St. Louis, MO), then precipitated in 20% trichloroacetic acid, 100mM Tris-HCl, pH 8.5, overnight at 4°C. The dried protein pellets were denatured, reduced, alkylated and digested with endoproteinase LysC and trypsin (both from Roche Applied Science, Indianapolis, IN) as described previously.³¹ Peptide mixtures were pressure-loaded onto split-triphasic microcapillary columns, installed in-line with a Quaternary Agilent 1100 series HPLC pump coupled to Deca-XP ion trap tandem mass spectrometer (ThermoElectron, San Jose, CA) and analyzed via seven-step chromatography as described in Ref. 31.

The MS/MS datasets were searched using SEQUEST³² against a database of 171 YS40 predicted gene products, combined with 2224 protein sequences from *Thermus thermophilus*, strain HB8 (chromosome and large plasmid) downloaded from NCBI on 2005-08-01, as well as usual contaminants such as human keratins, IgGs, and proteases. In addition, to estimate background correlations, each sequence in the database was randomized (keeping the same amino acid composition and length) and the resulting “shuffled” sequences were concatenated to the “normal” sequences and searched at the same time (the total number of sequences searched was 5144).

DTASelect/CONTRAST program³³ was used to select spectra/peptide matches with normalized difference in cross-correlation score (DeltCn) of at least 0.11, a minimum cross-correlation score (XCorr) of 1.8 for singly-, 2.5 for doubly-, and 3.5 for triply-charged spectra, a maximum Sp rank of 10, and a minimal length of 7 amino acids. In addition, the peptides had to be fully tryptic. No peptides matching shuffled protein sequences passed this criteria set. Spectral counts are considered to be a good estimation of absolute protein abundance³⁴. To account for the fact that larger proteins tend to contribute more peptide/spectra, spectral counts are divided by protein length defining a Spectral Abundance Factor (SAF).³⁵ SAF values are normalized against the sum of all SAFs for each run (removing redundant proteins) allowing us to compare protein levels across different runs using the Normalized Spectral Abundance Factor (NSAF) value.

φY540 DNA polymerase

The gene encoding φY540 DNA polymerase was PCR amplified using appropriate primers annealing at the beginning and the end of φY540 gene 33 and containing engineered *NdeI* site CATATG overlapping with the initiating ATG codon of gene 33 and a *HindIII* site downstream of the termination codon (primer sequences are available from the authors upon request). The amplified fragment was treated with *NdeI* and *HindIII* and cloned into appropriately digested pet21d plasmid and transformed into the *E. coli* expression strain BL-21 pLysS. Cells were grown in 1 L of LB medium and induced with 1 mM IPTG. Cell pellet was dissolved in 15 ml of lysis buffer and centrifuged at 17000 rpm for 30 min (no heat treatment). Lysate was diluted to 0.25M NaCl, and applied on a Heparin Sepharose High-Trap column (GE Healthcare, Newark, NJ), equilibrated with 50 mM Tris pH 7.5, containing 0.25 M NaCl and 2 mM mercaptoethanol. After washing with the same buffer, φY540 DNA polymerase was eluted in about 0.3-0.35 M NaCl and appeared to be over 80% pure by SDS-PAGE. Assays of its enzymatic activities were done essentially as described by Pavlov et al., Ref. 36.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIH RO1 grant GM64530 (to KS) and NIH GM61898 to Seth Darst. The authors thank Galina Glazko and Frank Emmert-Streib (both from Stowers Institute) for assistance on the gene clustering analysis and the analysis on codon usage, respectively.

References

1. Palm P, Schleper C, Grampp B, Yeats S, McWilliam P, Reiter WD, Zillig W. Complete nucleotide sequence of the virus SSV1 of the archaeobacterium *Sulfolobus shibatae*. *Virology* 1991;185:242–250. [PubMed: 1926776]
2. Arnold HP, Zillig W, Ziese U, Holz I, Crosby M, Utterback T, Weidmann JF, Umayam LA, Teffera K, Kristjansson JK, Klenk HP, Nelson KE, Fraser CM. A novel lipothrixvirus, SIFV, of the extremely thermophilic crenarchaeon *Sulfolobus*. *Virology* 2000;267:252–266. [PubMed: 10662621]
3. Wiedenheft B, Stedman K, Roberto F, Willits D, Gleske AK, Zoeller L, Snyder J, Douglas T, Young M. Comparative genomic analysis of hyperthermophilic archaeal Fuselloviridae viruses. *J. Virol* 2004;78:1954–1961. [PubMed: 14747560]
4. Prangishvili D, Garrett RA, Koonin EV. Evolutionary genomics of archaeal viruses: Unique viral genomes in the third domain of life. *Virus Res* 2006;117:52–67. [PubMed: 16503363]
5. Hjorleifsdottir, S.; Hreggvidsson, GO.; Fridjonsson, OH.; Aevvarsson, A.; Kristjansson, JK. Bacteriophage RM 378 of a thermophilic host organism. Decode Genetics EHF. Patent: WO 0075335-A 14-DEC-2000. 2000.
6. Sakaki Y, Oshima T. Isolation and characterization of a bacteriophage infectious to an extreme thermophile, *Thermus thermophilus* HB8. *J. Virol* 1975;15:1449–1453. [PubMed: 1142476]
7. Durand D, Sankoff D. Tests for gene clustering. *J. Comput. Biol* 2003;10:453–482. [PubMed: 12935338]
8. Miller ES, Kutter EM, Mosig G, Arisaka F, Kunisawa T, Ruger W. Bacteriophage T4 genome. *Microbiol. Mol. Biol. Rev* 2003;67:86–156. [PubMed: 12626685]
9. Miller ES, Heidelberg JF, Eisen JA, Nelson WC, Durkin AS, Ciecko A, Feldblyum TV, White O, Paulsen IT, Nierman WC, Lee J, Szczypinski B, Fraser CM. Complete genome sequence of the broad-host-range vibriophage KVP40: comparative genomics of a T4-related bacteriophage. *J. Bacteriol* 2003;185:5220–5233. [PubMed: 12923095]
10. Mesyanzhinov VV, Robben J, Grymonprez B, Kostyuchenko VA, Bourkaltseva MV, Sykilinda NN, Krylov VN, Volckaert G. The genome of bacteriophage φKZ of *Pseudomonas aeruginosa*. *J. Mol. Biol* 2002;317:1–19. [PubMed: 11916376]

11. Matsugi J, Murao K, Ishikura H. Characterization of a *B. subtilis* minor isoleucine tRNA deduced from tDNA having a methionine anticodon CAT. *J. Biochem. (Tokyo)* 1996;119:811–816. [PubMed: 8743586]
12. Muramatsu T, Nishikawa K, Nemoto F, Kuchino Y, Nishimura S, Miyazawa T, Yokoyama S. Codon and amino-acid specificities of a transfer RNA are both converted by a single post-transcriptional modification. *Nature* 1988;336:179–181. [PubMed: 3054566]
13. Muramatsu T, Yokoyama S, Horie N, Matsuda A, Ueda T, Yamaizumi Z, Kuchino Y, Nishimura S, Miyazawa T. A novel lysine-substituted nucleoside in the first position of the anticodon of minor isoleucine tRNA from *Escherichia coli*. *J. Biol. Chem* 1988;263:9261–9267. [PubMed: 3132458]
14. Nureki O, Niimi T, Muramatsu T, Kanno H, Kohno T, Florentz C, Giege R, Yokoyama S. Molecular recognition of the identity-determinant set of isoleucine transfer RNA from *Escherichia coli*. *J. Mol. Biol* 1994;236:710–724. [PubMed: 8114089]
15. Filée J, Forterre P, Laurent J. The role played by viruses in the evolution of their hosts: a view based on informational protein phylogenies. *Res. Microbiol* 2003;154:237–243. [PubMed: 12798227]
16. Ponomarev VA, Makarova KS, Aravind L, Koonin EV. Gene duplication with displacement and rearrangement: origin of the bacterial replication protein PriB from the single-stranded DNA-binding protein Ssb. *J. Mol. Microbiol. Biotechnol* 2003;4:225–229. [PubMed: 12867746]
17. Hermoso JM, Méndez E, Soriano F, Salas M. Location of the serine residue involved in the linkage between the terminal protein and the DNA of phage ϕ 29. *Nucleic Acids Res* 1985;13:7715–7728. [PubMed: 3934646]
18. Garmendia C, Salas M, Hermoso JM. Site-directed mutagenesis in the DNA linking site of bacteriophage ϕ 29 terminal protein: isolation and characterization of a Ser232----Thr mutant. *Nucleic Acids Res* 1988;16:5727–5740. [PubMed: 3135531]
19. Garmendia C, Hermoso JM, Salas M. Functional domain for priming activity in the phage ϕ 29 terminal protein. *Gene* 1990;88:73–79. [PubMed: 2341040]
20. Washburn MP, Wolters D, Yates JR 3rd. Large-scale analysis of the yeast proteome by multidimensional protein identification technology. *Nat. Biotechnol* 2001;19:242–247. [PubMed: 11231557]
21. Makarova KS, Wolf YI, Koonin EV. Potential genomic determinants of hyperthermophily. *Trends Genet* 2003;19:172–176. [PubMed: 12683966]
22. Nechaev S, Severinov K. Bacteriophage-induced modifications of host RNA polymerase. *Annu. Rev. Microbiol* 2003;57:301–322. [PubMed: 14527281]
23. Slesarev AI, Mezhevaya KV, Makarova KS, Polushin NN, Shcherbinina OV, Shakhova VV, Belova GI, Aravind L, Natale DA, Rogozin IB, Tatusov RL, Wolf YI, Stetter KO, Malykh AG, Koonin EV, Kozyavkin SA. The complete genome of hyperthermophile *Methanopyrus kandleri* AV19 and monophyly of archaeal methanogens. *Proc. Natl. Acad. Sci. U.S.A* 2002;99:4644–4649. [PubMed: 11930014]
24. Polushin N, Malykh A, Morocho AM, Slesarev A, Kozyavkin S. High-throughput production of optimized primers (fimers) for whole-genome direct sequencing. *Methods Mol. Biol* 2005;288:291–304. [PubMed: 15333911]
25. Ewing B, Hillier L, Wendl MC, Green P. Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Res* 1998;8:175–815. [PubMed: 9521921]
26. Besemer J, Borodovsky M. Heuristic approach to deriving models for gene finding. *Nucleic Acids Res* 1999;27:3911–392. [PubMed: 10481031]
27. Altschul SF, Madden TI, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res* 1997;25:3389–3402. [PubMed: 9254694]
28. Felsenstein, J. PHYLIP (Phylogeny Inference Package) version 3.6. Department of Genome Sciences, University of Washington; Seattle: 2005. Distributed by the author
29. Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 1997;25:955–964. [PubMed: 9023104]
30. Walker DR, Koonin EV. SEALS: a system for easy analysis of lots of sequences. *Proc. Int. Conf. Intell. Syst. Mol. Biol* 1997;5:333–339. [PubMed: 9322058]

31. Tomomori-Sato C, Sato S, Parmely TJ, Banks CA, Sorokina I, Florens L, Zybaylov B, Washburn MP, Brower CS, Conaway RC, Conaway JW. A mammalian mediator subunit that shares properties with *Saccharomyces cerevisiae* mediator subunit Cse2. *J. Biol. Chem* 2004;279:5846–5851. [PubMed: 14638676]
32. Eng J, McCormack AL, Yates JR 3rd. An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. *J. Amer. Mass Spectrom* 1994;5:976–989.
33. Tabb DL, McDonald WH, Yates JR 3rd. DTASelect and Contrast: Tools for assembling and comparing protein identifications from shotgun proteomics. *J. Proteome Res* 2002;1:21–26. [PubMed: 12643522]
34. Liu H, Sadygov RG, Yates JR 3rd. A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal. Chem* 2004;76:4193–4201. [PubMed: 15253663]
35. Powell DW, Weaver CM, Jennings JL, McAfee KJ, He Y, Weil PA, Link AJ. Cluster analysis of mass spectrometry data reveals a novel component of SAGA. *Mol. Cell. Biol* 2004;24:7249–7259. [PubMed: 15282323]
36. Pavlov AR, Belova GI, Kozyavkin SA, Slesarev AI. Helix-hairpin-helix motifs confer salt resistance and processivity on chimeric DNA polymerases. *Proc. Natl. Acad. Sci. U. S. A* 2002;99:13510–13515. [PubMed: 12368475]




Figure 1. The ϕ YS40 genome. Bacteriophage ϕ YS40 genome is schematically presented with predicted ORFs indicated by arrows. Arrow direction indicates the direction of transcription. Several ORFs with clear functional predictions for their products are color-coded (see also Table 1 for more details).




Figure 2.
 Sequence alignment of the TP proteins.
 Multiple alignment of terminal proteins (TP) from ϕ 29 family phages and phage ϕ YS40 gp65. The stretch of * indicates a region of a predicted amphipathic alpha-helix in TP. Distances, in amino acid residues, from the ends of each sequence and between blocks, are shown in parentheses. A white font in blue indicates the residue identical in all sequences compared, yellow shading indicates the conservation of hydrophobic residues, grey shading indicates the conservation of polar and charged residues. The white font in red indicates the Ser₂₃₂ that is essential for TP priming activity.




Figure 3.
SDS-PAGE analysis of the ϕ YS40 virion proteins.
The SDS gel shows the protein composition of purified ϕ YS40 virions. The two major bands identified by mass-spectrometry are indicated.




Figure 4.
MudPIT analysis of ϕ YS40 lysates.

- A. Normalized Spectral Abundance Factor (NSAF) values measured for ϕ YS40 proteins detected in at least two of the three runs.
- B. NSAFs for contaminating *T. thermophilus* proteins detected in at least two of the three runs.
- C. All 33 ϕ YS40 genes for which products were detected are plotted along the genome as a function of the measured NSAF values (when proteins were identified in several runs, maximal NSAF values are reported). The arrows under the x axis represent the position of the leftward and rightward predicted transcription clusters.

Table 1

Gene products of phage ϕ YS40 and their predicted molecular functions.

ORF name	ORF strand/position ^a	ORF length (amino acids)	The best database match with validated similarity	Taxonomic origin of the best match	Function and other properties ^b
1	-/(7..1938)	643	34419532	<i>Vibrio</i> phage KVP40	distal tail fiber protein
2	-/(1941..4586)	881	48696430	<i>Staphylococcus</i> phage K	unknown
3	-/(4573..7410)	945	90591438	<i>Flavobacterium johnsoniae</i> UW101	portal protein
4	-/(7412..8068)	218	19924248	<i>Methanocaldococcus jannaschii</i>	TM, unknown
5	-/(8096..8530)	144	9631083	<i>Lymnaea dispar</i>	S-adenosylmethionine decarboxylase (adoMetDC)
6	-/(8564..8788)	74	33357605	nucleopolyhedrovirus	unknown
7	-/(8801..9412)	203	33860394	<i>Thermotoga maritima</i>	dUTPase
8	-/(9399..9941)	180	23029929	<i>Burkholderia cepacia</i> phage Bcep22	flavin-dependent thymidylate synthase
9	-/(9955..10782)	275	46200225	<i>Microbulbifer degradans</i>	unknown
10	-/(10816..11331)	171	22978288	<i>Thermus thermophilus</i> HB27	gp18, unknown function
11	-/(11310..11783)	157	23029305	<i>Ralstonia metallidurans</i>	RecA/RadA recombinase
12	-/(11776..12795)	339	23110678	<i>Microbulbifer degradans</i>	Rad52 strand-exchange protein
13	-/(12792..13367)	191	27262500	<i>Helicobacterium</i> sp.	DNA helicase DnaB
14	-/(13413..14756)	447	37526389	<i>Photobacterium luminescens</i>	unknown
15	-/(14743..15036)	97	15595102	<i>Borrelia burgdorferi</i>	IMP dehydrogenase / GMP reductase
16	15124..15453	109	9964625	<i>Streptococcus pneumoniae</i>	DNA binding HTH-domain protein, transcription regulator
17	15467..16576	369	52216967	<i>Microbulbifer degradans</i>	Major structural protein
18	16640..17050	136	29348669	<i>Novosphingobium aromattictvorans</i>	unknown
19	-/(17108..18343)	411	21229604	<i>Xanthomonas campestris</i>	unknown
20	-/(18400..18837)	145	23104360	<i>Acetobacter vinelandii</i>	unknown
21	-/(18834..19214)	126	20808702	<i>Thermoanaerobacter tengcongensis</i>	unknown
22	-/(19187..19960)	257	68		unknown
23	-/(19944..21620)	558	150		unknown
24	-/(21669..22277)	202	109		unknown
25	-/(22302..23015)	237	68		unknown
26	-/(22975..23901)	308	150		unknown
27	-/(23898..25247)	449	147		unknown
28	25396..26796	466	68		unknown
29	26822..27331	169	150		unknown
30	-/(27328..29085)	585	109		unknown
31	-/(29090..29803)	237	71		unknown
32	-/(29818..30291)	157	47		unknown
33	30387..32498	703	68		unknown
34	-/(32491..32781)	96	147		unknown
35	-/(32768..33034)	88	68		unknown
36	-/(33031..33309)	92	150		unknown
37	33381..33746	37	109		unknown
38	33730..34158	121	71		unknown
39	34188..34616	142	47		unknown
40	34631..35155	174	68		unknown
41	35201..37594	797	150		unknown
42	37607..38206	199	47		unknown
43	38240..38446	68	68		unknown
44	38459..38911	150	150		unknown
45	38898..39227	109	109		unknown
46	39224..39439	71	71		unknown
47	39441..39884	147	147		4 TMs, unknown

ORF name	ORF strand/position ^d	ORF length (amino acids)	The best database match with validated similarity	Taxonomic origin of the best match	Function and other properties ^b
48	39877..40185	102			unknown
49	40201..40548	115			unknown
50	40558..41013	151			unknown
51	41010..42482	490			unknown
52	42536..43408	290		<i>Mesorhizobium</i> sp. BNC1	UDP-3-O-[3-hydroxy-myristoyl] glucosamine N-acyltransferase
53	43411..43938	175			unknown
54	43940..44425	161			unknown
55	--/(44426..45127)	233			unknown
56	45187..46209	340		<i>Geobacter metallireducens</i>	conserved bacterial protein, unknown
57	46199..47536	466		<i>Symbiobacterium thermophilum</i>	spore cortex synthesis protein SpoVR
58	47564..49414	616		<i>Bdellovibrio bacteriovorus</i>	unknown
59	49453..51312	619		<i>Desulfotobacterium hafniense</i>	putative serine protein kinase
60	51410..51997	195		<i>Streptomyces</i> phage phi-BT1	putative dNMP Kinase
61	52035..52484	149			terminase large subunit
62	--/(52477..54345)	622		<i>Methanocaldococcus jannaschii</i>	unknown
63	--/(54320..55108)	262			unknown
64	--/(55105..55485)	126		<i>Bacillus</i> phage B103	terminal protein
65	--/(55466..56017)	183			unknown
66	56049..56315	88			unknown
67	--/(56362..57102)	246			unknown
68	--/(57104..57754)	216			tail sheath protein
69	--/(57775..59721)	648		<i>Chloroflexus aurantiacus</i>	unknown
70	--/(59782..60492)	236			unknown
71	--/(60495..61157)	220		<i>Staphylococcus</i> phage K	Zn ribbon, similar to archaeal transcription factor IIB
72	--/(61167..61682)	171			unknown
73	--/(61756..63168)	470			Major structural protein
74	--/(63204..64838)	544		<i>Staphylococcus</i> phage K	unknown, 3 coiled coil regions
75	--/(64838..65098)	86			unknown
76	--/(65085..69662)	1525			unknown
77	--/(69684..74918)	1744			unknown, 3 coiled coil regions
78	--/(74931..75296)	121			unknown
79	--/(75309..79883)	1524			helicase (DEAD motif replaced by DDAE)
80	--/(79880..80743)	287		<i>Aspergillus nidulans</i>	unknown
81	--/(80788..82740)	650			unknown
82	--/(82771..84609)	612			unknown
83	--/(84867..85094)	75			unknown
84	--/(85328..85558)	76			unknown
85	--/(85767..85919)	50			unknown
86	--/(86022..86273)	83			unknown
87	--/(86382..86618)	78			unknown
88	--/(86909..87154)	81			unknown
89	--/(87505..87990)	161		<i>Deinococcus radiodurans</i>	2 TMs, unknown
90	--/(88074..88529)	151			unknown
91	--/(88642..89250)	202			unknown
92	--/(89349..89783)	144			unknown
93	--/(89796..90221)	141			unknown
94	--/(90481..90927)	148			unknown
95	--/(91036..91212)	58			unknown
96	91231..91359	43			3 TMs, unknown
97	--/(91417..91824)	135			unknown
98	--/(91835..92380)	181			unknown
99	--/(92503..93045)	180			unknown
100	--/(93045..93635)	196			unknown
101	--/(93619..94131)	170			unknown
102	--/(94337..94873)	178			unknown

ORF name	ORF strand/position ^d	ORF length (amino acids)	The best database match with validated similarity	Taxonomic origin of the best match	Function and other properties ^b
103	--/(94885..95373)	162			unknown
104	--/(95510..96025)	171			unknown
105	--/(96096..96626)	176			unknown
106	--/(96833..97354)	173			unknown
107	--/(97575..99263)	562			unknown
108	--/(99280..100323)	347	15643692	<i>Thermotoga maritima</i>	ATPase
109	--/(100462..101157)	231			unknown
110	--/(101227..101973)	248			unknown
111	--/(102138..102530)	130			unknown
112	--/(102531..103076)	181			unknown
113	--/(103077..103616)	179			unknown
114	--/(103616..104107)	163			unknown
115	--/(104451..104693)	80	11992695	<i>Escherichia coli</i>	glycosyltransferase
116	--/(104803..105279)	158			unknown
117	--/(105422..105979)	185			unknown
118	--/(105969..106520)	183			unknown
119	--/(106510..107076)	188			unknown
120	--/(107090..107539)	149			unknown
121	--/(107552..108046)	164			unknown
122	--/(108141..108644)	167			unknown
123	--/(108772..109290)	172			unknown
124	--/(109328..109819)	163			unknown
125	--/(109998..110513)	171	18462664	<i>Shigella flexneri</i>	unknown
126	--/(110561..111145)	194			unknown
127	--/(111157..111654)	165			unknown
128	--/(111663..112133)	156			unknown
129	--/(112165..112677)	170			unknown
130	--/(112689..113195)	168			unknown
131	--/(113202..113630)	142			unknown
132	113852..114388	178			coiled coil, unknown
133	--/(114385..115032)	215			unknown
134	--/(115155..115724)	189			unknown
135	--/(115727..116299)	190			unknown
136	--/(116271..116693)	140			unknown
137	116815..117474	219			unknown
138	--/(117442..118005)	187			unknown
139	--/(118395..119999)	534	19552983	<i>Corynebacterium glutamicum</i>	unknown
140	--/(120226..120777)	183			unknown
141	120821..120994	58			unknown
142	--/(120953..123997)	1014			coiled coil, unknown
143	--/(124012..124536)	174			unknown
144	--/(124553..125593)	346	10956653	<i>Rhodococcus equi</i>	M27/M37 peptidase
145	--/(125598..126548)	316			unknown
146	--/(126553..126813)	86			3 TMs, unknown
147	126870..127055	61			unknown
148	127065..127460	131			unknown
	127471..127959	162			unknown
149	127979..129967	662	34762157	<i>Fusobacterium nucleatum</i>	putative baseplate assembly protein
150	129964..131859	631	903862	<i>Escherichia coli</i> phage K3	unknown
151	131870..134260	796			wac fibrin neck whisker
152	134253..136364	703			unknown
153	136388..137287	299			unknown
154	137294..137644	116			unknown
155	137634..138497	287			unknown
156	138469..139269	266			unknown

ORF name	ORF strand/position ^a	ORF length (amino acids)	The best database match with validated similarity	Taxonomic origin of the best match	Function and other properties ^b
158	139253..143296	1347			unknown
159	143322..143846	174			unknown
160	144155..144367	70			unknown
161	-/(144357..145424)	355	15674141	<i>Lactococcus lactis</i>	Radical SAM superfamily enzyme
162	-/(145421..146374)	317			unknown
163	-/(146390..147022)	210			unknown
164	147094..147639	181			unknown
165	147677..148306	209			unknown
166	148300..148689	129			unknown
167	148736..150229	497			unknown
168	150256..151341	361			unknown
169	151338..151907	189			unknown
170	151894..152157	87			unknown

^a φ YS40 virion proteins detected by MudPIT are indicated in red.

^a position of the ORFs in the phage YS40 genome; “-” indicates a leftwards transcription orientation.

^b presence of transmembrane domains (TM) and coiled coil regions are indicated.