

Stress Response in *Lactococcus lactis*: Cloning, Expression Analysis, and Mutation of the Lactococcal Superoxide Dismutase Gene

JAN WILLEM SANDERS, KEES J. LEENHOUTS,† ALFRED J. HAANDRIKMAN,‡
GERARD VENEMA,* AND JAN KOK

Department of Genetics, Biomolecular Sciences and Biotechnology Institute,
University of Groningen, 9751 NN Haren, The Netherlands

Received 21 March 1995/Accepted 5 July 1995

In an analysis of the stress response of *Lactococcus lactis*, three proteins that were induced under low pH culture conditions were detected. One of these was identified as the lactococcal superoxide dismutase (SodA) by N-terminal amino acid sequence analysis. The gene encoding this protein, designated *sodA*, was cloned by the complementation of a *sodA sodB Escherichia coli* strain. The deduced amino acid sequence of *L. lactis* SodA showed the highest degree of similarity to the manganese-containing Sod (MnSod) of *Bacillus stearothermophilus*. A promoter upstream of the *sodA* gene was identified by primer extension analysis, and an inverted repeat surrounding the –35 hexanucleotide of this promoter is possibly involved in the regulation of the expression of *sodA*. The expression of *sodA* was analyzed by transcriptional fusions with a promoterless *lacZ* gene. The induction of β -galactosidase activity occurred in aerated cultures. Deletion experiments revealed that a DNA fragment of more than 130 bp surrounding the promoter was needed for the induction of *lacZ* expression by aeration. The growth rate of an insertion mutant of *sodA* did not differ from that of the wild type in standing cultures but was decreased in aerated cultures.

Lactic acid bacteria (LAB) are widely used in fermented food production. Normal growth of LAB does not require strictly anaerobic environments. A number of stirring steps in the various production processes in which LAB are involved provide ample contact with oxygen, without obvious deleterious effects to these organisms. A number of LAB can use molecular oxygen or hydrogen peroxide to regenerate NAD⁺, by the action of NADH oxidase and NADH peroxidase (14). With the stepwise reduction of O₂ to H₂O the toxic intermediates O₂^{•-} and H₂O₂ are generated. Hydrogen peroxide was found to inhibit growth of lactococci (1), and exposure of lactococci to a sublethal dose of H₂O₂ induced an oxidative stress response (14), characterized by an increased survival after exposure to a lethal level of H₂O₂ compared with cells that were not pretreated.

Most LAB can deal with oxygen radicals by either a superoxide dismutase (Sod) or a high internal Mn²⁺ concentration (2). Sod dismutates oxygen radicals by catalyzing the reaction $2\text{O}_2^{\bullet-} + 2\text{H}^+ \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$ (20).

Sods are found in a wide variety of prokaryotic and eukaryotic organisms, and in a number of instances the genes encoding these enzymes have been cloned and characterized (5, 9, 12, 33). In prokaryotes three types of Sods can be distinguished depending on the metal cofactor contained in these enzymes (Cu-Zn, Fe, or Mn) (36, 40). A single organism can have two *sod* genes; the corresponding enzymes differ in their metal cofactor and in their expression pattern in response to oxygen.

The regulation of Sod expression in *Escherichia coli* has been studied extensively (13, 18, 25). Inactivation of *sod* genes can be growth inhibiting (in *E. coli* [10]) or even lethal (in *Legionella pneumophila* [40]).

All streptococci tested (including *Lactococcus lactis* subsp. *lactis*, formerly *Streptococcus lactis*) appear to carry a manganese-containing Sod (MnSod) (52). The MnSod of *L. lactis* is active under anaerobic conditions. Higher enzyme activity has been observed with increasing O₂ concentrations in the medium (24, 43). Although most organisms use catalase for the breakdown of H₂O₂, streptococci lack this activity; instead, they have NADH-peroxidase activity to decompose this compound (1).

One of the typical properties of LAB is their ability to produce large amounts of lactic acid, thereby causing a rapid acidification of their environment. Several microorganisms are known to adapt to medium with a low pH (19, 35). In the present paper we studied the effect of a pH downshift on the expression of proteins in *L. lactis*. An acid-induced protein was isolated, and by N-terminal amino acid sequence analysis as well as by activity assays, this protein appeared to be a Sod. The lactococcal *sodA* gene was cloned and sequenced, and its expression as well as the growth characteristics of a *sodA* mutant were studied.

MATERIALS AND METHODS

Bacterial strains, plasmids, and growth conditions. The bacterial strains and plasmids used in this study are listed in Table 1. *L. lactis* was grown at 30°C in M17 medium, with 0.5% glucose; solidified M17 medium contained 1.5% agar. Erythromycin (Em) and chloramphenicol (Cm) were used at final concentrations of 5 $\mu\text{g/ml}$. MRS broth (15) containing 0.5% glucose with or without 2% β -glycerophosphate was used for acid-induction experiments. *E. coli* was grown in TY broth (39) at 37°C with vigorous agitation or on TY medium supplemented with 1.5% agar. Ampicillin (Ap) and paraquat were used at 100 $\mu\text{g/ml}$ and 10 $\mu\text{g/ml}$, respectively.

Preparation of cell extracts, PAGE, and Sod activity detection. Cultures of *L. lactis* were harvested by centrifugation, and the pellets were resuspended in 10 mM Tris–1 mM EDTA (pH 7.4) and disrupted according to the method of Van de Guchte et al. (49). The cell extract was used for analysis by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) according to

* Corresponding author. Mailing address: Department of Genetics, Biomolecular Sciences and Biotechnology Institute, University of Groningen, Kerklaan 30, 9751 NN Haren, The Netherlands. Phone: 31-50-63-2092. Fax: 31-50-63-2348. Electronic mail address: Sandersj@BIOL.RUG.NL.

† Present address: International Centre for Genetic Engineering and Biotechnology, Department of Molecular Pathology, I-34012 Trieste, Italy.

‡ Present address: Hercules, European Research Centre, 3771 ME Barneveld, The Netherlands.

TABLE 1. Bacterial strains and plasmids

Strain or plasmid	Relevant characteristic(s)	Reference or source
Strains		
<i>L. lactis</i>		
MG1363	Plasmid-free derivative of NCDO712	21
LL108	<i>repA</i> ⁺ derivative of MG1363, Cm ^r	28a
MGSOD4	<i>sodA::lacZ</i> Em ^r	This study
MGSOD7	Δ <i>sodA</i> derivative of MG1363, Em ^r	This study
MGSOD8	Sod ⁺ derivative of MG1363, Em ^r	This study
<i>E. coli</i>		
NM522	<i>supE thi</i> Δ (<i>lac-proAB</i>) Δ <i>hsd5</i> ($r^- m^-$) [F' <i>proAB lacI^qZ</i> Δ M15]	23
OX326A	Δ <i>sodA</i> Δ <i>sodB</i>	45
EC101	<i>repA</i> ⁺ derivative of JM101, Km ^r	28
EC1000	<i>repA</i> ⁺ derivative of MC1000, Km ^r	27a
Plasmids		
pUC19	<i>lacZ'</i> Ap ^r	50
pORI13	Promoterless <i>lacZ</i> , Em ^r , Ori ⁺ of pWV01, Rep ⁻	41a
pORI19	<i>lacZ'</i> Em ^r , Ori ⁺ of pWV01, Rep ⁻	28
pVE6007	pWV01 derivative encoding a temperature-sensitive Rep protein, Cm ^r	30
pSOD1	Ap ^r , pUC19 with 7.6-kb <i>Sau3A</i> fragment	This study
pSOD2	Ap ^r , pUC19 with 0.8-kb + 0.9-kb <i>PvuII</i> fragment	This study
pSOD3	<i>sodA::lacZ</i> Em ^r	This study
pSOD4	<i>sodA::lacZ</i> Em ^r	This study
pSOD5	<i>sodA::lacZ</i> Em ^r	This study
pSOD6	<i>sodA::lacZ</i> Em ^r	This study
pSOD7	Internal <i>NdeI-EcoRI</i> fragment of <i>sodA</i> in pORI19	This study

the protocol of Laemmli (27). Nondenaturing PAGE was carried out similarly, except that SDS and mercaptoethanol were omitted. Prior to loading, samples were incubated at 37°C for 15 min. Polyacrylamide (PA) gels were stained with Coomassie brilliant blue for total protein detection. Sod activity in nondenaturing gels was determined according to the protocol of Beauchamp and Fridovich (4).

N-terminal amino acid analysis. Cell extracts of acid-shocked cells were separated on SDS-16% PA-piperazine diacrylamide gels (3). Proteins were transferred to polyvinylidene difluoride membrane (Millipore Corporation, Bedford, Mass.) which was subsequently stained with Coomassie brilliant blue, according to protocols of Eurosequence b.v. (Groningen, The Netherlands). After destaining with 50% methanol, the desired protein band was cut from the gel and subjected to automated Edman degradation with an Applied Biosystems 477A sequencer (Applied Biosystems, Inc., Foster City, Calif.). Phenylthiohydantoin amino acids were identified with an on-line high-pressure liquid chromatographer (model 120A; Applied Biosystems, Inc.).

Molecular cloning techniques. Molecular cloning techniques were performed essentially as described by Sambrook et al. (41). DNA was introduced in *E. coli* by electrotransformation (51). A genome bank of *L. lactis* in pUC19 (8) was used to clone the *sod* gene. DNA sequencing was done on double-stranded plasmid DNA by the dideoxy chain-termination method (42) and the T7 sequencing kit (Pharmacia LKB Biotechnology AB, Uppsala, Sweden) according to the manufacturer's instructions. Oligonucleotides were synthesized with an Applied Biosystems 381A DNA synthesizer.

DNA sequences were analyzed with the PC/Gene sequence analysis program (IntelliGenetics Inc., Geneva, Switzerland). Protein homology searches in the PIR bank (release 40) were carried out with the FASTA program (38). Protein sequence alignments were carried out with the PALIGN program of PC/Gene by using the structure genetic matrix or with the CLUSTAL program, both with standard settings.

Primer extension analysis. RNA was isolated from an exponentially growing standing *L. lactis* culture at an optical density at 600 nm of 0.5 as previously described (48). A synthetic oligonucleotide complementary to the mRNA from positions 90 to 127 (corresponding to coordinates 565 to 601 in Fig. 3) was used for primer extension. Ten picomoles of primer was annealed to 10 μ g of RNA, and then cDNA synthesis was performed as previously described (41). The product was analyzed on a sequencing gel next to a sequence reaction with the same primer, providing a size marker.

β -Galactosidase assays. Cell extracts were prepared from exponentially growing cultures, either in completely filled standing bottles or by shaking in 100-ml bottles with 15 ml of medium. β -Galactosidase activity was determined essen-

tially as described by Miller (32). Protein concentrations of the cell extracts were determined by the method of Bradford (6) with bovine serum albumin as a standard.

Construction of a Sod⁻ mutant. An internal *NdeI-EcoRI* fragment of the *sodA* gene was cloned in the integration insertion vector pORI19, using a Rep⁺ *E. coli* helper strain (EC101 [28]). This plasmid, pSOD7, was used to disrupt the *sodA* gene in *L. lactis* MG1363, as described before (22). The proper chromosomal location of the integrated plasmid was confirmed by Southern hybridization (not shown), and the strain was named MGSOD7.

Nucleotide sequence accession number. The sequence presented in Fig. 3 has been assigned the GenBank nucleotide sequence accession number U17388.

RESULTS

A 24-kDa acid-induced protein of *L. lactis* is a Sod. *L. lactis* MG1363 was subjected to acid stress by growth in MRS medium with or without 2% β -glycerophosphate as the buffering agent. After overnight incubation the medium had acquired a pH of 6.2 in the buffered and a pH of 4.8 in the unbuffered medium. Cell extracts were analyzed by SDS-PAGE (Fig. 1A). At least three proteins with estimated molecular sizes of 72, 64, and 24 kDa were expressed at a higher level in a medium with a low pH, although the effect, probably due to slight variations in the growth conditions, was not invariably observed. Proteins of 72 and 64 kDa were also expressed at a higher level in exponentially growing cells subjected to a heat shock of 42°C for 1 h (Fig. 1). Therefore, and on the basis of their sizes (16, 26), we speculate that these proteins represent the lactococcal DnaK and GroEL proteins, respectively. The 24-kDa protein was partially purified by SDS-PAGE, and the N-terminal amino acid sequence was determined to be (Thr or Ala)-Phe-Thr-Leu-Pro-Glu-Leu-Pro-Tyr-Ala-Pro-Asn-Ala-Leu-Glu-Pro-Phe. A computer homology search revealed that this amino acid sequence is similar to the N-terminal amino acid sequences of a number of Sods (see Fig. 5). To determine whether Sod was induced by low pH conditions, cell extracts were separated on a nondenaturing PA gel and assayed for Sod activity. A higher level of Sod activity was indeed detected in cells grown at the lower pH (Fig. 1B).

Cloning and sequencing of the *sodA* gene. To clone the *L. lactis* *sod* gene, use was made of the *E. coli* strain from which




FIG. 1. (A) SDS-12.5% PAGE analysis of cell extracts of acid-stressed *L. lactis* cells. Lanes: 1, cells grown overnight in buffered MRS; 2, cells grown overnight in unbuffered MRS; 3, cells taken from exponential growth phase and incubated 1 h at 42°C. (B) Nondenaturing (12.5%) PA gel assayed for Sod activity. Culture conditions for lanes 1 and 2 were the same as those for lanes 1 and 2 in panel A. The arrows indicate three acid-induced proteins. The pH values of the culture media after growth and the growth temperatures are indicated below the lanes. Molecular masses (in kilodaltons) are indicated at the left. Equal amounts of protein (30 μ g) were applied per lane.




FIG. 2. Nondenaturing (12.5%) PAGE of cell extracts; the gel was stained for Sod activity. Lanes: 1, *L. lactis* MG1363; 2, *E. coli* NMS22; 3, *E. coli* OX326A; 4, *E. coli* OX326A(pSOD2). Thirty micrograms of protein was applied per lane.

the *sodA* and *sodB* genes were deleted (OX326A). This strain lacks all Sod activity and is sensitive to 10 µg of paraquat per ml, in contrast to *E. coli* strains carrying one or both *sod* genes (45).

A genome bank of *L. lactis* MG1363 in pUC19 was used to transform *E. coli* OX326A, and transformants were selected for resistance to ampicillin and paraquat. Eight paraquat-resistant colonies were assayed for Sod activity on a nondenaturing PA gel. Six of these showed a band of Sod activity with a mobility similar to that of *L. lactis* Sod. The mobility pattern of Sod activity of one of these clones is shown in Fig. 2, lane 4. The Sod activity was obviously derived from lactococcal DNA, as it was clearly different from the two *E. coli* Sod activities (Fig. 2, lane 2) which are lacking in the cloning host (Fig. 2, lane 3).

Plasmid DNA of the Sod-expressing clones was isolated. Restriction enzyme analysis of three clones showed that all contained a 7.6-kb DNA fragment (data not shown). A partial *PvuII* digest of one of these clones (pSOD1) was ligated into the dephosphorylated *SmaI* site of pUC19. The ligation mixture was used to transform OX326A to paraquat resistance. Plasmid DNA of paraquat-resistant transformants was isolated. The smallest plasmid expressing active Sod, pSOD2, consisted of two adjacent *PvuII* fragments of 0.8 and 0.9 kb. The 0.8-kb fragment hybridized with a degenerate oligonucleotide probe designed from the N-terminal amino acid sequence of the isolated protein (data not shown).

Part of the nucleotide sequence of both strands of the insert in pSOD2 is presented in Fig. 3. The cloned fragment contained an open reading frame (ORF) of 618 bp that could encode a protein of 206 amino acids with a predicted molecular weight of 23,254 and a calculated pI of 4.8. The translation product of the first 18 codons of this ORF (except for Met) was identical to the N terminus of the 24-kDa acid-induced protein, indicating that the *L. lactis sod* gene had indeed been cloned. Accordingly, the ORF was designated *sodA*. The gene is preceded by a putative ribosome binding site with complementarity to the 3' end of the ribosomal 16S rRNA of *L. lactis* (11) with a ΔG° of -14 kcal/mol (-58.3 kJ/mol) (47). Upstream of the ribosome binding site a promoter-like structure was present, consisting of the -35 hexanucleotide TTGGCA, a spacing of 17 bp and the -10 hexanucleotide TATAAT. The -35 hexanucleotide is surrounded by an 8-bp inverted repeat ($\Delta G[25^\circ\text{C}] = -4.6$ kcal/mol [-19.2 kJ/mol]). To examine whether the putative promoter was active in vivo, primer extension was carried out. The results are presented in Fig. 4 and show that transcription started at an adenine 7 bp downstream of the -10 hexanucleotide. Downstream of *sodA*, a 13-bp in-

verted repeat ($\Delta G[25^\circ\text{C}] = -14.4$ kcal/mol [-60.2 kJ/mol]) followed by a stretch of T's could function as a transcription terminator.

One hundred fifty-three base pairs upstream of the *sodA* gene the stop codon of a second ORF is present. This ORF starts beyond the 5' end of the cloned fragment in pSOD2. The inverted repeat surrounding the -35 sequence of the *sodA* promoter may function as a terminator for this second ORF, although its level of free energy is low. The PIR protein database did not contain proteins homologous to the amino acid sequence deduced from the incomplete ORF.

L. lactis SodA is a MnSod. A homology comparison with other Sod proteins revealed that lactococcal SodA is similar to Sods of various organisms of both prokaryotic and eukaryotic origin (Fig. 5). The size of 206 amino acid residues of the lactococcal SodA is in agreement with the sizes of other Sods,




FIG. 3. Nucleotide sequence and deduced amino acid sequence of the *sodA* gene of *L. lactis* MG1363 and its surrounding regions. Facing arrows, inverted repeats; rbs, ribosome binding site (with the bases in lowercase and boldface type); -10, -35, putative promoter sequence (with the bases underlined and in boldface type); vertical arrow, transcription start site. Stop codons are given in boldface type and indicated with asterisks. The doubly underlined amino acid sequence is identical to that determined from the isolated 24-kDa protein. A number of relevant restriction enzyme sites are indicated.



FIG. 7. Nondenaturing PAGE of cell extracts of MG1363 (lane 1) and MGSOD7 (lane 2). The gel was stained for Sod activity; 30 μ g of protein was applied per lane.

Also in this situation, a twofold induction by aeration was observed.

To identify the DNA region responsible for oxygen-induced *sodA* expression, pSOD5 was constructed by cloning in pORI13 a 502-bp *PvuII-EcoRI* fragment (coordinates 1 to 502 in Fig. 3) immediately upstream of the *E. coli lacZ* gene devoid of its own promoter. This plasmid contains the *sodA* promoter and part of the upstream ORF but no *sodA* coding sequences. The level of *lacZ* expression from this plasmid replicating in strain LL108 was lower than that from pSOD4 in LL108 but was still inducible. However, with another subclone, pSOD6, carrying a 130-bp *BglII-EcoRI* fragment in front of *lacZ* (coordinates 370 to 502, Fig. 3), β -galactosidase expression was similar in standing and in aerated cultures and comparable to that from LL108(pSOD5) under aerated conditions. In pSOD5 and pSOD6 the same *lacZ* fusion point was used. This indicates that the 370-bp *PvuII-BglII* fragment (coordinates 1 to 370, Fig. 3) is involved in the regulation of *sodA* expression.

Growth of a Sod-negative *L. lactis* mutant is impeded in aerobic cultures. To examine possible phenotypic effects of SodA deficiency, the *sodA* gene was disrupted by the insertion of the integration vector pSOD7. Figure 7 shows that no SodA activity could be detected in the disruption mutant. The growth rate of MGSOD7 in aerated medium was lower than that of a SodA⁺ Em^r control strain, whereas the growth rates of both strains were comparable in standing cultures (Fig. 8). The lag phase of MGSOD7 was consistently somewhat longer than that of the SodA⁺ strain. Also the SodA-negative mutant grew only slowly on solid medium in the presence of O₂. Thus, although not essential, *L. lactis* SodA is important for optimal growth of the organism under aerobic conditions.

DISCUSSION

In the framework of a general interest in stress-induced protein expression in *L. lactis* we have examined the induction of proteins synthesized at a low pH and observed in overnight cultures the induction of three proteins with molecular sizes of 72, 64, and 24 kDa. The induction pattern resembles that of *Salmonella typhimurium* upon infection of macrophages, which have a low interior pH. In this case a 58-kDa protein was identified as GroEL, and a 68-kDa protein was identified as DnaK. In addition, an unknown 27-kDa protein was induced (7). The *L. lactis* DnaK and GroEL proteins are 65 and 57 kDa, respectively (16, 26). The 72- and 64-kDa lactococcal proteins identified here may, therefore, represent the products of *dnaK* and *groEL* in *L. lactis*.

The 24-kDa acid-induced protein was identified as SodA on the basis of its N-terminal amino acid sequence identity with the deduced amino acid sequence of the cloned *sodA* gene. The deduced amino acid sequence of SodA is highly similar to those of SodA from evolutionary closely and distantly related organisms. The observed homology of lactococcal SodA with the group of MnSods rather than with FeSods is in agreement with the previous observation that all streptococci, and *L. lactis*, possess MnSods (52).

The role of SodA under acid conditions is not clear. The protein does not play an obvious role in the growth of *L. lactis* in standing cultures, since growth of the SodA-negative mutant under these conditions was similar to that of the parent strain. In contrast, a negative effect of the *sodA* mutation on growth was observed in aerated cultures and on plates. Apparently, oxygen radicals inhibit growth under these conditions. Similar effects have been observed for Sod⁻ derivatives of *E. coli* (10) and *Streptococcus mutans* (33). A Sod-free *E. coli* strain grew slowly in rich medium and was unable to grow in minimal medium under aerobic conditions. An intracellular Sod appeared to be essential for *L. pneumophila*, a bacterium that needs oxygen for growth (40). The inactivation of Sod of the obligate anaerobe *Porphyromonas gingivalis* caused a rapid loss of viability upon exposure to oxygen (34). It was concluded that the latter organism lacks efficient alternative protection or repair systems to overcome oxidative damage, whereas *E. coli* and *S. mutans* do have such systems. The ability of SodA-deficient *L. lactis* to grow under aerated conditions indicates that also in this organism alternative protection systems operate. A candidate protection mechanism is a high intracellular glutathione concentration, both in *E. coli* and in *L. lactis* (17), since glutathione can also detoxify free radicals (31). The properties of Sod-deficient strains described thus far suggest that Sod is essential for obligate aerobic or anaerobic bacteria to withstand oxygen, whereas it is not essential for facultative (an)aerobic species.

In two independent studies (24, 43), a twofold higher level of Sod activity was observed in *L. lactis* upon aeration. We show here that under similar conditions *sodA* is induced twofold. Together with the results obtained with the *sodA* mutant, these results indicate that SodA is the only or, if not the only, the major Sod in *L. lactis*. The results of the *sodA::lacZ* transcrip-




FIG. 8. Growth of *L. lactis* MG1363 derivatives in liquid GM17 medium, either as aerated (dotted lines) or standing cultures (solid lines). Strains: filled box, MGSOD8 (SodA⁺ Em^r); filled triangle, MGSOD7 (SodA⁻ Em^r). A600, Absorbance at 600 nm.

tional fusions indicate that expression of *sodA* is, most likely, regulated at the transcriptional level. Primer extension revealed that the promoter upstream of *sodA* is active under standard growth conditions (standing culture). *lacZ* expression from pSOD5 indicates that the 500-bp DNA region upstream of *sodA* is required for inducibility under aerobic conditions. Since *lacZ* expression from pSOD6 was independent from aeration, the inducibility of *sodA* expression is probably specified by the 370-bp *PvuII-BglII* fragment. The fact that *lacZ* expression from pSOD6 is similar to that from pSOD5 under aerated conditions suggests that *sodA* expression is repressed under nonaerated conditions by a mechanism acting on DNA sequences 70 to 440 bp upstream of the *sodA* promoter.

Another possible target site for regulation is the putative inverted repeat sequence that encompasses the -35 hexanucleotide of the *sodA* promoter. A stem-loop structure is also present upstream of *E. coli sodA* (46). Studies of the expression of *E. coli sodA* showed that it is regulated by six global regulators: ArcA, Fnr, Fur, SoxRS, CfxB, and IHF (13). These regulators all act on a DNA fragment of less than 120 bp, making the expression of *sodA* dependent on growth conditions and on the presence of O₂, Fe, NO₃⁻, or xenobiotics (18).

Several differences in levels of expression exist between *E. coli* and *L. lactis sodA*. *E. coli* exhibits a basic level of Sod by the constitutive expression of *sodB*. Additional Sod is synthesized from *sodA* in response to environmental conditions. Under nonstress conditions the expression of *sodA* is repressed. In contrast, in *L. lactis*, *sodA* provides the cell with a basic level of SodA or with higher levels when required. It is conceivable that, as with *E. coli*, a number of regulatory systems are involved in the control of *sodA* expression, as Sod activity in *L. lactis* is not only oxygen pressure dependent, but also dependent on the carbon source (24). The cloning of the lactococcal *sodA* and the availability of a SodA-negative mutant form the basis for further studies to obtain more insight in the aerobic life of this fermentative organism.

ACKNOWLEDGMENTS

This work was supported by Unilever Research Laboratorium Vlaardingen, The Netherlands. J.K. was the recipient of a fellowship from the Royal Netherlands Academy of Arts and Sciences (KNAW).

We thank H. M. Steinman for providing the *sodA sodB E. coli* strain, Albert Bolhuis for the primer extension experiment, Rik van Veen for helpful discussions, and Henk Mulder for preparation of the figures.

REFERENCES

- Anders, R. F., D. M. Hogg, and G. R. Jago. 1970. Formation of hydrogen peroxide by group N streptococci and its effect on their growth and metabolism. *Appl. Microbiol.* **19**:608-612.
- Archibald, F. S., and I. Fridovich. 1981. Manganese, superoxide dismutase, and oxygen tolerance in some lactic acid bacteria. *J. Bacteriol.* **146**:928-936.
- Artoni, G., E. Gianazza, M. Zanoni, C. Gelfi, M. C. Tanzi, C. Barozzi, P. Ferruti, and P. G. Righetti. 1984. Fractionation techniques in a hydro-organic environment. II. Acryloyl-morpholine polymers as a matrix for electrophoresis in hydro-organic solvents. *Anal. Biochem.* **137**:420-428.
- Beauchamp, C., and I. Fridovich. 1971. Superoxide dismutase: improved assays and an assay applicable to acrylamide gels. *Anal. Biochem.* **44**:276-287.
- Bowler, C., L. Van Kaer, W. Van Camp, M. Van Montagu, D. Inze, and P. Dhaese. 1990. Characterization of the *Bacillus stearothermophilus* manganese superoxide dismutase gene and its ability to complement copper/zinc superoxide dismutase deficiency in *Saccharomyces cerevisiae*. *J. Bacteriol.* **172**:1539-1546.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein, utilizing the principle of protein-dye binding. *Anal. Biochem.* **12**:248-254.
- Buchmeier, N. A., and F. Heffron. 1990. Induction of *Salmonella* stress proteins upon infection of macrophages. *Science* **248**:730-732.
- Buist, G., J. Kok, K. J. Leenhouts, M. Dabrowska, G. Venema, and A. J. Haandrikman. 1995. Molecular cloning and nucleotide sequence of the gene encoding the major peptidoglycan hydrolase of *Lactococcus lactis*, a muramidase needed for cell separation. *J. Bacteriol.* **177**:1554-1563.
- Carlioz, A., M. L. Ludwig, W. C. Stallings, J. A. Fee, H. M. Steinman, and D. Touati. 1988. Iron superoxide dismutase. Nucleotide sequence of the gene from *Escherichia coli* K12 and correlations with crystal structures. *J. Biol. Chem.* **263**:1555-1562.
- Carlioz, A., and D. Touati. 1986. Isolation of superoxide dismutase mutants in *Escherichia coli*: is superoxide dismutase necessary for aerobic life? *EMBO J.* **5**:623-630.
- Chiaruttini, C., and M. Milet. 1993. Gene organization, primary structure and RNA processing analysis of a ribosomal RNA operon in *Lactococcus lactis*. *J. Mol. Biol.* **230**:57-76.
- Church, S. L. 1990. Manganese superoxide dismutase: nucleotide sequence and deduced amino acid sequence of a cDNA encoding a new human transcript. *Biochim. Biophys. Acta* **1087**:250-252.
- Compan, I., and D. Touati. 1993. Interaction of six global regulators in expression of manganese superoxide dismutase in *Escherichia coli* K-12. *J. Bacteriol.* **175**:1687-1696.
- Condon, S. 1987. Responses of lactic acid bacteria to oxygen. *FEMS Microbiol. Rev.* **46**:269-280.
- De Man, J. C., M. Rogosa, and M. E. Sharpe. 1960. A medium for the cultivation of lactobacilli. *J. Appl. Bacteriol.* **23**:130-135.
- Eaton, T., C. Shearman, and M. Gasson. 1993. Cloning and sequence analysis of the *dnkK* region of *Lactococcus lactis* subsp. *lactis*. *J. Gen. Microbiol.* **139**:3253-3264.
- Fahey, R. C., W. C. Brown, W. B. Adams, and M. B. Worsham. 1978. Occurrence of glutathione in bacteria. *J. Bacteriol.* **133**:1126-1129.
- Fee, J. A. 1991. Regulation of *sod* genes in *Escherichia coli*: relevance to superoxide dismutase function. *Mol. Microbiol.* **5**:2599-2610.
- Foster, J. W., and H. K. Hall. 1990. Adaptive acidification tolerance response of *Salmonella typhimurium*. *J. Bacteriol.* **172**:771-778.
- Fridovich, I. 1975. Superoxide dismutases. *Annu. Rev. Biochem.* **44**:147-159.
- Gasson, M. J. 1983. Plasmid complements of *Streptococcus lactis* NCDO712 and other lactic streptococci after protoplast-induced curing. *J. Bacteriol.* **154**:1-9.
- Godon, J. J., K. Jury, C. A. Shearman, and M. J. Gasson. 1994. The *Lactococcus lactis* sex-factor aggregation gene *cluA*. *Mol. Microbiol.* **12**:655-663.
- Gough, J. A., and N. E. Murray. 1983. Sequence diversity among related genes for recognition of specific targets in DNA molecules. *J. Mol. Biol.* **166**:1-19.
- Hansson, L., and M. H. Häggström. 1984. Effects of growth conditions on the activities of superoxide dismutase and NADH-oxidase/NADH-peroxidase in *Streptococcus lactis*. *Curr. Microbiol.* **10**:345-352.
- Hassan, H. M., and L. W. Schrum. 1994. Roles of manganese and iron in the regulation of the biosynthesis of manganese-superoxide dismutase in *Escherichia coli*. *FEMS Microbiol. Rev.* **14**:315-324.
- Kim, S. G., and C. A. Batt. 1993. Cloning and sequencing of the *Lactococcus lactis* subsp. *groESL* operon. *Gene* **127**:121-126.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (London)* **227**:680-685.
- Law, J. Unpublished data.
- Law, J., A. J. Haandrikman, G. Venema, and K. J. Leenhouts. Development of a general system for generating random chromosomal insertions. Submitted for publication.
- Leenhouts, K. J. Unpublished data.
- Leenhouts, K. J., and G. Venema. 1993. Lactococcal plasmid vectors, p. 65-94. In K. G. Hardy (ed.), *Plasmids, a practical approach*, 2nd ed. Oxford University Press, New York.
- Maguin, E., P. Duwat, T. Hege, D. Ehrlich, and A. Gruss. 1992. New thermosensitive plasmid for gram-positive bacteria. *J. Bacteriol.* **174**:5633-5638.
- Meister, A. 1988. Glutathione metabolism and its selective modification. *J. Biol. Chem.* **263**:17205-17208.
- Miller, J. H. 1972. *Experiments in molecular genetics*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Nakayama, K. 1992. Nucleotide sequence of *Streptococcus mutans* superoxide dismutase gene and isolation of insertion mutants. *J. Bacteriol.* **174**:4928-4934.
- Nakayama, K. 1994. Rapid viability loss on exposure to air in a superoxide dismutase-deficient mutant of *Porphyromonas gingivalis*. *J. Bacteriol.* **176**:1939-1943.
- Olson, E. R. 1993. Influence of pH on bacterial gene expression. *Mol. Microbiol.* **8**:5-14.
- Parker, M. W., and C. C. F. Blake. 1988. Iron- and manganese-containing superoxide dismutases can be distinguished by analysis of their primary structures. *FEBS Lett.* **229**:377-382.
- Parker, M. W., and C. C. F. Blake. 1988. Crystal structure of manganese superoxide dismutase from *Bacillus stearothermophilus* at 2.4 Å resolution. *J. Mol. Biol.* **199**:649-661.
- Pearson, W. R., and D. J. Lipman. 1988. Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**:2444-2448.
- Rottlander, E., and T. A. Trautner. 1970. Genetic and transfection studies with *Bacillus subtilis* phage SP50. *J. Mol. Biol.* **108**:47-60.
- Sadosky, A. B., J. W. Wilson, H. M. Steinman, and H. A. Shuman. 1994. The

- iron superoxide dismutase of *Legionella pneumophila* is essential for viability. *J. Bacteriol.* **176**:3790–3799.
41. **Sambrook, J., E. F. Fritsch, and T. Maniatis.** 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
 - 41a. **Sanders, J. W.** Unpublished data.
 42. **Sanger, F. S., S. Nicklen, and A. R. Coulson.** 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* **74**:5463–5467.
 43. **Smart, J. B., and T. D. Thomas.** 1987. Effect of oxygen on lactose metabolism in lactic streptococci. *Appl. Environ. Microbiol.* **53**:533–541.
 44. **Steinman, H. M.** 1978. The amino acid sequence of manganese superoxide dismutase from *Escherichia coli* B. *J. Biol. Chem.* **253**:8708–8720.
 45. **Steinman, H. M.** 1992. Construction of an *Escherichia coli* K-12 strain deleted for manganese and iron superoxide dismutase genes and its use in cloning the iron superoxide dismutase gene of *Legionella pneumophila*. *Mol. Gen. Genet.* **232**:427–430.
 46. **Takeda, Y., and H. Avila.** 1986. Structure and gene expression of the *E. coli* Mn-superoxide dismutase. *Nucleic Acids Res.* **14**:4577–4589.
 47. **Tinoco, I., Jr., P. N. Borer, B. Dengler, M. D. Levine, O. C. Uhlenbeck, D. M. Crothers, and J. Gralla.** 1973. Improved estimation of secondary structure in ribonucleic acids. *Nature (London)* **246**:40–41.
 48. **Van Asseldonk, M., A. Simons, H. Visser, W. M. de Vos, and G. Simons.** 1993. Cloning, nucleotide sequence, and regulatory analysis of the *Lactococcus lactis dnaJ* gene. *J. Bacteriol.* **175**:1637–1644.
 49. **Van de Guchte, M., T. Van der Lende, J. Kok, and G. Venema.** 1991. Distance-dependent translational coupling and interference in *Lactococcus lactis*. *Mol. Gen. Genet.* **227**:65–71.
 50. **Yanisch-Perron, C., J. Vieira, and J. Messing.** 1985. Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors. *Gene* **33**:103–119.
 51. **Zabarovsky, E. R., and G. Winberg.** 1990. High efficiency electroporation of ligated DNA into bacteria. *Nucleic Acids Res.* **18**:5912.
 52. **Zitzelsberger, W., F. Götz, and K. H. Schleifer.** 1984. Distribution of superoxide dismutases, oxidases, and NADH peroxidases in various streptococci. *FEMS Microbiol. Lett.* **21**:243–246.