

SSP2 and OSW1, Two Sporulation-Specific Genes Involved in Spore Morphogenesis in *Saccharomyces cerevisiae*

Jing Li,* Seema Agarwal*^{†,1} and G. Shirleen Roeder*^{†,‡,2}

*Department of Molecular, Cellular and Developmental Biology, [†]Howard Hughes Medical Institute and [‡]Department of Genetics, Yale University, New Haven, Connecticut 06520-8103

Manuscript received October 2, 2006
Accepted for publication October 27, 2006

ABSTRACT

Spore formation in *Saccharomyces cerevisiae* requires the synthesis of prospore membranes (PSMs) followed by the assembly of spore walls (SWs). We have characterized extensively the phenotypes of mutants in the sporulation-specific genes, *SSP2* and *OSW1*, which are required for spore formation. A striking feature of the *osw1* phenotype is asynchrony of spore development, with some spores displaying defects in PSM formation and others spores in the same ascus blocked at various stages in SW development. The *Osw1* protein localizes to spindle pole bodies (SPBs) during meiotic nuclear division and subsequently to PSMs/SWs. We propose that *Osw1* performs a regulatory function required to coordinate the different stages of spore morphogenesis. In the *ssp2* mutant, nuclei are surrounded by PSMs and SWs; however, PSMs and SWs often also encapsulate anucleate bodies both inside and outside of spores. In addition, the SW is not as thick as in wild type. The *ssp2* mutant defect is partially suppressed by overproduction of either *Spo14* or *Sso1*, both of which promote the fusion of vesicles at the outer plaque of the SPB early in PSM formation. We propose that *Ssp2* plays a role in vesicle fusion during PSM formation.

SPORULATION in the budding yeast, *Saccharomyces cerevisiae*, involves two overlapping processes—meiosis and spore morphogenesis. During meiosis, a diploid cell undergoes a single round of DNA replication, followed by two rounds of chromosome segregation, to generate four haploid nuclei. During spore morphogenesis, the four daughter nuclei are packaged into spores, which are contained within an ascus.

Spore morphogenesis (reviewed by NEIMAN 2005) initiates with enlargement of the outer surface of the spindle pole body (SPB) to form the meiotic outer plaque. As cells undergo meiosis II, vesicles coalesce on the meiotic plaque to form a novel intracellular membrane called the prospore membrane (PSM). The double-layered PSM extends outward from each SPB to surround the adjacent nucleus. When the leading edges of the PSM meet and become fused, the haploid nucleus is fully encapsulated to form a prospore. PSM closure triggers the final step of spore morphogenesis, spore wall (SW) formation. The SW is assembled between the two membranes of the PSM and consists of four layers. The two inner layers are composed of glucan and mannan, which are components of vegetative cell walls.

The two outer layers are composed of chitosan and dityrosine and provide much of the spore's resistance to environmental stress. The inner layer of the former PSM forms the plasma membrane underlying the SW; the outer layer of the PSM breaks down during SW formation.

Genetic analysis indicates that the vesicles that fuse to form the PSM are post-Golgi vesicles, closely related to those normally involved in secretion (NEIMAN 1998). A widely accepted model for vesicle fusion events in the secretory pathway is the soluble *N*-ethylmaleimide-sensitive factor attachment protein receptors (SNARE) hypothesis. According to this model, every vesicle carries on its surface a vesicular (*v*-)SNARE that can interact only with a target (*t*-)SNARE on the correct acceptor membrane (JAHN and SUDHOF 1999). In the case of secretion in budding yeast, vesicles destined for the plasma membrane carry on their surface the *v*-SNARE molecules, *Snc1* and *Snc2*. The *Snc* proteins interact specifically with the *t*-SNARE proteins, *Sec9*, and *Sso1* and/or *Sso2*, on the plasma membrane (GERST 1999).

The synthesis of PSMs during sporulation also depends on SNAREs, although the players differ somewhat from those used during secretion in vegetative cells. In vegetative cells, *Sso1* and *Sso2* function redundantly. Sporulation, however, specifically requires *Sso1* (but not *Sso2*) (JANTTI *et al.* 2002; ENYENIHI and SAUNDERS 2003). *Sec9* is essential for secretion in vegetative cells, but it plays only a minor role during sporulation, where its function is largely replaced by the

¹Present address: Department of Internal Medicine, Section of Medical Oncology, Yale University School of Medicine, New Haven, CT 06520.

²Corresponding author: Department of Molecular, Cellular and Developmental Biology, Howard Hughes Medical Institute, Yale University, P.O. Box 208103, New Haven, CT 06520-8103.
E-mail: shirleen.roeder@yale.edu

sporulation-specific Sec9 homolog, Spo20 (NEIMAN 1998). In the *ssol* mutant or in the *sec9 spo20* double mutant, secretory vesicles accumulate near the meiotic plaque, but fail to coalesce to form a membrane sheet (NAKANISHI *et al.* 2006).

Another protein important for PSM formation is Spo14, a phospholipase D enzyme. Spo14 is essential during sporulation, and it localizes to PSMs (ROSE *et al.* 1995). The phenotype of a *spo14* null mutant is similar to that of the *ssol* and *sec9 spo20* mutants, indicating that Spo14 also participates in vesicle fusion (RIEDEL *et al.* 2005; NAKANISHI *et al.* 2006). Studies of nonnull alleles of *SPO14* and of mutations affecting the Spo14-interacting protein, Sma1, indicate that Spo14 acts at two discrete steps in PSM formation: the initial fusion of vesicles on the meiotic plaque and subsequent enlargement of the PSM to surround the adjoining nucleus (RIEDEL *et al.* 2005; NAKANISHI *et al.* 2006).

Expansion of the PSM depends on a complex of proteins, called the leading-edge complex, which includes Don1, Ssp1, and Ady3 (MORENO-BORCHART *et al.* 2001; NICKAS and NEIMAN 2002). These proteins initially localize to the SPB; as the PSM expands to engulf the adjacent nucleus, they localize to a ring corresponding to the growing edge of the PSM. The leading-edge complex is believed to be important in determining the shape and size of the PSM and in regulating the amount of cytoplasm surrounding each encapsulated nucleus (NEIMAN 2005).

In a screen for genes required for sporulation, we identified the sporulation-specific (*SSP2*) and outer spore wall (*OSW1*) genes. Concurrent with our studies, *SSP2* and *OSW1* were also characterized by two other groups (SARKAR *et al.* 2002; COLUCCIO *et al.* 2004). Our results indicate that the *ssp2* and *osw1* null mutants are proficient in meiotic recombination and chromosome segregation, but defective in the subsequent processes of PSM and SW formation. The *osw1* mutant exhibits a heterogeneous phenotype with different spores in the same ascus arrested at different stages of PSM and SW formation. On the basis of the similarity of *osw1* to various kinase mutants, we suggest that the *Osw1* protein plays a role in regulating and coordinating spore morphogenesis. In the *ssp2* mutant, nuclei are surrounded by PSMs and SWs, but the SW is only half the thickness of its wild-type counterpart. In addition, anucleate spore-like bodies are generated, raising the possibility that PSM formation is not tightly coupled to SPBs. The *ssp2* defect is partially suppressed by overproduction of Spo14 or Sso1, suggesting that Ssp2 plays a role in vesicle fusion during PSM formation.

MATERIALS AND METHODS

Plasmids: Standard methods were used in plasmid constructions (SAMBROOK *et al.* 1989). To construct an *SSP2*

disruption, *SSP2* was first amplified by PCR from the *S. cerevisiae* λ -clone 6767 (ATCC 70786) to introduce a *Bam*HI site ~270 bp upstream of the *SSP2* ORF and a *Kpn*I site ~80 bp downstream. The *Bam*HI–*Kpn*I fragment containing *SSP2* was inserted between the *Bam*HI and *Kpn*I sites of Bluescript SK (+) (Stratagene, La Jolla, CA) to create pJS1. To construct pJS2 (*ssp2::URA3*), the ~1-kbp *Hind*III–*Sal*I fragment of *URA3* (CHUA and ROEDER 1998) was used to replace the *Hind*III–*Xho*I fragment of *SSP2* (nucleotides 71–727 of the coding region) in pJS1.

An *SSP2*–*GFP* fusion gene was constructed as follows. PCR was used to introduce a *Not*I site ~250 bp upstream of the *SSP2* ORF and an *Nhe*I site just before the stop codon. The *Not*I–*Nhe*I fragment (~1.4 kbp) containing *SSP2* was then subcloned into the *Not*I–*Nhe*I sites of pSA111 (AGARWAL and ROEDER 2000), thus creating pJ82, which contains the coding sequences for green fluorescent protein (*GFP*) fused to the 3' end of *SSP2*.

To construct an *OSW1* disruption, a 2.0-kbp *Eco*RI fragment containing *OSW1* from the *S. cerevisiae* λ clone 6933 (ATCC 71157) was first subcloned into pHSS6 (SEIFERT *et al.* 1986) to create pJS4. To construct pJS5 (*osw1::URA3*), the ~1-kbp *Hind*III–*Sma*I fragment of *URA3* (CHUA and ROEDER 1998) was used to replace the *Hinc*II–*Hind*III fragment of *OSW1* (nucleotides 70–721 of the coding region) in pJS4.

An *OSW1*–*GFP* fusion gene was constructed as follows. PCR was used to introduce a *Not*I site ~290 bp upstream of the *OSW1* ORF and an *Nhe*I site just before the stop codon. The *Not*I–*Nhe*I fragment (~1.1 kbp) containing *OSW1* was subcloned into the *Not*I–*Nhe*I sites of pSA111 (AGARWAL and ROEDER 2000) to create pJ83, which contains *GFP* fused to the 3' end of *OSW1*.

pSB8 is a centromere-containing, pRS314-based plasmid encoding Don1–*GFP* (TACHIKAWA *et al.* 2001). pME1086 is a multicopy YEp351-based plasmid encoding Spo14–*GFP* (RUDGE *et al.* 1998). The *Xba*I–*Sal*I fragment (~4.5 kbp) containing part of *SPO14* and the full sequence of *GFP* from the *SPO14*–*GFP* plasmid was subcloned into the *Xba*I–*Sma*I sites of the integrating plasmid pRS305 (SIKORSKI and HIETER 1989) to create pJ121. YEpSSO1 and YEpSSO2 are pMAC561-based plasmids containing the *SSO1* and *SSO2* genes (RUOHONEN *et al.* 1997); pMA56 was used as the control plasmid.

Yeast strains: Yeast manipulations were performed, and media were prepared as described previously (SHERMAN *et al.* 1986). Gene disruptions were confirmed by PCR analysis. All strains used are isogenic with BR2495 (ROCKMILL and ROEDER 1990), which has the following genotype: *MATa/MAT α his4-280/his4-260 leu2-27/leu2-3,112 arg4-8/ARG4 thr1-1/thr1-4 cyh10/CYH10 ade2-1/ade2-1 ura3-1/ura3-1 trp1-1/trp1-289*. Diploids used in this study were made by mating appropriate haploids, generated by transformation and/or genetic crosses. All strains used are diploids homozygous for the disruptions/fusions indicated.

ssp2::URA3 and *osw1::URA3* mutant strains were constructed by transforming with the *Bam*HI–*Kpn*I fragment of pJS2 and the *Eco*RI fragment of pJS5, respectively.

pAFS59 (STRAIGHT *et al.* 1996) containing 256 Lac operator (*LacO*) repeats was used to integrate *LacO* at the *LEU2* locus, which is located 22 kbp to the left of *CENIII*. pAFS152 (SHONN *et al.* 2000), containing the *GFP*–*Lac* repressor (*LacI*) fusion driven by the *CYC1* promoter, was integrated at the *URA3* locus.

pJ121 was digested with *Bsa*BI and then used to transform wild-type, *ssp2::URA3* and *osw1::URA3* haploids, creating strains in which *SPO14*–*GFP* is integrated at its genomic loci.

Measurements of sporulation and nuclear division: To measure sporulation efficiency, each saturated culture (1 ml) was washed once with distilled water and resuspended in 10 ml of sporulation medium (2% potassium acetate). After 72 hr at

30°, cells were observed by differential interference contrast (DIC) microscopy. Three cultures were examined for each strain, and at least 100 cells were counted for each culture. The percentages presented are averages.

To determine the frequency and kinetics of meiotic nuclear division, cells were fixed in 70% ethanol and frozen at -20° prior to staining with 4'-6-diamidino-2-phenylindole (DAPI) (HONG and ROEDER 2002). Cells were observed and counted under the fluorescence microscope (Nikon Eclipse E800, Plan Apo100×/1.4 oil objective).

Chromosome segregation was monitored in wild-type, *ssp2::URA3*, and *osw1::URA3* strains with the LacO-marked chromosome III (*LEU2*) bound with GFP-LacI (STRAIGHT *et al.* 1996). The cells were first fixed with formaldehyde, stained with DAPI (1 µg/ml), and then observed under the fluorescence microscope. All fluorescence images presented in this article were captured with a fluorescence microscope (Nikon Eclipse E800, Plan Apo100×/1.4 oil objective) equipped with an FITC HYQ filter to visualize the GFP signals. Images were captured with a Sensys CCD camera (Photometrics, Tucson, AZ). Each image presented represents a single focal plane.

Analysis of spread nuclei: Meiotic nuclei were spread and stained with antibodies as described (BAILIS and ROEDER 1998). Guinea pig anti-GFP antibodies (TSUBOUCHI and ROEDER 2002) and rat anti-tubulin antibodies (Sera-Lab) were used at a 1:100 dilution. Anti-rabbit antibodies conjugated with FITC and anti-rat antibodies conjugated with Texas Red (both from Jackson ImmunoResearch Labs, West Grove, PA) were used at a 1:200 dilution as secondary antibodies. Chromosomal DNA was stained with 1 µg/ml DAPI.

Staining of fixed whole cells: To observe the Ssp2-GFP fluorescent signal combined with DAPI staining in whole cells, ~20 µl of sporulating cells were scraped from sporulation plates and suspended in solution A that contains 3.7% formaldehyde, 1.2 M sorbitol, and 50 mM potassium phosphate buffer, pH 6.8. After fixation for 1 hr at room temperature with rotation, cells were washed three times with solution A and incubated at 37° for 1 hr in solution A supplemented with 0.1% β-mercaptoethanol and 25 µg/ml zymolyase. Cells were washed twice with solution A and resuspended in a small volume of phosphate-buffered saline (PBS). Cells were then added to Poly-L-lysine coated slides and rinsed once with PBS followed by PBS + 0.1% NP-40. Guinea pig anti-GFP antibody (TSUBOUCHI and ROEDER 2002) was diluted 1:100 in PBS + 1% BSA and used to stain cells. After overnight incubation at 4°, cells were rinsed with PBS, then PBS + 0.1% NP-40, and then PBS. Anti-guinea pig antibodies conjugated with FITC and diluted 1:200 in PBS + 1% BSA were used as secondary antibodies. After 2.5 hr of incubation at room temperature, the cells were rinsed in PBS as described above and then mounted with DAPI.

Electron microscopy: Samples for examination in the electron microscope (EM) were prepared by potassium permanganate (KMnO₄) fixation as described by Du and NOVICK (2002). Samples were sectioned using a Sorvall MT-2 Ultramicrotome (Waltham, MA) and then analyzed using a Zeiss EM 10A transmission EM (Zeiss, Jena, Germany). For measurements of SW thickness, 11 asci were examined for wild type and 13 for *ssp2*.

RESULTS

Identification of the *SSP2* and *OSW1* genes: We previously identified transposon insertions that generate in-frame lacZ fusion genes expressed specifically during sporulation (BURNS *et al.* 1994; ROSS-MACDONALD *et al.* 1999). Diploids homozygous for these insertions




FIGURE 1.—Ssp2 and Osw1 are essential for sporulation. Typical cells of wild type (A and B), *ssp2* (C and D), and *osw1* (E and F) are shown. Cells were sporulated and observed under DIC microscopy after 72 hr in sporulation medium. Bar, 2 µm.

were constructed and screened for sporulation competence. Diploids derived from strains M86 and M88 were unable to sporulate; these represent insertions in ORFs YOR242c and YOR255w, respectively. Both of these ORFs were identified in previous screens for sporulation-defective mutants (RABITSCH *et al.* 2001; SARKAR *et al.* 2002; ENYENIHI and SAUNDERS 2003). YOR242c has been termed *SSP2* by SARKAR *et al.* (2002), and YOR255w has been named *OSW1* by COLUCCIO *et al.* (2004). *SSP2* encodes a protein of 371 amino acids, which shares significant homology with IPF16670, a predicted gene product of *Candida albicans*. *OSW1* encodes a protein of 278 amino acids with no significant similarity to other proteins in databases.

Ssp2 and Osw1 are required for sporulation: Deletion/disruption mutants of *SSP2* and *OSW1* were made by replacing most of each coding region with the *URA3* gene (see MATERIALS AND METHODS). Homozygous *ssp2* and *osw1* diploids and an isogenic wild type were tested for sporulation efficiency by DIC microscopy (see MATERIALS AND METHODS). In wild type, 65% of cells produce mature four-spored asci (Figure 1, A

and B) after 72 hr in sporulation medium. However, only 5 and 1% of cells generate mature asci in *ssp2* and *osw1*, respectively. A significant fraction of cells (63% in *ssp2* and 57% in *osw1*) appear to contain compartments, which are irregular in size and shape, suggestive of immature spores (Figure 1, C–F). The Ssp2 and Osw1 proteins are therefore required for spore formation.

Meiotic division is not affected in *ssp2* and *osw1* mutants: To assess meiotic nuclear division, cells harvested at different time points after the induction of meiosis were examined by fluorescence microscopy after staining with DAPI, a DNA-binding dye. Cells that have completed the meiosis I division contain two discrete nuclei, whereas cells that have completed both meiotic divisions contain four nuclei. Wild-type, *ssp2*, and *osw1* cells undergo nuclear division with nearly identical kinetics (Figure 2A). There is no evidence of nuclear fragmentation in the mutants; four DAPI-stained bodies, equivalent in size, are present in nearly every ascus (data not shown). Thus, analysis of nuclear division suggests that the kinetics and efficiency of meiotic progression are normal in the *ssp2* and *osw1* mutants.

Commitment to meiotic recombination is normal in the *ssp2* and *osw1* mutants: To examine the effect of *ssp2* and *osw1* on meiotic recombination, the null mutants were used to measure commitment to gene conversion (intragenic recombination) in return-to-growth assays. Cells were introduced into sporulation medium and then returned to growth medium prior to the commitment to meiotic chromosome segregation (ESPOSITO and ESPOSITO 1974). The frequency of gene conversion was measured in *ssp2*, *osw1*, and an isogenic wild type by selecting for prototrophic recombinants at four different loci carrying heteroalleles conferring auxotrophy. Commitment to meiotic gene conversion is normal in *ssp2* and *osw1* mutants (Figure 3), with respect both to timing and to frequency.

Chromosome segregation is not affected in the *ssp2* and *osw1* mutants: To examine chromosome segregation, strains were constructed that produce a GFP–LacI fusion protein and are homozygous for a tandem array of LacO repeats inserted 22 kbp from *CENIII* (STRAIGHT *et al.* 1996) (see MATERIALS AND METHODS). Thus, segregation of the LacO repeats, and by implication the centromere of chromosome III, can be monitored by fluorescence microscopic detection of the GFP–LacI protein (STRAIGHT *et al.* 1996).

Wild-type, *ssp2*, and *osw1* cells were harvested after 24 hr of sporulation, when ~60% of wild-type cells had undergone one or both nuclear divisions (Figure 2A). Cells were fixed, stained with DAPI, and observed by fluorescence microscopy. In wild type, some cells show only one GFP signal, indicative of meiotic prophase (or a failure to enter meiosis). Others show two distinct GFP signals, indicative of meiosis I division products. Some show four distinct GFP dots (one per DAPI-stained body), indicating the end of meiosis II (Figure 2B). The




FIGURE 2.—Chromosome segregation occurs normally in *ssp2* and *osw1*. (A) Meiotic nuclear division was examined in wild-type, *ssp2::URA3*, and *osw1::URA3* strains. Both binucleate and tetranucleate cells were scored as having undergone nuclear division. (B and C) Chromosome segregation was monitored in a strain that produces GFP–LacI fusion protein and is homozygous for a tandem array of LacO repeats inserted 22 kbp from *CENIII*. (B) An example of a cell that has undergone normal segregation is shown. Four DAPI-stained bodies indicate that chromosomes have undergone both meiotic divisions. One GFP dot per DAPI-stained body indicates that GFP-tagged chromosomes have segregated correctly. Bar, 2 μ m. (C) Quantitation of GFP signals in wild-type, *ssp2::URA3*, and *osw1::URA3* strains. Cells were harvested after 24 hr of sporulation, stained with DAPI, and examined by fluorescence microscopy. The percentages of cells with one, two, three, or four GFP dots are indicated. At least 100 cells were scored for each strain. The experiments shown in A and C were performed twice with qualitatively similar results.




FIGURE 3.—Commitment to meiotic recombination is normal in *ssp2* and *osw1*. (A–D) Intragenic recombination was measured in wild-type, *ssp2::URA3*, and *osw1::URA3* strains sporulated at 30°. Prototroph formation was monitored at four different heteroallelic loci at different time points during sporulation as described by SYM *et al.* (1993). The frequencies of prototroph formation at *HIS4* (*his4-280/his4-260*), *THR1* (*thr1-1/thr1-4*), *LEU2* (*leu2-27/leu2-3,112*), and *TRP1* (*trp1-1/trp1-289*) are shown as a function of time.

ssp2 and *osw1* mutants show the same pattern of GFP signals as wild type (Figure 2C), indicating that chromosomes segregate correctly in the absence of Ssp2 or Osw1. Since crossing over is required for correct chromosome segregation in meiosis, these results indicate that the *ssp2* and *osw1* mutants are proficient in crossing over.

Localization of the Ssp2 and Osw1 proteins during meiosis: *SSP2-GFP* and *OSW1-GFP* fusion genes were constructed (see MATERIALS AND METHODS) and used to determine the intracellular locations of Ssp2 and Osw1. Single-copy plasmids carrying the fusion genes do not complement. However, when the fusion genes are present on a multicopy plasmid (carrying the 2 μ circle origin of DNA replication), they enable the corresponding null mutants to sporulate as efficiently as wild type, indicating that the fusion proteins are partially functional. The *ssp2* + 2 μ *SSP2-GFP* and *osw1* + 2 μ *OSW1-GFP* strains were introduced into sporulation medium, and cells harvested at various time points were observed by fluorescence microscopy.

Figure 4 shows images of living cells expressing *SSP2-GFP* during the course of sporulation. In vegetative cells and early in meiosis, no Ssp2-GFP signal is detected (data not shown). Starting around 18 hr, when many cells have completed the meiosis II division, the Ssp2-GFP signal is detected as small ring-like structures (Figure 4, A and B). By 25 hr, the rings in most cells have enlarged to occupy most of the ascus (Figure 4C). Each Ssp2-GFP ring surrounds an individual nucleus, as determined by staining fixed cells both with anti-GFP antibodies and with DAPI (Figure 4G). Thus, the Ssp2-GFP signals observed must be associated with PSMs and/or SWs. SARKAR *et al.* (2002) also reported Ssp2

localization to rings surrounding nuclei starting after the meiosis II division.

Osw1 localization shows different patterns at different times during sporulation. When living cells expressing *OSW1-GFP* are observed by fluorescence microscopy, Osw1-GFP first appears as two dots (Figure 5A) and later as four distinct foci (Figure 5B). Double staining of spread nuclei with anti-tubulin and anti-GFP antibodies indicates that these dots represent SPBs at meiosis I (two dots) and meiosis II (four dots) (Figure 5, G and H). After meiotic nuclear division is complete, the Osw1-GFP signals enlarge to fully surround nuclei, in addition to being present at the four SPBs (Figure 5C).

The leading edge of PSMs is affected by *osw1* but not by *ssp2*: Don1 was used as a marker to determine whether *ssp2* and/or *osw1* are defective in formation of the leading edge of the PSM. Living cells producing a Don1-GFP fusion protein were examined.

In wild type, Don1-GFP first appears as four distinct dots (Figure 6A). As the leading edge of the PSM moves away from the SPB to engulf the adjacent nucleus, Don1 staining appears as rings or bars (Figure 6B) depending on whether the leading edge of the membrane is viewed from the side (bars) or in cross section (rings). No obvious differences in Don1 localization were observed between the *ssp2* mutant (Figure 6, C and D) and wild type.

In the *osw1* mutant, a variable number of Don1 dots/rings (one to four) were observed (Figure 6, E–H). Some cells contain partially developed or aberrantly shaped Don1 rings (Figure 6G); other cells contain four Don1 dots or rings at different stages of development (Figure 6, F and H). These observations suggest that Don1 rings, and thus the leading edges of the PSMs,




FIGURE 4.—Localization of Ssp2-GFP in sporulating cells. (A–F) The *ssp2* mutant producing Ssp2-GFP was sporulated and then analyzed by observing GFP fluorescence in living cells. (A–C) Representative cells producing Ssp2-GFP. (D–F) Diagrams of the images in A–C. The diagram is based on the background GFP signal a cell displays when overexposed. Bar, 2 μ m. (G) *ssp2* cells containing 2 μ SSP2-GFP were fixed and then stained with DAPI (blue) and anti-GFP antibodies (green). The fourth DAPI-stained body is visible in a different plane of focus. Time points for each image: A–B, 20 hr; C, 24 hr; G, 24 hr.

develop aberrantly and asynchronously in the absence of Osw1.

Both Ssp2 and Osw1 are required for proper localization of Spo14: To examine PSM formation further, the Spo14 protein, a marker for the PSM, was localized in diploid cells in which both copies of the *SPO14* gene had been disrupted by insertion of a *SPO14-GFP* fusion gene (see MATERIALS AND METHODS). *SPO14-GFP* integration has no effect on sporulation efficiency in wild-type, *ssp2*, or *osw1* strains (data not shown). No specific GFP signal was observed in living cells. Thus, to detect Spo14-GFP, cells were fixed and stained with anti-GFP antibodies; cells were simultaneously stained with DAPI to visualize nuclei.

When sporulation is complete, ~70% of wild-type cells show four Spo14-GFP circles around four DAPI-stained nuclear bodies (Figure 7, A–C). However, none of the *ssp2* cells show a similar pattern. Instead, Spo14-GFP signals most often appear as punctate dots dispersed around DAPI signals (Figure 7, D–L). *ssp2* cells sometimes display linear stretches of continuous Spo14 staining (Figure 7, G and J, open arrows); occasionally, small Spo14-GFP circles centered around DAPI-stained material are observed (Figure 7D, open arrow). Large circles surrounded by Spo14-GFP staining (like the circles seen in wild type) were never observed. The




FIGURE 5.—Localization of Osw1-GFP in sporulating cells. (A–F) The *osw1* mutant producing Osw1-GFP was sporulated and then analyzed by observing GFP fluorescence in living cells. (A–C) Representative cells producing Osw1-GFP. (D–F) Diagrams of the images in A–C. In the case of the cell shown in C, three SPBs are evident (arrowheads), while the fourth SPB is evident in a different plane of focus. (G–H) Spread nuclei from *osw1* containing 2 μ *OSW1-GFP* were stained with DAPI (blue), anti-GFP (green), and anti-tubulin (red) antibodies. Bar, 2 μ m. Time points for each image: A, 17 hr; B, 18 hr; C, 23 hr; G–H, 24 hr.

predominantly punctate staining pattern and the absence of large circles suggest that PSMs do not mature properly in the *ssp2* mutant.

Spo14-GFP mislocalizes even more severely in the *osw1* mutant. Spo14-GFP staining often appears to be concentrated to one side of the ascus. There are no obvious circles of Spo14 staining, although short linear stretches are sometimes observed (Figure 7, M, P, and S, open arrows). DAPI-stained masses without any surrounding Spo14-GFP signal are often observed (Figure 7, N and T, arrowheads), suggesting the absence of a PSM.

Thus, Spo14-GFP mislocalizes in both mutants, indicating roles for Ssp2 and Osw1 in the formation of PSMs.

***ssp2* and *osw1* mutants form aberrant structures in prospores:** To investigate further the *ssp2* and *osw1* defects in spore formation, sectioned cells were examined in the EM. In wild-type cells, each PSM starts to develop at one side of a nucleus (Figure 8A), then expands to




FIGURE 6.—Don1 rings are normal in *ssp2*, but aberrant in *osw1*. Wild-type (A and B), *ssp2* (C and D), and *osw1* (E–H) strains carrying *DON1-GFP* on a centromeric plasmid were sporulated. They were then analyzed by observing GFP fluorescence in living cells. The cell in E has two Don1 rings. In F, the top left Don1 signal is discontinuous, suggestive of incomplete ring formation. In H, the lower Don1 bar is significantly longer than the others. Time points for each image: A, 21 hr; B, 25 hr; C, 21 hr; D, 25 hr; E, 21 hr; F, 23 hr; G, 23 hr; H, 25 hr. Bar, 2 μ m.

engulf the nucleus (Figure 8B), and eventually closes when the nucleus is fully surrounded. Later in spore morphogenesis, the SW materials are deposited between the two layers of the PSM (Figure 8C). In our EM preparations, the two innermost layers of the SW (composed of glucan and mannan) appear as one thick electron-translucent layer (Figure 8D), referred to hereafter as the inner SW (ISW). The two outer layers of the SW (composed of dityrosine and chitosan) appear as one thin darkly stained layer, referred to hereafter as the outer spore wall (OSW). The inner layer of the former PSM defines the plasma membrane underlying the mature SW.

During the early stage of PSM development in the *ssp2* mutant, PSMs are formed; however, short pieces of membrane-like structures are often seen close by the more fully developed PSMs (Figure 8E, arrows). The frequency of these membranous structures is 1.7 ± 0.9 /spore section. (Similar pieces of membrane were never observed in wild type.) By the time the PSM has successfully encapsulated an entire daughter nucleus, longer stretches of free membrane are frequently observed (Figure 8, F and G, arrows). Prospores do form and SWs start to develop in the *ssp2* mutant; however, PSMs and SWs often encapsulate unidentified electron-dense materials both inside and outside of spores (Figure 8, H and I, arrows). In addition, some spores are very irregular in shape (Figure 8J). The ISW does not reach the thickness seen in wild type (compare Figure 8, H–J, with 8D). The ISW is 0.22 ± 0.01 μ m thick in wild type compared to 0.09 ± 0.01 μ m in *ssp2*. The OSW appears to form normally in *ssp2*.

The *osw1* mutant displays various defects. The most obvious is the asynchronous development of the four spores within an ascus. At the stage when PSMs develop, sometimes only a subset of nuclei become surrounded by PSMs (Figure 9, A–C). At the SW maturation stage, some asci contain prospores with immature SWs of different thicknesses (Figure 9D). A second defect is that the OSW is rarely seen (Figure 9, C–F). In addition, electron-dense materials often accumulate around the surface of the SW (Figure 9, E and F, arrowheads). This accumulation has been observed in other SW-defective mutants (WAGNER *et al.* 1997; CHRISTODOULIDOU *et al.*

1999; TACHIKAWA *et al.* 2001). Last, the immature spores sometimes are very irregular in shape (Figure 9, E and F), perhaps due to the absence of the OSW for stabilization.

Overproduction of Spo14 partially suppresses the sporulation defect of *ssp2*: In the process of immunolocalizing the Spo14 protein, we discovered that overproduction of the Spo14–GFP protein partially suppresses the sporulation defect of the *ssp2* mutant. When *SPO14-GFP* was introduced on a multicopy plasmid, 21% of *ssp2* cells formed mature four-spored asci (as assessed by DIC microscopy), compared to only 5% of cells carrying an empty vector. Overproduction of Spo14–GFP does not alter the sporulation efficiency of wild type or *osw1* (RUDGE *et al.* 1998) (data not shown).

Both wild-type and *ssp2* strains overproducing Spo14–GFP were sporulated, and living cells were examined for GFP fluorescence. Around the time of the meiosis II division, many cells (both wild type and mutant) contain four small GFP dots, most likely representing Spo14 near SPBs, as reported previously (Figure 10, A and D) (RUDGE *et al.* 1998). (SPB staining was not observed in cells containing single-copy *SPO14-GFP*.) At later times during sporulation in wild type, circles of Spo14–GFP were observed; these eventually enlarged to fill the entire ascus (Figure 10, B and C). In *ssp2* overproducing Spo14–GFP, a subset of cells contained large rings and were indistinguishable from the mature asci observed in wild type (Figure 10E). The frequency of cells that exhibit normal Spo14–GFP staining ($\sim 23\%$) is similar to the frequency of cells that completed sporulation ($\sim 21\%$). The remaining cells contained rings that were aberrant in several respects. First, the rings did not fill the entire ascus. Second, some rings had aberrant protrusions (Figure 10, F and G, arrowheads), reminiscent of the anucleate spore-like structures observed in the EM. Third, some of the cells were irregular in shape (Figure 10G, arrow), similar to the foot-shaped spores observed in the EM (Figure 8J). In every case, Spo14–GFP staining was more continuous in *ssp2* cells containing multiple copies of the *SPO14-GFP* gene than in cells carrying an integrated version of *SPO14-GFP*.

Overproduction of Sso1 partially suppresses the sporulation defect of *ssp2*: To determine whether the




FIGURE 7.—The localization of Spo14–GFP is aberrant in *ssp2* and *osw1*. Wild-type, *ssp2*, and *osw1* strains with *SPO14–GFP* integrated (see MATERIALS AND METHODS) were sporulated for 3 days before cells were fixed and stained with DAPI (blue) and rabbit anti-GFP antibodies (green). Arrows, small Spo14–GFP circles or short linear stretches of Spo14–GFP signals. Arrowheads, DAPI-stained bodies without surrounding Spo14–GFP signals. Bar, 2 μ m. The fourth Spo14–GFP circle in A and C and the fourth DAPI-stained body in T and U were evident in another focus plane. No specific GFP signals were observed in a control strain lacking Spo14–GFP (data not shown).

Sso proteins can function as suppressors, multicopy plasmids carrying *SSO1* or *SSO2* (RUOHONEN *et al.* 1997) were introduced into wild-type, *ssp2*, and *osw1* strains. Overproduction of Sso1 increases sporulation in *ssp2* (from 5 to 22%), but has no effect on sporulation efficiency in wild type or *osw1*. Overproduction of Sso2 had no effect on sporulation in wild type, *ssp2*, or *osw1*. The differential effect of *SSO1* and *SSO2* overproduction is consistent with the finding that Sso1, but not Sso2, functions in meiotic cells (JANTTI *et al.* 2002).

DISCUSSION

SSP2 and OSW1—two genes involved in spore morphogenesis: We have characterized extensively the phenotypes of the *ssp2* and *osw1* mutants. Our data demonstrate that the Ssp2 and Osw1 proteins are not required for early events in meiosis, including recombination, chromosome segregation, and nuclear division. These results are consistent with those of SARKAR *et al.* (2002), who examined nuclear division and recombination in *ssp2*. Our data indicate that Ssp2 and Osw1 are required for the normal formation of PSMs and mature SWs. This defect in spore morphogenesis is consistent with the timing of expression of the *SSP2* and *OSW1* genes (CHU *et al.* 1998; NAG and AXELROD 1998), which are expressed at the same time as genes involved in PSM formation.

Osw1 may play a regulatory role in spore morphogenesis: A striking feature of the *osw1* null mutant phenotype is the asynchrony of spore morphogenesis among different spores in the same ascus. Some nuclei are never encapsulated by PSMs even at very late times in sporulation medium, although well-defined PSMs encapsulate other nuclei in the same ascus. Consistent with the heterogeneity observed in the EM, fluorescence microscopy demonstrates variation in the number and morphology of Don1 rings. Some asci contain spores with SWs of different thicknesses, providing another indication of a lack of coordination among spores within an ascus. OSWs are rarely seen in *osw1*; the aberrant shapes of some prospores might result from lack of the OSW for stabilization. COLUCCIO *et al.* (2004) also observed an absence of OSWs in *osw1*; they did not report any other defects.

Several other mutants have been described that display heterogeneous defects in the development of PSMs and SWs (FRIESEN *et al.* 1994; KRISAK *et al.* 1994; WAGNER *et al.* 1997; STRAIGHT *et al.* 2000; COLUCCIO *et al.* 2004). These include mutations in four genes encoding protein kinases, *SMK1*, *SPS1*, *CAK1* and *MPS1*, as well as mutations in the *SPO75* gene, which encodes an integral membrane protein. The spectrum of phenotypes displayed by the *smk1* null mutant is very similar to that of the *osw1* mutant (FRIESEN *et al.* 1994; KRISAK *et al.* 1994). It is generally assumed that the Smk1 kinase acts in a signaling pathway to promote and coordinate the




FIGURE 8.—*ssp2* forms aberrant PSMs and SWs. Wild-type (A–D) and *ssp2* (E–J) strains were incubated in sporulation medium at 30°. (A and E) Cells were harvested at 24 hr, when ~60% of cells have undergone one or both nuclear divisions. (B–D and F–J) Cells were harvested at 36 hr, when spore morphogenesis is complete in wild type. Arrows in E–G indicate aberrant membrane-like structures; arrows in H and I indicate aberrant anucleated spore-like bodies. N, nucleus; PSM, prospore membrane; ISW, inner spore wall; OSW, outer spore wall; imSW, immature spore wall. Bar, 500 nm.

different stages in spore wall morphogenesis (WAGNER *et al.* 1999). The similarity in phenotypes between the *osw1* and *smk1* mutants suggests that *Osw1* also performs a regulatory function.

The location of the *Osw1* protein is compatible with its hypothesized function in the regulation of PSM and SW development. *Osw1* is found at SPBs around the time that the meiotic plaque is active in initiating formation of the PSM. At subsequent stages, it surrounds developing prospores, consistent with it being a component of the PSM.

***Ssp2* affects PSMs and SWs:** Our study reveals two aspects of the *ssp2* mutant phenotype. First, there is a defect in PSM development. PSMs are often observed encapsulating materials both inside spores and in the spaces between spores. Furthermore, the PSM component, *Spo14*, fails to localize properly. Second, the ISW is only half the thickness of its wild-type counterpart. This aberration presumably explains why the developing spores are not refractile to light and therefore not observed in the light microscope. Unlike the *osw1* mutant in which different spores develop to different extents,

the *ssp2* mutant phenotype is fairly homogeneous with almost all spores exhibiting the same phenotype.

There are two possible interpretations of the *ssp2* mutant phenotype. First, the *Ssp2* protein might act at two different steps in spore morphogenesis, first during PSM development and later during SW assembly. A more economical interpretation is that the primary defect in *ssp2* is in PSM formation and the alteration in the SW is a secondary consequence of the defect in the PSM. SW materials might not be properly deposited between the two layers of the PSM when the PSM is aberrant. A role for *Ssp2* in PSM development is consistent with the localization of *Ssp2* to rings surrounding prospores even at very early stages in spore morphogenesis.

The *ssp2* mutant phenotype has been characterized previously by two groups (SARKAR *et al.* 2002; COLUCCIO *et al.* 2004). Both groups reported that the PSM develops properly. However, the development of anucleate spore-like bodies is a subtle phenotype that might easily have escaped detection, and the localization of PSM components was not examined previously. Neither group reported any difference in the thickness of the ISW,




FIGURE 9.—*osw1* forms aberrant PSMs and SWs. *osw1* cells were analyzed as described. Open arrowheads in E and F, electron-dense materials around the surface of the SW. N, nucleus (numbered in A–C); PSM, prospore membrane; ISW, inner spore wall. (A) N1 and N2 are being engulfed by a PSM, while N3 does not have any PSM. (B) N1, N3, and N4 have PSMs, while N2 does not. (C) N2 is fully encapsulated by a PSM and an incipient SW structure is discernible, while N1 does not show any PSM formation. Bar, 500 nm.

although no measurements were presented. The most striking difference between our results and those published previously is that we do see an OSW in *ssp2*, whereas both SARKAR *et al.* (2002) and COLUCCIO *et al.*

(2004) described the failure to develop OSWs as the sole defect in *ssp2*. The difference between our results and those published previously might be due to differences in yeast strain background and/or to differences in the methods used to prepare samples for examination in the EM. Our results indicate that Ssp2 acts at a much earlier stage than previously thought.

Ssp2 acts in the vesicle fusion pathway of PSM formation: Formation of the PSM involves two distinct processes (NEIMAN 2005). The first process, membrane initiation, is a homotypic fusion event in which post-Golgi vesicles fuse with each other. This event occurs at the meiotic plaque of the SPB, and the plaque is essential for the coalescence of vesicles to initiate a PSM. The second process, membrane expansion, involves heterotypic fusion events in which vesicles fuse with the developing PSM until the nucleus is fully encapsulated.

The *ssp2* mutant defects are partially suppressed by overproduction of either Spo14 or Sso1. Since these proteins are involved in vesicle fusion during PSM formation (RIEDEL *et al.* 2005; NAKANISHI *et al.* 2006), it is likely that Ssp2 also plays a role in vesicle fusion. As noted in the Introduction, Spo14 in both initiation and expansion of the PSM (RIEDEL *et al.* 2005; NAKANISHI *et al.* 2006). Similarly, Ssp2 could act at both steps. In support of a role in PSM initiation, the *ssp2* mutant produces pieces of membrane that are not associated with SPBs, as if vesicle fusion events are not obligatorily coupled to the meiotic outer plaque in the absence of Ssp2.

Other observations suggest that Ssp2 plays a role in membrane expansion. Although fully developed PSMs are often observed in the EM in *ssp2* cells, large circles of Spo14–GFP staining are never observed by fluorescence microscopy. This discrepancy suggests that Spo14–GFP localization is aberrant in PSMs, with Spo14 localizing in a punctate pattern instead of uniformly throughout the PSM. The aberrant localization of Spo14 in the PSMs




FIGURE 10.—Overproduction of Spo14 partially suppresses *ssp2*. Wild-type (A–C) and *ssp2* (D–G) strains overexpressing *SPO14-GFP* were sporulated. They were then analyzed by observing GFP fluorescence in living cells. Bar, 2 μ m. Open arrowheads in F and G point to aberrant protrusions also observed in the EM (Figure 8, H and I, arrows). Arrow in G points to a spore that is aberrant in shape, reminiscent of images seen in the EM (*e.g.*, Figure 8J).

that surround nuclei implies that PSM development is perturbed subsequent to the initiation stage. In addition, it is noteworthy that the PSMs that form in the *ssp2* mutant do not always capture a nucleus (Figure 8, H and I), suggestive of a failure at the expansion stage (NEIMAN 1998).

We thank Aaron Neiman, JoAnne Engebrecht, and Sirkka Keranen for providing plasmids. Oligonucleotides were synthesized by the W. M. Keck Foundation Biotechnology Resource Laboratory at Yale University. We are grateful to members of the Roeder lab for helpful discussions and comments on the manuscript. In addition, we thank Li-Lin Du, JoAnne Engebrecht, and Aaron Neiman for helpful discussions and sharing unpublished data and protocols. We are grateful to Barry Piekos for his able assistance with electron microscopy. This work was supported by grant GM28904 from the United States Public Health Service to G.S.R. and by the Howard Hughes Medical Institute.

LITERATURE CITED

- AGARWAL, S., and G. S. ROEDER, 2000 Zip3 provides a link between recombination enzymes and synaptonemal complex proteins. *Cell* **102**: 245–255.
- BAILIS, J. M., and G. S. ROEDER, 1998 Synaptonemal complex morphogenesis and sister-chromatid cohesion require Mek1-dependent phosphorylation of a meiotic chromosomal protein. *Genes Dev.* **12**: 3551–3563.
- BURNS, N., B. GRIMWADE, P. B. ROSS-MACDONALD, E.-Y. CHOI, K. FINBERG *et al.*, 1994 Large-scale analysis of gene expression, protein localization and gene disruption in *Saccharomyces cerevisiae*. *Genes Dev.* **8**: 1087–1105.
- CHRISTODOULIDOU, A., P. BRIZA, A. ELLINGER and V. BOURIOTIS, 1999 Yeast ascospore wall assembly requires two chitin deacetylase isozymes. *FEBS Lett.* **460**: 275–279.
- CHU, S., J. DERISI, M. EISEN, J. MULHOLLAND, D. BOTSTEIN *et al.*, 1998 The transcriptional program of sporulation in budding yeast. *Science* **282**: 699–703.
- CHUA, P. R., and G. S. ROEDER, 1998 Zip2, a meiosis-specific protein required for the initiation of chromosome synapsis. *Cell* **93**: 349–359.
- COLUCCIO, A., E. BOGENGRUBER, M. N. CONRAD, M. E. DRESSER, P. BRIZA *et al.*, 2004 Morphogenetic pathway of spore wall assembly in *Saccharomyces cerevisiae*. *Eukaryot. Cell* **3**: 1464–1475.
- DU, L., and P. NOVICK, 2002 Pag1p, a novel protein associated with protein kinase Cbk1p, is required for cell morphogenesis and proliferation in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* **13**: 503–514.
- ENYENIHI, A. H., and W. S. SAUNDERS, 2003 Large-scale functional genomic analysis of sporulation and meiosis in *Saccharomyces cerevisiae*. *Genetics* **163**: 47–54.
- ESPOSITO, R. E., and M. S. ESPOSITO, 1974 Genetic recombination and commitment to meiosis in *Saccharomyces*. *Proc. Natl. Acad. Sci. USA* **71**: 3172–3176.
- FRIESEN, H., R. LUNZ, S. DOYLE and J. SEGALL, 1994 Mutation of the *SPS1*-encoded protein kinase of *Saccharomyces cerevisiae* leads to defects in transcription and morphology during spore formation. *Genes Dev.* **8**: 2162–2175.
- GERST, J. E., 1999 SNAREs and SNARE regulators in membrane fusion and exocytosis. *Cell. Mol. Life Sci.* **55**: 707–734.
- HONG, E.-J. E., and G. S. ROEDER, 2002 A role for Ddc1 in signaling meiotic double-strand breaks at the pachytene checkpoint. *Genes Dev.* **16**: 363–376.
- JAHN, R., and T. C. SUDHOF, 1999 Membrane fusion and exocytosis. *Annu. Rev. Biochem.* **68**: 863–911.
- JANTTI, J., M. K. AALTO, M. OYEN, L. SUNDQVIST, S. KERANEN *et al.*, 2002 Characterization of temperature-sensitive mutations in the yeast syntaxin 1 homologues Sso1p and Sso2p, and evidence of a distinct function for Sso1p in sporulation. *J. Cell Sci.* **115**: 409–420.
- KRISAK, L., R. STRICH, R. S. WINTERS, J. P. HALL, M. J. MALLORY *et al.*, 1994 *SMK1*, a developmentally regulated MAP kinase, is required for spore wall assembly in *Saccharomyces cerevisiae*. *Genes Dev.* **8**: 2151–2161.
- MORENO-BORCHART, A. C., K. STRASSER, M. G. FINKBEINER, A. SHEVCHENKO and M. KNOP, 2001 Prospore membrane formation linked to the leading edge protein (LEP) coat assembly. *EMBO J.* **20**: 6946–6957.
- NAG, D. K., and J. AXELROD, 1998 Identification of yeast meiosis-specific genes by differential display. *Methods* **16**: 423–433.
- NAKANISHI, H., M. MORISHITA, C. L. SCHWARTZ, A. COLUCCIO, J. ENGBRECHT *et al.*, 2006 Phospholipase D and the SNARE Sso1p are necessary for vesicle fusion during sporulation in yeast. *J. Cell Sci.* **119**: 1406–1415.
- NEIMAN, A. M., 1998 Prospore membrane formation defines a developmentally regulated branch of the secretory pathway in yeast. *J. Cell Biol.* **140**: 29–37.
- NEIMAN, A. M., 2005 Ascospore formation in the yeast *Saccharomyces cerevisiae*. *Microbiol. Mol. Biol. Rev.* **69**: 565–584.
- NICKAS, M. E., and A. M. NEIMAN, 2002 *Ady3p* links spindle pole body function to spore wall synthesis in *Saccharomyces cerevisiae*. *Genetics* **160**: 1439–1450.
- RABITSCH, K. P., A. TOTH, M. GALOVA, A. SCHLEIFFER, G. SCHAFFNER *et al.*, 2001 A screen for genes required for meiosis and spore formation based on whole-genome expression. *Curr. Biol.* **11**: 1001–1009.
- RIEDEL, C. G., M. MAZZA, P. MAIER, R. KORNER and M. KNOP, 2005 Differential requirement for phospholipase D/Spo14 and its novel interactor Smal for regulation of exocytotic vesicle fusion in yeast meiosis. *J. Biol. Chem.* **280**: 37846–37852.
- ROCKMILL, B., and G. S. ROEDER, 1990 Meiosis in asynaptic yeast. *Genetics* **126**: 563–574.
- ROSE, K., S. A. RUDGE, M. A. FROHMAN, A. J. MORRIS and J. ENGBRECHT, 1995 Phospholipase D signaling is essential for meiosis. *Proc. Natl. Acad. Sci. USA* **92**: 12151–12155.
- ROSS-MACDONALD, P., P. S. COELHO, T. ROEMER, S. AGARWAL, A. KUMAR *et al.*, 1999 Large-scale analysis of the yeast genome by transposon tagging and gene disruption. *Nature* **402**: 413–418.
- RUDGE, S. A., A. J. MORRIS and J. ENGBRECHT, 1998 Relocalization of phospholipase D activity mediates membrane formation during meiosis. *J. Cell Biol.* **140**: 81–90.
- RUOHONEN, L., J. TOIKKANEN, V. TIEAHO, M. OUTOLA, H. SODERLUND *et al.*, 1997 Enhancement of protein secretion in *Saccharomyces cerevisiae* by overproduction of Sso protein, a late-acting component of the secretory machinery. *Yeast* **13**: 337–351.
- SAMBROOK, J., E. F. FRITSCH and T. MANIATIS, 1989 *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- SARKAR, P. K., M. A. FLORCZYK, K. A. McDONOUGH and D. K. NAG, 2002 *SSP2*, a sporulation-specific gene necessary for outer spore wall assembly in the yeast *Saccharomyces cerevisiae*. *Mol. Genet. Genomics* **267**: 348–358.
- SEIFERT, H. S., E. Y. CHEN, M. SO and F. HEFFRON, 1986 Shuttle mutagenesis: a method of transposon mutagenesis for *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **83**: 735–739.
- SHERMAN, F., G. R. FINK and J. B. HICKS, 1986 *Methods in Yeast Genetics: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- SHONN, M. A., R. MCCARROLL and A. W. MURRAY, 2000 Requirement of the spindle checkpoint for proper chromosome segregation in budding yeast meiosis. *Science* **289**: 300–303.
- SIKORSKI, R., and P. HIETER, 1989 A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**: 19–27.
- STRAIGHT, A. F., A. S. BELMONT, C. C. ROBINETT and A. W. MURRAY, 1996 GFP tagging of budding yeast chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion. *Curr. Biol.* **6**: 1599–1608.
- STRAIGHT, P. D., T. H. GIDDINGS and M. WINEY, 2000 *Mps1p* regulates meiotic spindle pole body duplication in addition to having novel roles during sporulation. *Mol. Biol. Cell* **11**: 3525–3537.
- SYM, M., J. ENGBRECHT and G. S. ROEDER, 1993 Zip1 is a synaptonemal complex protein required for meiotic chromosome synapsis. *Cell* **72**: 365–378.
- TACHIKAWA, H., A. BLOECHER, K. TATCHELL and A. M. NEIMAN, 2001 A Gip1p-Glc7p phosphatase complex regulates septin organization and spore wall formation. *J. Cell Biol.* **155**: 797–808.

TSUBOUCHI, H., and G. S. ROEDER, 2002 The Mnd1 protein forms a complex with Hop2 to promote homologous chromosome pairing and meiotic double-strand break repair. *Mol. Cell. Biol.* **22**: 3078–3088.

WAGNER, M., P. BRIZA, M. PIERCE and E. WINTER, 1999 Distinct steps in yeast spore morphogenesis require distinct *SMK1* MAP kinase thresholds. *Genetics* **151**: 1327–1340.

WAGNER, M., M. PIERCE and E. WINTER, 1997 The CDK-activating kinase *CAK1* can dosage suppress sporulation defects of *smk1* MAP kinase mutants and is required for spore wall morphogenesis in *Saccharomyces cerevisiae*. *EMBO J.* **16**: 1305–1317.

Communicating editor: R. S. HAWLEY