

## Vitamin B<sub>12</sub> deficiency in a breast fed infant

A J McPHEE,\* G P DAVIDSON,† M LEAHY,‡ AND T BEARE‡

\*Department of Paediatrics, Queen Victoria Hospital, Rose Park, South Australia and Departments of †Gastroenterology and ‡Paediatrics, Adelaide Children's Hospital, North Adelaide, South Australia

**SUMMARY** We report the case of a 5 month old breast fed infant who presented with a history of vomiting, pallor, and failure to thrive. Investigations showed severe nutritional vitamin B<sub>12</sub> deficiency with a megaloblastic pancytopenia. This deficiency was due to low vitamin B<sub>12</sub> concentrations in the maternal breast milk, and subsequent investigations showed maternal pernicious anaemia. Treatment of the infant with vitamin B<sub>12</sub> resulted in a rapid clinical and haematological improvement. This case represents an unusual presentation of pernicious anaemia.

Dietary vitamin B<sub>12</sub> deficiency in infancy is rare, and most reported cases are breast fed infants of mothers who themselves are deficient in vitamin B<sub>12</sub>, usually on the basis of deficient (particularly vegetarian) diets.<sup>1-4</sup> The development of haematologic, neurologic, and metabolic abnormalities in the breast fed offspring of these mothers is usually the presenting feature of the maternal deficiency, which itself may be mild. The case reported here is that of a breast fed infant whose vitamin B<sub>12</sub> deficiency presented as vomiting, failure to thrive, and megaloblastic pancytopenia at 5-6 months of age. Although maternal serum vitamin B<sub>12</sub> concentrations and blood film were normal, concentrations of vitamin B<sub>12</sub> in the breast milk were very low. Subsequent studies showed maternal vitamin B<sub>12</sub> deficiency due to subclinical pernicious anaemia. Three similar cases, two of whom had prominent neurologic abnormalities, have been reported.<sup>5-7</sup>

### Case report

A girl, the first born infant of healthy unrelated parents (birth weight 3080 g), was exclusively breast fed and maintained the 25-50th weight percentile until 3-4 months of age when vomiting became a problem. This did not respond to the introduction of solids and in the three months before admission she gained only 250 g. Pallor was first noted by her local physician at 5 months of age, and anaemia was diagnosed (haemoglobin concentration 74 g/l). Treatment with oral iron was without effect, and at the time of her referral to the Adelaide Children's Hospital a complete blood count showed pancytopenia with a haemoglobin of 70 g/l, white

cell count  $7.5 \times 10^9/l$ , absolute neutrophil count  $530 \times 10^9/l$ , and platelets  $55 \times 10^9/l$ .

On admission, she was found to be a pale placid infant, with a weight of 5640 g (<3rd percentile), length 66 cm (10th-25th percentile), and head circumference 43 cm (10th percentile). There was no lymphadenopathy and no organomegaly. Neurodevelopmental assessment was consistent with postnatal age, and maternal dietary history was normal. Subsequent investigations of mother and infant are shown in the table.

Table Results of investigations on infant and mother

Investigation	Infant	Mother
Haemoglobin (g/l)	52	136
White cell count ( $\times 10^9/l$ )	3.1	6.6
Absolute neutrophil count ( $\times 10^6/l$ )	400	4550
Platelet count ( $\times 10^9/l$ )	46	>200
Serum vitamin B <sub>12</sub> (pmol/l)*	<37	273
Serum folate (nmol/l)†	49	9.8
Breast milk vitamin B <sub>12</sub> (pmol/l)‡	—	44.3
Schilling test	Normal	Abnormal
Barium meal and follow through	Normal	Not tested
Gastric parietal cell antibody	Negative	Positive +
Thyroid cytoplasmic antibody	Not tested	Positive ++
Intrinsic factor antibody	Negative	Positive +++
Free thyroxine index§	Not tested	55
Thyroid stimulating hormone (IU/l)	Not tested	126

\*Serum vitamin B<sub>12</sub> normal range=221-885 pmol/l.

†Serum folate normal range=6.8-47.7 nmol/l.

‡Breast milk vitamin B<sub>12</sub> normal range=207-1549.

§Free thyroxine index normal range 75-150.

||Thyroid stimulating hormone normal result <10 IU/l.

The infant's blood film showed appreciable poikilocytosis and anisocytosis with macrocytic changes. The reticulocyte count was 1.4% and there were two nucleated red cells per 100 white cells. Occasional myelocytes and metamyelocytes were seen. A bone marrow biopsy specimen showed appreciable megakaryoblastic changes with considerable disparity between nuclear and cytoplasmic development in all cell lines, and there was a myeloid: erythroid ratio of 2:1. The low serum vitamin B<sub>12</sub> concentrations prompted further studies in the infant, including a Schilling test and gastric acid secretion studies all of which gave normal results.

Breast milk vitamin B<sub>12</sub> concentrations were determined using the same competitive protein binding assay used to determine serum vitamin B<sub>12</sub> concentrations. Although the validity of this assay system for breast milk B<sub>12</sub> was not determined, the concentration of 44.3 pmol/l was considerably lower than concentrations determined in our laboratories, using the same assay system, on breast milk samples from healthy lactating women (206.6–1549.4 pmol/l).<sup>4</sup> Also, the concentration of 44.3 pmol/l is low compared with other published normal ranges of breast milk vitamin B<sub>12</sub> determined by other methods.<sup>8,9</sup> Overall, the low breast milk vitamin B<sub>12</sub> concentrations suggested subclinical maternal vitamin B<sub>12</sub> deficiency, and subsequent investigations showed classical (Addisonian) pernicious anaemia and hypothyroidism.

The infant was transfused and vitamin B<sub>12</sub> (250 µg) was administered as part of her Schilling test. Serial blood counts showed a rapid improvement in neutrophil and platelet counts (see figure). Vomiting stopped coincident with the administration of vitamin B<sub>12</sub>, and a considerable 'character' change, manifested by increased activity and responsiveness, was noted in the child by both parents and hospital staff. The mother was treated with thyroxine and parenteral vitamin B<sub>12</sub>. Follow up of the infant at 12 and 18 months showed a clinically normal child with




Figure Neutrophil and platelet counts before and after treatment with 250 µg vitamin B<sub>12</sub>.

normal haematology and serum vitamin B<sub>12</sub> concentrations. Now, at 8 years of age she is functioning normally in an age appropriate school setting.

## Discussion

The estimated vitamin B<sub>12</sub> requirements of the growing infant are 0.06–0.10 µg/day and the normal neonatal vitamin B<sub>12</sub> stores are of the order of 20–25 µg.<sup>10</sup> Therefore the normal newborn infant has sufficient vitamin B<sub>12</sub> stores to last for six to eight months, even in the presence of inadequate dietary intake or defective vitamin B<sub>12</sub> absorption.<sup>10</sup> On the other hand, the vitamin B<sub>12</sub> stores of the infant of a deficient mother may be as low as 2–5 µg<sup>10</sup> and although normal breast milk has considerable vitamin B<sub>12</sub>,<sup>8,9</sup> the vitamin B<sub>12</sub> content of the breast milk of deficient mothers is low, as shown by this and other cases.<sup>1–7</sup> Overall, the vitamin B<sub>12</sub> state of the breast fed infant of a vitamin B<sub>12</sub> deficient mother is precarious, with marginal stores being aggravated by inadequate dietary intake.

In the present case the maternal complete blood picture and serum vitamin B<sub>12</sub> concentration were normal. The low breast milk vitamin B<sub>12</sub> concentration was the only clue to the mother's aberrant vitamin B<sub>12</sub> state. A similar situation has been described in one other case.<sup>2</sup>

Based on the estimated daily requirements presented above, adequate vitamin B<sub>12</sub> intake with a breast milk B<sub>12</sub> concentration of 44.3 pmol/l would require an intake of 1–1½ litres of milk per day. While this would appear achievable, we speculate that the onset of vomiting at 3 months of age in the present case was critical in limiting vitamin B<sub>12</sub> deficiency. Also, as discussed below, the vomiting itself may have been caused by vitamin B<sub>12</sub> deficiency. Alternatively, because our normal range for breast milk vitamin B<sub>12</sub> is considerably higher than other published normal ranges,<sup>8,9</sup> it is possible that our assay method may have overestimated breast milk vitamin B<sub>12</sub> and that inadequate intake was present from an early age.

The clinical features of vitamin B<sub>12</sub> deficiency in infancy are predominantly neurologic and haematologic. Neurologic features include an acquired movement disorder, developmental regression, torpor, and even coma.<sup>1–4,6,7</sup> The appreciable character change noted in our patient after the administration of vitamin B<sub>12</sub> suggests that her placidity on presentation was an early neurologic feature. Other case reports,<sup>6,7</sup> including one from this institution,<sup>4</sup> have reported long term developmental and neurologic sequelae of vitamin B<sub>12</sub> deficiency in infancy. In general, such sequelae seem to be associated with profound neurologic abnormalities at the time of

presentation and presumably the absence of such abnormalities in the present case may explain the good long term outcome.

The haematologic features of vitamin B<sub>12</sub> deficiency—namely, a megaloblastic pancytopenia—were well illustrated in our case. Other reported features such as mild hepatosplenomegaly, diarrhoea, and a curious palmar pigmentation appear variable, and were not seen in our patient. Vomiting has not been reported previously in association with vitamin B<sub>12</sub> deficiency in infancy, but was the presenting complaint in our case. The absence of any structural cause, and the prompt resolution of vomiting coincident with the administration of vitamin B<sub>12</sub> suggest that the vomiting was a symptom of the vitamin B<sub>12</sub> deficiency.

Three other cases of occult maternal pernicious anaemia presenting as symptomatic vitamin B<sub>12</sub> deficiency in a breast fed child have been reported.<sup>5-7</sup> Haematologic features were prominent in the case described by Lampkin *et al.*,<sup>5</sup> while developmental regression was the presenting complaint in the case of Sadowitz *et al.*,<sup>6</sup> and obtundation with hypothermia was seen in the case of Johnson and Roloff.<sup>7</sup>

In summary, a case of vitamin B<sub>12</sub> deficiency in the breast fed infant of a mother with occult pernicious anaemia is presented. Vomiting and a megaloblastic pancytopenia were features of the infant's presentation. A low breast milk vitamin B<sub>12</sub> concentration was the only clue to the maternal deficiency. This case serves to emphasise that vitamin B<sub>12</sub> deficiency presenting at less than 6 months of age is almost exclusively seen in breast fed infants of vitamin B<sub>12</sub> deficient mothers. In the absence of a deficient maternal diet (particularly a

strict vegetarian diet), occult pernicious anaemia should be considered as the reason for the maternal deficiency.

#### References

- <sup>1</sup> Jadhav M, Webb JKG, Vaishnava S, Baker SJ. Vitamin B<sub>12</sub> deficiency in Indian infants. A clinical syndrome. *Lancet* 1962;ii:903-7.
- <sup>2</sup> Lampkin BC, Saunders EF. Nutritional vitamin B<sub>12</sub> deficiency in an infant. *J Pediatr* 1969;75:1053-5.
- <sup>3</sup> Higginbottom MC, Sweetman L, Nyhan WL. A syndrome of methylmalonic aciduria, homocystinuria, megaloblastic anemia and neurologic abnormalities in a vitamin B<sub>12</sub> deficient breast fed infant of a strict vegetarian. *N Engl J Med* 1978;229:317-23.
- <sup>4</sup> Wighton MC, Manson JI, Speed I, Robertson E, Chapman E. Brain damage in infancy and dietary vitamin B<sub>12</sub> deficiency. *Med J Aust* 1979;2:1-3.
- <sup>5</sup> Lampkin BC, Shore NA, Chadwick D. Megaloblastic anemia of infancy secondary to maternal pernicious anemia. *N Engl J Med* 1966;274:1168-71.
- <sup>6</sup> Sadowitz PD, Livingston A, Cavanaugh RM. Developmental regression as an early manifestation of vitamin B<sub>12</sub> deficiency. *Clin Pediatr (Phila)* 1986;25:369-71.
- <sup>7</sup> Johnson PR, Roloff JS. Vitamin B<sub>12</sub> deficiency in an infant strictly breast-fed by a mother with latent pernicious anaemia. *J Pediatr* 1982;100:917-9.
- <sup>8</sup> Sneed SM, Zane C, Thomas MR. The effects of ascorbic acid, vitamin B<sub>6</sub>, vitamin B<sub>12</sub> and folic acid supplementation on the breast milk and maternal nutritional status of low socioeconomic lactating women. *Am J Clin Nutr* 1981;34:1338-46.
- <sup>9</sup> Bijur AM, Desai AG. Composition of breast milk with reference to vitamin B<sub>12</sub> and folic acid in Indian mothers. *Indian J Pediatr* 1985;52:147-50.
- <sup>10</sup> Willoughby MLN. Vitamin B<sub>12</sub> metabolism and deficiency. In: Willoughby MLN, ed. *Paediatric haematology*. Edinburgh: Churchill Livingstone, 1977:35-42.

Correspondence to Dr G P Davidson, Gastroenterology Department, Adelaide Children's Hospital, North Adelaide, South Australia 5006, Australia.

Accepted 22 February 1988