

Research

Open Access

Comparison of plants used for skin and stomach problems in Trinidad and Tobago with Asian ethnomedicine

Cheryl Lans*

Address: BCICS, University of Victoria, British Columbia, V8W 2Y2, Canada

Email: Cheryl Lans* - cher2lans@netscape.net

* Corresponding author

Published: 05 January 2007

Received: 16 October 2006

Accepted: 05 January 2007

Journal of Ethnobiology and Ethnomedicine 2007, 3:3 doi:10.1186/1746-4269-3-3

This article is available from: <http://www.ethnobiomed.com/content/3/1/3>

© 2007 Lans; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This paper provides a preliminary evaluation of fifty-eight ethnomedicinal plants used in Trinidad and Tobago for skin problems, stomach problems, pain and internal parasites for safety and possible efficacy. Thirty respondents, ten of whom were male were interviewed from September 1996 to September 2000 on medicinal plant use for health problems. The respondents were obtained by snowball sampling, and were found in thirteen different sites, 12 in Trinidad and one in Tobago. The uses are compared to those current in Asia. *Bambusa vulgaris*, *Bidens alba*, *Jatropha curcas*, *Neurolaena lobata*, *Peperomia rotundifolia* and *Phyllanthus urinaria* are possibly efficacious for stomach problems, pain and internal parasites. Further scientific study of these plants is warranted.

Background

Trinidad and Tobago is one country consisting of two adjacent islands located just northeast of the Venezuelan coast with a combined area of 5070 km² [1]. The human population of 1.25 million is multi-ethnic, multi-religious and multicultural and increases at 1% annually. In Trinidad, the major population centres are concentrated along the west coast and along an east-west transportation corridor in the north of the island [1].

The multi-ethnic population of Trinidad and Tobago is reflected in its folk medicinal use. Previous research has indicated that the folk medicines used by hunters are derived from ancient Amerindian practices [2]. This paper will continue to explore the cultural origins of Caribbean folk medicine by investigating the contribution of the Chinese to Caribbean folk medicine. Chinese medicine has been described as a complex and holistic system of medical practice with its own philosophy, diagnosis, treatment systems and pharmacology which also includes acu-

puncture, moxibustion and *Qi Gong*. However in this paper I will focus on 'Ben Cao' (Herbalism) [3].

The Chinese were the first Asian immigrants, arriving before the original East Indians who arrived in 1845. Chinese Tartars (192 men and one woman) were brought to Trinidad in the fall of 1806. These men from Macao, Penang and Canton were brought to cultivate tea but most were dissatisfied with local conditions and returned on the same ship [4,5]. The twenty-three who stayed made a living as entrepreneurs (butchers, shopkeepers, carpenters and market gardeners) and creolised (integrated into the local population).

Prominent sugarcane planters believed that the emancipation of Caribbean slaves in 1838 would create a labour shortage. In the 1840s, the British "opened" a labor market of displaced or impoverished peasantry in southern China to fill this shortage and 2,500 mainly-male Chinese were brought legitimately to Trinidad as indentured work-

ers, or were 'shanghaied' (abducted by European traders) [6]. After the first Opium War (1840–42), and second Opium War, the British (as well as French and Americans) occupied twelve major ports (and colonized Hong Kong) [6]. China's defeats in the Opium Wars led to the deregulation of Chinese immigration. This combined with the unrest, rebellion, and war in China, facilitated the organized labour traffic of one million southern Chinese to the West from 1840 to 1875 [6].

Three vessels brought 1,100 Chinese indentured labourers to Trinidad in 1853 and 600 more came in 1865 and 1866. In 1862, 467 immigrants came from Hong Kong. Most of the immigrants arriving between 1853 and 1866 came from the southern Guangdong province (Macao, Hong Kong and Canton). In the last 5 trips, a total of 2837 emigrants came from Macao, Amoy, Canton and Hong Kong. Chinese migration after 1911 was driven by the Chinese revolution. Punti traders described Hakka prisoners as pigs on the bills of lading and shipped them to the Caribbean and South America [4,5]. Between 1920s and 1940s new immigrants consisted of the families and friends of earlier migrants. They came as merchants, peddlers, traders and shopkeepers, not indentured labour [4]. Almost 9,000 more Chinese immigrants came voluntarily from British Guyana to Trinidad over the next century, after having served their indenture ship [5]. Chinese people now constitute approximately 1% of the Trinidad and Tobago population as an ethnic group but are also present in the large mixed-raced population of 18 – 25%.

There is one publication that describes the use of medicinal plants by the Chinese community in Trinidad [7]; it contained no plants in common with those in this research [1]. Nevertheless in the discussion section of this paper, comparisons will be made of the uses of the plants in Trinidad and Tobago and those current in Asia and South-east Asia. The ethnomedicinal literature available from Asia will be used in the non-experimental validation.

Fifty-eight plants used in ethnomedicine in Trinidad and Tobago for skin problems, stomach problems, pain and internal parasites are described in this paper and a non-experimental validation of them is presented. The recent publication of high-quality studies and clinical trials on the ethnomedicinal plants in this paper has enhanced the non-experimental validation of the plants presented in the discussion section.

Methods

Study design

This study adhered to the research guidelines and ethical protocols of Wageningen University in the Netherlands. Thirty respondents, ten of whom were male were inter-

viewed from September 1996 to September 2000. The respondents were obtained by snowball sampling, and were found in thirteen different sites, 12 in Trinidad and one in Tobago. Snowball sampling was used because there was no other means of identifying respondents. The chief objective of the sampling method was to identify knowledgeable respondents.

Twenty respondents were interviewed once, the other ten (who were healers) were interviewed three or four times. Healers were also asked to reconstruct the circumstances and contexts of the plant uses so that the means of administration of the plants could be identified. No interview schedule of questions was used but a more qualitative, conversational technique. Plants were collected when available to verify that the common names used by each respondent were the same in each ethnic group as those recorded in the literature. The majority of the plants were identified at the Herbarium of the University of the West Indies but voucher samples were not deposited. This ethnomedicinal study was part of a larger research project on ethnoveterinary medicine; other data collecting techniques were used in the larger study [1].

Non-experimental validation

The ethnomedicinal plants used in Trinidad and Tobago for skin problems, stomach problems, pain and internal parasites are presented in Tables 1 and 2.

The plant-based remedies were evaluated for safety and efficacy with a non-experimental method. Published sources such as journal articles and books and databases on pharmacology and ethnomedicine available on the Internet were searched to identify the plants' chemical compounds and clinically tested physiological effects. This data was incorporated with data on the reported folk uses, and their preparation and administration in Latin America, the Caribbean, Asia and Africa. For each species or genus the ethnomedicinal uses in other countries are given if available; then follows a summary of chemical constituents, in addition to active compounds if relevant (Tables 3 and 4). This type of ethnopharmacological review and evaluation has been previously published [2]. The plant uses in China are then given (Table 5) and a comparison of the uses in Trinidad and China is made in the discussion.

Results

The ethnomedicinal plants used in Trinidad and Tobago for skin problems, stomach problems, pain and internal parasites are presented in Tables 1 and 2.

Plants used for skin problems

Twelve plants are used for skin problems including one for the rash caused by measles plus one for shingles. The

Table 1: Ethnomedicinal plants used for skin problems in Trinidad and Tobago

	Scientific name	Family	Common name	Plant part used	Use
1	<i>Achyranthes indica</i>	Amaranthaceae	Man better man		Skin problems
2	<i>Acnistus arborescens</i>	Solanaceae	Wild tobacco	Leaves	Bathe babies for eczema
3	<i>Azadirachta indica</i>	Meliaceae	Neem	Leaves	Measles
4	<i>Bidens alba/Bidens pilosa</i>	Asteraceae	Needle grass/Railway daisy	Leafy branch	Bathe children
5	<i>Cassia alata</i>	Fabaceae- Caesalpiniaceae	Senna	Leaves	Skin problems
6	<i>Chamaesyce hirta/hypericifolia</i>	Euphorbiaceae	Malomay	Flower	Skin rashes, measles
7	<i>Croton gossypifolius</i>	Euphorbiaceae	Blood bush/Bois sang	Leaves	Bathe babies for eczema
8	<i>Eclipta prostrata</i>	Asteraceae	Congolala		Bathe for children's malnutrition for 9 days & woodlice nest
9	<i>Manihot esculenta</i>	Euphorbiaceae	Cassava	Leaves	Bathe babies for eczema
10	<i>Origanum vulgare</i>	Lamiaceae	Majoram		Bathe babies
11	<i>Sida carpinifolia</i> (syn. <i>Sida acuta</i>)	Malvaceae	Garaba broom	Leaf	Eczema
12	<i>Solanum americanum</i>	Solanaceae	Agouma, gouma	Plant	Bathe for children's malnutrition
13	<i>Spondias mombin</i>	Anacardiaceae	Hogplum	Leaves	Eczema

majority of the plants were being used for children including babies. The thirteen plants belong to nine plant families. Eight plants are used to bathe babies. *Acnistus arborescens*, *Croton gossypifolius* and *Manihot esculenta* are used to bathe babies for eczema. *Bidens alba/Bidens pilosa* and *Origanum vulgare* are used to bathe babies and older children. *Eclipta prostrata* is combined with a non-plant material and used to bathe children for malnutrition. *Solanum americanum* is also used to bathe children for malnutrition. *Azadirachta indica* and *Chamaesyce hirta/hypericifolia* are used for measles. *Sida carpinifolia* (syn. *Sida acuta*) and *Spondias mombin* are used for eczema. *Achyranthes indica*, *Cassia alata* and *Chamaesyce hirta/hypericifolia* are used for skin rashes and other skin problems.

Plants used for stomach problems, pain, internal parasites

The medicinal plants used for stomach problems, injuries, endoparasites, arthritis and bites are combined in Table 2. This grouping partially reflects the analgesic activity of many of the plants used. Eighteen plants are used for stomach problems including diarrhoea. Another fifteen plants are used for various kinds of pain including cuts, bites, sprains and arthritis. Four plants are used as anthelmintics. Other plants in the table are used for dropsy. Twenty-seven plant families are represented in Table 2.

The following plants are used as carminatives: *Cecropia peltata*, *Aframomum melegueta*, *Ferula asafoetida* and *Tournefortia hirsutissima*.

The following plants are used for stomach problems: *Ambrosia cumanensis*, *Aristolochia rugosa/trilobata*, *Capraria biflora*, *Dorstenia contrajerva*, *Cajanus cajan*, *Momordica charantia*, *Punica granatum*, *Brownea latifolia* and *Cocos nucifera*.

Diarrhoea is treated with the following plants: *Chamaesyce hirta*, *Eleusine indica*, *Peperomia rotundifolia*, *Phyllanthus urinaria* and *Scoparia dulcis*.

The plants used as anthelmintics are *Citharexylum spinosum*, *Cucurbita maxima*, *Portulaca oleraceae*, *Tagetes patula* and *Eupatorium triplinerve*.

Plants used specifically for pain are: *Brownea latifolia*, *Abelmoschus moschatus*, *Eupatorium macrophyllum*, *Morinda citrifolia* and *Cola nitida*.

Arthritis is treated with the following plants: *Nicotiana tabacum*, *Petiveria alliacea*, *Rosmarinus officinalis* and *Neuro-laena lobata*.

Plants used for cuts, injuries and swellings are: *Solanum melongena*, *Jatropha curcas/gossypifolia*, *Bidens alba/Bidens pilosa*, *Cucurbita pepo*, *Tournefortia hirsutissima*, *Bambusa vulgaris*, *Bixa orellana* and *Cocos nucifera*.

Scorpion and snake bites are treated with *Tamarindus indica*, *Nopalea cochinellifera*, *Centropogon cornutus* and *Rosmarinus officinalis*.

Table 2: Plants used for stomach problems, pain and internal parasites in Trinidad and Tobago

	Scientific name	Family	Common name	Part used	Use
1.	<i>Abelmoschus moschatus</i>	Malvaceae	Gumbo musque	Seeds	Grind in rum for foot cramp
2.	<i>Aframomum melegueta</i>	Zingiberaceae	Guinea pepper	Seeds	Carminative
3.	<i>Ambrosia cumanenesis</i>	Asteraceae	Altamis	Bark	Stomach pain, 2*3 inch piece bark in urine for 3 days use to wash foot for 3 days for arthritis
4.	<i>Aristolochia rugosa, trilobata</i>	Aristolochiaceae	Mat root, anico	Root	Stomach pain, colic, poisoning
5.	<i>Bambusa vulgaris</i>	Poaceae	Bamboo	Leaves	Poultice
6.	<i>Bidens alba/Bidens pilosa</i>	Asteraceae	Needle grass	Leafy branch	Cuts
7.	<i>Bixa orellana</i>	Bixaceae	Roucou	Root	Dropsy
8.	<i>Brownea latifolia</i>	Fabaceae	Cooper hoop	Flower, leaves	Gripe, pain
9.	<i>Cajanus cajan</i>	Fabaceae	Pigeon pea	Leaves	Food poisoning, colic, constipation
10.	<i>Capraria biflora</i>	Scrophulariaceae	Du thé pays	Leaves	Flavour for purgative
11.	<i>Cecropia peltata</i>	Cecropiaceae	Bois canôt	Stem	3 'Ridges' from inside stem boiled as a carminative
12.	<i>Centropogon cornutus</i>	Campanulaceae	Deer meat, crepe coq	Leaves	Snake, scorpion bite
13.	<i>Chamaesyce hirta</i>	Euphorbiaceae	Malomay		Diarrhoea
14.	<i>Citharexylum spinosum</i>	Verbenaceae	Bois côtelette	Leaf	Anthelmintic
15.	<i>Cocos nucifera</i>	Arecaceae	Coconut	Root- 7 inches, Shell	Dropsy, Hernia
16.	<i>Cola nitida</i>	Sterculiaceae	Obie seed	Seed	Any kind of pain
17.	<i>Cucurbita maxima</i>	Cucurbitaceae	Pumpkin	Seeds	Anthelmintic
18.	<i>Cucurbita pepo</i>	Cucurbitaceae	Pumpkin		Sprains, breaks
19.	<i>Dorstenia contrajerva</i>	Moraceae	Refriyau		Food poisoning
20.	<i>Eleusine indica</i>	Poaceae	Pied poule		Diarrhoea
21.	<i>Eupatorium macrophyllum</i>	Asteraceae	Z'herbe chatte		Pain
22.	<i>Eupatorium triplinerve</i>	Asteraceae	Ayapana, japanne	Leaves	Stomach problems (worms)
23.	<i>Ferula asafoetida</i>	Apiaceae	Asafoetida		Carminative
24.	<i>Jatropha curcas/gossypifolia</i>	Euphorbiaceae	White/Red Physic Nut	Leaf	Clean sores
25.	<i>Momordica charantia</i>	Cucurbitaceae	Caraaili	Vine	Stomach problems
26.	<i>Morinda citrifolia</i>	Rubiaceae	Noni	Leaves	Pains
27.	<i>Neurolaena lobata</i>	Asteraceae	Z'herbe á pique	Leaves	Tincture for arthritis
28.	<i>Nicotiana tabacum</i>	Solanaceae	Tobacco	Leaves	Arthritis
29.	<i>Nopalea cochinellifera</i>	Cactaceae	Rachette	Joint	Snake bites
30.	<i>Peperomia rotundifolia</i>	Piperaceae	Mowon		Diarrhoea
31.	<i>Petiveria alliacea</i>	Phytolaccaceae	Mapourite		Arthritis and rheumatism
32.	<i>Phyllanthus urinaria</i>	Euphorbiaceae	Red seed under leaf	Plant	Diarrhoea
33.	<i>Portulaca oleraceae</i>	Portulacaceae	Pussley	Plant	Anthelmintic
34.	<i>Punica granatum</i>	Punicaceae	Pome-granate	Seeds	Stomach problems
35.	<i>Rosmarinus officinalis</i>	Lamiaceae	Rosemary	Leaf	Arthritis, Snake bites
36.	<i>Scoparia dulcis</i>	Scrophulariaceae	Sweet broom	Root	Diarrhoea
37.	<i>Solanum melongena</i>	Solanaceae	Melongene	Fruit	Breaks
38.	<i>Tagetes patula</i>	Asteraceae	Marigold		Anthelmintic
39.	<i>Tamarindus indica</i>	Fabaceae	Tamarind		Scorpion bite
40.	<i>Tournefortia hirsutissima</i>	Boraginaceae	Chigger bush	Leaves	Tea, carminative, chiggers

Table 3: Non-experimental validation of plants used for skin problems in Trinidad and Tobago

Scientific name	Validation	Reference
<i>Achyranthes aspera</i>	<i>Achyranthes bidentata</i> is a commonly used Chinese medicinal plant and is used in Nepal and in Mauritius and Rodrigues for skin diseases. <i>Achyranthes bidentata</i> polysaccharide can inhibit non-enzyme glycation in D-galactose induced mouse aging model <i>in vivo</i> . <i>Achyranthes aspera</i> leaf extract and the non-alkaloid fraction containing mainly non-polar compounds have chemo-preventive activity.	8–10
<i>Azadirachta indica</i>	A paste made of <i>Azadirachta indica</i> and <i>Curcuma longa</i> used to treat 814 people with scabies cured 97% of them within three to five days of treatment. <i>Azadirachta indica</i> (leaves, bark, fruit, flowers, oil, and gum) have the following properties: antimicrobial effects, <i>in vitro</i> antiviral activity, and antibacterial activity. Some active principles of <i>Azadirachta indica</i> are azadirachtin, salannin nimbin, and 6-desacetylnimbin. Clinical symptoms associated with toxocariasis in 1009 Trinidadian schoolchildren (aged 5–12 years) included eczema.	11–14
<i>Bidens pilosa</i>	<i>Bidens pilosa</i> is a commonly used traditional Chinese medicine. <i>Bidens pilosa</i> contains ethyl caffeate, a natural phenolic compound. Extracts of dried aerial parts of <i>Bidens pilosa</i> showed some antimicrobial activity as do components of the extract such as phenylheptatriene, linolic acid and linolenic acid. The triterpenes as well as several flavonoids (aurones, chalcones) are antiinflammatory agents. The chloroform fractions from the roots of <i>Bidens aurea</i> are anti-parasitical <i>in vitro</i> . The constituents of <i>Bidens pilosa</i> explain the use of this plant in traditional medicine in the treatment of wounds, against inflammations and against bacterial infections of the gastrointestinal tract.	15–17
<i>Cassia alata</i>	"Jue ming zi" (<i>Cassia tora</i> L. and <i>Cassia occidentalis</i> L.) has traditionally been used to improve visual acuity and to remove "heat" from the liver in Chinese medicine. Modern physicians use "Jue ming zi" to treat hypercholesterolemia and hypertension. "Jue ming zi" contains chrysophenol, emodin, and rhein. Roasted "Jue ming zi" is given as a health drink tea. The antioxidant activity of the methanolic extracts of "Jue ming zi" (<i>Cassia tora</i> L. and <i>Cassia occidentalis</i> L.) was established. <i>Cassia alata</i> is used for skin problems in the Caribbean, India, in traditional East Asian medicine and in the Ivory Coast (West Africa) to treat bacterial infections caused by <i>Escherichia coli</i> , and fungal infections caused by <i>Candida albicans</i> and dermatophytes. <i>Cassia alata</i> L. possesses anti-inflammatory, analgesic, laxative and antiplatelet aggregating activity and it contains kaempferol-3-O-gentiobioside. <i>Cassia alata</i> has antifungal activity that may be attributed to chrysophanol. When <i>Cassia alata</i> extracts were evaluated relative to a standard antibacterial agent chloramphenicol and antifungal agent amphotericin B the extracts had therapeutic potential for the treatment of opportunistic infections of AIDS patients. A 10-year human study indicated that a <i>Cassia alata</i> leaf extract can be reliably used as a folk medicine to treat <i>Pityriasis versicolor</i> . The leaf extract contains anthraquinones, flavonoids, quinones and sterols and had no side-effects.	18–21
<i>Chamaesyce hirta</i>	<i>Chamaesyce hirta</i> is used in West Bengal for ringworm. Antibacterial effects of <i>Chamaesyce hirta</i> leaves were found by several investigators. An aqueous extract of <i>Chamaesyce hirta</i> strongly reduced the release of prostaglandins I ₂ , E ₂ , and D ₂ . Additionally <i>Chamaesyce hirta</i> extracts exerted an inhibitory effect on platelet aggregation and depressed the formation of carrageenin-induced rat paw oedema.	22
<i>Croton gossypifolius</i>	<i>Croton cascarilloides</i> wood has been used historically to blacken teeth in Asia. <i>Croton cascarilloides</i> wood soot has limited antimicrobial activity against <i>Mutans streptococci</i> . <i>Croton</i> species are used in Thailand to treat dysmenorrhea, gastric ulcers, gastric cancers, and dysentery. <i>Croton kongensis</i> Gagnep., is known in Thailand as "Plao Ngeon" or "Plao Noi". A crude CH ₂ Cl ₂ extract of <i>Croton kongensis</i> showed antimalarial and antimycobacterial activities. <i>Croton sylvaticus</i> showed 5-lipoxygenase inhibitory activity with IC(50) values <61 ppm. A review of papers published in 2003, found that <i>in vitro</i> and <i>in vivo</i> studies supported the use of <i>Croton lechleri</i> Mull. Arg. for wounds, tumors, herpes infection, the itching, pain and swelling of insect bites and other conditions.	23–26
<i>Eclipta prostrata</i>	<i>Eclipta prostrata</i> is commonly used as self medication by AIDS patients in southern Thailand and showed potential as a therapeutic agent against <i>Giardia intestinalis</i> infections. The hydroalcoholic extract of <i>Eclipta prostrata</i> plant showed antinociceptive, immunomodulatory and antiinflammatory effects.	27
<i>Origanum vulgare</i>	<i>Origanum</i> volatile oil has potential efficacy against the infection of dysentery bacteria (<i>Shigella sonne</i> (Sh. sonnei) and <i>Shigella flexneri</i>). The carvacrol constituent has the most effective antimicrobial activity in <i>Origanum vulgare</i> . Diarrheic children in Trinidad were positive for <i>Shigella</i> (33 or 14.0%), 4 for <i>Salmonella</i> , and 1 for Enteropathogenic <i>E. coli</i> . Two fecal samples were positive for <i>Campylobacter jejuni</i> , and 1 was positive for hookworm ova.	28–30
<i>Sida acuta</i>	<i>Sida acuta</i> contains ecdysterone, ephedrine, hentriacontane, hypolaetin-8-glucoside, beta sitosterol, stigmaterol and campesterol. These chemicals may be responsible for the plant's reported narcotic analgesic, anti-inflammatory and analgesic activity.	31
<i>Solanum americanum</i>	<i>Solanum americanum</i> extracts were active against <i>Microsporium gypseum</i> and <i>Cryptococcus neoformans</i> and showed intra-peritoneal subacute toxicity in mice. Alpha-solamargine isolated from the fresh fruits of <i>Solanum americanum</i> is a glycoalkaloid with biological activity against <i>Herpes simplex I</i> , <i>Herpes zoster</i> and genital <i>Herpes</i> and <i>Trypanosoma cruzi</i> . <i>Solanum melongena</i> contains an anthocyanin, delphinidin, which inhibits the collagenolytic ability of matrix metalloproteinases.	32, 33

Table 4: Non-experimental validation of plants used for stomach problems, pain and internal parasites in Trinidad and Tobago

Scientific name	Validation	#
<i>Aframomum melegueta</i>	A decoction of the leaves of <i>Aframomum melegueta</i> is used for rheumatism and as an anti-emetic agent and a decoction of the fruits for dysenteric conditions. The methanol extracts of the seeds were significantly active against Gram (+) and Gram (-) bacteria (<i>S.aureus</i> , <i>B.subtilis</i> , <i>E.coli</i> , <i>P.aeruginosa</i>) and fungi (<i>C. albicans</i> , <i>A.niger</i>). <i>Aframomum melegueta</i> has antimicrobial properties against <i>E. coli</i> and <i>Bacillus cereus</i> . The antioxidant extracts of <i>Aframomum melegueta</i> was attributed to its phenolic components. Scabies and acute poststreptococcal glomerulonephritis (the latter can be caused by several bacterial and viral infections) are frequently associated with <i>S. aureus</i> in Trinidad.	34–36
<i>Ambrosia cumanensis</i>	The ambrosanolid-type sesquiterpene lactone cumenin (from <i>Ambrosia psilostachya</i>) showed a potent inhibitory effect in NO production (IC ₅₀ = 9.38 ± 0.38 µM) with low cytotoxicity.	37
<i>Aristolochia</i> species	The Chinese herb "Mu Tong" has included <i>Aristolochia manshuriensis</i> only since the 1950s. The classical Chinese herbal literature until the mid 17th century identifies "Mu Tong" as several <i>Akebia</i> species. From the 17th until the early 20th century "Mu Tong" was based on <i>Clematis</i> species. Renal failure due to ingestion of large doses of <i>Aristolochia manshuriensis</i> has been reported in China and other countries while no toxicity was recorded in traditional Chinese herbal texts. <i>Aristolochia</i> 's topical anti-inflammatory activity has been recently described. Aristolochic acids, isolated from <i>Aristolochia longa</i> inhibited <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Streptococcus faecalis</i> , <i>Staphylococcus aureus</i> and <i>Staphylococcus epidermidis</i> . The chloroform and hexane extracts of <i>Aristolochia trilobata</i> leaves and bark were active against <i>Escherichia coli</i> and <i>Pseudomonas aeruginosa</i> and <i>Staphylococcus aureus</i> .	38, 39
<i>Bambusa vulgaris</i>	The anti-inflammatory effect of the methanol extract of the leaves of <i>Bambusa arundinacea</i> was significant when compared to standard drugs validating its use in Ayurvedic medicine. The methanol extract of <i>Bambusa arundinacea</i> also showed antihypersensitivity activity, immunosuppressive activity, wound healing property and antibacterial activity experimentally.	40
<i>Bidens pilosa</i>	The "Shidachuan" which was originally recorded in "Ben Cao Gang Mu Shi Yi" (A Supplement to the Compendium of Materia Medica) is "Longyacao" (<i>Agrimonia pilosa</i>). "Shijianchuan" should be "Guizhencao" (<i>Bidens bipinata</i>). Bioactive polyacetylenes were found in the methanolic extract of <i>Bidens pilosa</i> (whole plant). The anti-inflammatory effect of aqueous extracts of the three plants <i>Bidens pilosa</i> var. <i>minor</i> (Blume) Sherff, <i>Bidens pilosa</i> and <i>Bidens chilensis</i> DC was significant. The immuno-suppressive activity of <i>Bidens pilosa</i> is attributed to the polyacetylene isolated from leaves. The water extract of <i>Bidens pilosa</i> showed a higher activity against <i>Bacillus cereus</i> and <i>Escherichia coli</i> than gentamycin sulphate. In one study diarrheic children in Trinidad were found to be positive for enteropathogenic <i>E. coli</i> .	41–43
<i>Bixa orellana</i>	<i>Bixa orellana</i> exhibited antimicrobial activity with a low MIC against <i>Escherichia coli</i> (0.8 microg/ml) compared to gentamycin sulfate (0.9 8 g/ml). <i>Bixa orellana</i> exhibited a better MIC against <i>Bacillus cereus</i> (0.2 microg/ml) than gentamycin sulfate (0.5 microg/ml).	43
<i>Cajanus cajan</i>	Extracts of roots and leaves of <i>Cajanus cajan</i> yielded 8 compounds: betulinic acid, biochanin A, cajanol, genistein and 2'-hydroxygenistein, longistylin A and C, and pinostrobin. The stilbenes, longistylin A and C, and betulinic acid showed moderate <i>in vitro</i> activity against chloroquine-sensitive <i>Plasmodium falciparum</i> . A protein was purified from the leaves and may enhance body immunosurveillance. <i>Cajanus indicus</i> protein possesses both a preventive and curative role against chloroform-induced hepatotoxicity and may act by an anti-oxidative defence mechanism.	44–46
<i>Capraria biflora</i>	The dried leaves of <i>Capraria biflora</i> (aqueous extract (50–200 mg kg(-1))) produced a moderate analgesic effect.	47
<i>Cecropia peltata</i>	<i>Cecropia pachystachya</i> has antioxidant properties. The two flavonoids orientin and iso-orientin, isolated from the active butanolic fraction could be responsible for the observed anxiolytic-like effect of <i>C. glaziovii</i> . Steroids and amino acids in <i>C. peltata</i> may account for the antimicrobial activity exhibited against <i>E. coli</i> .	48, 49
<i>Centropogon cornutus</i>	<i>Centropogon cornutus</i> has a synonym <i>Lobelia cornuta</i> . Three new piperidine alkaloids were isolated from stems, leaves and flowers of <i>Lobelia laxiflora</i> . The residues obtained from the ethanol extracts from stems, leaves, and flowers showed anti-inflammatory potential.	50
<i>Chamaesyce hirta</i> syn. <i>Euphorbia hirta</i>	<i>Euphorbia hirta</i> aqueous extract is used for dysentery, colic, bronchial infections and to treat ulcers. The plant contains eucocyanidol, quercitol, camphol, quercetrin, dihydroellagitannins and dimeric hydrolysable tannins – euphorbins. Ethanol extracts of the aerial parts of the plant showed antimicrobial activity against <i>Escherichia coli</i> (enteropathogen), <i>Proteus vulgaris</i> , <i>Pseudomonas aeruginosa</i> and <i>Staphylococcus aureus</i> .	51–53
<i>Citharexylum spinosum</i>	Six new iridoid glucosides and one known iridoid glucoside were isolated from the fruits and other parts of <i>Citharexylum caudatum</i> . The aerial parts of <i>Citharexylum spinosum</i> L., contain five iridoid glucosides, and one known lignan glucoside. When formulated in jojoba oil and applied to mice tails followed by infection with <i>Schistosoma mansoni cercariae</i> , the iridoid mixture from leaves of <i>Citharexylum quadrangular</i> blocked cercarial penetration and caused significant reduction (94%; P < 0.05) in worm burden in treated mice in comparison to controls.	54–56
<i>Cocos nucifera</i>	Coconut kernel fiber can protect cells from loss of oxidative capacity with the administration of the procarcinogen 1,2-dimethylhydrazine (DMH). The alcoholic extract of ripe dried coconut shell of <i>Cocos nucifera</i> showed antifungal activity against all dermatophytes tested with twice the concentration needed against <i>E. floccosum</i> (200 µg/ml). <i>Cocos nucifera</i> fruit exocarp has significant activity against all enteropathogens tested. All the strains tested were resistant to chloramphenicol; the two <i>Escherichia coli</i> species, the two <i>Shigella flexneri</i> species and the two <i>Salmonella</i> sp. species were sensitive to trimethoprim, and the two <i>Shigella sonnei</i> species were resistant. The authors concluded that coconut could be used as an alternative method to treat drug resistant enteric infections.	57, 58
<i>Cola nitida</i>	In Nigeria, <i>Cola accuminata</i> , <i>Cola nitida</i> and <i>Cola milleni</i> are used in ethnobotany for the treatment of diarrhea and dysentery. <i>Cola</i> species contain caffeine, koletein and kolatin alkaloids, proanthocyanin, magnesium, sodium, potassium bromide, cobalt, caesium, zinc and selenium. The <i>Mycobacterium bovis</i> was susceptible at 1000 µg/ml of methanol extract root bark of both <i>Cola nitida</i> and <i>Cola milleni</i> but insensitive to methanol extracts of both the leaves and stem-bark of the three <i>Cola</i> sp. tested. The MIC of the methanol root extract of <i>Cola nitida</i> against <i>Mycobacterium bovis</i> is 125 µg/ml. The MIC of methanol root extract of <i>Cola nitida</i> against the six ATCC strain of <i>Mycobacterium vaccae</i> ranged from 500 µg/ml to above 1000 µg/ml. The control Rifampicin is active against <i>M. bovis</i> at 5 µg and 10 µg/ml.	59
<i>Cucurbita</i> species	The minimum inhibitory concentration (MIC) of 23 gr. of pumpkin seed (+/- 73 seeds) (<i>Cucurbita maxima</i>) in 100 ml. distilled water as an antiparasitic agent using canine tapeworms with an intestinal isolation of 5 to 6 hours was determined. Alterations in helminthic motility were found at a dose of > 23 gr. There is a protheolithic effect with an average survival time of 38.4 minutes. The anthelmintic effect is increased at 30 and 32 gr.	60
<i>Dorstenia contrajerva</i>	<i>Dorstenia</i> species contain furanocoumarins with analgesic, anti-inflammatory, antibacterial, antiviral, anticoagulant, and photosensitizing activities. Prenylated chalcones are also found and may have anti-carcinogenic and antiproliferative properties. <i>Dorstenia contrajerva</i> was active toward <i>Giardia lamblia</i> with IC(50) < 38 mug/ml. This antiprotozoal activity supports the popular use to treat diarrhoea and dysentery.	61–63
<i>Eleusine indica</i>	<i>Eleusine indica</i> ethanol extract showed activity against vesicular stomatitis virus. The plant contains hydrocyanic acid.	64

Table 4: Non-experimental validation of plants used for stomach problems, pain and internal parasites in Trinidad and Tobago (Continued)

<i>Eupatorium macrophyllum</i>	The ethanol extract of the leaves of <i>Eupatorium adenophorum</i> (100, 200 and 300 mg/kg, po) showed significant analgesic activity, compared to standard drugs diclofenac sodium and pentazocine. Petroleum ether and methanolic extracts of leaves of <i>Eupatorium ayapana</i> showed broad spectrum antibacterial activity at the tested concentration (250–1000 µg/ml) except against <i>Shigella dysenteriae</i> . The petroleum ether extract also showed antifungal activity. Two extracts (dichloromethane and methanol), of the dried stems and leaves of <i>Eupatorium inulaefolium</i> , the S2 fraction of the hexane extract and neurolelin B from the dichloro-methane extract, showed statistically significant antiplasmodial activity.	65–67
<i>Ferula a safoetida</i>	A <i>Ferula asafoetida</i> gum extract (3 mg/ml), decreased the average amplitude of spontaneous contractions of the isolated guinea-pig ileum to 54 +/- 7% of control. <i>Ferula asafoetida</i> gum extract (0.3–2.2 mg/100 g body weight) reduced the mean arterial blood pressure in anaesthetised rats.	68
<i>Jatropha curcas</i>	Two deoxyprossomerins were isolated from stems of <i>Jatropha curcas</i> . Two compounds had antibacterial constituents. <i>Jatropha curcas</i> crude bark extract accelerates the healing process of wounds on Wistar albino rats by increasing the skin breaking strength, granulation tissue breaking strength, wound contraction, dry granulation tissue weight and hydroxyproline levels. A significant decrease in epithelization period was also observed.	69, 70
<i>Momordia charantia</i>	<i>Momordia charantia</i> may induce both intestinal and also systemic anti-inflammatory responses and may have antiviral activity.	71, 72
<i>Morinda citrifolia</i>	The lyophilised aqueous extract of roots of <i>Morinda citrifolia</i> produced a dose-related, central analgesic activity in mice. The analgesic efficacy of the Noni extract was less strong than morphine but non-addictive and had no side effects. <i>Morinda citrifolia</i> fruit powder demonstrated over 70% COX-1 inhibition. The extracts from <i>Morinda citrifolia</i> leaf (45%) showed moderate inhibition on COX-1. The extracts from <i>Morinda citrifolia</i> bark (27%) and <i>Morinda citrifolia</i> fresh fruit juice (38%) presented low inhibition on COX-1. The extract from <i>Morinda officinalis</i> root was inactive (9.87%) at a concentration of 3.4 mg/ml.	73–75
<i>Neurolaena lobata</i>	<i>Neurolaena lobata</i> has antinociceptive and antibacterial effects. When tested against <i>Brugia pahangi</i> , a lymphatic dwelling filarial worm, the ethanol extract of <i>Neurolaena lobata</i> showed potential macro- and micro-filaricidal activity.	76
<i>Nicotiana tabacum</i>	The lack of nicotine-induced analgesia assessed by the tail flick reflex test in female rats is consistent with human studies showing that nicotine reduces pain elicited by brief noxious cutaneous stimulation in male but not female subjects.	77
<i>Peperomia rotundifolia</i>	In south-east Asia, <i>Peperomia pellucida</i> is used for wounds, skin problems, abdominal pain and other pains and for headache. <i>Peperomia pellucida</i> is reported to have analgesic activity in mice, antibacterial activity against <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> and <i>Staphylococcus aureus</i> , and antifungal activity. <i>Peperomia pellucida</i> ethyl-acetate soluble extracts and crude methanolic extracts were active against Gram-positive and Gram-negative bacteria.	78
<i>Petiveria alliacea</i>	<i>Petiveria alliacea</i> extract showed an antinociceptive effect which account for its popular use as an analgesic. The oral administration of <i>Petiveria alliacea</i> root crude lyophilized extract at the highest dose of extract tested (43.9 mg/kg body wt.) significantly reduced the number of migrating neutrophils, mononuclear cells and eosinophils. The <i>Petiveria alliacea</i> root extract also showed a significant analgesic effect. Thiosulfates, trisulfides and benzylsulfonic acid are antimicrobial compounds, with the benzyl-containing thiosulfates having the broadest spectrum of antimicrobial activity.	79–81
<i>Portulaca oleracea</i>	The ingestion of purslane (<i>Portulaca oleracea</i>) leaves may have a protective effect against oxidative stress caused by vitamin A deficiency.	82
<i>Punica granatum</i>	<i>Punica granatum</i> was used by Egyptians in ancient times as a treatment for tapeworm and other parasites. A pomegranate extract at a low extract concentration (0.01% v/v) delayed bacterial growth of <i>Staphylococcus aureus</i> FRI 722, while a higher concentration (1% v/v) eliminated bacterial growth.	83
<i>Rosmarinus officinalis</i>	<i>Rosmarinus officinalis</i> has historically been used as an analgesic and antirheumatic herb. The aqueous and ethanol extracts of <i>Rosmarinus officinalis</i> L. aerial parts induced a significant antinociceptive activity. In an observational study, a combination of reduced iso-alpha-acids from hops, rosemary extract and oleanolic acid decreased pain in patients suffering from rheumatic conditions and osteoarthritis.	84, 85
<i>Solanum melongena</i>	<i>Solanum melongena</i> contains significant quantities of histamine and serotonin.	86
<i>Scoparia dulcis</i>	<i>Scoparia dulcis</i> has traditionally been used to treat stomach troubles, inflammation, hemorrhoids, and hepatitis and as an analgesic. Biologically active substances from <i>Scoparia dulcis</i> include scoparic acid A, scoparic acid B, scopadulcic acid A and B, scopadulciol and scopadulin. The chloroform/methanol fractions <i>Scoparia dulcis</i> showed antimicrobial activity against the human pathogenic bacteria <i>Salmonella typhi</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Bacillus subtilis</i> , <i>Pseudomonas aeruginosa</i> , and <i>Proteus vulgaris</i> and the plant pathogenic fungi <i>Alternaria macrospora</i> , <i>Candida albicans</i> , <i>Aspergillus niger</i> , and <i>Fusarium oxysporum</i> .	87
<i>Tagetes patula</i>	<i>Tagetes erecta</i> callus cultures produce ascorbic acid as well as insecticidal pyrethrins. <i>Tagetes patula</i> oil contains several compounds with the major ones being limonene, (Z) and (E)-β-ocimene, dihydrotagetone, terpinolene, piperitone, peperitenone, E-caryophyllene and trans-sesquibabinene hydrate. The fourth instar larvae of <i>Aedes aegypti</i> (LC ₅₀ 13.57, LC ₉₀ 37.91) was most susceptible to <i>Tagetes patula</i> essential oil followed by <i>Anopheles stephensi</i> (LC ₅₀ 12.08, LC ₉₀ 57.62) and <i>Culex quinquefasciatus</i> (LC ₅₀ 22.33, LC ₉₀ 71.89).	88, 89
<i>Tamarindus indica</i>	In Thai traditional medicine, the fruit of <i>Tamarindus indica</i> is considered to be as a digestive, carminative, laxative, expectorant and a blood tonic. A crude <i>Tamarindus indica</i> seed extract inhibited the PLA2, protease, hyaluronidase, L-amino acid oxidase and 5'-nucleotidase enzyme activities of <i>Vipera russelli</i> venom in a dose-dependent manner. Mice that received the extract 10 min after the injection of venom were protected from venom-induced toxicity. The seed coat extract of <i>Tamarindus indica</i> has antioxidant activity. The extract is composed of flavonoids including tannins, polyphenols, anthocyanidin, and oligomeric proanthocyanidins. These flavonoids may produce vasorelaxant activity, increase capillary permeability and protection from oxidative stress. Excess nitric oxide production is associated with diseases such as autoimmunity, rheumatoid arthritis, inflammatory bowel disease and septic shock. <i>In vitro</i> studies demonstrated that the crude seed coat extract of <i>Tamarindus indica</i> suppressed nitric oxide production while producing no adverse effects.	90, 91
<i>Tournefortia hirsutissima</i>	In Taiwan, <i>Tournefortia sarmentosa</i> Lam. is used as a detoxicant, an anti-inflammatory agent, and a circulation promoter to remove blood stasis. Alkaloids, flavones, triterpenoids, and cinnamates are found in the genus <i>Tournefortia</i> . The stems of <i>Tournefortia sarmentosa</i> contain five phenolic compounds as well as salicylic acid and allantoin. <i>Tournefortia rufosericeae</i> leaves contain pyrrolizidine alkaloids (5% of dry weight).	92, 93

Table 5: Chinese ethnomedicinal uses for the Chinese-origin plants or closely related species used in Trinidad

Trinidad ethnomedicinal plant	Chinese ethnomedicinal plant and practice
<i>Abelmoschus moschatus</i>	Geographical origin S.E. Asia. Myricetin a flavonol, is found in tea, berries, fruits, and the herb of <i>Abelmoschus moschatus</i> . This flavonol has both antioxidative and cytoprotective properties and has been used successfully to treat depression and anxiety in traditional Chinese medicine [94].
<i>Achyranthes aspera</i> , <i>Achyranthes indica</i>	<i>Achyranthes bidentata</i> is grown in the tropical parts of China, Korea and Vietnam. Its roots ("Niu Xi", Radix <i>Achyranthes Bidentatae</i>) are used in traditional Chinese medicine as a tonic, emmenagogue, antiarthritic, diuretic, and antifertility agent to nourish the liver and kidneys, strengthen bones and muscles and invigorate circulation [95].
<i>Aristolochia rugosa</i> , <i>A. trilobata</i>	The stem of <i>Aristolochia manshuriensis</i> (AMA, Guanmuton) is a traditional Chinese medicinal herb largely harvested from the Northeast of China. It is used as a diuretic, anti-inflammatory, to alleviate swelling and to treat rheumatism [96].
<i>Bidens alba</i> / <i>Bidens pilosa</i>	<i>Bidens parviflora</i> (Xiaohua-Guizhencao) is used as a traditional antipyretic, anti-inflammatory and anti-rheumatic medicine in China [97]. <i>Bidens pilosa</i> was introduced into Asia and is common in Taiwan.
<i>Cajanus cajan</i>	In Chinese folk medicine pigeon pea leaves are used to staunch blood, as an analgesic and to kill parasites [98].
<i>Cassia alata</i>	<i>Cassia obtusifolia</i> seed, called "Juemingzi", is used to treat eye infections, headache, and dizziness [99]. <i>Cassia alata</i> can be purchased in herb shops in Thailand.
<i>Croton gossypifolius</i>	There are 21 species of <i>Croton</i> distributed throughout the southern part of China. Several species including <i>C. kongensis</i> are used in traditional Chinese medicine to alleviate dysmenorrhea (fruits), as a purgative (seeds), and to treat dyspepsia (bark) and malaria (leaves) [100].
<i>Eclipta prostrata</i>	In Chinese medicine this plant is called "Eclipta Prostrata Herba" (Yetbadetajo Hert) [101]. It is also used in Taiwanese folk medicine.
<i>Eupatorium macrophyllum</i>	<i>Eupatorium chinense</i> grows in the south of China and is used for colds, snakebite and inflammation [102].
<i>Momordica charantia</i>	<i>Momordica charantia</i> seeds are known in Chinese medicine as "Ku guazi". They are used for infections and immune disorders [103].
<i>Morinda citrifolia</i>	Chinese traditional tonic herbal medicine "BajiTian" (<i>Morinda officinalis</i>) has been prescribed in China for about two thousand years, for tonifying kidney, strengthening Yang-qi and relieving rheumatism [104].
<i>Phyllanthus urinaria</i>	<i>Phyllanthus urinaria</i> grows widely in China. It is used to treat jaundice, hepatitis B, nephrolithiasis, and painful disorders [105].
<i>Portulaca oleraceae</i>	<i>Portulaca oleracea</i> (Ma-Chi-Xian), grows widely in China, and is used traditionally for alleviating pain and swelling. It has anti-bacterial, anti-viral, anti-diabetic, and immuno-modulating activity [106].
<i>Sida acuta</i>	This medicinal plant is named "Huanghuaren" [107].
<i>Tamarindus indica</i>	In Thai traditional medicine, the fruit of <i>T. indica</i> is used as a digestive, laxative, expectorant and blood tonic. The seeds of <i>T. indica</i> are used as an anthelmintic, antidiarrheal, and an emetic, and the seed coat is used to treat burns and aid in wound healing as well as against dysentery. [90], [91]

Non-experimental validation of plants used for skin problems in Trinidad and Tobago

For each species or genus the ethnomedicinal uses in other countries, particularly Asian countries, are given if available; then follows a summary of chemical constituents, in addition to active compounds if relevant to the condition being treated (Tables 3 and 4).

Comparative evaluation of plants used for skin problems, stomach problems, pain and internal parasites

Table 5 contains a preliminary listing of the ethnomedicinal plants discussed in this paper that are used similarly in Chinese ethnomedicine. If the specific plant was not found in the literature search the closely related species that are used similarly in Chinese traditional medicine are listed.

The commonalities between Chinese traditional medicine and Trinidad and Tobago "bush medicine" are provided below.

Abelmoschus moschatus is used to treat depression and anxiety in traditional Chinese medicine [94]. In Trinidad and Tobago it is used for pain.

Achyranthes bidentata ("Niu Xi" in Chinese medicine, Radix *Achyranthes Bidentatae*) is used as a tonic, to nourish the liver and kidneys, and invigorate circulation [95]. *Achyranthes indica* is used in Trinidad and Tobago for skin rashes and other skin problems.

Aristolochia manshuriensis (AMA, "Guanmuton") is used in China as a diuretic and anti-inflammatory [96]. *Aristolochia rugosa/trilobata* are used in Trinidad and Tobago for stomach problems. Zhu claims that the Chinese herb "Mu Tong" has been based on *Aristolochia manshuriensis* only since the 1950s. The classical Chinese herbal literature until the mid 17th century identifies "Mu Tong" as several *Akebia* species and no toxicity related to "Mu Tong" was recorded in these traditional Chinese herbal texts.

Bidens parviflora ("Xiaohua-Guizhencao") is used as a traditional antipyretic, anti-inflammatory and anti-rheu-

matic medicine in China [97]. Plants used for cuts, injuries and swellings in Trinidad and Tobago include *Bidens alba*/*Bidens pilosa*.

During the ethnomedicinal research one of the respondents claimed that the use of *Cajanus cajan* for internal parasites was a recent addition to Trinidad folk medicine. This ethnomedicinal practice in Trinidad is the same as that reported for the folk medicine of China (to kill parasites) [98] but no definitive statements about its origins can be made at this time. *Momordica charantia* seeds or "Ku guazi" are used for infections and immune disorders [103]; in Trinidad and Tobago the plant is used for stomach problems.

"BaJiTian" (*Morinda officinalis*) has been prescribed in China for about two thousand years, for tonifying the kidney, strengthening "Yang-qi" and relieving rheumatism [104]. Plants used for pain in Trinidad and Tobago include *Morinda citrifolia*.

Phyllanthus urinaria is extensively grown in China. It is used to treat jaundice, hepatitis B, nephrolithiasis, and painful disorders [106]. Diarrhoea is treated with *Phyllanthus urinaria* in Trinidad and Tobago.

Portulaca oleracea ("Ma-Chi-Xian") is grown widely in China, and is used traditionally for alleviating pain and swelling [106]. It is used as an anthelmintic in Trinidad and Tobago. *Tamarindus indica* fruit is used as a blood tonic and the seed coat of *Tamarindus indica* is used to treat burns and aid in wound healing in China. In Trinidad and Tobago, scorpion and snake bites are treated with *Tamarindus indica*.

Discussion and conclusion

Vincent Yáñez, the captain of the caravel Niña reportedly dug up *Morinda citrifolia* in Hispaniola on December 30, 1492 [1]; yet this plant was not considered special in Trinidad until the forces of globalisation made "Noni" ubiquitous as an "Australasian cure-all" and it was then sold on the streets of Trinidad by herbalists and other traders [1]. This story illustrates that since Caribbean folk medicine is a product of globalisation and colonisation, research into its origins and plant uses is complex. Attributing specific uses to Chinese folk medicine would necessitate access to the earliest Chinese herbals.

The ship that brought 467 Chinese men, women, and children (from an original 549) in 1862 was the first ship to bring Chinese women to Trinidad. In the last 5 voyages (1862–1866), of 367 females embarked, 309 landed. The immigrant gender imbalance may have affected the dissemination of Chinese folk medicine into the Caribbean culture. Two wars taking place in eastern China in 1862

facilitated the immigration or abduction of Hakka peoples to the Americas and presumably the Puntí peoples came in the later stages of immigration [108,109]. If any of these original immigrants had expertise in Chinese plants, besides knowledge of opium, they did not widely advertise this under the British colonial administration.

It may be the case that the Chinese contribution to Caribbean folk medicine has formed part of its earliest foundation and its provenance is not remembered. Research on the Chinese contribution to Trinidad is complicated by the fact that many of the Hakka research population have lived up to their migratory reputation – moving on to North America. Language is also a barrier.

Cuba and other Caribbean countries have not adopted the model of China's barefoot doctors. Cuba's medical diplomacy and investment in biotechnology generates symbolic capital: intangible qualities (like honour, prestige, and reputation) which appear opposed to strictly economic interests, are in fact convertible back into material capital [110]. The Cuban policy is to demonstrate that its socialist state can provide a modern health care system and need not settle for small-scale technologies like traditional medicine [110]. In contrast it has been estimated that 80% of medications used in Chinese rural areas are derived from Chinese materia medica and related products. These products are economical and therefore provide important cost savings [2,111,112].

Similarly to the process taking place in the Caribbean, younger people in Taiwan have been moving away from Chinese medicines because work pressures force them to seek faster cures from allopathic doctors [2]. However tonic herbs such as "Danggui" (*Radix Angelica sinensis*), "Huangqi" (*Radix Astragalii/Astragalus membranaceus*), "Gou Qi Zi" (*Fructus barbarum*) and "Renshen" (*Radix Panax ginseng /Panax notoginseng*), are used by Taiwanese families in slow-cooking winter meals. These herbs are also popular for postnatal care, for the elderly and for post-surgical therapy [2].

Non-experimental validation is a new approach that is designed to introduce cost effectiveness into medicinal plant research. The findings of the non-experimental validation suggest that the majority of the therapeutic applications of the plants used in Caribbean folk medicine listed in this paper are justified, and more studies are warranted to explore their efficacy. All of the plants used in Trinidad and Tobago for skin problems merit clinical trials. The plants used for stomach problems, pain and internal parasites that should take priority in clinical trials are *Bambusa vulgaris*, *Bidens alba*, *Jatropha curcas*, *Neurolaena lobata*, *Peperomia rotundifolia* and *Phyllanthus urinaria*.

Competing interests

The author(s) declare that they have no competing interests.

Acknowledgements

This data collection was part of a larger study for a Ph.D. at Wageningen UR, the Netherlands. The fellowship support provided is appreciated. The Herbarium staff of the University of the West Indies provided essential plant identification. Thanks to all of the respondents. This paper serves as a small recognition of the official Bicentennial of the Arrival of the Chinese in T&T, 1806 – 2006.

References

- Lans C: *Creole remedies of Trinidad and Tobago*, book self-published on Lulu.com 2006.
- Lans C, Harper T, Georges K, Bridgewater E: **Medicinal and ethnoveterinary remedies of hunters in Trinidad.** *BMC Complement Altern Med* 2001, **1**:10.
- Chan K: **Chinese medicinal materials and their interface with Western medical concepts.** *J Ethnopharmacol* 2005, **96**:1-18.
- Anon: **The Chinese in Trinidad and Tobago.** 2006 [<http://library2.nalis.gov.tt/Default.aspx?tabid=249>].
- Gerard Besson: *The 'Land of Beginnings'. A historical digest.* *Newsday Newspaper Sunday August 27 2000*.
- Yun L, Laremont R: **Chinese Coolies and African Slaves in Cuba, 1847-74.** *Journal of Asian American Studies* 2001, **4**:99-122.
- Harris R: *Local Herbs Used in the Chinese Way (Tonics).* Book 1. *The Traditional Chinese Medical Centre, Trinidad and Tobago, WI* 1991.
- Deng HB, Cui DP, Jiang JM, Feng YC, Cai NS, Li DD: **Inhibiting effects of *Achyranthes bidentata* polysaccharide and *Lycium barbarum* polysaccharide on nonenzyme glycation in D-galactose induced mouse aging model.** *Biomed Environ Sci* 2003, **16**(3):267-75.
- Zeng Y, Zhong JM, Ye SQ, Ni ZY, Miao XQ, Mo YK, Li ZL: **Screening of Epstein-Barr virus early antigen expression inducers from Chinese medicinal herbs and plants.** *Biomed Environ Sci* 1994, **7**:50-5.
- Chakraborty A, Brantner A, Mukainaka T, Nobukuni Y, Kuchide M, Konoshima T, Tokuda H, Nishino H: **Cancer chemopreventive activity of *Achyranthes aspera* leaves on Epstein-Barr virus activation and two-stage mouse skin carcinogenesis.** *Cancer Lett* 2002, **177**(1):1-5.
- Baboolal S, Rawlins SC: **Seroprevalence of toxocarasis in schoolchildren in Trinidad.** *Trans R Soc Trop Med Hyg* 2002, **96**(2):139-43.
- Charles V, Charles SX: **The use and efficacy of *Azadirachta indica* ADR ('Neem') and *Curcuma longa* ('Turmeric') in scabies. A pilot study.** *Tropical and Geographical Medicine* 1992, **44**:178-181.
- Dasgupta T, Banerjee S, Yadava PK, Rao AR: **Chemopreventive potential of *Azadirachta indica* (Neem) leaf extract in murine carcinogenesis model systems.** *J Ethnopharmacol* 2004, **92**:23-36.
- Aftab Saeed: **Medicinal, culinary and aromatic plants in Pakistan.** *Medicinal, Culinary and Aromatic plants in the Near East* [<http://www.fao.org/docrep/x5402e/x5402e15.htm>]. Proceedings of the International Expert Meeting organized by the Forest Products Division FAO Forestry Department and the FAO Regional Office for the Near East. Cairo, Egypt 19 – 21 May 1997
- Chiang YM, Lo CP, Chen YP, Wang SY, Yang NS, Kuo YH, Shyr LF: **Ethyl caffeate suppresses NF-kappaB activation and its downstream inflammatory mediators, iNOS, COX-2, and PGE2 in vitro or in mouse skin.** *Br J Pharmacol* 2005, **146**:352-63.
- Alvarez A, Pomar F, Sevilla J, Montero MJ: **Gastric antisecretory and antiulcer activities of an ethanolic extract of *Bidens pilosa* L. var. *radiata* Schult. Bip.** *J Ethnopharmacol* 1999, **67**:333-40.
- Geissberger P, Sequin U: **Constituents of *Bidens pilosa* L.: do the components found so far explain the use of this plant in traditional medicine?** *Acta Trop* 1991, **48**:251-61.
- Crockett CO, Guede-Guina F, Pugh D, Vangah-Manda M, Robinson TJ, Olubadewo JO, Ochillo RF: ***Cassia alata* and the preclinical search for therapeutic agents for the treatment of opportunistic infections in AIDS patients.** *Cell Mol Biol* 1992, **38**:505-11. Erratum in: *Cell Mol Biol* 1992; 38: 615.
- Damodaran S, Venkataraman S: **A study on the therapeutic efficacy of *Cassia alata*, Linn. leaf extract against *Pityriasis versicolor*.** *J Ethnopharmacol* 1994, **42**:19-23.
- Yen GC, Chen HW, Duh PD: **Extraction and identification of antioxidative component from *Jue Ming Zi* (*Cassia tora* L.).** *J Agric Food Chem* 1998, **46**:820-824.
- Cuellar MJ, Giner RM, Recio MC, Manez S, Rios JL: **Topical anti-inflammatory activity of some Asian medicinal plants used in dermatological disorders.** *Fitoterapia* 2001, **72**:221-9. *cassia anti-infla*
- Mukhopadhyay SK, Buddhadeb D, Duary B, Dasgupta MK, (Ed), Ghosh DC, (Ed), Gupta DD, (Ed), Majumdar DK, (Ed), Chattopadhyay GN, (Ed), Ganguli PK, (Ed), Muni PS, (Ed), Bhattacharya D: . In *Ethnobotany of some common crop field weeds in a sub-humid agricultural tract of West Bengal* Proceedings of the national symposium on sustainable agriculture in sub-humid zone, Sriniketan, West Bengal, India:272-277. 3 – 5 March 1995
- Tayanin GL, Bratthall D: **Black teeth: beauty or caries prevention? Practice and beliefs of the Kammu people.** *Community Dent Oral Epidemiol* 2006, **34**:81-6. *croton*
- Frum Y, Viljoen AM: **In vitro 5-lipoxygenase and anti-oxidant activities of South African medicinal plants commonly used topically for skin diseases.** *Skin Pharmacol Physiol* 2006, **19**:329-335.
- Thongtan J, Kittakoop P, Ruangrunsi N, Saenboonrueng J, Thebtaranonth Y: **New antimycobacterial and antimalarial 8,9-secoaurane diterpenes from *Croton kongensis*.** *J Nat Prod* 2003, **66**:868-70.
- Jones K: **Review of sangre de drago (*Croton lechleri*) – a South American tree sap in the treatment of diarrhea, inflammation, insect bites, viral infections, and wounds: traditional uses to clinical research.** *J Altern Complement Med* 2003, **9**:877-96.
- Sawangaroen N, Subhadhirasakul S, Phongpaichit S, Siripanth C, Jangjaroen K, Sawangaroen K: **The in vitro anti-giardial activity of extracts from plants that are used for self-medication by AIDS patients in southern Thailand.** *Parasitol Res* 2005, **95**:17-21.
- Liao F, Huang Q, Yang Z, Xu H, Gao Q: **Experimental study on the antibacterial effect of origanum volatile oil on dysentery bacilli in vivo and in vitro.** *J Huazhong Univ Sci Technolog Med Sci* 2004, **24**:400-3.
- Santoyo S, Cavero S, Jaime L, Ibanez E, Senorans FJ, Reglero G: **Supercritical carbon dioxide extraction of compounds with antimicrobial activity from *Origanum vulgare* L.: determination of optimal extraction parameters.** *J Food Prot* 2006, **69**:369-75.
- Khan-Mohammed Z, Adesiyun AA, Swanston WH, Chadee DD: **Frequency and characteristics of selected enteropathogens in fecal and rectal specimens from childhood diarrhea in Trinidad, 1998-2000.** *Rev Panam Salud Publica* 2005, **17**:170-7.
- Malairajan P, Geetha Gopalakrishnan, Narasimhan S, Jessi Kala Veni K: **Analgesic activity of some Indian medicinal plants.** *J Ethnopharmacol* 2006, **106**:425-8. *sida acuta*
- Al Chami L, Mendez R, Chataing B, O'Callaghan J, Usubillaga A, LaCruz L: **Toxicological effects of alpha-solamargine in experimental animals.** *Phytother Res* 2003, **17**:254-8.
- Nagase H, Sasaki K, Kito H, Haga A, Sato T: **Inhibitory effect of delphinidin from *Solanum melongena* on human fibrosarcoma HT-1080 invasiveness in vitro.** *Planta Medica* 1998, **64**:216-9.
- Konning GH, Agyare C, Ennison B: **Antimicrobial activity of some medicinal plants from Ghana.** *Fitoterapia* 2004, **75**:65-7. *afmomomum*
- Adegoke GO, Skura BJ: **Nutritional profile and antimicrobial spectrum of the spice *Aframomum danielli* K. Schum.** *Plant Foods Hum Nutr* 1994, **45**:175-82.
- Suite M: **Cutaneous infections in Trinidad.** *Int J Dermatol* 1990, **29**:31-4.
- Lastra AL, Ramirez TO, Salazar L, Martinez M, Trujillo-Ferrara J: **The ambrosanolid cumanin inhibits macrophage nitric oxide synthesis: some structural considerations.** *J Ethnopharmacol* 2004, **95**:221-7.
- Zhu YP: **Toxicity of the Chinese herb mu tong (*Aristolochia manshuriensis*). What history tells us.** *Adverse Drug React Toxicol Rev* 2002, **21**:171-7.

39. Camporese A, Balick MJ, Arvigo R, Esposito RG, Morsellino N, De Simone F, Tubaro A: **Screening of anti-bacterial activity of medicinal plants from Belize (Central America).** *J Ethnopharmacol* 2003, **87**:103-7.
40. Muniappan M, Sundararaj T: **Anti-inflammatory and antiulcer activities of *Bambusa arundinacea*.** *J Ethnopharmacol* 2003, **88**:161-7.
41. Chih HW, Lin CC, Tang KS: **Anti-inflammatory activity of Taiwan folk medicine "ham-hong-chho" in rats.** *American Journal of Chinese Medicine* 1995, **23**:273-8.
42. Xie ZW: **Textural research on "Shidachuan" and "Shijianchuan" in "Ben Cao Gang Mu Shi Yi" (a supplement to the compendium of materia medica).** *Zhongguo Zhong Yao Za Zhi* 2000, **25**:49-51.
43. Rojas JJ, Ochoa VJ, Ocampo SA, Munoz JF: **Screening for antimicrobial activity of ten medicinal plants used in Colombian folkloric medicine: a possible alternative in the treatment of non-nosocomial infections.** *BMC Complement Altern Med* 2006, **6**:2.
44. Duker-Eshun G, Jaroszewski JW, Asomaning WA, Oppong-Boachie F, Brogger Christensen S: **Antiplasmodial constituents of *Cajanus cajan*.** *Phytother Res* 2004, **18**:128-30.
45. Datta S, Sinha S, Bhattacharyya P: **Effect of a herbal protein, CI-I, isolated from *Cajanus indicus* on immune response of control and stressed mice.** *J Ethnopharmacol* 1999, **67**:259-267.
46. Ghosh A, Sarkar K, Sil PC: **Protective effect of a 43 kD protein from the leaves of the herb, *Cajanus indicus* L. on chloroform induced hepatic-disorder.** *J Biochem Mol Biol* 2006, **39**:197-207.
47. Acosta SL, Muro LV, Sacerio AL, Pena AR, Okwei SN: **Analgesic properties of *Capraria biflora* leaves aqueous extract.** *Fitoterapia* 2003, **74**:686-8.
48. Consolini AE, Ragone MI, Migliori GN, Conforti P, Volonte MG: **Cardiotonic and sedative effects of *Cecropia pachystachya* Mart. (ambay) on isolated rat hearts and conscious mice.** *J Ethnopharmacol* 2006, **106**:90-6.
49. Rocha FF, Lapa AJ, De Lima TC: **Evaluation of the anxiolytic-like effects of *Cecropia glazioui* Sneth in mice.** *Pharmacol Biochem Behav* 2002, **71**:183-90.
50. Philipov S, Istatkova R, Ivanovska N, Denkova P, Tosheva K, Navas H, Villegas J: **Phytochemical study and anti-inflammatory properties of *Jobelia laxiflora* L.** *Z Naturforsch [C]* 1998, **53**:311-7.
51. Vijaya K, Ananthan S, Nalini R: **Antibacterial effect of theaflavin, polyphenol 60 (*Camellia sinensis*) and *Euphorbia hirta* on *Shigella* spp. – a cell culture study.** *J Ethnopharmacol* 1995, **49**:115-8.
52. Tona L, Kambu K, Ngimbi N, Mesia K, Penge O, Lusakibanza M, Cimanga K, De Bruyne T, Apers S, Totte J, Pieters L, Vlietinck AJ: **Antiamoebic and spasmolytic activities of extracts from some antiarrhoeal traditional preparations used in Kinshasa, Congo.** *Phytomedicine* 2000, **7**:31-8.
53. Wang YC, Huang TL: **Screening of anti-Helicobacter pylori herbs deriving from Taiwanese folk medicinal plants.** *FEBS Immunol Med Microbiol* 2005, **43**:295-300.
54. Ayers S, Sneden AT, Caudatosides A-F: **New iridoid glucosides from *Citharexylum caudatum*.** *J Nat Prod* 2002, **65**:1621-6.
55. Balazs B, Toth G, Duddeck H, Soliman HS: **Iridoid and lignan glycosides from *Citharexylum spinosum* L.** *Nat Prod Res* 2006, **20**:201-5.
56. Bahgat M, Shalaby NM, Ruppel A, Maghraby AS: **Humoral and cellular immune responses induced in mice by purified iridoid mixture that inhibits penetration of *Schistosoma mansoni* cercariae upon topical treatment of mice tails.** *J Egypt Soc Parasitol* 2005, **35**:597-613.
57. Pillai MG, Thampi BS, Menon VP, Leelamma S: **Influence of dietary fiber from coconut kernel (*Cocos nucifera*) on the 1,2-dimethylhydrazine-induced lipid peroxidation in rats.** *J Nutr Biochem* 1999, **10**:555-60.
58. Alanis AD, Calzada F, Cervantes JA, Torres J, Ceballos GM: **Antibacterial properties of some plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders.** *J Ethnopharmacol* 2005, **100**:153-7.
59. Adeniyi BA, Groves MJ, Gangadharam PR: **In vitro anti-mycobacterial activities of three species of *Cola* plant extracts (Sterculiaceae).** *Phytother Res* 2004, **18**:414-8.
60. Diaz Obregon D, Lloja Lozano L, Carbajal Zuniga V: **Preclinical studies of *Cucurbita maxima* (pumpkin seeds) a traditional intestinal antiparasitic in rural urban areas.** *Rev Gastroenterol Peru* 2004, **24**:323-7. [Article in Spanish]
61. Tovar-Miranda R, Cortés-García R, Santos-Sánchez NF, Joseph-Nathan P: **Isolation, total synthesis, and relative stereochemistry of a dihydrofurocoumarin from *Dorstenia contrajerva*.** *J Nat Prod* 1998, **61**:1216-20.
62. Ngameni B, Touaibia M, Patnam R, Belkaid A, Sonna P, Ngadjui BT, Annabi B, Roy R: **Inhibition of MMP-2 secretion from brain tumor cells suggests chemopreventive properties of a furanocoumarin glycoside and of chalcones isolated from the twigs of *Dorstenia turbinata*.** *Phytochemistry* 2006, **67**:2573-9.
63. Calzada F, Yopez-Mulia L, Aguilar A: **In vitro susceptibility of *Entamoeba histolytica* and *Giardia lamblia* to plants used in Mexican traditional medicine for the treatment of gastrointestinal disorders.** *J Ethnopharmacol* in press. 2006, Jun 2; dorstenia
64. Ali Abdul M, Mackeen MM, El-Sharkawy S, Hamid J, Ismail N, Ahmad F, Lajis N: **Antiviral and cytotoxic activities of some plants used in Malaysian indigenous medicine.** *Pertanika Journal of Tropical Agricultural Science* 1996, **19**:129-136.
65. Mandal SK, Boominathan R, Parimaladevi B, Dewanjee S, Mandal SC: **Analgesic activity of methanol extract of *Eupatorium adenophorum* Spreng. leaves.** *Indian J Exp Biol* 2005, **43**:662-3.
66. Gupta M, Mazumder UK, Chaudhuri I, Chaudhuri RK, Bose P, Bhattacharya S, Manikandan L, Patra S: **Antimicrobial activity of *Eupatorium ayapana*.** *Fitoterapia* 2002, **73**:168-70.
67. Blair S, Mesa J, Correa A, Carmona-Fonseca J, Granados H, Saez JL: **Antimalarial activity of neurolepin B and derivatives of *Eupatorium inulaefolium* (Asteraceae).** *Pharmazie* 2002, **57**:413-5.
68. Fatehi M, Farifteh F, Fatehi-Hassanabad Z: **Antispasmodic and hypotensive effects of *Ferula asafoetida* gum extract.** *J Ethnopharmacol* 2004, **91**:321-4.
69. Ravindranath N, Reddy MR, Mahender G, Ramu R, Kumar KR, Das B: **Deoxypreussomerins from *Jatropha curcas*: are they also plant metabolites?** *Phytochemistry* 2004, **65**:2387-90.
70. Shetty S, Udupa SL, Udupa AL, Vollala VR: **Wound healing activities of bark extract of *Jatropha curcas* Linn in albino rats.** *Saudi Med J* 2006, **27**:1473-6.
71. Manabe M, Takenaka R, Nakasa T, Okinaka O: **Induction of anti-inflammatory responses by dietary *Momordica charantia* L. (bitter melon).** *Biosci Biotechnol Biochem* 2003, **67**:2512-7.
72. Grover JK, Yadav SP: **Pharmacological actions and potential uses of *Momordica charantia*: a review.** *J Ethnopharmacol* 2004, **93**:123-32.
73. Wang MY, West BJ, Jensen CJ, Nowicki D, Su C, Palu AK, Anderson G: ***Morinda citrifolia* (Noni): a literature review and recent advances in Noni research.** *Acta Pharmacol Sin* 2002, **23**:1127-41.
74. Younos C, Rolland A, Fleurentin J, Lanhers MC, Misslin R, Mortier F: **Analgesic and behavioural effects of *Morinda citrifolia*.** *Planta Med* 1990, **56**:430-4.
75. Li RW, Myers SP, Leach DN, Lin GD, Leach G: **A cross-cultural study: anti-inflammatory activity of Australian and Chinese plants.** *J Ethnopharmacol* 2003, **85**:25-32.
76. Fujimaki Y, Kamachi T, Yanagi T, Caceres A, Maki J, Aoki Y: **Macrophilicidal and microfilaricidal effects of *Neurolaena lobata*, a Guatemalan medicinal plant, on *Brugia pahangi*.** *J Helminthol* 2005, **79**:23-8.
77. Carstens E, Anderson KA, Simons CT, Carstens MI, Jinks SL: **Analgesia induced by chronic nicotine infusion in rats: differences by gender and pain test.** *Psychopharmacologia* 2001, **157**:40-45.
78. Khan MR, Omoloso AD: **Antibacterial activity of *Hygrophila stricta* and *Peperomia pellucida*.** *Fitoterapia* 2002, **73**:251-4.
79. Kim S, Kubec R, Musah RA: **Antibacterial and antifungal activity of sulfur-containing compounds from *Petiveria alliacea* L.** *J Ethnopharmacol* 2006, **104**:188-92.
80. Lopes-Martins RA, Pegoraro DH, Woisky R, Penna SC, Sertie JA: **The anti-inflammatory and analgesic effects of a crude extract of *Petiveria alliacea* L. (Phytolaccaceae).** *Phytomedicine* 2002, **9**:245-8.
81. Kubec R, Kim S, Musah RA: **S-Substituted cysteine derivatives and thiosulfinate formation in *Petiveria alliacea* – part II.** *Phytochemistry* 2002, **61**:675-80.
82. Malek F, Boskabady MH, Borushaki MT, Tohidi M: **Bronchodilatory effect of *Portulaca oleracea* in airways of asthmatic patients.** *J Ethnopharmacol* 2004, **93**:57-62.

83. Braga LC, Shupp JW, Cummings C, Jett M, Takahashi JA, Carmo LS, Chartone-Souza E, Nascimento AM: **Pomegranate extract inhibits *Staphylococcus aureus* growth and subsequent enterotoxin production.** *J Ethnopharmacol* 2005, **96**:335-9.
84. Hosseinzadeh H, Nourbakhsh M: **Effect of *Rosmarinus officinalis* L. aerial parts extract on morphine withdrawal syndrome in mice.** *Phytother Res* 2003, **17**:938-41.
85. Lukaczer D, Darland G, Tripp M, Liska D, Lerman RH, Schiltz B, Bland JS: **A pilot trial evaluating Meta050, a proprietary combination of reduced iso-alpha acids, rosemary extract and oleonic acid in patients with arthritis and fibromyalgia.** *Phytother Res* 2005, **19**:864-9.
86. Pramod SN, Venkatesh YP: **Allergy to eggplant (*Solanum melongena*).** *J Allergy Clin Immunol* 2004, **113**:171-3.
87. Latha M, Ramkumar KM, Pari L, Damodaran PN, Rajeshkannan V, Suresh T: **Phytochemical and antimicrobial study of an anti-diabetic plant: *Scoparia dulcis* L.** *J Med Food* 2006, **9**:391-4.
88. Sarin R: **Insecticidal activity of callus culture of *Tagetes erecta*.** *Fitoterapia* 2004, **75**:62-4.
89. Dharmagadda VS, Naik SN, Mittal PK, Vasudevan P: **Larvicidal activity of *Tagetes patula* essential oil against three mosquito species.** *Bioresour Technol* 2005, **96**:1235-40.
90. Ushanandini S, Nagaraju S, Harish Kumar K, Vedavathi M, Machiah DK, Kemparaju K, Vishwanath BS, Gowda TV, Girish KS: **The anti-snake venom properties of *Tamarindus indica* (leguminosae) seed extract.** *Phytother Res* 2006, **20**:851-8.
91. Komutarin T, Azadi S, Butterworth L, Keil D, Chitsomboon B, Suttajit M, Meade BJ: **Extract of the seed coat of *Tamarindus indica* inhibits nitric oxide production by murine macrophages *in vitro* and *in vivo*.** *Food Chem Toxicol* 2004, **42**:649-58.
92. Lin YL, Tsai YL, Kuo YH, Liu YH, Shiao MS: **Phenolic compounds from *Tournefortia sarmentosa*.** *J Nat Prod* 1999, **62**:1500-3.
93. Roque-Abelo L: **Chemical defense and aposematism: the case of *Utetheisa galapagensis*.** *Chemoecology* 2002, **12**:153-157.
94. Liu IM, Liou SS, Cheng JT: **Mediation of beta-endorphin by myricetin to lower plasma glucose in streptozotocin-induced diabetic rats.** *J Ethnopharmacol* 2006, **104**:199-206. abelmoschus
95. Sun HX: **Adjuvant effect of *Achyranthes bidentata* saponins on specific antibody and cellular response to ovalbumin in mice.** *Vaccine* 2006, **24**:3432-9.
96. Hu SL, Zhang HQ, Chan K, Mei QX: **Studies on the toxicity of *Aristolochia manshuriensis* (Guanmuton).** *Toxicology* 2004, **198**:195-201.
97. Wang N, Yao X, Ishii R, Kitanaka S: **Bioactive sucrose esters from *Bidens parviflora*.** *Phytochemistry* 2003, **62**:741-6.
98. Zu Yg, Fu Yj, Liu W, Hou Cl: **Simultaneous determination of four flavonoids in *Pigeonpea* [*Cajanus cajan* (L.) Millsp.] leaves using RP-LC-DAD.** *Chromatographia* 2006, **63**:499.
99. Jiang TF, Lv ZH, Wang YH: **Separation and determination of anthraquinones in *Cassia obtusifolia* (Leguminosae) by micellar electrokinetic capillary electrophoresis.** *J Sep Sci* 2005, **28**:2225-9.
100. Wei Chen, Xiao-Dong Yang, Jing-Feng Zhao, Jing-Hua Yang, Hong-Bin Zhang, Zi-Yan Li, Liang Li: **Three New, 1-Oxygenated ent-8,9-Secokaurane Diterpenes from *Croton kongensis*.** *Helvetica Chimica Acta* **89**:537-541.
101. Du A, Hu S: **Effects of a herbal complex against *Eimeria tenella* infection in chickens.** *J Vet Med B Infect Dis Vet Public Health* 2004, **51**:194-7. eclipta
102. Yang SP, Cheng JG, Huo J, Jiang HL, Chen KX, Yue JM: **Seven New Sesquiterpene Lactones from *Eupatorium chinense*.** *Chinese Journal of Chemistry* 2005, **23**:1530-1536.
103. Fong WP, Poon YT, Wong TM, Mock JW, Ng TB, Wong RN, Yao QZ, Yeung HW: **A highly efficient procedure for purifying the ribosome-inactivating proteins alpha- and beta-momorcharins from *Momordica charantia* seeds, N-terminal sequence comparison and establishment of their N-glycosidase activity.** *Life Sci* 1996, **59**:901-9.
104. Li YF, Gong ZH, Yang M, Zhao YM, Luo ZP: **Inhibition of the oligosaccharides extracted from *Morinda officinalis*, a Chinese traditional herbal medicine, on the corticosterone induced apoptosis in PC12 cells.** *Life Sci* 2003, **72**:933-42.
105. Wanxing Wei, Yuanjiang Pan, Yaozu Chen, Cuiwu Lin, Tengyou Wei, Shukai Zhao: **Carboxylic Acids from *Phyllanthus urinaria*.** *Chemistry of Natural Compounds* 2005, **41**:17-21.
106. Li Y, Ooi LS, Wang H, But PP, Ooi VE: **Antiviral activities of medicinal herbs traditionally used in southern mainland China.** *Phytother Res* 2004, **18**:718-22. portulaca
107. Cao JH, Qi YP: **Studies on the chemical constituents of the herb huanghuaren (*Sida acuta* Burm. f.).** *Zhongguo Zhong Yao Za Zhi* 1993, **18**:681-2. [Article in Chinese]
108. Helen Atteck, Philip Atteck: **Stress of Weather. A Collection of Original Source Documents Relating to a voyage from China to Trinidad, West Indies in 1862 in conjunction with a family chronicle** Wanata Enterprises, Ontario, Canada; 2000.
109. Millett Trevor M: **The Chinese in Trinidad Port of Spain, Trinidad:** Inprint Caribbean; 1993.
110. Brodwin P: **The cultural politics of biomedicine in the Caribbean. Review article.** *New West Indian Guide/Nieuwe West-Indische Gids* 1998, **72**:101-109.
111. Gauri V, Cercone J, Briceno R: **Separating financing from provision: evidence from 10 years of partnership with health cooperatives in Costa Rica.** *Health Policy Plan* 2004, **19**:292-301.
112. Fischer K: **On Building Alliances: Credit Union Service Organizations.** *The Anthill* 2005, **5**(1): [<http://bcics.uvic.ca/anthill/v5/i1/cuso.htm>].

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

