



## Application of response surface methodology in medium optimization for pyruvic acid production of *Torulopsis glabrata* TP19 in batch fermentation

ZHANG Jian<sup>†</sup>, GAO Nian-fa

(Tianjin Key Lab of Industrial Microbiology, College of Biotechnology, Tianjin University of Science and Technology, Tianjin 300457, China)

<sup>†</sup>E-mail: zj96sk@163.com

Received Apr. 12, 2006; revision accepted July 26, 2006

**Abstract:** Response surface methodology (RSM) was used to optimize the fermentation medium for enhancing pyruvic acid production by *Torulopsis glabrata* TP19. In the first step of optimization, with Plackett-Burman design, ammonium sulfate, glucose and nicotinic acid were found to be the important factors affecting pyruvic acid production significantly. In the second step, a 2<sup>3</sup> full factorial central composite design and RSM were applied to determine the optimal concentration of each significant variable. A second-order polynomial was determined by the multiple regression analysis of the experimental data. The optimum values for the critical components were obtained as follows: ammonium sulfate 0.7498 (10.75 g/L), glucose 0.9383 (109.38 g/L) and nicotinic acid 0.3633 (7.86 mg/L) with a predicted value of maximum pyruvic acid production of 42.2 g/L. Under the optimal conditions, the practical pyruvic acid production was 42.4 g/L. The determination coefficient ( $R^2$ ) was 0.9483, which ensures adequate credibility of the model. By scaling up fermentation from flask to jar fermentor, we obtained promising results.

**Key words:** Response surface methodology, *Torulopsis glabrata*, Pyruvic acid, Fermentation, Medium optimization  
**doi:**10.1631/jzus.2007.B0098      **Document code:** A      **CLC number:** TQ922

### INTRODUCTION

Pyruvic acid, also known as 2-oxopropanoic acid,  $\alpha$ -ketopropionic acid or acetylformic acid, is the most important  $\alpha$ -oxocarboxylic acid. It plays a central role in energy metabolism in living organisms. Industrially, it is used mainly as a starting material in the biosynthesis of pharmaceuticals, such as L-tryptophan, L-tyrosine, L-dihydroxyphenylalanine (Uchio *et al.*, 1976), L-phenylacetylcarbinol (Rosche *et al.*, 2002), and N-acetyl-D-neuraminic acid (Mahmoudian *et al.*, 1997). It is also employed in the production of crop protection agents, antioxidant, polymers, cosmetics and food additives. Calcium pyruvate also has a strong effect in reducing fat because it can accelerate the metabolism of fatty acids in the human body (Roufs, 1996), and presently it serves as a fat burner in the food industry. As it is widely used in drug, agrochemical, chemical and food industries, the commercial demand for pyruvic acid has

been expanding (Yonehara and Miyata, 1994).

Basically, there are four different known approaches to produce pyruvic acid: (1) chemical synthesis and oxidation (Tsujino *et al.*, 1992; Ai and Ohdan, 1995), (2) enzymes (Burdick and Schaeffer, 1987; Eisenberg *et al.*, 1997), (3) resting cells (Izumi *et al.*, 1982; Schinschel and Simon, 1993; Ogawa *et al.*, 2001), and (4) fermentation processes (Yokota *et al.*, 1994; Li *et al.*, 2001a; 2001b; Causey *et al.*, 2004). In comparison with these approaches, the fermentation processes is regarded as one of the most promising routes for the production of pyruvic acid.

Recent research efforts have focused on further process development (optimization) and scale-up of pyruvic acid production. As we all known, medium components play a very important role in enhancing the pyruvic acid accumulation, therefore, medium optimization study is very important. In general, medium optimization by the traditional 'one-factor-at-a-time' technique was used (Gokhade *et al.*, 1991). This

method is not only laborious and time consuming but also often leads to an incomplete understanding of the system behaviour, resulting in confusion and a lack of predictive ability.

Response surface methodology (RSM) is a powerful and efficient mathematical approach widely applied in the optimization of fermentation process, e.g. media components on enzyme production (Adinarayana and Ellaiah, 2002; Park *et al.*, 2002; Pui *et al.*, 2002), production of other metabolites (Zhang *et al.*, 1996; Sunitha *et al.*, 1998; Sadhukhan *et al.*, 1999; Hujanen *et al.*, 2001), spore production (Yu *et al.*, 1997) and biomass production optimization (Lhomme and Roux, 1991). It can give information about the interaction between variables, provide information necessary for design and process optimization, and give multiple responses at the same time.

In this study, optimization of fermentation media of *Torulopsis glabrata* TP19 was investigated using RSM to increase the pyruvic acid production. In the first optimization step, a Plackett-Burman (PB) design was used to determine the likely effects of medium components on pyruvic acid production. In the second step, the factors that had significant effects were optimized using a central composite design (CCD) and response surface analysis. Based on optimal medium, scale-up was carried out in a 5-L jar fermentor.

## MATERIALS AND METHODS

### Microorganism

*Torulopsis glabrata* TG-06 wild type was obtained from Tianjin Key Lab of Industrial Microbiology, China. From this a mutation *Torulopsis glabrata* TP19 (NA<sup>-</sup>+Bio<sup>-</sup>+TPP<sup>-</sup>+Pdx<sup>-</sup>), which was used as a pyruvic acid producer in this research, was developed in the authors' laboratory. It was maintained on agar slants containing (per liter): 30 g glucose, 10 g peptone, 20 g agar, 1 g KH<sub>2</sub>PO<sub>4</sub> and 0.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O. The organism was subcultured over the interval of 2 months and stored at 4 °C.

### Medium

The seed medium consisted of (per liter): 30 g glucose, 5 ml corn steep liquor, 10 g peptone, 1 g

KH<sub>2</sub>PO<sub>4</sub> and 0.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O. The components in the fermentation medium were added according to the design of each experiment. Trace element solution containing (per liter of 2 mol/L HCl): 2 g CaCl<sub>2</sub>·2H<sub>2</sub>O, 2 g FeSO<sub>4</sub>·7H<sub>2</sub>O, 5 g ZnCl<sub>2</sub>, 0.2 g MnCl<sub>2</sub>·4H<sub>2</sub>O and 0.5 g CuSO<sub>4</sub>·5H<sub>2</sub>O. The initial pH of all media was adjusted to 5.0. In the fermentor, the pH was automatically controlled at 5.0 with 8 mol/L NaOH solution. All vitamins were sterilized by microfiltration; and CaCO<sub>3</sub> was sterilized by dry-heat sterilization at 160 °C for 30 min before being added to the medium.

### Cultivation

A loopful of cells from the slant were transferred into a 250-ml conical flask containing 30-ml seed medium and incubated for 20 h on a rotary shaker operating at 200 r/min at 30 °C. It was then inoculated either into a 500-ml flask containing 50 ml fermentation medium or into a 5-L jar fermentor (Biostat M.B. Braun Co., Germany) containing 3 L fermentation medium. The inoculum size was 10% (v/v). In shake flask culture, it was incubated on a rotary shaker operating at 200 r/min at 30 °C. In fermentor culture, it was incubated at 30 °C, the agitation speed was controlled at 400 r/min. The air flow rate was 1.2 L/min. Foam formation was prevented by adding sterile polypropylene glycol solution (10% aqueous PPG 2000).

### Sample preparation

After 60 h, fermentation broth was collected. The cells were removed by centrifugation at 12000 r/min for 10 min, the supernatant was collected and further diluted by a factor of 100 with Milli-Q water. This diluent was then filtered through a 0.45 µm millipore filter prior to injection into the chromatographic column.

### HPLC

Pyruvic acid concentration was measured by HPLC (agilent 1100 series) using a Bio-Rad HPX-87H Aminex column 7.8 mm×300 mm and a UV detector at 210 nm with 0.005 mol/L H<sub>2</sub>SO<sub>4</sub> as the eluent at a flow rate of 0.6 ml/min at 55 °C.

### Experimental designs and data analysis

#### 1. Plackett-Burman (PB) design

PB design, a very useful tool, was used to screen

' $n$ ' variables in just ' $n+1$ ' number of experiments (Plackett and Burman, 1946; Rama *et al.*, 1999; Ghanem *et al.*, 2000). There will be a tremendous decrease in total experiments. In this part, the PB design was used to evaluate the relative importance of various nutrients for pyruvic acid production in batch fermentation. This design does not consider the interaction effects among the variables and is used to screen the important variables affecting the pyruvic acid production. The experimental design for screening of medium components is shown in Tables 1 and 2. Each variable was set at two levels, that is, high level and low level. The high level of each variable was set far enough from the low level to identify which ingredients of the media have significant influence on the pyruvic acid production. Trace element solution was constantly maintained at 5 ml/L. The design matrix (Table 2) was developed using an SAS package, version 8.01.

**Table 1 Medium components for screening using a PB design**

Variables	Levels	
	Low (-1)	High (+1)
A Glucose (g/L)	80	100
B Ammonium sulfate (g/L)	8	10
C $\text{KH}_2\text{PO}_4$ (g/L)	1.0	1.5
D $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (g/L)	0.5	1.5
E Nicotinic acid (mg/L)	6.0	7.5
F Thiamine-HCl ( $\mu\text{g/L}$ )	20	25
G Biotin ( $\mu\text{g/L}$ )	10	15
H Pyridoxine-HCl (mg/L)	0.4	0.6

**Table 2 The experimental design using the PB method for screening of medium components**

Run	Variables								Pyruvic acid yield (g/L)
	A	B	C	D	E	F	G	H	
1	1	-1	1	-1	-1	-1	1	1	32.5
2	1	1	-1	1	-1	-1	-1	1	36.3
3	-1	1	1	-1	1	-1	-1	-1	36.9
4	1	-1	1	1	-1	1	-1	-1	35.0
5	1	1	-1	1	1	-1	1	-1	38.5
6	1	1	1	-1	1	1	-1	1	40.2
7	-1	1	1	1	-1	1	1	-1	36.8
8	-1	-1	1	1	1	-1	1	1	33.4
9	-1	-1	-1	1	1	1	-1	1	32.6
10	1	-1	-1	-1	1	1	1	-1	35.3
11	-1	1	-1	-1	-1	1	1	1	34.1
12	-1	-1	-1	-1	-1	-1	-1	-1	31.8

## 2. Central composite design (CCD)

CCD is one of the response surface methodologies (Chakravarti and Sahai, 2002). After the identification of the components affecting the pyruvic acid yield significantly, a CCD was adopted to optimize the major variables (ammonium sulfate, glucose and nicotinic acid), which were selected through PB design. Coded levels for independent variables are presented in Table 3.

**Table 3 Levels of the variables tested in CCD**

Variables	Coded levels				
	-1.682	-1	0	1	1.682
Ammonium sulfate, $X_1$	8.3	9.0	10.0	11.0	11.7
Glucose, $X_2$	83.2	90.0	100.0	110.0	116.8
Nicotinic acid, $X_3$	5.8	6.5	7.5	8.5	9.2

$$X_1=(x_1-10)/1; X_2=(x_2-100)/10; X_3=(x_3-7.5)/1$$

For statistical calculations the variables  $X_i$  were coded as  $x_i$  according to Eq.(1)

$$X_i=(x_i-x_0)/\Delta x_i, i=1, 2, 3, \dots, K, \quad (1)$$

where  $X_i$  is (dimensionless) coded value of the real variable  $x_i$ ,  $x_0$  is the real value of  $X_i$  at the center point (zero) level, and the  $\Delta x_i$  is the step change value.

A  $2^3$ -factorial CCD, with six axial points ( $a=1.682$ ) and six replications at the centre points ( $n_0=6$ ) leading to a total number of 20 experiments was employed (Table 4) for the optimization of the three chosen medium variables. Pyruvic acid yield ( $Y$ , g/L) was used as the dependent output variable.

Second degree polynomials, Eq.(2), which includes all interaction terms, were used to calculate the predicted response.

$$Y = \beta_0 + \sum \beta_i X_i + \sum \beta_{ii} X_i^2 + \sum \beta_{ij} X_i X_j, \quad (2)$$

where  $Y$  represents response variable,  $\beta_0$  is the interception coefficient,  $\beta_i$ , coefficient of the linear effect,  $\beta_{ii}$ , the coefficient of quadratic effect and  $\beta_{ij}$ , the coefficient of interaction effect.

The experiments were performed in duplicate with the mean values taken for analysis. An SAS package, version 8.01, was used for multiple regression analysis of the experimental data obtained.  $F$ -test was employed to evaluate the statistical significance

**Table 4** Experimental design matrix and results of CCD

Run	Factor			Pyruvic acid yield, $Y$ (g/L)
	$X_1$	$X_2$	$X_3$	
1	-1	-1	-1	37.9
2	-1	-1	1	38.2
3	-1	1	-1	39.8
4	-1	1	1	40.5
5	1	-1	-1	39.3
6	1	-1	1	40.0
7	1	1	-1	40.2
8	1	1	1	41.7
9	-1.682	0	0	39.8
10	1.682	0	0	41.8
11	0	-1.682	0	38.2
12	0	1.682	0	42.4
13	0	0	-1.682	39.5
14	0	0	1.682	39.8
15	0	0	0	41.3
16	0	0	0	41.2
17	0	0	0	41.5
18	0	0	0	41.6
19	0	0	0	41.4
20	0	0	0	41.4

of the quadratic polynomial. The multiple coefficients of correlation  $R$  and the determination coefficient of correlation  $R^2$  were calculated to evaluate the performance of the regression equation. The optimum levels of the selected variables were obtained by solving the regression equation using a multi-stage Monte-Carlo optimization (Conley, 1984) program and also by analysing the response-surface plots (Khuri and Cornell, 1987).

## RESULTS AND DISCUSSION

### Plackett-Burman (PB) design

The first optimization step was using a 12-run PB design to identify the significant factors for pyruvic acid production by *Torulopsis glabrata* TP19. According to the resulting effects of these eight variables on pyruvic acid concentration and the associated significant levels presented in Table 5, it can be seen that with the increase in the concentration of ammonium sulfate, glucose, nicotinic acid,  $\text{KH}_2\text{PO}_4$ , thiamine-HCl and  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , all have positive

effects on pyruvic acid production. With increase in the pyridoxine-HCl levels, biotin has negative effects on pyruvic acid production. With the help of relative ranking, ammonium sulfate, glucose and nicotinic acid within the tested limits were selected for further optimization, which had the most significant effects on pyruvic acid production.

**Table 5** Coefficients,  $t$  values and significance levels calculated from the pyruvic acid yield obtained in the screening experiments

Code	Coefficient	$t$ value	$P >  T $	Ranking
A	1.0167	4.1157	0.0260	2*
B	1.8500	7.4893	0.0049	1**
C	0.5167	2.0916	0.1276	4
D	0.1500	0.6072	0.5866	8
E	0.8667	3.5085	0.0392	3*
F	0.3833	1.5518	0.2185	6
G	-0.1833	-0.7422	0.5118	7
H	-0.4333	-1.7543	0.1777	5

\* Statistically significant at 95% of probability level; \*\* Statistically significant at 99% of probability level

The positive effects of ammonium sulfate and glucose were, probably, caused by the requirement of a large quantity of substrate to synthesize cells. It is possible that the high concentration of nicotinic acid could contribute to the improvement of the glucose consumption rate. This may be due to the fact that nicotinic acid is a component of NAD, a cofactor of glycolysis (Miyata and Yonehara, 1996), resulting in both good cell growth and satisfactory pyruvic acid production.

The PB design was proved to be a powerful tool to rapidly determine the effects of medium constituents on pyruvic acid production of *Torulopsis glabrata* TP19. However the optimal concentrations of medium components that significantly affect pyruvic acid production could not be obtained. Further work need to be done to find out this information.

### Central composite design (CCD)

This is a very useful tool to determine the optimal level of medium constituents and their interaction. Based on the PB design, where ammonium sulfate, glucose and nicotinic acid were selected for their significant effects on the pyruvic acid production, a CCD was used for further optimization. Table 3 gave the variation levels at which these components were

supplemented to pyruvic acid production. Other nutrients concentrations were set at their centre point tested in the PB design. Table 4 gives the design and results of experiments carried out by the CCD design. The results obtained were submitted to analysis of variance on SAS package, with the regression model given as

$$Y = 41.416 + 0.598X_1 + 1.015X_2 + 0.27X_3 - 0.31X_1^2 - 0.2X_1X_2 + 0.15X_1X_3 - 0.49X_2^2 + 0.15X_2X_3 - 0.72X_3^2, \quad (3)$$

where  $Y$  is the response value, that is, the pyruvic acid production, and  $X_1$ ,  $X_2$  and  $X_3$  are the coded levels of ammonium sulfate, glucose and nicotinic acid, respectively.

The analysis of variance of the quadratic regression model demonstrated that Eq.(3) was a highly significant model, as was evident from the Fisher's  $F$ -test with a very low probability value [ $(P \text{ model} > F) = 0.0001$ ]. The model's goodness of fit was checked by determination coefficient ( $R^2$ ). In this case, the value of the determination coefficient ( $R^2 = 0.9483$ ) indicated that only 5.17% of the total variations were not explained by the model. The value of the adjusted determination coefficient [ $\text{Adj}(R^2) = 0.9019$ ] was also very high in supporting the high significance of the model. Among the model terms,  $X_1$ ,  $X_2$ ,  $X_2^2$  and  $X_3^2$  were significant with a probability of 99%;  $X_3$  and  $X_1^2$  were significant with a probability of 95% (Table 6). The interaction between  $X_1$ ,  $X_2$  and  $X_3$ , however,


**Table 6** Analysis of variance for the fitted quadratic polynomial model

Term	Effect	SS	DF	F ratio	P>F
$X_1$	0.598	4.8799	1	28.6470	0.0003**
$X_2$	1.015	14.0734	1	82.6170	0.0001**
$X_3$	0.270	1.0049	1	5.8992	0.0355*
$X_1^2$	-0.310	1.4161	1	8.3134	0.0163*
$X_1X_2$	0.200	0.3200	1	1.8786	0.2005
$X_1X_3$	0.150	0.1800	1	1.0567	0.3282
$X_2^2$	-0.490	3.4637	1	20.3340	0.0011**
$X_2X_3$	0.150	0.1800	1	1.0567	0.3282
$X_3^2$	-0.720	7.4721	1	43.8650	0.0001**
Model	—	31.2741	9	20.3990	0.0001
Error	—	1.7034	10	—	—
Total	—	32.9775	19	—	—

\* Statistically significant at 95% of probability level; \*\* Statistically significant at 99% of probability level

had no significant influence on pyruvic acid production.

The fitted response for the above regression model was plotted in Fig.1. 3D graphs were generated for the pair-wise combination of the three factors while keeping the other one at its optimum levels for pyruvic acid production. Graphs are given here to highlight the roles played by various factors and also to emphasize the roles played by the physical constraints vis-a-vis the biosynthetic aspects in the final yield of the pyruvic acid.



**Fig.1** Effects of (a) ammonium sulfate ( $X_1$ ) and glucose ( $X_2$ ), (b) ammonium sulfate ( $X_1$ ) and nicotinic acid ( $X_3$ ), (c) nicotinic acid ( $X_3$ ) and glucose ( $X_2$ ) and their interactive effect on the pyruvic acid production ( $Y$ ) with other nutrient set at centre level

The predicted optimum levels of the tested variables, namely, ammonium sulfate, glucose and nicotinic acid were obtained by applying regression analysis on Eq.(3) using SAS package software, version 8.01. The optimal levels were as follows:  $X_1=0.7498$  (10.75 g/L),  $X_2=0.9383$  (109.38 g/L),  $X_3=0.3633$  (7.86 mg/L) with the corresponding  $Y=42.2$  g/L. Verification of the predicted values was conducted by using optimal medium in inoculation experiments. The practical corresponding response was 42.4 g/L. This result corroborated the validity and the effectiveness of this model.

In an attempt to approximate industrial conditions for pyruvic acid production, scale-up was carried out in a jar fermentor by using optimal medium. In this way, we achieved a volumetric productivity of 0.911 g pyruvate/(L·h), a pyruvate/glucose yield of 0.53 g/g, and pyruvic acid production of 54.6 g/L. These results are encouraging for optimization under pilot scale or industrial scale conditions.

## CONCLUSION

RSM was performed to optimize the medium components for pyruvic acid production of *Torulopsis glabrata* TP19. A highly significant quadratic polynomial obtained by the CCD was very useful for determining the optimal concentrations of constituents that have significant effects on pyruvic acid production.

The optimal supplementary components (per liter) consisted of 109.38 g glucose, 10.75 g ammonium sulfate, 1.25 g  $\text{KH}_2\text{PO}_4$ , 1 g  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ , 7.86 mg nicotinic acid, 22.5  $\mu\text{g}$  thiamine·HCl, 12.5  $\mu\text{g}$  biotin, 0.5 mg pyridoxine·HCl and 5 ml trace element solution. Under the optimal condition, 42.2 g/L pyruvic acid could be produced in theory and 42.4 g/L pyruvic acid in practical experiment. Scale-up in jar fermentor produced adequate results, although accurate moisture and temperature control, optimized  $\text{O}_2$  mass transfer, as well as fed batch studies, promise further process improvement.

## References

Adinarayana, K., Ellaiah, P., 2002. Response surface optimization of the critical medium components for this production of alkaline protease by a newly isolated *Bacillus* sp. *J. Pharm. Pharmaceut. Sci.*, **5**(3):272-278.

- Ai, M., Ohdan, K., 1995. Formation of pyruvic acid by oxidative dehydrogenation of lactic acid. *Chem. Lett.*, **24**(5): 405. [doi:10.1246/cl.1995.405]
- Burdick, B.A., Schaeffer, J.R., 1987. Co-immobilized coupled enzyme systems on nylon mesh capable of gluconic and pyruvic acid production. *Biotechnol. Lett.*, **9**(4):253-258. [doi:10.1007/BF01027159]
- Causey, T.B., Shanmugam, K.T., Yomano, L.P., Inram, L.O., 2004. Engineering *Escherichia coli* for efficient conversion of glucose to pyruvate. *Proc. Natl. Acad. Sci. USA*, **101**(8):2235-2240. [doi:10.1073/pnas.0308171100]
- Chakravarti, R., Sahai, V., 2002. Optimization of compactin production in chemically defined production medium by *Penicillium citrinum* using statistical methods. *Process Biochem.*, **38**(4):481-486. [doi:10.1016/S0032-9592(02)00138-3]
- Conley, W.C., 1984. Computer Optimization Techniques. Petrocelli Books, Princeton, NJ, p.147-163.
- Eisenberg, A., Seip, J.E., Gavagan, J.E., Payne, M.S., Anton, D.L., DiCosimo, R., 1997. Pyruvic acid production using methylotrophic yeast transformants as catalyst. *J. Mol. Catal. B: Enzymatic*, **2**(4-5):223-232. [doi:10.1016/S1381-1177(96)00021-5]
- Ghanem, N.B., Yusef, H.H., Mahrouse, H.K., 2000. Production of *Aspergillus terreus* xylanase in solid-state cultures: application of the Plackett-Burman experimental design to evaluate nutritional requirements. *Bioresour. Technol.*, **73**(2):113-121. [doi:10.1016/S0960-8524(99)00155-8]
- Gokhade, D.V., Patil, S.G., Bastawde, K.B., 1991. Optimization of cellulase production by *Aspergillus niger* NCIM 1207. *Appl. Biochem. Biotechnol.*, **30**(2):99-109.
- Hujanen, M., Linko, S., Linko, Y.Y., Leisola, M., 2001. Optimization of media and cultivation conditions for L-(+)-lactic acid production by *Lactobacillus casei* NRRL B-441. *Appl. Microbiol. Biotechnol.*, **56**(1-2): 126-130. [doi:10.1007/s002530000501]
- Izumi, Y., Matsumura, Y., Tani, Y., Yamada, H., 1982. Pyruvic acid production from 1,2-propanediol by thiamin-requiring *Acinetobacter* sp. 80-M. *Agric. Biol. Chem.*, **46**(3):2673-2679.
- Khuri, A.I., Cornell, J.A., 1987. Response Surfaces: Design and Analyses. Dekker, New York.
- Lhomme, B., Roux, J.C., 1991. Utilization of experimental designs for optimization of *Rhizopus arrhizus* culture. *Bioresour. Technol.*, **35**(3):301-312. [doi:10.1016/0960-8524(91)90129-8]
- Li, Y., Chen, J., Lun, S.Y., 2001a. Efficient pyruvate production by a multi-vitamin auxotroph of *Torulopsis glabrata*: key role and optimization of vitamin levels. *Appl. Microbiol. Biotechnol.*, **55**(6):680-685. [doi:10.1007/s002530100598]
- Li, Y., Chen, J., Lun, S.Y., 2001b. Biotechnological production of pyruvic acid. *Appl. Microbiol. Biotechnol.*, **57**(4):451-459. [doi:10.1007/s002530100804]
- Mahmoudian, M., Noble, D., Drake, C.S., Middleton, R.F., Montgomery, D.S., Picrcey, J.E., Ramlakhan, D., Todd, M., Dawson, M., 1997. An efficient process for produc-

- tion of N-acetylneuraminic acid using N-acetylneuraminic acid aldolase. *Enzyme Microb. Technol.*, **20**(5):393-400. [doi:10.1016/S0141-0229(96)00180-9]
- Miyata, R., Yonehara, T., 1996. Improvement of fermentative production of pyruvate from glucose by *Torulopsis glabrata* IFO 0005. *J. Ferment. Bioeng.*, **82**(5):475-479. [doi:10.1016/S0922-338X(97)86986-3]
- Ogawa, J., Soong, C.L., Masashi, I., Shimizu, S., 2001. Enzymatic production of pyruvate from fumarate—an application of microbial cyclic-imide-transforming pathway. *J. Mol. Catal. B: Enzymatic*, **11**(4-6):355-359. [doi:10.1016/S1381-1177(00)00024-2]
- Park, Y.S., Kang, S.W., Lee, J.S., Hong, S.I., Kim, S.W., 2002. Xylanase production in solid state fermentation by *Aspergillus niger* mutant using statistical experimental designs. *Appl. Microbiol. Biotechnol.*, **58**(6):761-766. [doi:10.1007/s00253-002-0965-0]
- Plackett, R.L., Burman, J.P., 1946. The design of optimum multifactorial experiments. *Biometrika*, **33**(4):305-325. [doi:10.2307/2332195]
- Puri, S., Beg, Q.K., Gupta, R., 2002. Optimization of alkaline protease from *Bacillus* sp. by response surface methodology. *Curr. Microbiol.*, **44**(4):286-290. [doi:10.1007/s00284-001-0006-8]
- Rama Mohan Reddy, P., Reddy, G., Seenayya, G., 1999. Production of thermostable  $\beta$ -amylase and pullulanase by *Clostridium thermosulfurogenes* SV2 in solid-state fermentation: screening of nutrients using Plackett-Burman design. *Bioprocess Eng.*, **21**(2):175-179. [doi:10.1007/s004490050659]
- Rosche, B., Leksawasdi, N., Sandford, V., Breuer, M., Hauer, B., Rogers, P., 2002. Enzymatic (R)-phenylacetylcarbinol production in benzaldehyde emulsions. *Appl. Microbiol. Biotechnol.*, **60**(1-2):94-100. [doi:10.1007/s00253-002-1084-7]
- Roufs, J.B., 1996. Pyruvate: does it amp endurance and burn more fat? *Muscle Fitness*, **57**(2):195-197.
- Sadhukhan, A.K., Ramana Murthy, M.V., Ajaya Kumar, R., Mohan, E.V.S., Vandana, G., Bhar, C., Venkateswara Rao, K., 1999. Optimization of mycophenolic acid production in solid-state fermentation using response surface methodology. *J. Ind. Microbiol. Biotechnol.*, **22**(1):33-38. [doi:10.1038/sj.jim.2900597]
- Schinschel, C., Simon, H., 1993. Preparation of pyruvate from (R)-lactate with *Proteus* species. *J. Biotechnol.*, **31**(2):191-203. [doi:10.1016/0168-1656(93)90160-O]
- Sunitha, I., Subba Rao, M.V., Ayyanna, C., 1998. Optimization of medium constituents and fermentation conditions for the production of L-glutamic acid by the co-immobilized whole cells of *Micrococcus glutamicus* and *Pseudomonas reptilivora*. *Bioprocess Eng.*, **18**(5):353-359. [doi:10.1007/PL00008995]
- Tsujino, T., Ohigashi, S., Sugiyama, S., Kawashiro, K., Hayashi, H., 1992. Oxidation of propylene glycol and lactic acid to pyruvic acid in aqueous phase catalyzed by lead modified alladium-on-carbon and related systems. *J. Mol. Catal.*, **71**(1):25-35. [doi:10.1016/0304-5102(92)80005-2]
- Uchio, R., Kikuchi, K., Hirose, Y., 1976. Process for Producing Pyruvic Acid by Fermentation. US Patent, No. 3993 543.
- Yokota, A., Shimizu, H., Terasawa, Y., Takaoka, N., 1994. Pyruvic acid production by a lipoic acid auxotroph of *Escherichia coli* W1485. *Appl. Microbiol. Biotechnol.*, **41**(6):638-643. [doi:10.1007/BF00167278]
- Yonehara, T., Miyata, R., 1994. Fermentative production of pyruvate from glucose by *Torulopsis glabrata*. *J. Ferment. Bioeng.*, **78**(2):155-159. [doi:10.1016/0922-338X(94)90255-0]
- Yu, X., Hallet, S.G., Sheppard, J., Watson, A.K., 1997. Application of the Plackett-Burman experimental design to evaluate nutritional requirements for the production of *Colletotrichum coccodes* spores. *Appl. Microbiol. Biotechnol.*, **47**(3):301-305. [doi:10.1007/s002530050930]
- Zhang, J., Marcin, C., Shifflet, M.A., Salmon, P., Brix, T., Greasham, R., Buokland, B., Chartrain, M., 1996. Development of a defined medium fermentation process for physotigmine production by *Streptomyces griseofuscus*. *Appl. Microbiol. Biotechnol.*, **44**(5):568-575. [doi:10.1007/s002530050601]



Editors-in-Chief: Wei YANG & Peter H. BYERS  
ISSN 1673-1581 (Print); ISSN 1862-1783 (Online), monthly

*Journal of Zhejiang University*

SCIENCE B

www.zju.edu.cn/jzus; www.springerlink.com

jzus@zju.edu.cn

**JZUS-B focuses on "Biomedicine, Biochemistry & Biotechnology"**

**JZUS-B online in PMC:** <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=371&action=archive>