

Remodelling

Estimation of the age of a fracture by this criterion is very difficult because initial deformity, the amount of callus produced, and age of the child are major variables. In the young infant with a stable, undisplaced fracture the remodelling process may be complete at three months while the older child with an angular deformity or a markedly displaced fracture may continue to remodel for two years.

Intracranial haemorrhage

Computed tomography is the primary diagnostic method for the evaluation of head trauma and its value in the management of the abused child is well documented.²⁵⁻²⁹ The appearance of haematoma on computed tomography changes slowly with time and is well understood.³⁰ Freshly clotted blood is of higher attenuation (whiter) than brain, becoming the same density as brain from two to four weeks, and less dense than brain after this time, approaching the density of cerebrospinal fluid by six weeks. Chronic subdural haematomas associated with ventricular dilatation may be difficult to distinguish from cerebral atrophy on computed tomography but magnetic resonance imaging (MRI) easily identifies it as chronic haematoma. MRI has been shown to be superior to computed tomography in the investigation of subacute or chronic head injury³¹ but the MRI appearances during acute haemorrhage are more complex, change rapidly during the first week, and have a wide differential diagnosis.³² Nevertheless, once haemorrhage has been identified on computed tomography, MRI can be used to assess its age more accurately if that is appropriate.

Department of Radiology,
The Children's Hospital,
Ladywood Middleway,
Birmingham B16 8ET

S CHAPMAN

- O'Connor JF, Cohen J. Dating fractures. In: Kleinman PK, ed. *Diagnostic imaging of child abuse*. Baltimore: Williams and Wilkins, 1987:103-13.
- Ogden JA. Injury to the immature skeleton. In: Touloukian RJ, ed. *Pediatric trauma*. St Louis: Mosby Year Book, 1990:399-439.
- Crilly R. Longitudinal overgrowth of the chicken radius. *J Anat* 1972;112:11-8.
- Alexander CJ. Effect of growth rate on the strength of the growth plate shaft junction. *Skeletal Radiol* 1976;1:67-76.
- Keith A. The foundation of our knowledge of bone growth by Duhamel and Hunter. *Br J Surg* 1917;5:685-93.

- Stacher G, Firschein HE. Collagen and mineral kinetics in bone after fracture. *Am J Physiol* 1967;213:863-6.
- Herndon WA. Child abuse in a military population. *J Pediatr Orthop* 1983;3:73-6.
- Ebbin AJ, Gollub MH, Stein AM, Wilson MG. Battered child syndrome at the Los Angeles County General Hospital. *Am J Dis Child* 1969;118:660-7.
- Akbarnia B, Torg JS, Kirkpatrick J, Sussman S. Manifestations of the battered-child syndrome. *J Bone Joint Surg* 1974;56A:1159-66.
- Kleinman PK, Marks SC, Blackbourne B. The metaphyseal lesion in abused infants: a radiologic-histopathologic study. *AJR* 1986;146:895-905.
- Trueta J, Amato VP. The vascular contribution to osteogenesis. III. Changes in the growth cartilage caused by experimentally induced ischaemia. *J Bone Joint Surg [Br]* 1960;42B:571-87.
- Kleinman PK, Marks SC, Spevak MR, Belanger PL, Richmond JM. Extension of growth plate cartilage into the metaphysis: a sign of healing fracture in abused infants. *AJR* 1991;156:775-9.
- Maresh MM. Linear growth of long bones of extremities from infancy through adolescence. *Am J Dis Child* 1955;89:725-42.
- Cumming WA. Neonatal skeletal fractures. Birth trauma or child abuse? *Journal of the Canadian Association of Radiologists* 1979;30:30-3.
- Kleinman PK. *Diagnostic imaging of child abuse*. Baltimore: Williams and Wilkins, 1987:7-8.
- Graif M, Stahl-Kent V, Ben-Ami T, Strauss S, Amit Y, Itzhak Y. Sonographic detection of occult bone fractures. *Pediatr Radiol* 1988;18:383-5.
- Smeets AJ, Robben SGF, Meradji M. Sonographically detected costochondral dislocation in an abused child. *Pediatr Radiol* 1990;20:566-7.
- Sty JR, Starshak RJ. The role of bone scintigraphy in the evaluation of the suspected abused child. *Radiology* 1983;146:369-75.
- Merten DF, Radkowski MA, Leonidas JC. The abused child: a radiological reappraisal. *Radiology* 1983;146:377-81.
- Haase GM, Ortiz VN, Stakianakis GN, Morse TS. The value of radionuclide bone scanning in the early recognition of deliberate child abuse. *J Trauma* 1980;20:873-5.
- Jaudes PK. Comparison of radiography and radionuclide bone scanning in the detection of child abuse. *Pediatrics* 1984;73:166-8.
- Smith FW, Gilday DL, Ash JM, Green MD. Unsuspected costo-vertebral fractures demonstrated by bone scanning in the child abuse syndrome. *Pediatr Radiol* 1980;10:103-6.
- Rosenthal L, Hill RO, Chuang S. Observation on the use of ^{99m}Tc-phosphate imaging in peripheral bone trauma. *Radiology* 1976;119:637-41.
- Salter RB. Special features of fractures and dislocation in children. In: Heppenstall RB, ed. *Fracture treatment and healing*. Philadelphia: WB Saunders, 1980:190.
- Zimmerman RA, Bilaniuk LT, Genneralli T. Computed tomography of shearing injuries of the cerebral white matter. *Radiology* 1978;127:393-6.
- Zimmerman RA, Bilaniuk LT, Bruce D, Schut L, Uzzell B, Goldberg HI. Computed tomography of craniocerebral injury in the abused child. *Radiology* 1979;130:687-90.
- Tsai FY, Zee CS, Apthorp JS, Dixon GH. Computed tomography in child abuse head trauma. *CT: The Journal of Computed Tomography* 1980;4:277-86.
- Merten DF, Osborne DRS, Radkowski MA, Leonidas JC. Craniocerebral trauma in the child abuse syndrome: radiological observations. *Pediatr Radiol* 1984;14:272-7.
- Cohen RA, Kaufman RA, Myers PA, Towbin RB. Cranial computed tomography in the abused child with head injury. *AJR* 1986;146:97-102.
- Scotti G, Terbrugge K, Melançon D, Bélanger G. Evaluation of the age of subdural haematomas by computerized tomography. *J Neurosurg* 1977;47:311-5.
- Sato Y, Yuh WTC, Smith WL, Alexander RC, Kao SCS, Ellerbroek CJ. Head injury in child abuse: evaluation with MR imaging. *Radiology* 1989;173:653-7.
- Zimmerman RD, Heier LA, Snow RB, Liu DPC, Kelly AB, Deck MDF. Acute intracranial haemorrhage: intensity changes on sequential MR scans at 0.5T. *AJR* 1988;150:651-61.

Fungal skin infections

Fungal skin infections or superficial mycoses are common causes of skin disease in most age groups. They include three main diseases—dermatophyte or ringworm infections, candidosis, and pityriasis versicolor. There are significant differences, though in the epidemiology and clinical behaviour of these infections when they present in childhood. This is in part due to variations in rates of exposure in the case of dermatophytosis but in other examples may reflect real differences in the expression of host resistance.

Resistance to skin infection

The factors that determine the outcome of skin infection are divided into innate mechanisms and those that require the expression of immunological memory.^{1,2} Innate mechanisms include the capacity for epidermal cells to respond to damage to the stratum corneum by increased proliferation. Although scientific proof is lacking there is no reason to suppose that this capacity varies with age, except perhaps in photoaged skin. Likewise fungal inhibition due to reversible binding by unsaturated transferrin,³ present in sweat and serum, is not known to be affected by age. However the

presence of free fatty acids (FFAs) on the skin surface is known to affect fungi as those of medium chain length inhibit the growth of dermatophytes while facilitating the growth of pityrosporum yeasts. The composition of sebum containing these FFAs changes at puberty when the inhibitory moieties of medium chain length come to dominate. This is believed to be the explanation of the comparative rarity of scalp ringworm, tinea capitis, in postpubertal children whereas pityrosporum infections, such as pityriasis versicolor, are seen more frequently. FFAs appear to act by direct inhibition of fungal growth but they may also interfere with adhesion,⁴ the interaction between fungal cell wall receptors with keratinocytes, which is essential for subsequent penetration of the skin.

The other main processes determining the outcome of fungal invasion are antigen reception by Langerhan's cells,⁵ the accumulation of effector cells, principally neutrophils, the site of infection via the production of chemotactic factors by the fungi and the activation of T lymphocytes, predominantly helper cells.^{6,7} The means by which the latter effect fungal clearance is also not understood. Production of cytokines which amplify neutrophil killing or

epidermal growth as well as the activation of keratinocytes themselves to express HLA-DR and elaborate more cytokines such as interleukin-1 are probably equally important. The net effect is that patients with defective T lymphocyte function including AIDS patients are especially susceptible to most superficial mycoses, although unless this is accompanied by appropriate levels of exposure this defect will manifest itself by altered clinical expression of disease rather than by increased incidence. Dermatophytosis in the AIDS patient is an example where bizarre or extensive infections can occur but the diseases are not more common than in the appropriate control groups.⁸

Susceptibility to superficial mycoses may, in some cases, be genetically determined. There is evidence, for instance, for HLA linkage in the candida endocrinopathy syndrome⁹ and for an autosomal recessive mediated trait in the tropical infection, tinea imbricata.¹⁰

Dermatophytosis

The dermatophyte or ringworm fungi are mycelial organisms that produce long chains of cells, hyphae, in order to penetrate the stratum corneum.¹¹ They do this by the production of proteases which can break down keratin. These fungi cause exogenous infections originating either from other humans or animals or, more rarely, soil. They produce a characteristic group of infections known collectively as tineas whose clinical expression is largely determined by the site of infection, the identity of the organism, and the level of host response.

There are major differences in the epidemiology of infection caused by dermatophytes in children and adults. Foot infection including nail disease (onychomycosis) is rare in childhood¹² whereas it is the commonest pattern in adults. The reasons for this are not clear. Childhood nail infections, for instance, are seen occasionally, indicating that the nail is susceptible to attack. One possible reason is the lowered opportunity for exposure to a suitable environment for transmission of infection which is provided ideally by communal changing rooms. Tinea pedis begins to peak in incidence in the teens and early adult life.

The converse is true of tinea capitis, scalp ringworm, which with few exceptions, is a disease of childhood.¹³ There are two main sources of infection, animals (zoophilic infections) and other children (anthropophilic infections).¹⁴ With the former the main cause in most countries is the cat or dog ringworm, *Microsporium canis*, although cattle ringworm, *Trichophyton verrucosum* may also occur. Here exposure is sporadic depending on contact with an infected animal. Increased susceptibility in children, as mentioned previously, is the main reason for the development of the infection in childhood compared to adult life. In Europe, the Middle East and some US and Latin American cities *M canis* is now the dominant organism. By contrast ringworm infections caused by anthropophilic fungi are more common in Africa, India, and the USA.^{15 16} While host susceptibility may explain the prevalence of childhood infections to some extent exposure to an infected population is important. Carriage rates of organisms in clinically normal scalps in endemic areas may be as high as 4%.¹⁷ Some of these children may subsequently develop ringworm or the carrier status may be reversed. The combination of high rates of endemicity, mild clinical signs of infection, and inability to mount effective control measures has led to the perpetuation of hyperendemic foci for scalp ringworm. Typical causative organisms are *Trichophyton tonsurans* (USA), *Trichophyton soudanense* and *Microsporium audouini* (Africa), and *Trichophyton violaceum* (North Africa, Middle East, and the Indian subcontinent).

Scalp ringworm is easily treated with griseofulvin; most

topical antifungal treatments are ineffective on their own.¹⁸ Newer antifungal drugs such as itraconazole or terbinafine are effective, although their limitations are still not clear. Treatment of *M canis* infections with ketoconazole, however, is often unsuccessful.¹⁹ Griseofulvin is normally given in doses of 10 mg/kg daily for six to eight weeks. Though with zoophilic infections it is not necessary to keep children off school, some measure of caution is advisable with the anthropophilic fungi. Here the traditional method of control is griseofulvin combined with exclusion of infected children from school. There is no study that has examined the combined use of oral antifungals and topical applications such as ketoconazole shampoo as methods of limiting spread. With anthropophilic infections it is comparatively easy to screen for spread within classrooms as even though the organisms of the trichophyton unlike microsporium species do not usually fluoresce under ultraviolet (Wood's) light they can be grown from screening scalp samples. One method is the isolation of fungi from scalp massage brushes passed through children's hair.²⁰ Our practice is to treat infected individuals with griseofulvin and carriers, that is those with positive culture but no clinical or microscopic evidence of hair shaft invasion, with ketoconazole shampoo.

Cost in developing countries is often a contraindication to effective control. While the cost of griseofulvin is low compared with newer alternatives the price of a six week course for large numbers of children is high where resources are limited. However there is evidence that the use of single doses of up to 1 g of griseofulvin may be effective methods of eradicating infection in the majority of infected children.¹³ This method is also highly appropriate as the drug can be administered under supervision by a medical assistant or nurse.

Candidosis

Superficial infections affecting the skin and mucous membranes caused by species of the genus candida are very common in clinical practice. These fungi are normal commensals in healthy hosts, although disease is more likely where there is a defect in individual susceptibility. The carriage rates of candida in normal hosts range from 18% (range 4–66) in the mouth to 12% (range 0–33) in the gastrointestinal tract.² In children carriage rates vary considerably. For instance in newborn infants they may reach over 70% in those in neonatal intensive care units.

The commonest superficial infection is oral candidosis, which is seen in infants. Chronic oropharyngeal infection is much less common. Where it occurs it is important to consider the possibility of inherited or acquired immunodeficiency including AIDS and the rare infection chronic mucocutaneous candidosis (see below).

Chronic mucocutaneous candidosis (CMC) is a persistent superficial infection with candida species affecting principally the mouth, the skin, and nails.²¹ While originally it was hoped that a deeper understanding of the immunological basis of this syndrome would provide a potential answer to our knowledge of the nature of resistance to infection, the true nature of the immunological defect(s) in this condition is still elusive. It has become apparent though that some of the supposed defects such as reduced delayed type hypersensitivity and neutrophil killing of candida are reversible with successful antifungal treatment. This suggests that some form of immunomodulation by the organism itself may be involved and, for instance, an immunosuppressive effect of circulating cell wall mannan has been proposed as one possible mechanism.²² The expression of a human iC3b receptor-like epitope by *Candida albicans* is of interest in this context as the presence of antibodies to this substance in the serum of CMC patients has been shown to block iC3b

rosetting.²³ It has also been found that patients with CMC are susceptible to other cutaneous infections notably dermatophytosis and human papilloma virus infections and so the defect is not specific to candida.

The classification of CMC has also changed with time. The best known variant is one associated with the polyendocrinopathy syndrome, usually hypoparathyroidism and hypoadrenalism. This is inherited as an autosomal recessive trait.²⁴ Patients with this form generally have evidence of autoantibodies to other organs and some have involvement of other systems, for example pernicious anaemia or primary ovarian failure. In addition to this type some patients may have underlying hypothyroidism, inherited as an autosomal recessive trait. Other patients have no clear predisposition and are said to have sporadic type CMC. In addition in some families without endocrinopathy there is evidence of either autosomal recessive or autosomal dominant susceptibility.²⁵ All forms are usually clinically indistinguishable and range from mild to severe, although those with the sporadic form appear more likely to have extensive disease. Rare adult forms of the disease may also occur. It is important to screen children periodically for endocrinopathy because they may develop endocrine disease years after their first presentation with CMC. One further risk factor is the appearance of leukoplakia on the tongue that may progress to squamous carcinoma. However progression appears to be a particular problem in smokers. Other complications include bronchiectasis and potentially fatal bacterial septicaemia.²¹

Treatment of CMC is now easier, although relapse of infection is common. The drug ketoconazole¹⁹ is very effective and itraconazole²⁶ or fluconazole²⁷ are alternatives. In each case relapse of oral infection is common and it is important to stop treatment once the mouth is clear, instituting further treatment should this be necessary. The reason for this cautious approach is the risk of drug resistance developing where an azole is used continuously. It is not clear whether this will occur with all azoles but resistant candidosis has been described with both ketoconazole and fluconazole. In future studies it will be important to combine monitoring of changes in the patterns of drug sensitivity with genetic typing (karyotyping, restriction fragment length polymorphisms) to check that resistance has truly evolved in the original strains.²⁸ For severe infections other drugs such as intravenous amphotericin B can be used instead.

Pityrosporum infections

The pityrosporum yeasts are common skin commensals, although their emergence is usually delayed until after puberty.²⁹ They may cause disease in childhood in two ways. Firstly they have been isolated in large numbers from cradle cap eczema giving rise to the suggestion that they may be responsible for this condition.³⁰ The situation is analogous to that seen with adult seborrhoeic dermatitis. In addition these fungi have also been found to be a cause of septicaemia in premature neonates; intravenous feeding with intralipid appears to be a risk factor for infection.³¹ The route of infection is not clear but is thought to be via an intravenous line. The commonest disease caused by the pityrosporum yeasts, pityriasis versicolor, is rare in childhood.

Conclusions

While superficial mycoses are common in adults, infections in children present specific challenges. In the healthy child scalp ringworm remains a problem although in industrialised countries, with the exception of the USA, it tends to be a

sporadically occurring disease. In the developing world, however, the prevalence of infection is likely to persist in the face of other more pressing financial priorities. In the immunocompromised patient, including those with HIV infection and CMC, there are still a number of challenges, not only in treatment but also in our knowledge of basic immunopathological mechanisms. The new understanding of the function of epidermis as an immunologically active organ may provide some of the answers that have been so elusive.

R J HAY

St John's Institute of Dermatology,
Guy's Hospital,
London SE1 9RT

- Sohnle PG. Dermatophytosis. In: Cox RA, ed. *Immunology of fungal diseases*. Florida: CRC Press, 1989:1-27.
- Odds FC. *Candida and candidosis*. London: Baillière Tindall, 1988.
- King RD, Khan HA, Foye JC, et al. Transferrin, iron and dermatophytes 1. Serum dermatophyte inhibitory component definitely identified as unsaturated transferrin. *J Lab Clin Med* 1975;86:204-12.
- Zurita J, Hay RJ. The adherence of dermatophyte microconidia and arthroconidia to human keratinocytes in vitro. *J Invest Dermatol* 1987;89:529-34.
- Braathen LR, Kaaman T. Human epidermal Langerhans cells induce cellular immune responses to trichophytin in dermatophytosis. *Br J Dermatol* 1983;109:295-9.
- Jones HE, Reinhardt JH, Rinaldi MG. Model dermatophytosis in naturally infected subjects. *Arch Dermatol* 1974;110:369-74.
- Calderon RA, Hay RJ. Cell-mediated immunity in experimental murine dermatophytosis. II Adoptive transfer of immunity to dermatophyte infection by lymphoid cells from donors with acute or chronic infections. *Immunology* 1984;53:405-10.
- Torssänder J, Karlsson A, Morfeldt-Mason L, et al. Dermatophytosis and HIV infection—study in homosexual men. *Acta Derm Venereol (Stokh)* 1988;68:53-9.
- Ahonen P, Koskimies S, Lokki ML, et al. The expression of autoimmune polyglandular disease type I appears associated with several HLA-A antigens but not with HLA-DR. *J Clin Endocrinol Metab* 1988;66:1152-7.
- Serjeantson S, Lawrence G. Autosomal recessive inheritance of susceptibility to tinea imbricata. *Lancet* 1977;ii:13-5.
- Rebell G, Taplin D. *Dermatophytes: their recognition and identification*. 2nd Ed. Florida: University of Miami Press, 1976.
- English MP, Gibson MD. Studies in the epidemiology of tinea pedis I and II. Tinea pedis in school children. *BMJ* 1959;ii:1442-8.
- Clayton YM. Scalp ringworm (tinea capitis). In: Verbov JL, ed. *Superficial fungal infections*. Manchester: MTP Press, 1986:1-8.
- Rippon JW. Epidemiology and emerging patterns of dermatophyte species. *Curr Top Med Mycol* 1985;1:208-34.
- Verhagen AR. Distribution of dermatophytes causing tinea capitis in Africa. *Trop Geogr Med* 1973;26:101-20.
- Krowchuk DP, Lucky AW, Primmer SI. Current status of the identification and management of tinea capitis. *Pediatrics* 1983;72:625-31.
- Branson DM, Desai DR, Barsky S, Foley SM. An epidemic of infection with Trichophyton tonsurans revealed in a 20 year old survey of fungal infections in Chicago. *J Am Acad Dermatol* 1983;3:322-30.
- Wright S, Robertson VJ. An institutional survey of tinea capitis in Harare, Zimbabwe and a trial of miconazole cream versus Whitfield's ointment in its treatment. *Clin Exp Dermatol* 1986;11:371-7.
- Jones HE, ed. *Ketoconazole today. A review of clinical experience* Manchester: ADIS Press, 1987.
- MacKenzie DWR. Hairbrush technique in detection and eradication of non-fluorescent scalp ringworm. *BMJ* 1963;iii:363-5.
- Herrod HG. Chronic mucocutaneous candidiasis in childhood and complications of non-candida infection: report of the Pediatric Immunodeficiency Collaborative Group. *J Pediatr* 1990;116:377-82.
- Durandy A, Fischer A, Le Deist F, et al. Mannan specific and mannan induced T-cell suppressive activity in patients with chronic mucocutaneous candidosis. *J Clin Immunol* 1987;7:400-10.
- Ollert MW, Wadsworth E, Calderone RA. Reduced expression of the functionally active complement receptor for iC3b but not for C3d on an avirulent mutant of *Candida albicans*. *Infect Immun* 1990;58:909-13.
- Ahonen P, Myllarniemi S, Sipila I, et al. Clinical variation of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) in a series of 68 patients. *N Eng J Med* 1990;322:1829-36.
- Sams WM, Jorizzo JL, Snyderman R, et al. Chronic mucocutaneous candidiasis; immunological studies of three generations of a single family. *Am J Med* 1979;67:948-59.
- Grant SM, Clissold SP. Itraconazole: a review of its pharmacodynamic and pharmacokinetic properties and therapeutic use in superficial and systemic mycoses. *Drugs* 1989;37:310-44.
- Grant SM, Clissold SP. Fluconazole: a review of its pharmacodynamic and pharmacokinetic properties, and therapeutic potential in superficial and systemic mycoses. *Drugs* 1990;39:877-916.
- Pfaller MA. Epidemiologic typing of candida species. *Curr Top Med Mycol* 1992;4:43-63.
- Faergemann J. Lipophilic yeasts in skin disease. *Semin Dermatol* 1985;4:173-84.
- Broberg A, Faergemann J. Infantile seborrhoeic dermatitis and pityrosporum ovale. *Br J Dermatol* 1989;120:359-62.
- Dankner WM, Spector SA, Fierer J, et al. Malassezia fungemia in neonates and adults; complication of hyperalimentation. *Rev Infect Dis* 1987;9:743-53.