

ing the importance of the data in this type of survey being primarily recorded by general practitioners.

There is an impression among those dealing with a large number of "cot death" cases that these infants tend to be significantly overweight for their age (J. L. Emery, personal communication, 1970). One child in our series suffered a cot death during the period of the trial, and this infant was in the group of children who were over the 90th percentile for age; necropsy gave the cause of death as resulting from acute bronchiolitis. This may well be a chance finding in a relatively large series of children, but it points the need for further investigation in this field.

If it be assumed that the case is proved, and that obese infants do have a greater liability to respiratory infection than those of normal weight, the question arises why this should be the case. Two possibilities emerge as indications for further investigation. Firstly, that obesity leads to underventilation of the lungs, with possible impairment of coughing and clearing of secretions from the respiratory tree, leading to a tendency for minor respiratory infections to become more serious and prolonged. The second possible mechanism by which obesity might predispose to an increased incidence of

respiratory infection could be due to a defect in the immune defences of the body, associated either directly with the obesity or indirectly with the early introduction of artificial feeding. The early abandonment of breast-feeding may also play a part in the reduction of the immune response at this age. Further work on both of these possibilities would be valuable and interesting, and we hope that our clinical survey will stimulate more detailed biochemical and respiratory function studies in this field.

We are grateful to local general practitioners for their help and co-operation, without which this paper could not have been written; to Dr. W. J. McQuillan for advice and encouragement, and to the health visitors in his department for recording weights and other details of babies and collection of data cards from general practitioners; to the department of statistics, University of Birmingham, for advice on the statistical aspects of the trial; and to Dr. Brian Williams, who also gave helpful criticism and advice.

#### References

- British Medical Journal*, 1970, 2, 64.  
Hutchinson-Smith, B. (1970). *Medical Officer*, 123, 257.  
Tanner, J. M., Whitehouse, R. H., and Takaiishi, M., (1966). *Archives of Disease in Childhood*, 41, 613.

## Infectious Complications in Bone Marrow Transplant Patients

C. O. SOLBERG, H. J. MEUWISSEN, R. N. NEEDHAM, R. A. GOOD, J. M. MATSEN

*British Medical Journal*, 1971, 1, 18-23

#### Summary

In 11 patients receiving transplants of allogeneic bone marrow, the graft was successful in six. Nine patients developed infections, and six died—five of septicaemia and one of *Pneumocystis carinii* pneumonia. Fifty individual infections occurred. Predisposing factors included severe underlying diseases, long-term exposure to resistant hospital organisms, heavy immunosuppressive therapy, and graft-versus-host disease. Gram-negative bacilli and *Candida albicans* were the most common causative organisms. In every instance of septicaemia identical organisms were isolated from blood cultures and simultaneously obtained stool cultures. Infection with exogenous organisms often occurred in patients occupying conventional isolation rooms. Isolation of one patient for 45 days in a laminar air flow room prevented infection with exogenous organisms.

#### Introduction

Infection remains a major problem in all types of internal organ transplantation (Rifkind *et al.*, 1964; Kelly *et al.*, 1967;

Fulginiti *et al.*, 1968; Montgomery *et al.*, 1969). Bone marrow transplantation is still in its early stages (Pegg, 1966; Mathé *et al.*, 1967), and studies of associated infections are limited (Mathé *et al.*, 1965). It is apparent, however, that candidates for bone marrow transplantation are particularly susceptible to infection owing to the nature of their underlying disease processes (Mathé *et al.*, 1967; Bergsma and Good, 1968), the heavy immunosuppressive therapy often required to obtain a take of the graft, and the subsequent occurrence of graft-versus-host disease (Blaese *et al.*, 1964; Mathé *et al.*, 1965).

The present study reports the infectious complications in bone marrow transplant patients at the University of Minnesota Medical Center from 1968 to 1970. Results of extensive microbiological monitoring of patients established the origin of several infections and provided information for improved prophylaxis and therapy in future bone marrow transplantation.

#### Patients and Methods

Eleven patients received transplants of allogeneic\* bone marrow. All suffered from a variety of serious life-threatening diseases. Details of the patients' diagnoses, age, sex, and bone marrow transplantation are presented in Table I. The methods of histocompatibility matching and the technique of bone marrow transplantation have already been described (Meuwissen *et al.*, 1969).

Demonstrable cell-mediated immune functions were completely absent in Cases 1-4, deficient in Cases 5-7, and unimpaired in Cases 8-11. Case 4 received small doses of methotrexate to facilitate the take of the bone marrow while Cases 1-3 received no immunosuppressive therapy. In addi-

\* Allogeneic: of same species but not of identical genetic constitution—for example, father and son.

University of Minnesota Medical Center, Minneapolis, Minnesota 5545  
C. O. SOLBERG, M.D., Associate Professor of Medicine. (Present address: School of Medicine, Medical Department B, University of Bergen, Bergen, Norway)

H. J. MEUWISSEN, M.D., Lecturer

R. N. NEEDHAM, B.S., Research Fellow

R. A. GOOD, M.D., PH.D., Regent's Professor of Paediatrics and Microbiology

J. M. MATSEN, M.D., Assistant Professor of Paediatrics and Laboratory Medicine

TABLE I—Bone Marrow Transplant Patients

Case No.	Age and Sex	Disease	Bone Marrow Donor	Immunosuppressive Therapy	Outcome
1	5 months. Male	Sex-linked lymphopenic hypogammaglobulinaemia	Sister, MLC compatible, CTA incompatible	None	Immunological reconstitution
2	7 months. Female	Autosomal recessive lymphopenic hypogammaglobulinaemia	Sister, HL-A identical	None	Immunological reconstitution; chronic GVHD
3	13 months. Male	Sex-linked lymphopenic hypogammaglobulinaemia	Sister, HL-A identical	None	Beginning immunological reconstitution. Died of infection 27 days after TP
4	10 months. Male	Sex-linked lymphopenic hypogammaglobulinaemia	Father, MLC and CTA incompatible	Antihuman lymphocyte serum. Methotrexate 0.5 mg./kg with 3-6 day intervals after TP	Beginning immunological reconstitution. Died of infection 42 days after TP
5	15 years. Male	Hodgkin's disease, stage IV B	Brother, HL-A identical	Cyclophosphamide 50 mg./kg daily for 2 days before TP	Beginning immunological reconstitution. Died of infection and GVHD 27 days after TP
6	15 years. Male	Hodgkin's disease, stage IV B	Brother, HL-A identical	Long-term chemotherapy discontinued 2 weeks before TP	No immunological reconstitution. Died of Hodgkin's disease of stomach
7	7 years. Female	Chronic mucocutaneous candidiasis	Sister, MLC and CTA incompatible	None	Improvement of candidiasis; chronic GVHD
8	8 years. Male	Non-lymphopenic hypogammaglobulinaemia (Bruton's type)	Sister, HL-A identical	None	No immunological reconstitution
9	9 years. Female	Idiopathic aplastic anaemia	Brother, HL-A identical	Before 1st TP: antilymphocyte serum, prednisone and azathioprine. 2nd TP: cyclophosphamide 60 mg./kg daily for 3 days before TP	Died of infection 16 days after TP
10	21 years. Female	Aplastic anaemia (drug-induced?)	Brother, HL-A identical	Long-term prednisone. Cyclophosphamide 75 mg./kg daily for 2 days before TP	Died of infection 3 days after TP
11	25 years. Female	Acute lymphoblastic leukaemia	Sister, MLC compatible?	Long-term prednisone. Cyclophosphamide 37.5 mg./kg daily for 2 days before TP	Recovery of bone marrow and transient remission of leukaemia. Died of infection 122 days after TP

MLC = Mixed leucocyte culture. CTA = Cytotoxic assay. TP = Transplantation. GVHD = Graft-versus-host disease.

tion to other drugs high doses of cyclophosphamide were given to Case 5 and to Cases 9-11. Seven patients received HL-A identical sibling bone marrow. Graft-versus-host disease occurred in six patients (Table I).

The principal markers for the presence and function of donor cells in the recipient were as follows: (1) sex chromosome markers, (2) red blood cell antigenic markers, (3) gammaglobulin allotype markers (kindly performed by Dr. Arthur Steinberg), (4) functional criteria such as increased gammaglobulin production in hypogammaglobulinaemic patients or the appearance of lymphocytes and in-vitro lymphocyte function in previously lymphopenic unresponsive patients, and (5) transfer of delayed hypersensitivity from donor to recipient.

### Immunological Reconstitution

In Cases 1 and 2 immunological reconstitution was documented by persistent production of immunoglobulins, antibodies, and the in-vitro lymphocyte response to phytohaemagglutinin and allogeneic cells. In Case 1 (a male) peripheral blood lymphocytes, dividing under the influence of phytohaemagglutinin, were of female karyotype, whereas spontaneously dividing bone marrow cells showed a mixture of male and female karyotypes. Eight weeks after transplantation this patient developed immunologically induced pancytopenia attributed to a reaction of the grafted lymphoid cells against host constituents. One month later treatment of the pancytopenia with a second marrow transplant was as successful in restoring the red-cell, platelet, and granulocyte populations as the first had been in establishing the lymphocyte and plasma cell lines (Meuwissen *et al.*, 1969). Cases 1 and 2 were still in good clinical condition and immunologically reconstituted 23 and 16 months after transplantation, respectively.

In Cases 3 and 4 the onset of immunological reconstitution was shown by a rise in phytohaemagglutinin-stimulated blastoid cells in peripheral blood lymphocyte cultures, and by a positive response to a challenging dose of dinitrofluorobenzene, 1/1,000, and to intradermal IgE. In Case 3 necropsy also showed proliferating bone marrow in the subcutaneous fat around the site of the peritoneal catheter used for bone marrow transplantation, and in Case 4 donor type red blood cells occurred in the peripheral blood one week after transplantation, while these cells had previously been absent.

In Case 5 necropsy showed infiltration of the lymph nodes and the spleen by numerous plasma cells and large pale blastoid cells characteristic of graft-versus-host disease. Lesions typical of this disease were also present in the skin. Case 11 had a second relapse of acute lymphoblastic leukaemia and showed no response to prolonged treatment with various antileukaemic drugs. Complete recovery of the marrow occurred after bone marrow transplantation. Three months later a third relapse occurred and the patient died. In the absence of markers, donor type cells could not be found in the blood. In the remaining patients no evidence of take of bone marrow was found.

### Bacteriological Tests

Microbiological sampling and culturing methods and biochemical and serological organism identification techniques have been described (Solberg *et al.*, 1970). Sensitivity determinations were done with either the twofold tube dilution method (using Mueller-Hinton broth (BBL) and an inoculum of  $10^5$ - $10^6$  organisms per ml) or the single high-potency disc technique (Bauer *et al.*, 1966). Identity of bacterial strains was documented by antibacterial spectra in addition to the biochemical and serological techniques indicated above.

A diagnosis of septicaemia was based on positive blood cultures together with other evidence of septicaemia—that is, a primary site of infection, spiking fever, shock, and/or gastrointestinal symptoms. The presence of pulmonary infection was judged by typical clinical and roentgenological findings together with a predominant growth of micro-organisms in repeated deep cough sputum cultures or deep tracheal aspiration cultures. Urinary tract infection was diagnosed when urine catheter or midstream specimens contained  $10^5$  or more organisms per ml.

### Results

#### INCIDENCE OF INFECTIONS

A total of 50 individual infections occurred in 9 of the 11 transplant patients. Cases 6 and 8 did not develop infection related to bone marrow transplantation. Each failed to have a definite take of the bone marrow and no evidence of graft-versus-host disease was observed. In the remaining patients,

nine infections occurred during the three weeks before transplantation; eight were cured and one infection was carried over to the post-transplantation period. After transplantation 41 additional infections developed.

#### INFECTING ORGANISMS

Twenty-eight infections in eight patients were caused by Gram-negative bacilli (Table II). *Candida albicans* accounted for nine infections in four patients, *Staphylococcus aureus* for five infections in four patients, *Haemophilus influenzae* for two infections in two patients, and, finally, *Bacillus* sp., *Bacteroides* sp., *Clostridium perfringens*, enterococcus, and *Pneumocystis carinii* accounted for or were involved in 12 infections in five patients.

#### TYPE OF INFECTION

Septicaemia occurred eight times in six patients; two patients had two episodes, the organisms isolated during the two episodes being different (Chart, Table II). Seven of the eight septic episodes occurred after transplantation. Enteric Gram-negative bacilli were involved six times, and *C. albicans* and *Bacteroides* sp. once each. Infection with multiple organisms, from two to five pathogens verified by two or three consecutive blood cultures, occurred in four instances. In all septic episodes blood cultures grew out strains also isolated from simultaneously obtained stool cultures. Septicaemia was the primary cause of death in five patients, three of these also having internal organ abscesses (Table III).

Pneumonia occurred 11 times in six patients (Chart, Table II). The principal causative organisms were *Klebsiella* sp., *Pn. carinii*, and *Staph. aureus*. Three infections occurred before transplantation and were eradicated. Pneumonia was the primary cause of death of one patient, and a contributing cause in another patient (Table III). *Pn. carinii* was the infecting organism in both patients, and in each instance the organism, most probably, was carried over from the pre-transplant period.

Urinary tract infections occurred 11 times in seven patients. Gram-negative bacilli were present in 10 cases and *C. albicans* in one. Cure was achieved in eight instances. In two

TABLE III—Terminal Infectious Diseases in Six Bone Marrow Transplant Patients

Case No.	Survival (Days)	Infectious Disease	Aetiology
3	27	Pneumonia	<i>Pn. carinii</i>
4	42	Septicaemia Pneumonia	<i>E. coli</i> , <i>Ps. aeruginosa</i> <i>Pn. carinii</i>
5	27	Septicaemia Lung and spleen abscesses	<i>Ps. aeruginosa</i> <i>E. coli</i> , <i>Pr. mirabilis</i>
9	16	Septicaemia Multiple metastatic abscesses	<i>Bacillus</i> sp., <i>Cl. perfringens</i> , enterococcus, <i>E. coli</i> , <i>Klebsiella</i> sp., <i>Bacillus</i> sp., <i>Cl. perfringens</i> , enterococcus, <i>E. coli</i> , <i>Klebsiella</i> sp.
10	3	Septicaemia Urinary tract infection	<i>E. coli</i> <i>E. coli</i>
11	122	Septicaemia Multiple metastatic abscesses	<i>C. Albicans</i> <i>C. Albicans</i>

patients who died of septicaemia the organisms were never eradicated from the urine. In these patients identical organisms were isolated from urine and blood cultures.

Cases 5, 9, and 11 had multiple metastatic abscesses in the brain, gastrointestinal wall, kidneys, liver, lungs, and/or spleen in addition to septicaemia. Identical organisms were isolated from the abscesses and blood cultures. Case 7 developed lung abscesses after pneumonia with enterococci.

Seven patients had miscellaneous infections—that is, mucocutaneous candidiasis, otitis media, draining pustules, and wounds. Case 7 had suffered from mucocutaneous candidiasis for several years and improved significantly after transplantation. The other patients were successfully treated with antibiotics.

#### RELATION TO IMMUNOSUPPRESSIVE THERAPY AND GRAFT-VERSUS-HOST DISEASE


Owing in part to immunosuppressive therapy before transplantation, Cases 5 and 9-11 had granulocyte counts less than 100/mm<sup>3</sup> for several days after transplantation. Each developed septicaemia, and three patients (Cases 5, 9, and 10) with Hodgkin's disease and aplastic anaemia died 27, 16 and 3 days after transplantation, respectively. Necropsy of these three patients showed diffuse necrosis and patchy ulcerations of the gastrointestinal mucosa.

TABLE II—Infections in Bone Marrow Transplant Patients

Diagnosis	No. of Cases	No. of Infections	Aetiology	Results		
				Cured	Suppressed	Not Cured
Septicaemia	6	8	<i>Bacteroides</i> sp.	1	—	—
			<i>Candida albicans</i>	—	—	1
			<i>Escherichia coli</i>	—	—	1
			<i>Pseudomonas aeruginosa</i>	—	—	1
			<i>E. coli</i> , <i>Proteus mirabilis</i>	2	—	—
			<i>E. coli</i> , <i>Ps. aeruginosa</i>	—	—	1
Pneumonia	6	11	<i>Bacillus</i> sp., <i>Clostridium perfringens</i> , enterococcus, <i>E. coli</i> , <i>Klebsiella</i> sp.	—	—	1
			Enterococcus	1	—	—
			<i>Haemophilus influenzae</i>	1	—	—
			<i>Klebsiella</i> sp.	3	—	—
			<i>Pneumocystis carinii</i>	—	—	2
			<i>Staphylococcus aureus</i>	2	—	—
Urinary tract infection	7	11	<i>Ps. aeruginosa</i>	1	—	—
			Enterococcus, <i>E. coli</i> , <i>Pr. mirabilis</i>	1	—	—
Abscesses (brain, gastrointestinal wall, kidney, liver, lung, spleen)	4	12	<i>C. albicans</i>	—	1	—
			Gram-negative bacilli*	8	—	2
Miscellaneous infections	7	8	<i>C. albicans</i>	—	—	5
			Enterococcus	1	—	—
			<i>E. coli</i> , <i>Pr. mirabilis</i>	—	2	—
			<i>Bacillus</i> sp., <i>Clostridium perfringens</i> , <i>E. coli</i> , enterococcus, <i>Klebsiella</i> sp.	—	—	4
			<i>C. albicans</i>	1	—	1
			Enterococcus	1	—	—
<i>H. influenzae</i>	1	—	—			
<i>Staph. aureus</i>	2	1	—			
<i>E. coli</i> , <i>Pr. mirabilis</i>	1	—	—			

\**E. coli*, *Klebsiella* sp., *Pr. mirabilis*, *Ps. aeruginosa*.

Cases 1-5 and 7 developed graft-versus-host disease from one to five weeks after bone marrow transplantation. In these patients 21 infections occurred after transplantation—three in the period between transplantation and the onset of graft-versus-host disease and 18 during the first five weeks of graft-versus-host disease (see Chart).



Cases 1-4 with absence of demonstrable cell-mediated immune functions all developed graft-versus-host disease. Cases 1-3 received HL-A identical marrow, and the graft-versus-host disease symptoms were less severe than in Case 4, who received HL-A non-identical paternal marrow. Two patients (Cases 1 and 2) recovered from their graft-versus-host disease and did not develop life-threatening symptoms. One (Case 3) developed severe *Pn. carinii* pneumonia and died 27 days after transplantation, and one (Case 4) died of massive septicaemia, pneumonia, and multiple gastrointestinal ulcerations, probably caused by his severe graft-versus-host disease.

#### ORIGIN OF INFECTION

In cases 1, 3, 4, 5, and 9 bacteriological samples from the nose, throat, axillae, groin, perineum, urine, and stool were obtained regularly at intervals of two to four days. On admission Case 4 was kept in a laminar air flow room under strict isolation procedures (Solberg *et al.*, 1970), while the other four were isolated in non-laminar air flow hospital rooms, but under the same isolation procedures as in the

laminar air flow room. Case 4 occupied the laminar air flow room for 45 days. This patient maintained his own endogenous bacterial flora during his entire stay in the laminar air flow room, and no contamination or infection occurred with organisms from the environment or from the personnel entering the room. The terminal septicaemia was

Infections in relation to bone marrow transplantation and graft-versus-host disease. \*Later, the patient received one more bone marrow transplant without evidence of graft-versus-host disease and infectious complications. †Previously the patient had received a bone marrow transplant with little or no evidence of immunological reconstitution, no graft-versus-host disease and no infectious complications.

caused by *Klebsiella* sp. and *Ps. aeruginosa* identical to strains repeatedly isolated from his stool.

In the other four patients colonization of the nose, throat, and gut with exogenous organisms (*Ps. aeruginosa*, *Klebsiella*, *Moraxella*, and/or *Bacillus* sp.) occurred from 5 to 29 days after admission to hospital. Initially the organisms were isolated in small numbers, but within two to seven days large numbers were usually recovered. Cases 1 and 5 developed infections with these organisms eight and five days after colonization, respectively. Case 1 developed pneumonia with a *Klebsiella* strain identical to that which had recently colonized his throat. Case 5 developed septicaemia with a *Pseudomonas* strain identical to that which had colonized his gut. Case 9 also developed septicaemia, and several blood cultures grew out five different organisms, identical to strains being isolated from simultaneously obtained stool cultures. One of these strains (*Klebsiella* sp.) had colonized the gut 20 weeks previously. Two patients (Cases 5 and 9) died of septicaemia. Case 3 had two episodes of septicaemia and had pneumonia three times. There was no evidence of exogenous organisms causing these infections. The first episode of septicaemia was caused by *E. coli* and *Pr. mirabilis* and the second by *Bacteroides* sp., organisms which were repeatedly isolated in large numbers from the patient's stools.

## Discussion

Infection has been one of the most serious complications of organ transplantation. In renal transplantation infections have contributed to more than two-thirds of the deaths (Rifkind *et al.*, 1964; Kelly *et al.*, 1967), and in 17 recipients of liver homografts Fulginiti *et al.* (1968) reported that 10 of the 15 deaths were due to infectious complications. Montgomery *et al.* (1969) found that 10 out of 28 cardiac transplant patients died of infections complicating the surgery and subsequent immunosuppressive therapy. The principal reasons for the severe infections in solid organ transplantation include heavy immunosuppressive therapy, complicated surgical procedures, vascular post-surgery complications, and long-term exposure of debilitated patients to resistant hospital organisms.

As documented in our study, infection is also a serious complication in bone marrow transplantation. Nine of the 11 patients in our series experienced a total of 41 infections after transplantation neither received immunosuppressive therapy. Two who did not develop infection in relation to transplantation, neither received immunosuppressive therapy nor developed graft-versus-host disease. The principal reasons for the increased susceptibility to infection in our patients include severe underlying medical problems, long-term exposure to antibiotic-resistant hospital organisms, heavy immunosuppressive therapy, and, finally, frequent occurrence of graft-versus-host disease.

All patients reported here had severe underlying diseases, including hypogammaglobulinaemia, chronic mucocutaneous candidiasis, neoplasia, and aplastic anaemia. In all of these conditions there is a predisposition to either a specific infectious process (in mucocutaneous candidiasis) or to opportunistic infections in general. Before transplantation our patients had experienced repeated infections resulting in frequent admission to hospital with the inherent risk of exposure to antibiotic-resistant hospital organisms. Infections due to hospital-adapted strains are usually more difficult to treat, and the persistence of the organisms appears to be enhanced (McDermott, 1957).

In addition to severe underlying diseases and long-term exposure to resistant hospital organisms, immunosuppressive therapy greatly enhanced the susceptibility of our patients to infection. Six patients in the present series received cyclophosphamide, methotrexate, prednisone, azathioprine, and/or antihuman lymphocyte serum to suppress the immune response in order to obviate transplant rejection. Two patients with aplastic anaemia, one with Hodgkin's disease, and one with acute leukaemia—all with pre-existing granulocytopenia—had fewer than 100 granulocytes per mm<sup>3</sup> for several days after administration of cyclophosphamide. The three patients with aplastic anaemia and Hodgkin's disease also developed widespread necrosis of the intestinal mucosa. Despite heroic measures, including repeated leucocyte transfusions, each of these four patients developed septicaemia with organisms present in the gut flora, and the three patients with aplastic anaemia and Hodgkin's disease died within a few days. High doses of cyclophosphamide are known to produce intravascular coagulation in all tissues in monkeys (van Bekkum *et al.*, 1970) and haemorrhagic gastroenteritis in dogs (Epstein *et al.*, 1969). Therefore the gut necrosis which occurred in our patients may have been related to the high doses of cyclophosphamide. The pronounced neutropenia was in all likelihood a direct result of the administration of this drug.

Finally, graft-versus-host disease is associated with a pronounced degree of immunological incompetence due to production of lymphoid atrophy and depression of bone marrow (Blaese *et al.*, 1964; Mathé *et al.*, 1967; van Bekkum and de Vries, 1967). Furthermore, graft-versus-host disease may cause severe damage to the intestinal mucosa, resulting in continuous seeding of enteric organisms into the blood and

persistence of septicaemia despite what might appear to be adequate therapy by in-vitro testing (de Vries *et al.*, 1961). Six of our patients developed graft-versus-host disease. In these patients 18 separate infections occurred within four to five weeks after the onset of graft-versus-host disease, and three died—one of pneumonia and two of septicaemia. The latter two patients had multiple intestinal ulcerations, and identical organisms were isolated from the blood and stool cultures.

In the eight septic episodes which occurred in our patients, blood cultures always grew out strains isolated from simultaneously obtained stool cultures, and four of the five patients who died of septicaemia had multiple intestinal ulcerations, most likely related to graft-versus-host disease and/or administration of cyclophosphamide. Therefore, in future patients intestinal decontamination with non-absorbable antibacterial and antifungal agents should be considered. Decontamination of the gut flora may prevent fatal septicaemia caused by seeding of enteric organisms into the blood. Furthermore, the gastrointestinal damage which occurs after bone marrow transplantation may be less severe and healing may occur more rapidly if the load of intestinal organisms is reduced. Finally, evidence is accumulating that intestinal decontamination may reduce the severity of graft-versus-host disease (Keast, 1968; Ledney, 1969).

## INFECTION WITH EXOGENOUS ORGANISMS

The principal drawback of intestinal decontamination remains the risk of superinfection with exogenous organisms resistant to the antimicrobial agents used. It should be possible, however, to minimize this risk in the future by using strict aseptic techniques, by sterilizing all food, medications, and utensils used for the patients, and by use of laminar air flow rooms (Solberg *et al.*, 1970).

The importance of infection with exogenous organisms in these highly susceptible patients should not be overlooked. Four of our patients occupying conventional isolation rooms were the subjects of extensive bacteriological monitoring. Each was colonized with exogenous organisms 5 to 29 days after admission to hospital. Later, three of these patients developed documented infections due to those organisms, and two died. In these three patients organisms originally exogenous became part of the resident flora and greatly increased in numbers for several days to weeks before precipitating the infectious episodes. The other bone marrow transplant patients who occupied conventional isolation rooms were not the subjects of extensive bacteriological monitoring. Most probably some of the infections which occurred in these patients were also due to exogenous organisms. In contrast, one of our patients was kept in a laminar air flow room for 45 days, under strict isolation procedures. Extensive microbiological monitoring of the patient, the laminar air flow room, and the personnel entering the isolation facility showed that the patient maintained his endogenous bacterial flora during his entire stay in the room, and no colonization occurred with organisms from the environment or from any of the personnel entering the room. To prevent infection with exogenous organisms in future bone marrow transplantation, admission of the patients to laminar air flow rooms under strict isolation seems indicated, especially if intestinal decontamination is attempted.

Despite the frequent occurrence of fatal infections in bone marrow transplantation, aggressive attempts at diagnosis are of far more than academic interest since several of these infections, including those caused by fungi and *Pn. carinii*, can be treated by large doses of antimicrobial agents and immune globulins (Rifkind *et al.*, 1966). The rapid onset and fulminant course of infections in these patients, especially during periods of heavy immunosuppressive therapy or graft-versus-host disease, requires close observation by skilled per-

sonnel and early institution of therapy. Continuous monitoring of the patients' flora may often help to ensure specific therapy before results of appropriate cultures are available, as shown in our septicaemia patients who always developed infections with organisms present in the intestinal flora. To secure a specific bacteriological diagnosis, however, the value of blood cultures, endotracheal aspirations, and urine and wound cultures should be emphasized in view of the repeatedly significant findings in these samples.

In future bone marrow transplantations, admission of patients to laminar air flow rooms under strict isolation conditions, and decontamination of the intestinal flora, may be important steps in ensuring survival of the patients. Improved immunosuppressive therapy and methods to ameliorate the graft-versus-host disease, however, seem more important in preventing fatal infections.

This work was supported in part by U.S. Public Health Service Grants 1F05-TW-01494-01, 1-R01-CA11505-01, AI-08677, and HE-06314, and by grants from the National Foundation-March of Dimes, and the American Heart Association.

Requests for reprints should be addressed to Dr. John M. Matsen, Box 198, University of Minnesota Hospital, Minneapolis, Minnesota 55455, U.S.A.

## References

- Bauer, A. W., Kirby, W. M. M., Sherris, J. C., and Turck, M. (1966). *American Journal of Clinical Pathology*, **45**, 493.
- Bergsma, D., and Good, R. A. (editors) (1968). *Immunologic Deficiency Diseases in Man*. New York, National Foundation.
- Blaese, R. M., Martinez, C., and Good, R. A. (1964). *Journal of Experimental Medicine*, **119**, 211.
- de Vries, M. J., Crouch, B. G., van Putten, L. M., and van Bekkum, D. W. (1961). *Journal of the National Cancer Institute*, **27**, 67.
- Epstein, R. B., Storb, R., Clift, R. A., and Thomas, E. D. (1969). *Cancer Research*, **29**, 1072.
- Fulginiti, V. A., et al. (1968). *New England Journal of Medicine*, **279**, 619.
- Keast, D. (1968). *Immunology*, **15**, 237.
- Kelly, W. D., et al. (1967). *Surgery*, **62**, 704.
- Ledney, G. D. (1969). *Transplantation*, **8**, 127.
- McDermott, W. (1957). *Yale Journal of Biology and Medicine*, **30**, 257.
- Mathé, G., et al. (1965). *European Journal of Cancer*, **1**, 75.
- Mathé, G., et al. (1967). *Scandinavian Journal of Haematology*, **4**, 193.
- Meuwissen, H. J., Gatti, R. A., Terasaki, P. I., Hong, R., and Good, R. A. (1969). *New England Journal of Medicine*, **281**, 691.
- Montgomery, J. R., et al. (1969). Abstract: Ninth Interscience Conference on Antimicrobial Agents and Chemotherapy, Washington D.C., October.
- Pegg, D. E. (1966). *Bone Marrow Transplantation*, p. 71. London, Lloyd-Luke.
- Rifkind, D., Faris, T. D., and Hill, R. B., jun. (1966). *Annals of Internal Medicine*, **65**, 943.
- Rifkind, D., Marchioro, T. L., Waddell, W. R., and Starzl, T. E. (1964). *Journal of the American Medical Association*, **189**, 397.
- Solberg, C. O., et al. (1970). *Applied Microbiology*. In press.
- van Bekkum, D. W., and de Vries, M. J. (1967). *Radiation Chimaeras*. London, Logos Press.
- van Bekkum, D. W., Dicke, K. A., Balner, H., Hollander, C. F., and van Putten, L. M. (1970). *Experimental Hematology*, **20**, 27.

# Measles Immunoglobulins in Subacute Sclerosing Panencephalitis

J. H. CONNOLLY, MARGARET HAIRE, DIANA S. M. HADDEN

*British Medical Journal*, 1971, **1**, 23-25

## Summary

Normal responses of measles specific immunoglobulins M and G (IgM and IgG) were defined in 10 children with measles. Abnormal responses of measles IgM and IgG were found in both sera and cerebrospinal fluids from three cases of subacute sclerosing panencephalitis. In two patients the serum titres of measles IgM and IgG were abnormally high. The measles IgM was present during prolonged illnesses in serum and cerebrospinal fluid, which suggested a correlation with the known persistence of measles virus antigen in the brain of the three patients. It was concluded that both measles IgM and IgG may be produced within the central nervous system in subacute sclerosing panencephalitis.

## Introduction

The viral aetiology of subacute sclerosing panencephalitis was suggested by the finding of Cowdry type A inclusions (Dawson, 1933) and paramyxovirus-like filaments in brain (Bouteille, Fontaine, Vedrenne, and Delarue, 1965). Measles antibody in high or increasing titre was found in serum and cerebrospinal fluid (C.S.F.) and measles virus antigen was found in the brain of the same patients (Connolly, Allen,

Hurwitz, and Millar, 1967; Freeman, Magoffin, Lennette, and Herndon, 1967). Further evidence of the relationship between measles virus and subacute sclerosing panencephalitis was reported at a conference (Sever and Zeman, 1968) and measles virus has now been isolated from cultured brain cells of patients with this disease (Baublis and Payne, 1968; Horta-Barbosa, Fuccillo, Sever, and Zeman, 1969).

Measles IgM and IgG were titrated in 10 children with measles and in three previously reported patients with subacute sclerosing panencephalitis to assess the diagnostic value of the immunoglobulin responses in both acute and persistent measles virus infections.

## Patients, Materials, and Methods

The childhood measles group consisted of nine children with uncomplicated measles and one child with measles-associated encephalitis who recovered completely. Serum samples were obtained from all children, and a C.S.F. sample from the child with encephalitis. "Normal" C.S.F. samples were obtained during diagnostic lumbar pneumoencephalography from four patients with epilepsy and one patient with a head injury. The clinical, pathological, epidemiological, and virological data on the three cases of subacute sclerosing panencephalitis have already been reported (Connolly, Allen, Hurwitz, and Millar, 1968).

All specimens were stored at  $-20^{\circ}\text{C}$ , except the normal C.S.F. samples, which were stored at  $-70^{\circ}\text{C}$ , and were tested for measles complement-fixing antibody to confirm the diagnosis of measles where necessary. Virus specific IgM and IgG were detected with sheep antihuman IgM and IgG (Wellcome Reagents Ltd.) by the indirect immunofluorescence technique (Haire and Hadden, 1970). Serum and C.S.F.

Department of Microbiology, The Queen's University of Belfast, Belfast BT12 6BN

J. H. CONNOLLY, M.D., M.R.C.PATH., Consultant Virologist  
MARGARET HAIRE, M.D., Senior Experimental Officer  
DIANA S. M. HADDEN, M.B., D.C.H., Research Assistant