ERRATA

Contrasts Between Phagocyte Antibiotic Uptake and Subsequent Intracellular Bactericidal Activity

W. LEE HAND AND NEVA L. KING-THOMPSON

Veterans Administration Medical Center (Atlanta), Decatur, Georgia 30033, and the Department of Medicine, Emory University School of Medicine, Atlanta, Georgia 30303

Volume 29, no. 1, p. 137, Table 2: No. of viable S. aureus at 1 h for PMN plus clindamycin should read " $5.18 (\pm 1.06) \times 10^{5}$ ".

Antimicrobial Susceptibility of Five Subgroups of Mycobacterium fortuitum and Mycobacterium chelonae

JANA M. SWENSON, RICHARD J. WALLACE, JR., VELLA A. SILCOX, AND CLYDE THORNSBERRY

Antimicrobics and Infection Mechanisms Branch, Hospital Infections Program, and Mycobacteriology Reference Branch, Bacterial Diseases Division, Center for Infectious Diseases, Centers for Disease Control, Atlanta, Georgia 30333, and the University of Texas Health Center at Tyler, Tyler, Texas 75710

Volume 28, no. 6, p. 810, Table 4: The MIC ranges and modal MICs, respectively, should be $\leq 0.5-1.0$ and $\leq 0.5,1.0$ for amikacin, $\leq 0.25-4$ and 0.5,1.0 for doxycycline, >16 and >16 for erythromycin, and $\leq 2-16$ and ≤ 2 for sulfamethoxazole.

An Animal Source for the ROB-1 β-Lactamase of *Haemophilus influenzae* Type b

ANTONE A. MEDEIROS, ROGER LEVESQUE, AND GEORGE A. JACOBY

Miriam Hospital, Brown University, Providence, Rhode Island 02906, and Massachusetts General Hospital, Boston, Massachusetts 02114

Vol. 29, no. 2, p. 214: In paragraph 2 of the Discussion, the second sentence should begin "An ampicillin-resistant strain of *Pasteurella multocida*. . .".