

Characterization of Tocopherol Cyclases from Higher Plants and Cyanobacteria. Evolutionary Implications for Tocopherol Synthesis and Function¹

Scott E. Sattler, Edgar B. Cahoon², Sean J. Coughlan³, and Dean DellaPenna*

Department of Biochemistry and Molecular Biology, Biochemistry Building, Michigan State University, East Lansing, Michigan 48824-1319 (S.E.S., D.D.P.); and DuPont Crop Genetics, Experimental Station, Wilmington, Delaware 19880-0402 (E.B.C., S.J.C.)

Tocopherols are lipophilic antioxidants synthesized exclusively by photosynthetic organisms and collectively constitute vitamin E, an essential nutrient for both humans and animals. Tocopherol cyclase (TC) catalyzes the conversion of various phytyl quinol pathway intermediates to their corresponding tocopherols through the formation of the chromanol ring. Herein, the molecular and biochemical characterization of TCs from *Arabidopsis* (*VTE1* [VITAMIN E 1]), *Zea mays* (*SXD1* [Sucrose Export Deficient 1]) and *Synechocystis* sp. PCC6803 (slr1737) are described. Mutations in the *VTE1*, *SXD1*, or slr1737 genes resulted in both tocopherol deficiency and the accumulation of 2,3-dimethyl-6-phytyl-1,4-benzoquinone (DMPBQ), a TC substrate. Recombinant *SXD1* and *VTE1* proteins are able to convert DMPBQ to γ -tocopherol in vitro. In addition, expression of maize *SXD1* in a *Synechocystis* sp. PCC6803 slr1737 knockout mutant restored tocopherol synthesis, indicating that TC activity is evolutionarily conserved between plants and cyanobacteria. Sequence analysis identified a highly conserved 30-amino acid C-terminal domain in plant TCs that is absent from cyanobacterial orthologs. *vte1-2* causes a truncation within this C-terminal domain, and the resulting mutant phenotype suggests that this domain is necessary for TC activity in plants. The defective export of Suc in *sxd1* suggests that in addition to presumed antioxidant activities, tocopherols or tocopherol breakdown products also function as signal transduction molecules, or, alternatively, the DMPBQ that accumulates in *sxd1* disrupts signaling required for efficient Suc export in maize.

Tocopherols (α -, β -, γ -, and δ -tocopherol) are lipophilic antioxidants that collectively constitute vitamin E, an essential nutrient for both humans and animals. Tocopherol synthesis has only been observed in photosynthetic organisms (plants, algae, and some cyanobacteria), a distribution that suggests the pathway evolved in cyanobacteria to aid in protecting the cell from reactive oxygen species generated by photosynthesis. Plant tocopherol biosynthetic enzymes are nuclear encoded and were presumably acquired from the endosymbiotic cyanobacteria that gave rise to plastids (Goksoyr, 1967). The localization of tocopherols and most of the tocopherol biosynthetic enzymes in plastid membranes supports the cyanobacterial origins of the pathway in plants (Soll et al., 1980, 1985; Lichtenthaler et al., 1981; Fryer, 1992; Kruk and Strzalka, 1995; Arango and Heise, 1998a).

Although comparatively little is known about tocopherol functions in photosynthetic organisms, the physiological importance of these molecules in human and other animal systems has been studied extensively. The complete absence of dietary tocopherols, for example, results in chronic wasting, death, and fetal reabsorption in rats (Bramley et al., 2000). Less severe tocopherol dietary deficiencies in humans and animal models are associated with numerous degenerative diseases such as atherosclerosis, arthritis, some cancers, vision maladies, weakened immune system, and neuromuscular abnormalities (Brigelius-Flohe and Traber, 1999; Bramley et al., 2000; Ricciarelli et al., 2001).

Among the best characterized functions of tocopherols in cells is their ability to scavenge and quench reactive oxygen species and lipid-soluble by-products of oxidative stress (Brigelius-Flohe and Traber, 1999; Bramley et al., 2000; Ricciarelli et al., 2001). In addition to being lipophilic, tocopherols are capable of donating a single electron to form the resonance-stabilized tocopheroxyl radical (Liebler, 1993; KamalEldin and Appelqvist, 1996). Tocopherols are unique in this regard to other phenolic antioxidants, such as hydroxyquinones, which must donate two electrons to attain a stable structure. Tocopherols can also donate two electrons, which results in opening of the chromanol ring to form the corresponding tocoquinone derivative. These com-

¹ This work was supported in part by the Michigan State University Center for Novel Plant Products.

² Present address: U.S. Department of Agriculture-Agricultural Research Service Plant Genetics Research Unit, Donald Danforth Plant Science Center, 975 N. Warson Rd., St. Louis, MO 63132.

³ Present address: Agilent Technologies Inc., Little Falls Site, 2850 Centreville Rd, Wilmington, DE 19808.

* Corresponding author; e-mail dellapenna@msu.edu; fax 517-353-9334.

Article, publication date, and citation information can be found at www.plantphysiol.org/cgi/doi/10.1104/pp.103.024257.

bined molecular characteristics allow tocopherols to protect polyunsaturated fatty acids from lipid peroxidation by scavenging lipid peroxy radicals that propagate lipid peroxidation chain reactions in membranes (Burton et al., 1986; Liebler, 1993). Though direct evidence is lacking, tocopherols are thought to play similar roles in protecting the polyunsaturated fatty acid-rich plastid membrane from lipid peroxidation.

Recent studies in mammalian systems have demonstrated additional biological activities of tocopherols that are independent of their antioxidant functions. The underlying mechanisms for these effects are the modulation of signal transduction pathways by specific tocopherols and, in some instances, transcriptional activation of gene expression mediated by tocopherol-binding proteins (Brigelius-Flohe and Traber, 1999; Sen et al., 2000; Chan et al., 2001; Ricciarelli et al., 2001; Yamauchi et al., 2001; Clement et al., 2002; Nobata et al., 2002). Modulation of the protein kinase C signaling cascade and eicosanoid synthesis are two well-characterized examples of the antioxidant-independent effects of tocopherols in mammalian systems (Greenberg et al., 1993; Tran et al., 1996; Azzi et al., 2002). Although direct experimental evidence is lacking for antioxidant-independent tocopherol activities in plants, these data raise the possibility that tocopherols may also have roles in plants that extend beyond their proposed antioxidant functions.

Though the functions of tocopherols in plants remain an open question, much has been learned about tocopherol synthesis and the pathway enzymes during the past 5 years (Norris et al., 1998; Shintani and DellaPenna, 1998; Collakova and DellaPenna, 2001; Schledz et al., 2001; Porfirova et al., 2002; Savidge et al., 2002; Shintani et al., 2002). Tocopherol synthesis draws substrates from two separate metabolic pathways, aromatic amino acid metabolism and isoprenoid synthesis. Homogentisic acid, an intermediate in aromatic amino acid degradation and the head group of tocopherols, is produced from *p*-hydroxyphenylpyruvate by the cytosolic enzyme *p*-hydroxyphenylpyruvate dioxygenase (HPPD; Garcia et al., 1997, 1999; Norris et al., 1998; Dahnhardt et al., 2002). The isoprenoid-derived phytyl tail of tocopherols is a product of the plastid-localized 1-deoxy-D-xylulose-5-phosphate pathway (Eisenreich et al., 1998; Lichtenthaler, 1998). The remaining steps in tocopherol synthesis occur within the inner envelope of the chloroplast and include a phytyl transferase, two different methyltransferases, and a ring-producing enzyme, the tocopherol cyclase (TC; Soll et al., 1980, 1985; Arango and Heise, 1998a).

The TC adds a second oxygen-containing ring at the junction between the aromatic head group and phytyl tail to create a two-ring structure known as a chromanol ring (Fig. 1), which is essential for resonance stabilization of tocopheroxyl radicals after single-electron transfer. Previous work has characterized TC activity




Figure 1. Tocopherol biosynthetic pathway. This figure represents the enzymatic reactions and intermediates that are involved in tocopherol synthesis. 1, HPPD. 2, Homogentisate phytyl transferase (HPT). 3, 2-Methyl-6-phytyl-1,4-benzoquinone (MPBQ) methyltransferase. 4, TC. 5, γ -Tocopherol methyltransferase (γ -TMT). HPP, *p*-Hydroxyphenylpyruvate; HGA, homogentisic acid; SAM, S-adenosyl L-Met.

in chloroplasts and chromoplasts of higher plants and in cyanobacteria (Soll, 1979; Soll and Schultz, 1980; Soll et al., 1985; Stocker et al., 1993, 1994, 1996; Arango and Heise, 1998b). The primary substrate of the TC is reduced (quinol form) 2,3-dimethyl-6-phytyl-1,4-benzoquinone (DMPBQ), which is converted to γ -tocopherol by TC (Soll, 1979; Soll and Schultz, 1980; Soll et al., 1985). However, the enzyme characterized from *Anabaena* sp. PCC7120 has also been shown to cyclize other 6-prenyl-1,4-benzoquinol substrates in vitro (Stocker et al., 1996). In this report, we describe the isolation and functional characterization of TCs from *Arabidopsis*, maize (*Zea mays*), and *Synechocystis* sp. PCC6803 and discuss the evolutionary implications of tocopherol cyclization for both tocopherol synthesis and function. The identification and characterization of TC from *Arabidopsis* were recently reported by Porfirova et al. (2002).

RESULTS

Isolation and Characterization of *vte1* Mutants

To further understand the tocopherol pathway in higher plants, an HPLC-based screen of *Arabidopsis*

leaf tissue was developed to isolate mutants with tocopherol profiles that differ from wild type. Arabidopsis leaves accumulate approximately 10 ng α -tocopherol mg^{-1} fresh weight and 0.2 ng γ -tocopherol mg^{-1} fresh weight under standard growth conditions (see "Materials and Methods"). Numerous mutants were identified from an ethyl methanesulfonate (EMS)-mutagenized population, including two mutants that were devoid of tocopherols in leaf tissue (Fig. 2A). Genetic complementation tests confirmed that the mutants were allelic (data not shown). The mutants were designated *vte1-1* and *vte1-2* (vitamin e).

The visible phenotypes of both *vte1* mutants did not significantly differ from wild type when grown




Figure 2. HPLC analysis of tocopherols in wild-type and mutant Arabidopsis, maize, and *Synechocystis* sp. PCC6803. Tocopherols present in Arabidopsis, maize, and *Synechocystis* sp. PCC6803 lipid extracts were separated by normal phase HPLC and detected using a fluorescence detector with 290-nm excitation and 325-nm emission. Tocol, a synthetic tocopherol, was used as an internal recovery standard. A, Arabidopsis leaf tissue: solid line, Columbia wild type; dotted line, *vte1-1*; gray line, *vte1-2*. B, Maize leaf tissue: solid line, wild type; dotted line, *sxd1*. C, *Synechocystis* sp. PCC6803: solid line, wild type; dotted line, Δ *slr1737* insertional mutant; gray line, SXD1 expressed in the Δ *slr1737* insertional mutant. Retention times of α -, β -, δ -, and γ -tocopherol and tocol were determined by HPLC analysis of tocopherol standards. LU, Luminescence units.

under normal laboratory conditions (see "Materials and Methods"). Although several possibilities could result in a tocopherol-deficient phenotype, the two most likely are a loss of HPT activity or a loss of TC activity (Fig. 1). Assuming no genetic redundancy, a mutation disrupting either gene would result in a tocopherol-deficient phenotype, but the two classes of mutations should be readily distinguishable by the intermediates that accumulate. A defect in the TC should result in the accumulation of the DMPBQ, whereas a mutation in HPT would not accumulate tocopherol pathway prenyl quinone intermediates (Fig. 1). To understand the biochemical basis of *vte1-1* and *vte1-2*, prenyl quinones were isolated from each mutant and analyzed by HPLC (Fig. 3A). A novel peak with a retention time and spectrum consistent with the prenyl quinone DMPBQ (Hutson and Threlfall, 1980; Marshall et al., 1985; Johnson et al., 2000) was observed in the *vte1-1* and *vte1-2* mutants but not in wild type (Fig. 3A). This compound was purified by HPLC and a mass of 415 D, the mass of DMPBQ, was determined by fast atom bombardment mass spectroscopy (data not shown). These combined characteristics indicate that the novel compound that accumulates in *vte1-1* and *vte1-2* mutants is DMPBQ, the substrate of the TC.

In addition to green tissues, seeds also contain tocopherols, but instead of α -tocopherol predominating as in leaves, γ -tocopherol accumulates due to low γ -TMT activity (Shintani and DellaPenna, 1998). Lipids were extracted from the seeds of wild-type and *vte1* mutants, and the tocopherols were analyzed by HPLC (Fig. 4). Tocopherols were absent in *vte1-1* seed, but *vte1-2* seeds contained approximately 25% of the tocopherol level in wild type (84.9 ± 0.8 versus 322.5 ± 7.4 ng total tocopherols mg^{-1} seed in *vte1-2* and wild type, respectively), suggesting *vte1-2* is a weaker allele than *vte1-1*.

A map-based cloning approach was undertaken to isolate the gene encoding the TC from Arabidopsis. *vte1-1* was crossed to Landsberg *erecta*, and 1,100 individuals from an F₂ population were used to map the *VTE1* locus to a 140-kb interval on the bottom of chromosome 4. Analysis of the genes within this interval identified At4g32770, encoding an unknown protein of 488 amino acids that contains a putative N-terminal chloroplast transit peptide of 68 amino acids. Previously, At4g32770 and the *Synechocystis* sp. PCC6803 protein *slr1737* were identified as homologs of SXD1 (Suc Export Deficient 1) from Maize (Provencher et al., 2001). *slr1737* is a protein of unknown function and is in the same operon as *slr1736*, which encodes HPT (Collakova and DellaPenna, 2001; Schledz et al., 2001; Savidge et al., 2002), another enzyme of the tocopherol biosynthetic pathway (Fig. 1). At4g32770 was fully sequenced in both mutants, and each was found to contain a nonsense mutation. The *vte1-1* mutation creates a premature




Figure 3. HPLC analysis of the prenyl quinones from wild-type and mutant *Arabidopsis*, maize, and *Synechocystis* sp. PCC6803. Lipids were extracted from *Arabidopsis*, maize, and *Synechocystis* sp. PCC6803, and total prenyl quinones were isolated by thin-layer chromatography (TLC) and then analyzed by normal phase HPLC (see "Materials and Methods") A, *Arabidopsis*. Solid line, Columbia wild type; dotted line, *vte1-1*; gray line, *vte1-2*. B, Maize. Solid line, Wild type; dotted line, *sxd1*. C, *Synechocystis* sp. PCC6803. Solid line, Wild type; dotted line, Δ *slr1737* insertional mutant; gray line, *SXD1*cDNA expressed in the Δ *slr1737* mutant background. Insets, Spectra of the peak labeled DMPBQ, Phyllo, Phylloquinone; PQ, Plastoquinone.

stop codon at amino acid 237 (Trp to stop), whereas the *vte1-2* mutation creates a premature stop codon at amino acid 465 (Gln to stop). Both mutations are base transitions, G to A and C to T, respectively, which is consistent with the mutagenic properties of EMS. The identification of At4g32700 as the *Arabidopsis* TC concurs with a recent report by Porfirova et al. (2002).

Sequence Analysis and Evolutionary Origin

BLAST searches revealed that *SXD1*, *VTE1*, and *slr1737* share a high degree of amino acid sequence similarity (Table I) with other proteins in the nonredundant GenBank database: three proteins of unknown function in the cyanobacteria *Anabaena* sp. PCC7120, *N. punctiforme*, and *Synechococcus* sp. PCC7002. *SXD1* is a chloroplast-targeted protein of unknown function that had been identified previously based on a mutation causing a defect in symplastic photosynthate transport near the site of phloem loading within the minor veins of maize leaves (Russin et al., 1996; Provencher et al., 2001). When the *SXD1* and *VTE1* chloroplast transit peptides are removed, *SXD1*, *VTE1*, and the four cyanobacterial proteins share long stretches of amino acid identity.

The four cyanobacterial proteins are assumed to be orthologs of *VTE1* because the cyanobacterial genomes each contain obvious orthologs of the four other known genes of the tocopherol pathway: HPPD, HPT, MPBQ methyltransferase, and γ -TMT. *VTE1* and the cyanobacterial orthologs exist as single genes within their respective sequenced genomes. There are several other cyanobacteria whose genomes also have been sequenced: *Prochlorococcus ma-*




Figure 4. HPLC analysis of seed tocopherols in wild-type *Arabidopsis*, *vte1-1*, and *vte1-2*. Total seed lipids were extracted, and the tocopherols present were separated by reverse phase HPLC and detected using a fluorescence detector; 290-nm excitation and 325-nm emission. Tocol, a synthetic tocopherol, was used as an internal recovery standard. Solid line, Columbia wild type; dotted line, *vte1-1*; gray line, *vte1-2*. Retention times of α -, δ -, and γ -tocopherol and tocol were determined by HPLC analysis of tocopherol standards.

Table I Pairwise comparisons of VTE1 orthologs from plants and cyanobacteria

Pair-wise comparisons were performed using ClustalW and are expressed as percentage amino acid similarity. The predicted mature plant protein sequences were used for alignments. *Anabaena*, *Anabaena* sp. PCC7120 (all0245); *Nostoc*, *Nostoc punctiforme* (506-74); *Synecho*, *Synechococcus* sp. PCC7002.

	SXD1	VTE1	<i>Medicago truncatula</i>	Barley (<i>Hordeum vulgare</i>)	slr1737	<i>Synecho</i>	<i>Nostoc</i>
				%			
VTE1	79						
<i>M. truncatula</i>	80	84					
Barley	92	80	83				
slr1737	46	46	48	48			
<i>Synecho</i>	47	49	49	48	63		
<i>Nostoc</i>	55	55	54	55	59	67	
<i>Anabaena</i>	55	54	54	55	61	67	85

rinus MED4, *P. marinus* MIT9313, *Synechococcus* sp. PCC7002 WH8102, and *Thermosynechococcus elongatus* BP-1. The genomes of these organisms lack obvious VTE1 orthologs and obvious orthologs for HPPD, HPT, and γ -TMT (refer to pathway in Fig. 1). Thus, it appears likely that only a subgroup of cyanobacteria have evolved the ability to synthesize tocopherols.

VTE1, SXD1, and the four cyanobacterial orthologs lack any previously described protein motifs. There are numerous plant expressed sequence tags (ESTs) in the public database that share high similarity with VTE1 and SXD1, and full-length sequences of the *M. truncatula* and barley VTE1 orthologs were obtained from EST assemblies. These four representative plant sequences are more conserved than the four cyanobacterial sequences (Table I). Sequence alignment of the plant and the cyanobacterial protein sequences identified a highly conserved 30-amino acid carboxyl domain in the plant VTE1 orthologs (starting at Thr-458 of Arabidopsis VTE1) that is absent from the cyanobacteria proteins (Fig. 5). The last five amino acids of this carboxyl domain (KPPGL) are invariant among the plants represented, which include the bryophyte *P. patens*, monocots, and dicots. With the exception of vascular and nonvascular VTE1 orthologs, this 30-amino acid domain was not found in other proteins in the nonredundant database. Interestingly, the *C. reinhardtii* VTE1 ortholog has a short-

ened version of the carboxyl domain (Fig. 5) and lacks the last 12 amino acids (starting at Leu-477 of Arabidopsis VTE1), including the invariant KPPGL motif. The *vte1-2* mutation causes premature termination of VTE1 and deletion of 24 amino acids of the conserved carboxyl domain.

TC Function in Arabidopsis, Maize, and *Synechocystis* sp. PCC6803

To confirm that the VTE1 orthologs are required for tocopherol synthesis in plants other than Arabidopsis, lipids were isolated from leaves of the *sxd1* mutant and analyzed for tocopherols by HPLC. As with *vte1-1* and *vte1-2*, leaves of the *sxd1* mutant lack tocopherols, whereas wild-type maize leaves contain both α - and γ -tocopherols (Fig. 2B). Prenyl quinones from leaves of wild-type maize and the *sxd1* mutant were also analyzed by HPLC. This analysis indicated that like the *vte1* mutants, *sxd1* contained a prenyl quinone that was absent from wild type (Fig. 3B) with an absorbance spectrum and retention time identical to the DMPBQ that accumulated in *vte1* mutants.

To show that this gene family has an identical function in cyanobacteria and plants, an insertional mutant, Δ slr1737, was created in the slr1737 open reading frame (ORF) of *Synechocystis* sp. PCC6803




Figure 5. Alignment of the carboxyl termini of TC orthologs from plants and cyanobacteria. The asterisk above the At4g32770 protein sequence denotes the position of the *vte1-2* mutation. *Anabaena*, *Anabaena* sp. PCC7120 (all0245); *Chlamy*, *Chlamydomonas reinhardtii*; *Medicago*, *M. truncatula*; *Nostoc*, *N. punctiforme* (506-74); *Physcomitrella*, *Physcomitrella patens*; *Synecho*, *Synechococcus* sp. PCC7002. The *P. patens*, rice (*Oryza sativa*), and wheat (*Triticum aestivum*) sequences are partial sequences obtained from ESTs. Dark shading, Amino acid identity; light shading, Amino acid similarity. The threshold for amino acid consensus identity or similarity is 51%.




Figure 6. TC activity of proteins expressed in *E. coli*. *E. coli* cell lysates from cells overexpressing the empty pET vector or pET engineered to express TC proteins from Arabidopsis, maize, and *Synechocystis* sp. PCC6803 were incubated with radiolabeled 2,3-methyl-6-phytyl-1,4-benzenequinol (3 methyl ^{14}C) for 4 h as described in "Materials and Methods." Total lipids were extracted, separated by TLC, and radiolabeled products were detected by phosphor imager analysis. Products were identified by comigration with standards. The ^{14}C incorporation into γ -tocopherol was quantified densitometrically and expressed as pixels per microgram of total protein.

through homologous recombination. The growth rate of $\Delta\text{slr1737}$ was indistinguishable from wild type (data not shown). Lipids were extracted from $\Delta\text{slr1737}$ and wild-type cells and analyzed for tocopherol composition by HPLC. $\Delta\text{slr1737}$ lacked α -tocopherol, similar to the *vte1* and *sxd1* mutants (Fig. 2). The two unknown peaks detected in $\Delta\text{slr1737}$ immediately after α -tocopherol are also present in the tocopherol-deficient mutants $\Delta\text{slr1736}$ and $\Delta\text{slr0090}$ (Dahnhardt et al., 2002) and do not correspond to tocopherols or tocopherol pathway intermediates. HPLC analysis of prenyl quinones in $\Delta\text{slr1737}$ and wild type showed that, like *vte1* and *sxd1*, $\Delta\text{slr1737}$ accumulates DMPBQ (Fig. 3C).

As further proof that the cyanobacterial and plants genes are functionally equivalent, a SXD1 cDNA expression cassette was transformed into the $\Delta\text{slr1737}$ line. Lipids were isolated from the cells and analyzed by HPLC for tocopherols. The transformed cells contained α -tocopherol (Fig. 2C) and did not accumulate DMPBQ (Fig. 3C). Thus, expression of SXD1 restored a wild-type tocopherol profile to $\Delta\text{slr1737}$, indicating

that SXD1 is able to functionally complement $\Delta\text{slr1737}$. Hence, these cyanobacterial and plant genes have not only high sequence similarity but also functional equivalency, suggesting a common evolutionary ancestry.

To determine the activity of the VTE1 protein and its maize and cyanobacterial orthologs, we expressed VTE1, SXD1, and *slr1737* in *Escherichia coli* using the pET expression system. Lysates from *E. coli* expressing either VTE1 or SXD1 were able to convert [^{14}C]2,3-dimethyl-6-phytyl-1,4-benzoquinol into γ -tocopherol (Fig. 6). This result conclusively demonstrates that both genes encode an enzyme with TC activity. Activity was not observed with the *slr1737* protein expressed in *E. coli* for reasons that are unknown.

Carbohydrate Assimilation in *vte1*

Although SXD1 and VTE1 have similar enzymatic activities (Fig. 6) and primary biochemical phenotypes (tocopherol deficiency and DMPBQ accumulation, Figs. 2 and 3), *sxd1* was initially isolated because of a secondary phenotype, a Suc transport defect (Russin et al., 1996; Provencher et al., 2001). To determine whether *vte1* caused a similar Suc transport defect, Glc, Suc, and starch levels were analyzed in mature leaves from 4-week-old *vte1-1* plants. The leaves were sampled at the end and at the beginning of the photoperiod. There were no significant differences between wild-type and *vte1-1* leaves for Glc, Suc, and starch at the beginning or end of the photoperiod (Table II). Thus, unlike the *sxd1* mutant, carbohydrate metabolism in mature leaves appears unaffected by the *vte1* mutation. The molecular and biochemical similarities of the VTE1 and SXD1 proteins suggest the difference in *sxd1* and *vte1* carbohydrate metabolism phenotypes in maize and Arabidopsis reflects additional roles of tocopherols or the tocopherol pathway beyond antioxidant chemistry, rather than simply a difference in the enzymatic activity of the SXD1 and VTE1 proteins.

DISCUSSION

In this report, we have shown that the three proteins, VTE1, SXD1, and *slr1737* from a dicot, monocot,

Table II. Analysis of Glc, Suc, and starch in wild-type Arabidopsis and *vte1-1*

Glc, Suc, and starch levels were analyzed spectrophotometrically using the enzyme-coupled assays described in "Materials and Methods." Glc and Suc are expressed as nanomoles per milligram fresh wt ($n = 4$).

Carbohydrates	Beginning of Photoperiod		End of Photoperiod	
	Columbia	<i>vte1-1</i>	Columbia	<i>vte1-1</i>
Glc	2.32 \pm 0.62	2.46 \pm 0.58	2.36 \pm 0.79	2.63 \pm 0.83
Suc	0.07 \pm 0.07	0.16 \pm 0.16	1.56 \pm 0.60	1.75 \pm 0.28
Starch ^a	13 \pm 3	13 \pm 4	86 \pm 14	85 \pm 11

^a Starch is expressed as nanomoles of Glc monomers per milligram fresh wt.

and cyanobacterium, respectively, function as TCs. Mutations in the TC gene from each organism result in identical primary biochemical phenotypes, a block in tocopherol synthesis, and accumulation of DMPBQ, the endogenous substrate for the TC. In addition, the SXD1 and VTE1 proteins expressed in *E. coli* were able to convert DMPBQ to γ -tocopherol. Finally, expression of maize SXD1 was sufficient to complement the tocopherol-deficient phenotype of the *Synechocystis* sp. PCC6803 slr1737 deletion mutant (Δ slr1737). This result demonstrates that slr1737 and SXD1 are functionally equivalent and that the biochemical activity of TCs has been evolutionarily conserved between plants and cyanobacteria. Our finding that At4g32770 encodes a functional TC in *Arabidopsis* concurs with a recent report by Porfirova et al. (2002).

Sequence Analysis and Evolutionary Implications of the TC Family

The TCs (VTE1, SXD1, and slr1737) share significant amino acid similarity with each other and define an evolutionarily conserved gene family that includes putative orthologs in a large number of other plants and cyanobacteria. VTE1 orthologs were not identified in databases of fungal, animal, or non-photosynthetic bacterial species, none of which are known to produce tocopherols. Full-length sequences of two additional VTE1 orthologs from plants (barley and *M. truncatula*) and three from cyanobacteria (*N. punctiforme*, *Anabaena* sp. PCC7120, and *Synechococcus* sp. PCC7002) were identified in the public databases. Although all VTE1 proteins share a high degree of amino acid similarity, they are devoid of any previously described protein motifs, with the exception of ubiquitous phosphorylation and myristylation motifs. All VTE1 orthologs are hydrophobic proteins with low pIs and a high number of conserved Trp residues (Provencher et al., 2001). These characteristics are consistent with the TC activity characterized in *Anabaena variabilis* being membrane associated (Stocker et al., 1993, 1994, 1996).

Although plant and cyanobacterial TCs exhibit a high degree of protein sequence similarity, plant orthologs have additional N- and C-terminal domains that are absent in the four cyanobacterial TCs. The N-terminal domains of plant VTE1 orthologs are poorly conserved and are predicted to encode chloroplast transit peptides that would target each protein to the chloroplast. The N-terminal sequence of SXD1 has been demonstrated experimentally to be required for import into plastids (Provencher et al., 2001). HPT and γ -TMT, two other tocopherol biosynthetic enzymes, are also predicted to be chloroplast targeted (Shintani and DellaPenna, 1998; Collakova and DellaPenna, 2001; Schledz et al., 2001; Savidge et al., 2002), and chloroplastic localization of the TC is consistent with the reported localization of TC activ-

ity and tocopherol synthesis in plastids (Soll et al., 1985; Arango and Heise, 1998a).

In contrast to the N-terminal domain of plant TCs, the 30-amino acid C-terminal domain is highly conserved between angiosperms and the moss *P. patens*. This evolutionary conservation suggests an important function for this domain in tocopherol synthesis in plants, whereas the absence of the sequence from the four cyanobacterial VTE1 orthologs suggests that the domain is not an absolute requirement for TC enzymatic activity per se. The restriction of this C-terminal domain to vascular and nonvascular plants suggests it arose relatively recently in progenitors of land plants rather than in the endosymbiotic cyanobacteria that gave rise to plastids (Goksoyr, 1967). The shortened C-terminal domain present in the *C. reinhardtii* TC further suggests evolution took place in an ancestor common to *C. reinhardtii* and plants (i.e. before the split of Chlorophyta and Charophyta; Karol et al., 2001). The *vte1-2* mutation causes deletion of the majority of the conserved C-terminal domain, and tocopherols fail to accumulate in *vte1-2* leaf tissue but reach 25% of wild-type levels in seeds, indicating the truncated protein retains at least partial activity in vivo. The *vte1-2* phenotype suggests the C-terminal domain plays a more significant role in TC activity/function in leaf chloroplasts than in the plastids of seeds. The complementation of Δ slr1737 by SXD1 suggests that the presence of the C-terminal domain does not affect TC activity and function in cyanobacteria. The relevance of this highly conserved carboxyl domain for TC function in plants requires further investigation.

The *sxd1* and *vte1* Phenotypes

A surprising phenotype of the *sxd1* mutants is a block in Suc export from leaves and an accumulation of anthocyanins and starch in leaf blades (Russin et al., 1996). This pleiotropic phenotype results from aberrant plasmodesmata at the interface between the bundle sheath cells and the vascular parenchyma cells surrounding the minor veins. These defective plasmodesmata block symplastic transport of Suc to the phloem and, hence, cause the Suc export defect. Unlike *sxd1*, *vte1-1* and *vte1-2* do not accumulate anthocyanins in leaves, do not accumulate starch in cells surrounding the leaf veins, and do not have stunted growth (data not shown). In addition, the similar Suc, Glc, and starch levels in mature leaves of *vte1-1* and wild type (Table II) indicate that a functional Suc export pathway is present in *vte1*. However, this most obvious difference in phenotype between *sxd1* and *vte1* is not entirely unexpected considering the anatomical and physiological differences between C3 and C4 plants. Maize bundle sheath cells contain chloroplasts that differ morphologically and physiologically from the chloroplasts of mesophyll cells (Evert et al., 1977a; Fahn, 1990).

Maize bundle sheath plastids have few grana, little PSII activity, high NADP-malate decarboxylase activity, and accumulate starch (Evert et al., 1977a, 1977b, 1978; Fahn, 1990). Arabidopsis lacks a physiologically equivalent cell type to the C4 bundle sheath cell. Chloroplasts within the analogous cells surrounding the minor veins in Arabidopsis do not differ significantly from the chloroplasts of mesophyll cells (Haritatos et al., 2000). Mutants of the Arabidopsis *SUC2* gene, which encodes a Suc-H⁺ symporter required for apoplastic phloem loading, share striking similarities to *sxd1* mutants (Gottwald et al., 2000). *suc2* mutants accumulate anthocyanins and starch in their cotyledons, have stunted growth, and are seedling lethal in soil (Gottwald et al., 2000). Thus, the absence of the *sxd1* Suc export phenotype in *vte1* suggests that disruption of *VTE1* in Arabidopsis either does not affect Suc export, most likely because of the fundamentally different mechanisms of Suc export in maize and Arabidopsis, or that the effect is too small to be observable as an analogous whole-plant phenotype in Arabidopsis.

In *sxd1* mutants, the link between the production of aberrant plasmodesmata in the BS parenchyma cells and the defect in symplastic transport of Suc was straightforward and easy to rationalize (Provencher et al., 2001). However, the mechanistic link between the disruption of a gene encoding a chloroplast protein of unknown function (*SXD1*) and defective Suc transport was not so obvious. Provencher et al. (2001) raised the possibility that the *sxd1* mutation exerts its pleiotropic phenotype by disrupting or altering a signal from the chloroplast to the nucleus. The nature of this signal was unknown, and cloning of the *SXD1* locus did not provide further insight. Although the precise nature of this signal remains unclear, the demonstration that *SXD1* encodes a TC and determination of the primary *sxd1*, *vte1*, and Δ slr1737 biochemical phenotypes now greatly limits the possibilities.

The absence of tocopherols in *vte1* and Δ slr1737 does not cause a pleiotropic phenotype analogous to *sxd1*. *vte1* mutants are indistinguishable from wild type under normal growth conditions, and Δ slr1736 and Δ slr1737 mutants grow at rates identical to wild type (Collakova and DellaPenna, 2001). These data suggest that the *sxd1* phenotype is not simply due to the absence of tocopherols as lipophilic antioxidants because similar affects would be observed in the *vte1*, Δ slr1737, and Δ slr1736 mutants, which also lack tocopherols. In addition, the DMPBQ that accumulates in *vte1*, *sxd1*, and Δ slr1737 mutants can still act as an antioxidant by donating a pair of electrons and then being recycled by reduction back to the quinol form of DMPBQ (Kruk et al., 1994; Kruk and Strzalka, 1995; Liebler and Burr, 2000). Thus, even in the absence of tocopherols, the *sxd1* mutant is not entirely deficient in membrane-associated antioxidants.

Influencing membrane fluidity is another potential function of tocopherols, and this could be related to the plasmodesmatal defect in *sxd1* mutants. However, the consensus from cell fractionation studies is that tocopherols are localized exclusively in plastid membranes (Lichtenthaler et al., 1981; Fryer, 1992; Kruk and Strzalka, 1995), and although it cannot be excluded, it is unlikely that tocopherols or DMPBQ would be present in plasma membranes to directly impact plasmodesmatal development. However, the bundle sheath cell is a specialized cell type, and direct physical connections between plasmodesmata and chloroplasts through the endoplasmic reticulum have been reported in maize bundle sheath cells (Evert et al., 1977a, 1977b). This potential association between the chloroplast and plasmodesmata would provide a means for tocopherols or prenyl quinones to impact membrane fluidity at the plasma membrane. Still, it is difficult to envision a mechanism whereby membrane fluidity would alter plasmodesmata development. If diminished antioxidant capacity or altered membrane fluidity is not the cause of the *sxd1* phenotype, the question still remains: How does disruption of TC activity result in the pleiotropic *sxd1* phenotype?

In addition to the well-defined role of tocopherols as antioxidants, specific tocopherols, tocotrienols, and their oxidized products have been demonstrated to have biological activities in mammalian systems that are independent of their antioxidant functions. The unifying theme for these antioxidant-independent activities is the modification or modulation of various signal transduction pathways (Brigelius-Flohe and Traber, 1999; Ricciarelli et al., 2001; Clement et al., 2002). The effects of tocopherols on the protein kinase C signaling cascade and the synthesis of eicosanoids in mammals have been well characterized. α -Tocopherol posttranslationally inhibits the activity of protein kinase C in several mammalian systems (Chan et al., 2001; Azzi et al., 2002; Clement et al., 2002). Plant genomes also contain protein kinase C homologs and other components of this signaling pathway. Tocopherols also have been shown to posttranslationally inhibit the activity of phospholipase A₂ (Tran et al., 1996; Chandra et al., 2002), cyclooxygenase (COX-2; Jiang et al., 2000; Wu et al., 2001), and lipoxygenase-5 (Greenberglevy et al., 1993; Wang et al., 2000; Ricciarelli et al., 2002). These enzymes are involved in the production of eicosanoid signaling molecules (prostaglandins, prostacyclins, thromboxanes, and leukotrienes) from polyunsaturated fatty acids in mammals (Ohuchi and Levine, 1980; Tran et al., 1996; Kim et al., 2001). Like eicosanoids in animals, jasmonic acid and other oxylipins in plants are synthesized from polyunsaturated lipids by the action of lipoxygenase(s) and phospholipase(s) (Blee, 2002; Howe and Schillmiller, 2002). A third example of α -tocopherol-dependent, antioxidant-independent signal transduction in mammals is the transcriptional activator TAP

(tocopherol-associated protein). Upon the binding of α -tocopherol by TAP, the complex is translocated into the nucleus, where it has been shown to activate the transcription of transgenes (Yamauchi et al., 2001).

Although tocopherols have not yet been shown to affect protein kinase C signaling, transcriptional regulation, or the synthesis of jasmonic acid or other oxylipins in plants, studies in mammalian systems suggest plausible mechanisms whereby the absence of TC activity (and, hence, tocopherols) could affect signaling and result in the pleiotropic *sxd1* phenotype. Thus, it is possible that many tocopherol functions will be universal, including the roles tocopherols play in modulating signal transduction pathways or acting as signals themselves. Although the downstream events of signal transduction pathways would not necessarily be evolutionarily conserved between plants and mammals, many of the core components and biochemical motifs of signal transduction pathways are. We suggest that the *sxd1* phenotype is the first evidence that tocopherols act as signaling molecules or modulators of signaling in plants. Tocopherols, tocopherol derivatives, or tocopherol pathway intermediates may provide or modulate signals required for the development of maize bundle sheath vascular parenchyma plasmodesmata, analogous to the effects of tocopherols in mammalian signaling. Alternatively, the DMPBQ that accumulates in *sxd1* may interfere with an endogenous signaling pathway required for the process. Several groups have provided evidence that the redox status of the chloroplast, which is monitored through the plastoquinone (PQ) pool, regulates nuclear-encoded photosynthetic gene expression (Pfannschmidt et al., 1999a, 1999b, 2001; Alfonso et al., 2000; Allen and Pfannschmidt, 2000; Kujat and Owtrim, 2000; Li and Sherman, 2000; Pursiheimo et al., 2001; Trebitsh and Danon, 2001; Yang et al., 2001). DMPBQ has the same 2,3-dimethyl-1,4-benzoquinone head group as PQ and could interfere with redox signaling through the PQ pool. Another more remote possibility is that the SXD1 protein has an unknown substrate in addition to DMPBQ, and this product is required for signal transduction in maize. Although none of these models can be excluded based on the present data, they do provide a framework for future study.

The observation that Arabidopsis *vtel* mutants do not exhibit phenotypes analogous to *sxd1* suggests that the downstream signal transduction events impacted by tocopherol deficiency differ between monocots and dicots. The pathways leading to maize bundle sheath vascular parenchyma plasmodesmata formation may either be absent or not equivalent in Arabidopsis, or the effects are too subtle to be observed at the whole-organism level as in *sxd1*. Experiments to assess the whole-genome responses of Arabidopsis *vtel* mutants are under way.

MATERIALS AND METHODS

Growth Conditions and Seed Stocks

Arabidopsis plants were grown at 22°C under a 12-h photoperiod (120 μ E) in a vermiculite and potting soil mixture. M₃ EMS-mutagenized Arabidopsis seeds (Columbia ecotype) were purchased from Lehle Seed (Round Rock, TX). *vtel-1* was backcrossed to wild type three times, and *vtel-2* was backcrossed twice. Maize (*Zea mays*) plants were grown under greenhouse conditions in the same soil mixture and fertilized biweekly with 20-20-20 fertilizer. The maize *sxd1-2* allele used in this publication was isolated through the Trait Utility System for Corn (Pioneer Hybrids, Johnston, IA). The *sxd1-2* Mu insertion site and mutant phenotype were described previously (Provencher et al., 2001). *Synechocystis* sp. PCC6803 was grown on BG-11 media photoautotrophically or photomixotrophically (BG11 media containing 15 mM Glc) on plates or in liquid culture at 30°C and 50 to 70 μ E light.

Tocopherol Analysis

For tocopherol analyses, total lipids were extracted from 30 to 35 mg of Arabidopsis or maize leaf tissue or 15 to 20 mg of plate-grown *Synechocystis* sp. PCC6803 cells (Bligh and Dyer, 1959; Collakova and DellaPenna, 2001), and dissolved in 100 μ L of methanol or hexane. Methanol extracts (50 μ L) were subject to HPLC (Agilent 1100 series, Agilent, Wilmington, DE) on a Spherisorb ODS-2 5- μ m, 250- \times 4.6-mm reverse phase column (Column Engineering, Ontario, CA) at 28°C with a flow rate of 2 mL min⁻¹ with 95% (v/v) methanol and 5% (v/v) isopropanol. Hexane extracts (50- μ L volume) were subjected to HPLC on a ReliaSil Silica 5- μ m, 250- \times 4.6-mm normal phase column (Column Engineering) at 42°C with a flow rate of 2 mL min⁻¹ with 83% (v/v) hexane and 17% (v/v) isopropyl ether. Tocopherols were detected by fluorescence using 290-nm excitation and 325-nm emission.

Analysis of Prenyl Quinones

One gram of Arabidopsis or maize leaf tissue and a 500-mL culture of *Synechocystis* sp. PCC6803 (OD₇₃₀ = 0.8) were harvested and total lipids extracted (Collakova and DellaPenna, 2001). Prenyl quinones were purified by TLC as described by Pennock (1985) and eluted with diethyl ether. After drying, the samples were resuspended in 500 μ L of hexane. A 100- μ L aliquot was dried, resuspended in 75 μ L of isopropanol, and subjected to HPLC on a reverse phase column (described above) under conditions previously described by Johnson et al. (2000). Alternatively, 60 μ L of hexane extract was subjected to HPLC, on a normal phase column (described above), at 30°C with a 1 mL min⁻¹ flow rate using 0.1% (v/v) dioxane in hexane. Prenyl quinones were detected by A₂₅₄ using a diode array detector.

Map-Based Cloning of *vtel*

PCR-based markers were designed using INDEL or SNP from the Cereon Arabidopsis Polymorphism and Landsberg *erecta* Sequence Collection (Cereon Genomics LLC, Cambridge, MA; Jander et al., 2002). DNA was extracted from 1- to 2-mm developing leaves using Plant DNazol (Invitrogen, Carlsbad, CA) following the protocol of the manufacturer. Alternatively, DNA was isolated from 1- to 2-mm developing leaves in 1.1-mL tubes arrayed in a 96-well format. Two 4-mm glass beads were added to tubes along with 200 μ L of 10 mM Tris (pH 8.0) and 200 μ L of chloroform. The tubes were shaken with a commercial paint shaker for 5 min and centrifuged at 3,750 rpm for 10 min. One microliter of the aqueous phase or resuspended DNA from DNazol extractions was used in a 20- μ L PCR reaction.

Sequence Analysis

All DNA sequences other than *vtel* mutant alleles were obtained from public databases using BLAST: wheat (*Triticum aestivum*; BQ619591), rice (*Oryza sativa*; AU031770), *Physcomitrella patens* (BJ164574), *M. truncatula* (BF641171 and TC48011), barley (*Hordeum vulgare*; TC33553 and TC32886), Arabidopsis (AF302188), and maize (AF302187). A TC prefix denotes sequences obtained from The Institute for Genomic Research. All others have GenBank accession numbers. The cyanobacteria and algae sequences were

obtained from their respective genome sequencing projects. *Synechocystis* sp. PCC6803 and *Anabaena* sp. PCC7120 sequences were obtained from Cyanobase (<http://www.kazusa.or.jp/cyano/cyano.html>). *Chlamydomonas reinhardtii* and *Nostoc* sequences were obtained from the Joint Genome Institute (<http://www.jgi.doe.gov>). The *Synechococcus* sp. PCC7002 sequence was obtained from National Center for Biotechnology Information (www.ncbi.nlm.nih.gov). Alignments were performed using MacVector 7.0 (Oxford Publishing, London), which includes the ClustalW algorithm.

Construction of *slr1737*, *SXD1*, and *VTE1* Protein Expression Vectors

Primers 5'-CATATGACCCTAATTTATCTTCCTTTG-3' (F₁), 5'-CATATG-GACAAGATCTCCGTTAAACCTG-3' (F₂), 5'-CTCGAGTTACAGACCCGGT-GGCTTG-3' (R₁), and turbo *Pfu* (Stratagene, La Jolla, CA) were used to PCR amplify the *VTE1* cDNA from a seed cDNA library (a gift from Dr. John Ohlrogge, Michigan State University, East Lansing). F₁ was used to amplify the full-length cDNA, and F₂ was used to amplify a truncated version of the cDNA that encodes a version of the protein lacking chloroplast transit peptide. PCR products were cloned into the *Sma*I site of pBluescript II SK⁻. The two versions of the *VTE1* cDNA were cloned as *Nde*I-*Xho*I fragments (sites underlined in primers) into *Nde*I and *Xho*I sites of the pET30b expression vector (Novagen, Madison, WI).

Primers 5'-CATATGAAATTTCCGCCACAGTGGTTAC-3' (F₃) and 5'-GGATCCTAACGAATCAAACAAGGC-3' (R₂), and turbo *Pfu* were used to amplify the *slr1737* gene from *Synechocystis* sp. PCC6803 genomic DNA. The PCR product was cloned into the *Eco*RV site of pBluescript II SK⁻. The *slr1737* gene was cloned as *Nde*I-*Bam*HI fragments (sites underlined in primers) into *Nde*I and *Bam*HI sites of the pET30b expression vector (Novagen).

Primers 5'-TTCATATGGCAACGCCGCATAGCGGGTACCAC-3' (F₄) and 5'-TTGCGCGCGCTTCATTGTGACATTCGTTGG (R₃) were used to amplify the *SXD1* cDNA without a chloroplast transit peptide. The PCR product was cloned into *Nde*I and *Not*I sites (sites underlined in primers) of pET24d (Novagen). The fidelity of all constructs was confirmed by sequencing.

Generation of *Δslr1737*

Primers 5'-CTGTGTATTCTGACGGTGC-3' (F₅) and 5'-GGAGATTG-AGAGAAATTTATGATGC-3' (R₄) and *Pfu* polymerase (Stratagene) were used to PCR amplify the 5' region flanking *slr1737* from *Synechocystis* sp. PCC6803 genomic DNA. Primers 5'-ATAAATCTCTCAATCTCCGTACCGAATAA-CACTGCCTGTGTTTG-3' (F₆) and 5'-ACCTGTTCTTCTAACCAGTIG-3' (R₅) and *Pfu* polymerase (Stratagene) were used to PCR amplify the 3' region flanking *slr1737* from *Synechocystis* sp. PCC6803 genomic DNA (underlined nucleotides indicate an added *Bsi*WI restriction site). The 5'- and 3'-flanking PCR products were joined through re-amplification with *Pfu* polymerase using Primers F₅ and R₅ to generate a contiguous 1,041-bp fragment containing a *Bsi*WI site separating the 5'- and 3'-flanking portions. The PCR product was cloned into the pPCR-Script AMP vector (Stratagene) to generate pSLR1737-5'-3' flank. The *aadA* gene encoding spectinomycin adenyltransferase was inserted as a *Bsi*WI fragment into the corresponding site of pSLR1737-5'-3' flank. The resulting plasmid was then used to replace ORF *slr1737* by homologous recombination (Williams, 1988). Recombinant lines were selected by spectinomycin resistance, and replacement of the *slr1737* ORF with the *aadA* gene was confirmed by PCR.

Complementation of *Δslr1737* by Expression of the Maize *SXD1* cDNA

A vector, designated pSynExp-2, was used to express *SXD1* in *Synechocystis* sp. PCC6803. pSynExp-2 was derived from pPCR-Script AMP (Stratagene) and contained the *Synechocystis* sp. PCC6803 *psbA2* promoter linked by a multicloning site to Tn9, which encodes chloramphenicol acetyltransferase. To facilitate homologous recombination, the promoter and multicloning site were flanked by 512- and 429-bp sequences from the *slr2699* locus.

Primers 5'-TTTTTTTGTCTAGCACGCCGCATAGCGGGTACCAC-3' (F₇) and 5'-TTTTTTTGTCTAGCTGATCACATTCGTTGGTATCCTATAG-3' (R₆)

and *Pfu* polymerase were used to PCR amplify the coding sequence of the mature maize *SXD1* polypeptide from the *SXD1* cDNA. The PCR product was cloned into the *Nhe*I and *Bcl*I restriction sites of the multicloning site behind the *psbA2* promoter. The resulting plasmid was used to transform *Δslr1737* through homologous recombination of the *slr2699* locus. Recombination events were selected by chloramphenicol resistance, and the introduction of the maize *SXD1* cDNA into *slr1737* knockout mutants was confirmed by PCR analysis of genomic DNA isolated from chloramphenicol- and spectinomycin- (selection for *Δslr1737*) resistant cell lines.

TC activity from *Escherichia coli*-Expressed Proteins

C43 (DE3) cells containing the relevant pET vectors engineered to express TCs from the three organisms were grown at 30°C in 50 mL of Luria-Broth culture (50 μg mL⁻¹ kanamycin) to mid-log phase (0.4–0.6 OD_{600 nm}), then 1 mM isopropylthio-β-galactoside was added and grown at 15°C for 18 h (Miroux and Walker, 1996). Cells were harvested by centrifugation and washed with 100 mM KHPO₄ (pH 7.8) and 4 mM MgCl₂. The cells were resuspended in 1 mL of 100 mM KHPO₄ (pH 7.8), 4 mM MgCl₂, 4 mM dithiothreitol, and 4 mM glutathione, and sonicated six times for a duration of 10 s at 40 Hz. Dodecyl maltoside was added to a final concentration of 0.08 mM, and the cell lysate was shaken gently for 1 h at 4°C. Radiolabeled DMPBQ substrate was prepared by enzymatic labeling of MPBQ with [¹⁴C]-S-adenosyl L-Met (Amersham, Piscataway, NJ) using heterologously expressed *Synechocystis* sp. PCC6803 MPBQ methyl transferase (Shintani et al., 2002). The labeled DMPBQ was purified from the reaction by TLC (Pennock, 1985). Each 100-μL TC assay contained 4.5 × 10⁻⁴ μCi of 2,[[¹⁴C]-dimethyl-6-phytyl-1,4-benzoquinol. The substrate was reduced with 2 mg of NaBH₄ immediately before the assay. Each reaction contained 100 mM KHPO₄ (pH 7.8), 4 mM MgCl₂, 0.8 mM dithiothreitol, 0.8 mM glutathione, 0.8 mM dodecyl maltoside, 20 mM ascorbic acid, 1 mM cyclodextrin, and 20 μL of cell lysate from *E. coli* expressing *SXD1*, *VTE1*, or *slr1737*. The dodecyl maltoside and cyclodextrin were added first to solubilize the substrate. The reactions were sparged with nitrogen gas and gently shaken for 4 h at room temperature. Total lipids were extracted in extraction buffer containing 1 mg mL⁻¹ butylated hydroxytoluene and unlabeled 20 μg mL⁻¹ γ-tocopherol as a carrier. The prenyl lipids were separated by TLC, and the presence of the unlabeled prenyl quinone and tocopherol standards was determined by staining with Emmeric-Engel reagent (Pennock, 1985). [¹⁴C]-labeled compounds were detected by a 4-d exposure to a low-energy PhosphorImage screen (Molecular Dynamics, Piscataway, NJ) and analyzed using a PhosphorImager (Molecular Dynamics).

Analysis of Glc, Suc, and Starch in Leaves

Mature leaves from 4-week-old Arabidopsis plants were harvested at the end and the beginning of the photoperiod. Sugars were extracted from leaf tissue in 80% (v/v) ethanol at 80°C for 30 min. Starch was also extracted from the cleared leaf tissue using 0.2 M KOH at 95°C for 45 min and neutralized with 1 M acetic acid to pH 5.0. The sugars and starch levels were measured enzymatically through the conversion of NAD to NADH by Glc-6-phosphate dehydrogenase and observed at 340 nm as described (Stitt et al., 1989).

ACKNOWLEDGMENTS

We would like to acknowledge Chris Cook and Hiroshi Maeda for technical assistance in the map-based cloning of *vte1*, Eva Collakova for help with the analysis of prenyl quinones, Dr. Zigang Cheng for technical assistance in preparation of MPBQ substrate, and Maria Magallenes-Lundback for assistance with the HPLC analysis. We are also thankful to members of the Dr. DellaPenna laboratory for reviewing this manuscript and for their helpful discussions.

Received March 25, 2003; returned for revision May 2, 2003; accepted May 12, 2003.

LITERATURE CITED

Alfonso M, Perewoska I, Kirilovsky D (2000) Redox control of *psbA* gene expression in the cyanobacterium *Synechocystis* PCC 6803: involvement of the cytochrome b(6)/f complex. *Plant Physiol* 122: 505–515

- Allen JF, Pfannschmidt T (2000) Balancing the two photosystems: photosynthetic electron transfer governs transcription of reaction centre genes in chloroplasts. *Philos Trans R Soc Lond Ser B Biol Sci* **355**: 1351–1357
- Arango Y, Heise KP (1998a) Localization of alpha-tocopherol synthesis in chromoplast envelope membranes of *Capsicum annuum* L. fruits. *J Exp Bot* **49**: 1259–1262
- Arango Y, Heise KP (1998b) Tocopherol synthesis from homogentisate in *Capsicum annuum* L. (yellow pepper) chromoplast membranes: evidence for tocopherol cyclase. *Biochem J* **336**: 531–533
- Azzi A, Ricciarelli R, Zingg JM (2002) Non-antioxidant molecular functions of alpha-tocopherol (vitamin E). *FEBS Lett* **519**: 8–10
- Blee E (2002) Impact of phyto-oxylipins in plant defense. *Trends Plant Sci* **7**: 315–322
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* **37**: 911–917
- Bramley PM, Elmadfa I, Kafatos A, Kelly FJ, Manios Y, Roxborough HE, Schuch W, Sheehy PJA, Wagner KH (2000) Vitamin E. *J Sci Food Agric* **80**: 913–938
- Brigelius-Flohe R, Traber MG (1999) Vitamin E: function and metabolism. *FASEB J* **13**: 1145–1155
- Burton GW, Cheng SC, Webb A, Ingold KU (1986) Vitamin E in young and old human red blood cells. *Biochim Biophys Acta* **860**: 84–90
- Chan SS, Monteiro HP, Schindler F, Stern A, Junqueira VBC (2001) Alpha-tocopherol modulates tyrosine phosphorylation in human neutrophils by inhibition of protein kinase C activity and activation of tyrosine phosphatases. *Free Radic Res* **35**: 843–856
- Chandra V, Jasti J, Kaur P, Betzel C, Srinivasan A, Singh TP (2002) First structural evidence of a specific inhibition of phospholipase A(2) by alpha-tocopherol (vitamin E) and its implications in inflammation: crystal structure of the complex formed between phospholipase A(2) and alpha-tocopherol at 1.8 angstrom resolution. *J Mol Biol* **320**: 215–222
- Clement SA, Tan CC, Guo JL, Kitta K, Suzuki YJ (2002) Roles of protein kinase C and alpha-tocopherol in regulation of signal transduction for GATA-4 phosphorylation in HL-1 cardiac muscle cells. *Free Radic Biol Med* **32**: 341–349
- Collakova E, DellaPenna D (2001) Isolation and functional analysis of homogentisate phytyltransferase from *Synechocystis* sp. PCC 6803 and *Arabidopsis*. *Plant Physiol* **127**: 1113–1124
- Dahnhardt D, Falk J, Appel J, van der Kooij TAW, Schulz-Friedrich R, Krupinska K (2002) The hydroxyphenylpyruvate dioxygenase from *Synechocystis* sp. PCC 6803 is not required for plastoquinone biosynthesis. *FEBS Lett* **523**: 177–181
- Eisenreich W, Schwarz M, Cartayrade A, Arigoni D, Zenk MH, Bacher A (1998) The deoxyxylulose phosphate pathway of terpenoid biosynthesis in plants and microorganisms. *Chem Biol* **5**: R221–R233
- Evert RF, Eschrich W, Heyser W (1977a) Distribution and structure of plasmodesmata in mesophyll and bundle sheath cells of *Zea mays* L. *Planta* **136**: 77–89
- Evert RF, Eschrich W, Heyser W (1978) Leaf structure in relation to solute transport and phloem loading in *Zea mays* L. *Planta* **138**: 279–294
- Evert RF, Eschrich W, Neuberger DS, Eichhorn SE (1977b) Tubular extensions of plasmalemma in leaf cells of *Zea mays* L. *Planta* **135**: 203–205
- Fahn A (1990) *Plant Anatomy*. Butterworth-Heinemann Ltd., Oxford
- Fryer MJ (1992) The antioxidant effects of thylakoid vitamin E (alpha-tocopherol). *Plant Cell Environ* **15**: 381–392
- Garcia I, Rodgers M, Lenne C, Rolland A, Sailland A, Matringe M (1997) Subcellular localization and purification of a *p*-hydroxyphenylpyruvate dioxygenase from cultured carrot cells and characterization of the corresponding cDNA. *Biochem J* **325**: 761–769
- Garcia I, Rodgers M, Pepin R, Hsieh TF, Matringe M (1999) Characterization and subcellular compartmentation of recombinant 4-hydroxyphenylpyruvate dioxygenase from *Arabidopsis* in transgenic tobacco. *Plant Physiol* **119**: 1507–1516
- Goksoyr J (1967) Evolution of eukaryotic cells. *Nature* **214**: 1161
- Gottwald JR, Krysan PJ, Young JC, Evert RF, Sussman MR (2000) Genetic evidence for the in planta role of phloem-specific plasma membrane sucrose transporters. *Proc Natl Acad Sci USA* **97**: 13979–13984
- Greenberglevy SH, Budowski P, Grossman S (1993) Lipoygenase and other enzymes of arachidonic acid metabolism in the brain of chicks affected by nutritional encephalomalacia. *Int J Biochem* **25**: 403–409
- Haritatos E, Medville R, Turgeon R (2000) Minor vein structure and sugar transport in *Arabidopsis thaliana*. *Planta* **211**: 105–111
- Howe GA, Schillmiller AL (2002) Oxylin metabolism in response to stress. *Curr Opin Plant Biol* **5**: 230–236
- Hutson KG, Threlfall DR (1980) Synthesis of plastoquinone-9 and phytyl-plastoquinone from homogentisate in lettuce chloroplasts. *Biochim Biophys Acta* **632**: 630–648
- Jander G, Norris SR, Rounsley SD, Bush DF, Levin IM, Last RL (2002) *Arabidopsis* map-based cloning in the post-genome era. *Plant Physiol* **129**: 440–450
- Jiang Q, Elson-Schwab I, Courtemanche C, Ames BN (2000) gamma-tocopherol and its major metabolite, in contrast to alpha-tocopherol, inhibit cyclooxygenase activity in macrophages and epithelial cells. *Proc Natl Acad Sci USA* **97**: 11494–11499
- Johnson TW, Shen G, Zybailov B, Kolling D, Reategui R, Beuparlant S, Vassiliev IR, Bryant DA, Jones AD, Golbeck JH et al. (2000) Recruitment of a foreign quinone into the A(1) site of photosystem I: I. Genetic and physiological characterization of phylloquinone biosynthetic pathway mutants in *Synechocystis* sp. pcc 6803. *J Biol Chem* **275**: 8523–8530
- KamalEldin A, Appelqvist LA (1996) The chemistry and antioxidant properties of tocopherols and tocotrienols. *Lipids* **31**: 671–701
- Karol KG, McCourt RM, Cimino MT, Delwiche CF (2001) The closest living relatives of land plants. *Science* **294**: 2351–2353
- Kim KR, Kim MJ, Rhee SJ (2001) Effects of vitamin E on arachidonic acid cascade in platelets and aorta of acute cadmium-poisoned rats. *Nutr Res* **21**: 657–665
- Kruk J, Schmid GH, Strzalka K (1994) Antioxidant properties of plastoquinol and other biological prenylquinols in liposomes and solution. *Free Radic Res* **21**: 409–416
- Kruk J, Strzalka K (1995) Occurrence and function of alpha-tocopherol quinone in plants. *J Plant Physiol* **145**: 405–409
- Kujat SL, Owtrim GW (2000) Redox-regulated RNA helicase expression. *Plant Physiol* **124**: 703–713
- Li H, Sherman LA (2000) A redox-responsive regulator of photosynthesis gene expression in the cyanobacterium *Synechocystis* sp. strain PCC 6803. *J Bacteriol* **182**: 4268–4277
- Lichtenthaler HK (1998) The plants' 1-deoxy-D-xylulose-5-phosphate pathway for biosynthesis of isoprenoids. *Fett Lipid* **100**: 128–138
- Lichtenthaler HK, Prenzel U, Douce R, Joyard J (1981) Localization of prenylquinones in the envelope of spinach chloroplasts. *Biochim Biophys Acta* **641**: 99–105
- Liebler DC (1993) The role of metabolism in the antioxidant function of vitamin E. *Crit Rev Toxicol* **23**: 147–169
- Liebler DC, Burr JA (2000) Antioxidant reactions of alpha-tocopherolhydroquinone. *Lipids* **35**: 1045–1047
- Marshall PS, Morris SR, Threlfall DR (1985) Biosynthesis of tocopherols: a re-examination of the biosynthesis and metabolism of 2-methyl-6-phytyl-1,4-benzoquinol. *Phytochemistry* **24**: 1705–1711
- Miroux B, Walker JE (1996) Over-production of proteins in *Escherichia coli*: mutant hosts that allow synthesis of some membrane proteins and globular proteins at high levels. *J Mol Biol* **260**: 289–298
- Nobata Y, Urakaze M, Temaru R, Sato A, Nakamura N, Yamazaki K, Kishida M, Takata M, Kobayashi M (2002) alpha-tocopherol inhibits IL-8 synthesis induced by thrombin and high glucose in endothelial cells. *Hormone Metab Res* **34**: 49–54
- Norris SR, Shen X, DellaPenna D (1998) Complementation of the *Arabidopsis* pds1 mutation with the gene encoding *p*-hydroxyphenylpyruvate dioxygenase. *Plant Physiol* **117**: 1317–1323
- Ohuchi K, Levine L (1980) Alpha-tocopherol inhibits 12-*o*-tetradecanoylphorbol-13-acetate-stimulated deacylation of cellular lipids, prostaglandin production, and changes in cell morphology of Madin-Darby canine kidney Cells. *Biochim Biophys Acta* **619**: 11–19
- Pennock JF (1985) Biosynthesis of plastoquinone. *Methods Enzymol* **110**: 313–319
- Pfannschmidt T, Nilsson A, Allen JF (1999a) Photosynthetic control of chloroplast gene expression. *Nature* **397**: 625–628
- Pfannschmidt T, Nilsson A, Tullberg A, Link G, Allen JF (1999b) Direct transcriptional control of the chloroplast genes psbA and psaAB adjusts photosynthesis to light energy distribution in plants. *Plant Life* **48**: 271–276
- Pfannschmidt T, Schutze K, Brost M, Oelmüller R (2001) A novel mechanism of nuclear photosynthesis gene regulation by redox signals from the chloroplast during photosystem stoichiometry adjustment. *J Biol Chem* **276**: 36125–36130

- Porfirova S, Bergmuller E, Tropf S, Lemke R, Dormann P** (2002) Isolation of an Arabidopsis mutant lacking vitamin E and identification of a cyclase essential for all tocopherol biosynthesis. *Proc Natl Acad Sci USA* **99**: 12495–12500
- Provencher LM, Miao L, Sinha N, Lucas WJ** (2001) Sucrose export defective1 encodes a novel protein implicated in chloroplast-to-nucleus signaling. *Plant Cell* **13**: 1127–1141
- Pursiheimo S, Mulo P, Rintamaki E, Aro EM** (2001) Coregulation of light-harvesting complex II phosphorylation and lhcb mRNA accumulation in winter rye. *Plant J* **26**: 317–327
- Ricciarelli R, Zingg JM, Azzi A** (2001) Vitamin E: protective role of a Janus molecule. *FASEB J* **15**: 2314–2325
- Ricciarelli R, Zingg JM, Azzi A** (2002) The 80th anniversary of vitamin E: beyond its antioxidant properties. *Biol Chem* **383**: 457–465
- Russin WA, Evert RF, Vanderveer PJ, Sharkey TD, Briggs SP** (1996) Modification of a specific class of plasmodesmata and loss of sucrose export ability in the sucrose export defective1 maize mutant. *Plant Cell* **8**: 645–658
- Savidge B, Weiss JD, Wong YH, Lassner MW, Mitsky TA, Shewmaker CK, Post-Beittenmiller D, Valentin HE** (2002) Isolation and characterization of homogentisate phytyltransferase genes from *Synechocystis* sp. PCC 6803 and Arabidopsis. *Plant Physiol* **129**: 321–332
- Schledz M, Seidler A, Beyer P, Neuhaus G** (2001) A novel phytyltransferase from *Synechocystis* sp. PCC 6803 involved in tocopherol biosynthesis. *FEBS Lett* **499**: 15–20
- Sen CK, Khanna S, Roy S, Packer L** (2000) Molecular basis of vitamin E action: tocotrienol potently inhibits glutamate-induced pp60(c-Src) kinase activation and death of HT4 neuronal cells. *J Biol Chem* **275**: 13049–13055
- Shintani D, DellaPenna D** (1998) Elevating the vitamin E content of plants through metabolic engineering. *Science* **282**: 2098–2100
- Shintani DK, Cheng Z, DellaPenna D** (2002) The role of 2-methyl-6-phytylbenzoquinone methyltransferase in determining tocopherol composition in *Synechocystis* sp. PCC6803. *FEBS Lett* **511**: 1–5
- Soll J** (1979) Comparison of geranylgeranyl and phytol substituted methylquinols in the tocopherol synthesis of spinach chloroplasts. *Biochem Biophys Res Commun* **91**: 715–720
- Soll J, Kemmerling M, Schultz G** (1980) Tocopherol and plastoquinone synthesis in spinach chloroplast subfractions. *Arch Biochem Biophys* **204**: 544–550
- Soll J, Schultz G** (1980) 2-Methyl-6-phytylquinol and 2,3-dimethyl-5-phytylquinol as precursors of tocopherol synthesis in spinach chloroplasts. *Phytochemistry* **19**: 215–218
- Soll J, Schultz G, Joyard J, Douce R, Block MA** (1985) Localization and synthesis of prenylquinones in isolated outer and inner envelope membranes from spinach chloroplasts. *Arch Biochem Biophys* **238**: 290–299
- Stitt M, Lilley RM, Gerhardt R, Heldt HW** (1989) Metabolite levels in specific cells and subcellular compartments of plant leaves. *Methods Enzymol* **174**: 518–552
- Stocker A, Fretz H, Frick H, Ruttimann A, Woggon WD** (1996) The substrate specificity of tocopherol cyclase. *Bioorg Med Chem* **4**: 1129–1134
- Stocker A, Netscher T, Ruttimann A, Muller RK, Schneider H, Todaro LJ, Derungs G, Woggon WD** (1994) The reaction mechanism of chromanol ring formation catalyzed by tocopherol cyclase from *Anabaena variabilis* Kutzling (cyanobacteria). *Helv Chim Acta* **77**: 1721–1737
- Stocker A, Ruttimann A, Woggon WD** (1993) Identification of the tocopherol cyclase in the blue-green algae *Anabaena variabilis* Kutzling (cyanobacteria). *Helv Chim Acta* **76**: 1729–1738
- Tran K, Wong JT, Lee E, Chan AC, Choy PC** (1996) Vitamin E potentiates arachidonate release and phospholipase A(2) activity in rat heart myoblastic cells. *Biochem J* **319**: 385–391
- Trebitsh T, Danon A** (2001) Translation of chloroplast psbA mRNA is regulated by signals initiated by both photosystems II and I. *Proc Natl Acad Sci USA* **98**: 12289–12294
- Wang CX, Zien CA, Afithile M, Welti R, Hildebrand DF, Wang XM** (2000) Involvement of phospholipase D in wound-induced accumulation of jasmonic acid in Arabidopsis. *Plant Cell* **12**: 2237–2246
- Williams JGK** (1988) Construction of specific mutations in photosystem II photosynthetic reaction center by genetic engineering methods in *Synechocystis* 6803. *Methods Enzymol* **167**: 766–778
- Wu DY, Hayek MG, Meydani SN** (2001) Vitamin E and macrophage cyclooxygenase regulation in the aged. *J Nutr* **131**: 382S–388S
- Yamauchi J, Iwamoto T, Kida S, Masushige S, Yamada K, Esashi T** (2001) Tocopherol-associated protein is a ligand-dependent transcriptional activator. *Biochem Biophys Res Commun* **285**: 295–299
- Yang DH, Andersson B, Aro EM, Ohad I** (2001) The redox state of the plastoquinone pool controls the level of the light-harvesting chlorophyll a/b binding protein complex II (LHC II) during photoacclimation: cytochrome b(6)f deficient *Lemna perpusilla* plants are locked in a state of high-light acclimation. *Photosynth Res* **68**: 163–174