

## ActA Is Required for Crossing of the Fetoplacental Barrier by *Listeria monocytogenes*<sup>∇</sup>

Alban Le Monnier,<sup>1,2,3</sup> Nicolas Autret,<sup>1</sup> Olivier F. Join-Lambert,<sup>1,2,3</sup> Francis Jaubert,<sup>2,3,4</sup>  
Alain Charbit,<sup>1</sup> Patrick Berche,<sup>1,2,3</sup> and Samer Kayal<sup>1,2,3\*</sup>

U-570 INSERM,<sup>1</sup> University and Medical School Paris-Descartes,<sup>2</sup> and Assistance Publique—Hôpitaux de Paris,<sup>3</sup> 75730 Paris Cedex 15, France, and Département d'Anatomie Pathologique, Hôpital Necker-Enfants Malades, 75015 Paris, France<sup>4</sup>

Received 28 September 2006/Returned for modification 17 October 2006/Accepted 13 November 2006

**The facultative intracellular bacterial pathogen *Listeria monocytogenes* induces severe fetal infection during pregnancy. Little is known about the molecular mechanisms allowing the maternofetal transmission of bacteria. In this work, we studied fetoplacental invasion by infecting mice with various mutants lacking virulence factors involved in the intracellular life cycle of *L. monocytogenes*. We found that the placenta was highly susceptible to bacteria, including avirulent bacteria, such as an *L. monocytogenes* mutant with an *hly* deletion ( $\Delta$ LLO) and a nonpathogenic species, *Listeria innocua*, suggesting that permissive trophoblastic cells, trapping bacteria, provide a protective niche for bacterial survival. The  $\Delta$ LLO mutant, which is unable to escape the phagosomal compartment of infected cells, failed to grow in the trophoblast tissue and to invade the fetus. Mutant bacteria with *inlA* and *inlB* deletion ( $\Delta$ InlAB) grew in the placenta and fetus as well as did the wild-type virulent strain (EGDwt), indicating that in the murine model, internalins A and B are not involved in fetoplacental invasion by *L. monocytogenes*. Pregnant mice were then infected with an *actA* deletion ( $\Delta$ ActA) strain, a virulence-attenuated mutant that is unable to polymerize actin and to spread from cell to cell. With the  $\Delta$ ActA mutant, fetal infection occurs, but with a significant delay and restriction, and it requires a placental bacterial load 2 log units higher than that for the wild-type virulent strain. Definitive evidence for the role of ActA was provided by showing that a *actA*-complemented  $\Delta$ ActA mutant was restored in its capacity to invade fetuses. ActA-mediated cell-to-cell spreading plays a major role in the vertical transmission of *L. monocytogenes* to the fetus in the murine model.**

*Listeria monocytogenes* is a facultative intracellular fast-growing gram-positive bacterium widely spread in the environment. It is a food-borne pathogen responsible for severe and life-threatening infections in both humans and a large variety of animal species (13). Immunocompromised patients, including the elderly and pregnant women, represent high-risk groups for listeriosis (9). During pregnancy, listeriosis can be asymptomatic or can give rise to subclinical symptoms like a nonspecific fever despite the insidious development of fetoplacental infection resulting in abortion, stillbirth, or severe and disseminated neonatal infections markedly described as granulomatosis infantiseptica (9, 24). Little is known about molecular mechanisms implicated in the placental infection by *L. monocytogenes* and the subsequent vertical transmission to the fetus.

Most virulence factors involved in the intracellular growth and survival of *L. monocytogenes* have been identified and extensively studied (8, 10, 30). Adhesion and invasion of non-professional phagocytes are mainly dependent upon the expression of internalin A (InlA), which interacts with E-cadherin expressed on eukaryotic cells. After phagocytosis, bacteria produce listeriolysin O (LLO), a pore-forming cytolysin (1), allowing bacterial escape from the phagosomal compartment. Once in the cytoplasm, bacteria divide, move, and spread from cell to cell. This is due to ActA, a bacterial sur-

face-exposed protein, which induces actin cytoskeleton rearrangements and polymerization. Thus, bacteria protrude into and infect neighboring cells favoring the persistence of the intracellular life cycle of *L. monocytogenes* (28).

The placenta is a dynamic organ constituted of intricate maternal and fetal tissues, whose structure and function change throughout the pregnancy. The physiological barrier separating fetal and maternal blood in the placenta is mainly formed by fetally derived trophoblastic cells. Very few pathogens are capable of crossing the placental biological barrier. This includes some viruses (17); parasites such as *Toxoplasma gondii* (26) and *Plasmodium falciparum* (27); and very rare bacteria, including *Chlamydia psittaci*, *Coxiella burnetii*, and *L. monocytogenes* (6, 23, 25).

There is compelling evidence that the trophoblast plays a central role in vertical transmission of pathogens from mothers to the fetus (21). First of all, the trophoblast acts as a pregnancy-specific component of the innate immune system (14). During pregnancy, the trophoblast is responsive to CSF-1 which acts to organize the maternal immune response to bacterial infection at the uteroplacental interface through recruitment of polymorphonuclear neutrophils. These inflammatory cells are the main effector cells mobilized in the placenta to destroy *L. monocytogenes*, as opposed to macrophages that are mostly excluded from the murine placenta (14, 21).

Little is known about the pathophysiological process of placental invasion during listeriosis. Using a murine model, we recently demonstrated that placental invasion by *L. monocytogenes* is associated with bacterial growth within trophoblastic cells, thus allowing outward spreading from the initial foci to

\* Corresponding author. Mailing address: Department of Microbiology-INSERM, 156 rue de Vaugirard, 75730 Paris Cedex 15, France. Phone: 33 1 44 49 49 61. Fax: 33 1 44 49 49 60. E-mail: kayal@necker.fr.

<sup>∇</sup> Published ahead of print on 21 November 2006.

the adjacent structures (21). In human fetoplacental listeriosis, the involvement of InlA through its interaction with E-cadherin has been recently reported, using an in vitro model (19). Mouse E-cadherin compared with human E-cadherin has a single-amino-acid mutation which results in a decrease of its affinity for InlA (18, 20). However, although the InlA–E-cadherin interaction occurs with the same affinity in guinea pigs as in humans, it has been very recently published that an InlA mutant behaved as wild-type virulent strain in a pregnant guinea pig model (4). These authors provided evidence for a role of ActA in the vertical transmission of *L. monocytogenes* in this model (4).

In this work, using a murine model of pregnant mice (21). We studied the role of virulence factors (InlA, InlB, LLO, and ActA) involved in the intracellular life cycle of *L. monocytogenes*. To decipher the role of virulence factors for the crossing of the fetoplacental barrier, we systematically monitored for each mutant of *L. monocytogenes* the correlation between the infection of the placenta and its corresponding fetus. We show that ActA-dependent cell-to-cell spreading promotes fetal invasion. Final evidence for the crucial role of ActA was obtained by restoring fetal invasion in an *actA*-complemented ActA mutant.

#### MATERIALS AND METHODS

**Bacterial strains and cultures.** We used the wild-type virulent strain of *L. monocytogenes* EGD<sub>e</sub> (EGDwt) (12) various isogenic mutants (Table 1) and *L. innocua*. Bacteria were grown overnight in brain heart infusion broth (Difco Laboratories, Detroit, MI) at 37°C without antibiotics, reexpanded the next day, and collected at the end of the exponential phase and then were centrifuged at 5,000 × g for 30 min at 4°C, washed twice with lipopolysaccharide-free Hanks' balanced salt solution (Gibco, Long Island, NY), and resuspended in RPMI 1640 medium (Difco) before being stored at –80°C in 1-ml aliquots. Bacteria were titrated by serial dilution and plated on brain heart infusion agar. Before each experiment, an aliquot was thawed and diluted as convenient for the experiment.

**DNA techniques.** Obtaining of chromosomal DNA, plasmid isolation, restriction enzyme analyses, and PCR amplifications were performed as previously described by Autret et al. (3). Oligonucleotides were synthesized by Eurogentec (Paris, France). We used the AmpliTaq Gold DNA polymerase of *Thermus aquaticus* from Roche (Branchburg, NJ) and the pAT113/pAT145 system (29) as previously described (3).

**Cloning of *actA* and complementation of the ΔActA mutant strain.** The entire *actA* sequence including its promoter was amplified by PCR from EGDwt chromosomal DNA using primers actA-prom (5'-TGAAGCTTGGGAAGCAGTTGGGGT-3'), which contains a HindIII site (underlined), and actA-term (5'-TTG AATTCTGAATTCATATCATTACCTCACT-3'), which contains an EcoRI site (underlined). The fragment was subcloned into the pCR II plasmid (Invitrogen, Carlsbad, CA), and transferred to *Escherichia coli* TOP10 (Invitrogen, Cergy Pontoise, France). Recombinant bacteria were selected onto ampicillin-containing agar, and clones were checked by PCR. The plasmid of one clone was prepared and submitted to restriction by HindIII and EcoRI (New England Biolabs). The HindIII/EcoRI *actA* fragment was cloned into the HindIII/EcoRI-digested pAT113 plasmid, leading to the pAT113-*actA* plasmid.

Chromosomal integration of the pAT113 Tn1545 transposon requires the presence of its integrase, provided in *trans* by pAT145 plasmid. Competent EGD-ΔActA bacteria were thus prepared as described previously (3), and 2 μg of QIAGEN-purified plasmid pAT145 was used for electroporation. The resulting Kan<sup>r</sup> transformants were named ΔActA-145. Then *L. monocytogenes* ΔActA-145 cells were transformed by electroporation with 2 μg of purified pAT113-*actA* plasmid. Plasmid integration into chromosomal DNA was checked by PCR. The DNA sequences flanking the transposon carrying the wild-type *actA* allele were determined using ligation-mediated PCR, as described previously (3), and the site of transposon insertion into the genome was identified by sequence analysis. The transposon was inserted at position 75350 in open reading frame lmo0068, encoding a putative 107-amino-acid protein of unknown function. Protein secretion was checked by Western blot analysis.

TABLE 1. Bacterial strains used in this work

Bacterial strain	Characteristic(s)	Source or reference
<i>Listeria monocytogenes</i>		
Wild type (EGDwt)	Virulent EGD <sub>e</sub> strain	12
EGD mutants		
ΔInlAB	InlAB deleted	P. Cossart (Institut Pasteur, Paris)
ΔLLO	1,080-bp deletion in <i>hly</i> gene	15
ΔActA	1,752-bp deletion in <i>actA</i> gene	7
ΔActA+ <i>actA</i>	Insertion of <i>actA</i> gene in ΔActA mutant	This work
<i>Listeria innocua</i>		
<i>L. innocua</i> CIP 11254	Wild type	Institut Pasteur Collection

**Infection of mice.** Inbred BALB/c pregnant mice purchased from Elevage Janvier (Le Genest-St-Isle, France) were used for bacterial growth studies and histological staining. Couplings were carried out with 8- to 10-week-old BALB/c female mice. Mating was assessed by the appearance of a vaginal plug, denoting the first embryonic day of pregnancy. The gestation was checked at the 12th day and nonpregnant females were used as control mice. Mice were housed in wire-bottom cages, with free access to food and water, and held under these conditions for at least 24 h before infection. Animal experiments were approved by the Animal Welfare Committee of the University Paris-Descartes.

BALB/c female mice were inoculated intravenously (i.v.) at the 14th day of gestation via the lateral tail vein with 0.5 ml of a calibrated suspension of bacteria, extemporarily obtained by appropriate dilution into saline isotonic solution from a frozen stock. All mice were daily examined. At intervals after infection (1, 6, 24, 48, and 72 h), groups of mice were anesthetized by intramuscular injection of 200 μl of a mixture of ketamine at 200 mg kg<sup>-1</sup> (Imalgène 100; Merial, Lyon-France) and xylazine hydrochloride at 10 mg kg<sup>-1</sup> (Rampun, Bayer, Puteaux, France) and were sacrificed. The abdominal cavity was then aseptically opened, and each mouse was bled by intracardiac puncture with heparinized syringe (Heparin sodic; Sanofi-Wintrop, France). Organs (livers, spleens, and brains) and each fetoplacental unit were aseptically removed and homogenized for bacterial counts and histological studies. Each placenta and its respective fetus were independently dissected and analyzed. Bacterial counts were determined by plating serial 10-fold dilutions of each organ homogenate on BHI agar plates incubated at 37°C during 24 to 48 h. For each mouse, 100 μl of each placenta or fetus homogenate was separately pooled to determine the mean bacterial load. The results were expressed as a mean ± standard error expressed as log<sub>10</sub> CFU per organ (bacteria/organ). The 50% lethal doses (LD<sub>50</sub>) estimated by the probit method on groups of five mice were assessed in this work at 10<sup>4.3</sup> and 10<sup>4.6</sup> bacteria per mouse for EGDwt and *actA*-complemented ΔActA strains, respectively.

**Histology.** Histological studies were performed on placentas removed from mice at day 17 of gestation, 72 h after i.v. infection with 2 × 10<sup>5</sup> bacteria. Fetoplacental units were removed from the uterus horns by dissociation between implantation sites. One half of each placenta was used to quantify bacterial load, and the second half was used for histological analysis. For light microscopy studies (Nikon Eclipse E600; digital camera DXM1200), placentas were fixed overnight in 10% formalin, dehydrated with an alcohol gradient, and embedded in paraffin blocks. Sequential 5- to 7-μm placental sections were stained by Gram-Weigert techniques.

**Statistical analysis.** All values are given as the mean ± standard error of the mean. We used several statistical methods to determine the relationship between placental infection and the corresponding fetal infection. To compare the mean values at the indicated times between mutants, we used multifactorial analysis of variance in which the two analyzed factors were "time" (6 h and 1, 2, and 3 days postinfection) and "mutants" (EGD, ΔInlAB, ΔActA, and ΔActA+*actA*). The differences were considered significant for *P* ≤ 0.05. Placentas and fetuses were considered infected when bacterial counts were superior or equal to 1 bacterium/placenta or 50 bacteria/fetus. Receiver-operator characteristics analysis was used

to identify the cutoff values for placental bacterial load associated with the maximal probability of fetal infection, according to a method previously described (21).

## RESULTS

**Crossing of fetoplacental barrier requires the expression of virulence factors.** We recently reported that *L. monocytogenes* inoculated i.v. into pregnant mice can invade the placenta and cross the placental barrier to subsequently proliferate in the fetus (21). With wild-type virulent bacteria, fetal infection occurred when the placenta was infected with doses as low as  $1 \times 10^3$  bacteria. To study the role of virulence factors in the vertical transmission of *L. monocytogenes* from mother to fetus, we first i.v. infected pregnant mice (14 days) with a high dose ( $5 \times 10^7$  bacteria) of *L. innocua*, a nonvirulent, nonhemolytic species that does not display the *prfA*-dependent virulent genes. Bacterial growth was monitored in the blood, spleen, placenta, and fetuses from 1 h to 48 h after infection (Fig. 1). As expected, bacteria were rapidly eliminated from the blood and the organs (data in the liver are not shown). In contrast, we found that all placentas were infected by *L. innocua* as early as 1 h after inoculation. Bacteria survived in placental tissues for at least 2 days, at a low titer ( $\sim 1 \times 10^3$  bacteria), without any infection of the respective fetus. These data suggest that although the placentas become infected with *L. innocua*, the crossing of the fetoplacental barrier requires bacterial growth and the expression of virulence factors.

**LLO is required for bacterial growth in the placenta and fetal invasion.** We recently showed in pregnant mice that the virulent strain of *L. monocytogenes* (EGDwt) first targets the trophoblastic cells before crossing the fetoplacental barrier (21). Thus, fetal infection requires bacterial growth which is clearly associated with the intracellular life cycle of *L. monocytogenes* within trophoblast cells acting as phagocytes. To decipher the role of virulence factors in the placental invasion and the subsequent fetal infection, we studied *L. monocytogenes* mutants inactivated for the expression of various virulent factors.

We first tested the involvement of internalins A and B in placental infection. Pregnant mice were infected i.v. with  $5 \times 10^5$  cells of the  $\Delta$ InlAB mutant. The kinetics of bacterial growth was then observed for 3 days in the blood and organs (liver, spleen, and brain) and in all fetoplacental units. As compared to the EGDwt strain used as control, bacterial growth of the  $\Delta$ InlAB mutant was similar in blood and organs except for the liver, in which the growth of the  $\Delta$ InlAB mutant was moderately lower (Fig. 2A). Surprisingly, this held true for bacterial growth of  $\Delta$ InlAB bacteria in placenta and fetus (Fig. 2B). As wild-type bacteria, the  $\Delta$ InlAB mutant rapidly multiplied in these tissues, with a daily increase of about 2 log units. By day 3, all placentas were infected and the comparison between the EGDwt and  $\Delta$ InlAB mutant strains was not statistically different. We then determined statistically which bacterial load in the placenta was associated with a higher probability of fetal infection. Thus, the cutoff values were assessed at  $1.7 \times 10^3$  and  $1.0 \times 10^3$  CFU/placenta for the EGDwt and  $\Delta$ InlAB strains, respectively. The comparison of the means for fetal infection between wild-type EGDwt and  $\Delta$ InlAB mutant was not statistically different. These results suggest that in




FIG. 1. Infection of pregnant mice with *L. innocua*. Pregnant BALB/c mice were injected i.v. with  $5 \times 10^7$  bacteria by day 14 of gestation. Animals were monitored by quantifying bacterial growth at intervals. (A) Bacterial growth in spleen and blood of pregnant mice. The results shown are means  $\pm$  standard errors from groups of five mice (each experiment was repeated twice) and are expressed as the  $\log_{10}$  bacteria (CFU) per organ or  $\log_{10}$  bacteria per ml of blood. (B) The kinetics of bacterial growth are represented in the left panels by dot plots corresponding to individual bacterial counts for each placenta (top panel) and each fetus (lower panel) for the indicated times. In the right panels, means  $\pm$  standard errors for each time are given.

the murine model, InlA and InlB are not required for the invasion of placenta and the subsequent fetal infection.

We then tested the role of LLO, the major virulence factor of *L. monocytogenes* needed to escape phagosomes. Pregnant mice were i.v. infected with  $5 \times 10^5$  bacteria of a  $\Delta$ LLO mutant. As previously described, bacteria were rapidly eliminated from the blood and organs (spleen and liver), without brain infection (data not shown) (11). By day 1 and day 3 postinfection, about 10% of placentas (3/30 at day 1 and 4/36 at day 3) were infected at low levels (10 to 100 bacteria/placenta). Under these conditions, fetuses were never infected (Fig. 2). Thus, nonvirulent  $\Delta$ LLO bacteria can infect some placentas for at least 3 days, revealing that once infected, the placenta cannot easily eliminate bacteria, in contrast to the spleen and the liver. As for *L. innocua*,  $\Delta$ LLO bacteria were unable to grow in the placenta and to subsequently invade fetuses, because they are probably retained within the phagosomes of trophoblastic cells.

**ActA promotes the crossing of the fetoplacental barrier.** The murine fetoplacental barrier consists of two layers of trophoblastic cells and one of endothelial cells. We then studied the role of ActA, a virulence factor promoting cell-to-cell spreading. We first analyzed the infectious process in pregnant mice i.v. inoculated with  $5 \times 10^5$  bacteria of a  $\Delta$ ActA mutant. After infection, bacterial growth was monitored in the blood and organs (spleen, liver, and brain) for 3 days. No difference was observed for the bacterial growth in organs between pregnant and nonpregnant mice (data not shown). The  $\Delta$ ActA bacteria were completely eliminated from the blood within 6 h postinfection, which was correlated with the absence of brain infection, as previously de-




FIG. 2. Infection of pregnant mice with  $\Delta$ InlAB and  $\Delta$ LLO mutants. Pregnant BALB/c mice were injected i.v. at day 14 of gestation with  $5 \times 10^5$  of wild-type *L. monocytogenes* (EGD) and its isogenic *hly* or *inlAB* deletion mutants. Animals were monitored by quantifying bacterial growth at intervals. (A) Bacterial growth in organs (spleen, liver and brain) and in blood of pregnant mice for EGDwt and  $\Delta$ InlAB strains. Results shown are means  $\pm$  standard errors of the means from groups of three to five mice and are expressed as the  $\log_{10}$  bacteria (CFU) per organ or  $\log_{10}$  bacteria per ml of blood. (B) The kinetics of bacterial growth are represented on the left panels by dot plots corresponding to individual bacterial counts for each placenta (top panel) and each fetus (lower panel) for the indicated times and strains. In the right panels, means  $\pm$  standard errors for each time are given for wild-type,  $\Delta$ InlAB, and  $\Delta$ LLO strains. The numbers reported near each value correspond to the percentage of infected placenta among all analyzed placentas. When observed, fetal losses are symbolized by a cross. *P* values reported in each graph correspond to the comparison between EGDwt and the indicated mutant. NS, not statistically different.

scribed. After initial growth in the liver and the spleen by day 1 of infection, mutant bacteria then declined rapidly, indicating that the infectious process was well controlled in these organs. With this inoculum ( $5 \times 10^5$  bacteria), about 35% of placentas were infected by  $\Delta$ ActA bacteria as early as 6 h postinfection, compared to 54% of placentas being infected with EGDwt. The percentage of placentas infected by EGDwt reached 100% by day 2, whereas mutant bacteria infected only 50% and 90% of placentas by days 2 and 3 postinfection, respectively. Although the means for placental infection between EGDwt and the  $\Delta$ ActA mutant were statistically different, the rate of mutant growth in the placenta was similar to that of EGDwt, which might reflect rapid multiplication inside permissive trophoblastic cells (see below). As illustrated in Fig. 3, fetal infection was significantly delayed for 2 to 3 days in mice infected by  $\Delta$ ActA bacteria, as compared to EGDwt bacteria. The rate of mutant growth was lower in fetuses during the 3 first days postinfection (Fig. 3B). This was also quantified by calculating the cutoff values corresponding to the placental bacterial load associated with the higher probability to induce fetal infection. The cutoff value was estimated at  $1.2 \times 10^5$   $\Delta$ ActA bacteria per placenta, compared to  $1.7 \times 10^3$  for EGDwt bacteria (Table 2).

Final evidence for the role of the virulence factor ActA in the

crossing of fetoplacental barrier was obtained by inserting a new *actA* gene in the chromosome of the  $\Delta$ ActA mutant strain (see Materials and Methods). After *actA* insertion, the expression of this surface-exposed protein was restored. As illustrated in Fig. 3, the virulence of the *actA*-complemented  $\Delta$ ActA strain was almost completely restored. Complemented bacteria produced bacteremia and grew rapidly in organs (liver, spleen, and brain), as well as did EGDwt bacteria. This was also observed in placentas and fetuses, where complemented bacteria behaved similarly to EGDwt bacteria (Fig. 3B), with cutoff values in the placenta estimated at  $1.7 \times 10^3$  bacteria for EGDwt and  $2.0 \times 10^3$  for the *actA*-complemented  $\Delta$ ActA strain (Table 2).

**Bacterial cell-to-cell spreading is a key step for crossing the fetoplacental barrier.** Histologic examination of the placental labyrinthine zone was performed 48 h after infection with  $5 \times 10^4$  bacteria of the EGDwt,  $\Delta$ InlAB, or  $\Delta$ ActA strains (Fig. 4). After infection with EGDwt and  $\Delta$ InlAB, the lesions observed in placental villousities were similar, consisting of a centrifugal dissemination all over the syncytiotrophoblastic cells following the villous axis. In contrast,  $\Delta$ ActA bacteria induced infectious foci in the placenta, where bacteria mostly visible inside trophoblastic cells were unable to spread in an outward manner, as did EGDwt bacteria. These results clearly demonstrate that spreading within the laby-




FIG. 3. Role of ActA in the crossing of the fetoplacental barrier. Pregnant BALB/c mice were injected intravenously at day 14 of gestation with  $5 \times 10^5$  *L. monocytogenes actA* deletion mutant cells ( $\Delta$ ActA) or the  $\Delta$ ActA mutant complemented with *actA* ( $\Delta$ ActA+*actA*). As compared to the isogenic mutants, bacterial growth for the EGDwt strain is indicated by the dotted lines. Animals were monitored by quantifying bacterial growth at the indicated times. (A) Bacterial growth in organs (spleen, liver, and brain) and in blood of pregnant mice infected with  $\Delta$ ActA (left panels) and *actA*-complemented  $\Delta$ ActA+*actA* (right panels) strains. Results shown are means  $\pm$  standard errors from groups of three to five mice and are expressed as the log<sub>10</sub> bacteria (CFU) per organ or log<sub>10</sub> bacteria per ml of blood. (B) The kinetics of bacterial growth are represented on the left panels by dot plots corresponding to individual bacterial counts for each placenta (top panel) and each fetus (lower panel) for the indicated times and strains. In the right panels, the means  $\pm$  standard errors for each time and strain are given for  $\Delta$ ActA and the *actA*-complemented  $\Delta$ Act mutant and *P* values for the comparison between the two strains are also reported. The numbers reported near each value correspond to the percentage of infected placenta among all analyzed placentas. All data are representative of three independent experiments. *P* values reported in each graph correspond to the comparison between EGDwt and the indicated mutant. NS, not statistically different.

TABLE 2. Cutoff values for bacterial placental load associated with the highest probability of fetal infection

Bacterial strain	No. of fetoplacental units analyzed	Cutoff value (10 <sup>3</sup> CFU) <sup>a</sup>	Sensitivity (%)	Specificity (%)	PPV (%) <sup>b</sup>	NPV (%) <sup>c</sup>
EGDwt	185	1.7	91	99	99	91
$\Delta$ InlAB	48	1.0	60	92	85	73
$\Delta$ ActA	72	120.0	93	82	82	93
$\Delta$ ActA+ <i>actA</i>	36	2.0	89	100	100	87

<sup>a</sup> Cutoff values correspond to level of bacterial load of the placenta giving the highest probability of fetal infection.

<sup>b</sup> PPV, positive predictive value.

<sup>c</sup> NPV, negative predictive value.

rinthine zone is a crucial step for the crossing the fetoplacental barrier in the murine model.

DISCUSSION

Our results first show that the placenta is an advantageous and protective niche for any bacteria, even the nonvirulent  $\Delta$ LO *L. monocytogenes* mutant and *L. innocua*, which could survive at a stable and low level for several days in the placenta. Trophoblastic cells appear to play an immunological role against pathogens, protecting the fetuses from bacterial aggression, but are also the first placental target cells for intracellular pathogens as *L. monocytogenes* (14, 21).




FIG. 4. Histological examination of placental infectious foci obtained with wild-type *L. monocytogenes* and its  $\Delta$ InlAB and  $\Delta$ ActA isogenic mutants. Gram-Weigert staining at different magnifications (top and bottom panels) were performed on placental sections 72 h after intravenous infection with  $2 \times 10^5$  wild-type *L. monocytogenes* cells or with mutants with the *inlAB* locus ( $\Delta$ InlAB) or *actA* ( $\Delta$ ActA) deleted. Arrows indicate the important spreading to adjacent and distant cells. Arrowheads symbolized the well-delimited peripheral zones from foci induced by infection due to the  $\Delta$ ActA mutant.

Our data are in agreement with previous reports demonstrating that trophoblastic cells express an important phagocytic capacity, mainly during the two first trimesters of pregnancy in humans, a function still conserved in the late stage of pregnancy (2). The bacterial survival of nonvirulent bacteria in the placenta might be due to several causes, including the presence of latent bacteria confined in the phagosomal compartment; low intracellular growth due to the escape of few  $\Delta$ LLO bacteria by production of phospholipases (22); and extracellular replication in the placenta, which constitutes a protective environment for bacterial growth. Indeed, the immunological status of the placenta is characterized by a predominant Th2 anti-inflammatory response to prevent the rejection of the semiallogeneic fetus, thus restraining the recruitment of inflammatory cells at the early onset of the infection.

We observed that the survival of  $\Delta$ LLO mutants and *L. innocua* in the placenta was associated with neither bacterial growth nor fetal infection, as opposed to virulent wild-type *L. monocytogenes* as previously observed in the guinea pig model (5). This indicates that the virulence factor LLO is absolutely required in vivo for bacterial growth in trophoblastic cells and the subsequent fetal invasion. The vertical transmission of *L. monocytogenes* to the fetus is therefore dependent upon the expression of virulence factors promoting intracellular multiplication of this pathogen. Consequently, we first tested in pregnant mice a  $\Delta$ InlAB mutant of *L. monocytogenes*, which might be affected in its invasive capacity for trophoblastic cells. It has been published that internalization of *L. monocytogenes* in a human trophoblas-

tic cell line (BeWo) infected in vitro requires the expression of InlA (4), a surface-exposed protein interacting with high affinity with human and guinea pig E-cadherin. The affinity of mouse E-cadherin for InlA is lower than that of guinea pig E-cadherin (18). However, this single in vitro interaction does not account for the mechanism of in vivo infection of trophoblastic cells by *L. monocytogenes*, since the rate of placental infection with an InlA mutant was not reduced in the guinea pig model (4). Similarly, we found in the mouse model that a strain with InlA and InlB deletion invaded placentas and fetuses as well as did wild-type bacteria. Therefore, InlA and InlB are not implicated in the crossing of the murine fetoplacental barrier as for guinea pigs (4).

We then studied the role of ActA in placental infection by using a  $\Delta$ ActA mutant in pregnant mice. Growth of this virulence-attenuated mutant was restricted in vivo, as indicated by the absence of bacteremia and the rapid elimination of bacteria in organs, as previously described (16). However, we observed that the  $\Delta$ ActA strain retained the ability to grow in the placenta, although at a low rate compared to wild-type bacteria. As expected,  $\Delta$ ActA bacteria were strongly impaired in cell-to-cell spreading through placental tissues, bacteria often being seen packed inside trophoblastic cells, contrasting with the outward spreading and the rapid dissemination of wild-type bacteria to neighboring cells. This growth defect of  $\Delta$ ActA bacteria was correlated to a delayed and decreased rate of fetal infection. The placental bacterial load required for fetal infection with the  $\Delta$ ActA mutant was 2 log units higher than that seen with EGDwt. However, impairment of placental growth of the  $\Delta$ ActA mutant might also




FIG. 5. Model of crossing of the fetoplacental barrier by *L. monocytogenes*. Two layers of syncytiotrophoblastic cells (ScT) and one layer of endothelial cells (EC) surrounding a fetal vessel (FV) constitute the murine fetoplacental barrier. *L. monocytogenes* (LM) infects the syncytiotrophoblastic layer and crosses the fetoplacental barrier by ActA-dependent cell-to-cell spreading.

be partly explained by the lack of bacteremia, which was observed for wild-type bacteria in pregnant mice. Our results in the murine model strongly support the major role of cell-to-cell spreading in the vertical transmission of *L. monocytogenes*.

In conclusion, together with those of Barkadjiev et al. from guinea pigs (4, 5), our data demonstrate in the murine model (i) that the surface-exposed proteins InlA and InlB are not necessary for placental invasion and (ii) the crucial role of ActA-dependent cell-to-cell spreading in the vertical transmission of *L. monocytogenes* to the fetus, including direct evidence obtained by fully restoring fetal invasion in an *actA*-complemented  $\Delta$ ActA mutant. Therefore, the crossing of the murine fetoplacental barrier requiring ActA-dependent cell-to-cell spreading allows bacteria to cross the two layers of trophoblastic cells and endothelial cells from the fetal vessel, as illustrated in Fig. 5.

#### ACKNOWLEDGMENTS

We thank P. Cossart (Institut Pasteur, Paris) for the gift of various *Listeria* mutants and G. Pivert for the technical assistance.

This work was supported by University Paris-Descartes, Inserm (ANR-IRAP.2005), and Assistance Publique—Hopitaux de Paris.

#### REFERENCES

- Alouf, J. E. 2000. Cholesterol-binding cytolytic protein toxins. *Int. J. Med. Microbiol.* **290**:351–356.
- Amarante-Paffaro, A., G. S. Queiroz, S. T. Correa, B. Spira, and E. Bevilacqua. 2004. Phagocytosis as a potential mechanism for microbial defense of mouse placental trophoblast cells. *Reproduction* **128**:207–218.
- Autret, N., I. Dubail, P. Trieu-Cuot, P. Berche, and A. Charbit. 2001. Identification of new genes involved in the virulence of *Listeria monocytogenes* by signature-tagged transposon mutagenesis. *Infect. Immun.* **69**:2054–2065.
- Bakardjiev, A. I., B. A. Stacy, S. J. Fisher, and D. A. Portnoy. 2004. Listeriosis in the pregnant guinea pig: a model of vertical transmission. *Infect. Immun.* **72**:489–497.
- Bakardjiev, A. I., B. A. Stacy, and D. A. Portnoy. 2005. Growth of *Listeria monocytogenes* in the Guinea pig placenta and role of cell-to-cell spread in fetal infection. *J. Infect. Dis.* **191**:1889–1897.
- Buendia, A. J., R. M. De Oca, J. A. Navarro, J. Sánchez, F. Cuello, and J. Salinas. 1999. Role of polymorphonuclear neutrophils in a murine model of *Chlamydia psittaci*-induced abortion. *Infect. Immun.* **67**:2110–2116.
- Chakraborty, T., F. Ebel, E. Domann, K. Niebuhr, B. Gerstel, S. Pistor, C. J. Temm-Grove, B. M. Jockusch, M. Reinhard, U. Walter, et al. 1995. A focal adhesion factor directly linking intracellularly motile *Listeria monocytogenes* and *Listeria ivanovii* to the actin-based cytoskeleton of mammalian cells. *EMBO J.* **14**:1314–1321.
- Cossart, P. 2002. Molecular and cellular basis of the infection by *Listeria monocytogenes*: an overview. *Int. J. Med. Microbiol.* **291**:401–409.
- Doganay, M. 2003. Listeriosis: clinical presentation. *FEMS Immunol. Med. Microbiol.* **35**:173–175.
- Dussurget, O., J. Pizarro-Cerda, and P. Cossart. 2004. Molecular determinants of *Listeria monocytogenes* virulence. *Annu. Rev. Microbiol.* **58**:587–610.
- Gaillard, J. L., P. Berche, and P. Sansonetti. 1986. Transposon mutagenesis as a tool to study the role of hemolysin in the virulence of *Listeria monocytogenes*. *Infect. Immun.* **52**:50–55.
- Glaser, P., L. Frangeul, C. Buchrieser, C. Rusniok, A. Amend, F. Baquero, P. Berche, H. Bloecker, P. Brandt, T. Chakraborty, A. Charbit, F. Chetouani, E. Couve, A. de Daruvar, P. Dehoux, E. Domann, G. Dominguez-Bernal, E. Duchaud, L. Durant, O. Dussurget, K. D. Entian, H. Fsihi, F. Garcia-del Portillo, P. Garrido, L. Gautier, W. Goebel, N. Gomez-Lopez, T. Hain, J. Hauf, D. Jackson, L. M. Jones, U. Kaerst, J. Kreft, M. Kuhn, F. Kunst, G. Kurapat, E. Madueno, A. Maitournam, J. M. Vicente, E. Ng, H. Nedjari, G. Nordsiek, S. Novella, B. de Pablos, J. C. Perez-Diaz, R. Purcell, B. Remmel, M. Rose, T. Schlueter, N. Simoes, A. Tierrez, J. A. Vazquez-Boland, H. Voss, J. Wehland, and P. Cossart. 2001. Comparative genomics of *Listeria* species. *Science* **294**:849–852.
- Gray, M. L., and A. H. Killinger. 1966. *Listeria monocytogenes* and listeric infections. *Bacteriol. Rev.* **30**:309–382.
- Guleria, I., and J. W. Pollard. 2000. The trophoblast is a component of the innate immune system during pregnancy. *Nat. Med.* **6**:589–593.
- Guzman, C. A., M. Rohde, T. Chakraborty, E. Domann, M. Hudel, J. Wehland, and K. N. Timmis. 1995. Interaction of *Listeria monocytogenes* with mouse dendritic cells. *Infect. Immun.* **63**:3665–3673.
- Kocks, C., E. Gouin, M. Tabouret, P. Berche, H. Ohayon, and P. Cossart. 1992. *L. monocytogenes*-induced actin assembly requires the actA gene product, a surface protein. *Cell* **68**:521–531.
- Koi, H., J. Zhang, and S. Parry. 2001. The mechanisms of placental viral infection. *Ann. N. Y. Acad. Sci.* **943**:148–156.
- Lecuit, M., S. Dramsi, C. Gottardi, M. Fedor-Chaiken, B. Gumbiner, and P. Cossart. 1999. A single amino acid in E-cadherin responsible for host specificity towards the human pathogen *Listeria monocytogenes*. *EMBO J.* **18**:3956–3963.
- Lecuit, M., D. M. Nelson, S. D. Smith, H. Khun, M. Huerre, M. C. Vacher-Lavenu, J. I. Gordon, and P. Cossart. 2004. Targeting and crossing of the human maternofetal barrier by *Listeria monocytogenes*: role of internalin interaction with trophoblast E-cadherin. *Proc. Natl. Acad. Sci. USA* **101**:6152–6157.
- Lecuit, M., S. Vandormael-Pournin, J. Lefort, M. Huerre, P. Gounon, C. Dupuy, C. Babinet, and P. Cossart. 2001. A transgenic model for listeriosis: role of internalin in crossing the intestinal barrier. *Science* **292**:1722–1725.
- Le Monnier, A., O. F. Join-Lambert, F. Jaubert, P. Berche, and S. Kayal. 2006. Invasion of the placenta during murine listeriosis. *Infect. Immun.* **74**:663–672.
- Marquis, H., V. Doshi, and D. A. Portnoy. 1995. The broad-range phospholipase C and a metalloprotease mediate listeriolysin O-independent escape of *Listeria monocytogenes* from a primary vacuole in human epithelial cells. *Infect. Immun.* **63**:4531–4534.
- Maurin, M., and D. Raoult. 1999. Q fever. *Clin. Microbiol. Rev.* **12**:518–553.
- Mylonakis, E., M. Paliou, E. L. Hohmann, S. B. Calderwood, and E. J. Wing. 2002. Listeriosis during pregnancy: a case series and review of 222 cases. *Medicine (Baltimore)* **81**:260–269.
- Pappas, G., N. Akritidis, M. Bosilkovski, and E. Tsianos. 2005. Brucellosis. *N. Engl. J. Med.* **352**:2325–2336.
- Pfaff, A. W., S. Georges, A. Abou-Bacar, V. Letscher-Bru, J. P. Klein, M. Mousli, and E. Candolfi. 2005. *Toxoplasma gondii* regulates ICAM-1 mediated monocyte adhesion to trophoblasts. *Immunol. Cell Biol.* **83**:483–489.

27. **Scherf, A., B. Pouvelle, P. A. Buffet, and J. Gysin.** 2001. Molecular mechanisms of *Plasmodium falciparum* placental adhesion. *Cell Microbiol.* **3**:125–131.
28. **Tilney, L. G., and D. A. Portnoy.** 1989. Actin filaments and the growth, movement, and spread of the intracellular bacterial parasite, *Listeria monocytogenes*. *J. Cell Biol.* **109**:1597–1608.
29. **Trieu-Cuot, P., C. Carlier, C. Poyart-Salmeron, and P. Courvalin.** 1991. An integrative vector exploiting the transposition properties of Tn1545 for insertional mutagenesis and cloning of genes from gram-positive bacteria. *Gene* **106**:21–27.
30. **Vázquez-Boland, J. A., M. Kuhn, P. Berche, T. Chakraborty, G. Dominguez-Bernal, W. Goebel, B. Gonzalez-Zorn, J. Wehland, and J. Kreft.** 2001. *Listeria* pathogenesis and molecular virulence determinants. *Clin. Microbiol. Rev.* **14**:584–640.

---

*Editor:* D. L. Burns