

## Strain-Specific Single-Nucleotide Polymorphism Assays for the *Bacillus anthracis* Ames Strain<sup>†</sup>∇

Matthew N. Van Ert,<sup>1</sup> W. Ryan Easterday,<sup>1</sup> Tatum S. Simonson,<sup>1</sup> Jana M. U'Ren,<sup>1</sup> Talima Pearson,<sup>1</sup> Leo J. Kenefic,<sup>1</sup> Joseph D. Busch,<sup>1</sup> Lynn Y. Huynh,<sup>1</sup> Megan Dukerich,<sup>1</sup> Carla B. Trim,<sup>1</sup> Jodi Beaudry,<sup>1</sup> Amy Welty-Bernard,<sup>1</sup> Timothy Read,<sup>3</sup> Claire M. Fraser,<sup>3</sup> Jacques Ravel,<sup>3</sup> and Paul Keim<sup>1,2\*</sup>

Northern Arizona University, Department of Biological Sciences, Box 5640, Flagstaff, Arizona 86011<sup>1</sup>;  
Translational Genomics Research Institute, Pathogen Genomics Division, 445 N. Fifth Street,  
Phoenix, Arizona 85004<sup>2</sup>; and The Institute for Genomic Research, 9712 Medical Center Drive,  
Rockville, Maryland 20723<sup>3</sup>

Received 15 June 2006/Returned for modification 22 July 2006/Accepted 7 October 2006

**Highly precise diagnostics and forensic assays can be developed through a combination of evolutionary analysis and the exhaustive examination of genomic sequences. In *Bacillus anthracis*, whole-genome sequencing efforts revealed ca. 3,500 single-nucleotide polymorphisms (SNPs) among eight different strains and evolutionary analysis provides the identification of canonical SNPs. We have previously shown that SNPs are highly evolutionarily stable, and the clonal nature of *B. anthracis* makes them ideal signatures for subtyping this pathogen. Here we identified SNPs that define the lineage of *B. anthracis* that contains the Ames strain, the strain used in the 2001 bioterrorist attacks in the United States. Sequencing and real-time PCR were used to validate these SNPs across *B. anthracis* strains, including (i) 88 globally and genetically diverse isolates; (ii) isolates that were shown to be genetic relatives of the Ames strain by multiple-locus variable number tandem repeat analysis (MLVA); and (iii) several different lab stocks of the Ames strain, including a clinical isolate from the 2001 letter attack. Six SNPs were found to be highly specific for the Ames strain; four on the chromosome, one on the pX01 plasmid, and one on the pX02 plasmid. All six SNPs differentiated the *B. anthracis* Ames strain from the 88 unique *B. anthracis* strains, while five of the six separated Ames from its close genetic relatives. The use of these SNPs coupled with real-time PCR allows specific and sensitive (<100 fg of template DNA) identification of the Ames strain. This evolutionary and genomics-based approach provides an effective means for the discovery of strain-specific SNPs in *B. anthracis*.**

The 2001 anthrax letter attack illustrated the “real-world” efficacy of *Bacillus anthracis* as a bioterror agent. Forensic and epidemiological analysis of clinical samples and weaponized spores from the letter attack included the identification of the *B. anthracis* strain as Ames (3, 8, 17). This was initially accomplished using multiple-locus variable number tandem repeat analysis (MLVA) (3, 6, 8), which was one of the few technical approaches possible for this highly monomorphic pathogen (5). Identifying the strain and establishing its identity in attack locations were important in linking the dispersed anthrax cases and suggesting a possible source for the weaponized material. During the anthrax attack crisis, diagnostic speed, specificity, and sensitivity were often limiting factors, necessitating extraordinary efforts by public health officials and forensic labs (3). Further advancement of molecular diagnostics allows for more-efficient responses in disease outbreaks, whether natural or bioterrorist mediated.

Single-nucleotide polymorphisms (SNPs) are increasingly recognized as important markers for detecting and subtyping bacterial pathogens, including *B. anthracis* (1, 2, 4, 8, 16, 18,

19). Recent comparative full-genome sequencing allowed the discovery of about 3,500 SNPs among eight strains of *B. anthracis* (15, 17; J. Ravel, unpublished data) and represents a valuable resource for developing diagnostic signatures for this pathogen. SNPs are particularly attractive markers for subtyping efforts since they are evolutionarily stable (8, 15) and are amenable to high-throughput detection methods, such as real-time PCR, pyrosequencing, and mass spectrometry (1, 2, 4, 10, 20, 22, 23).

As binary markers, SNPs would seem to have limited subtyping power, requiring a large number of SNPs to be interrogated for subtyping purposes. However, researchers have demonstrated that a small number of SNPs can be used to efficiently define genetic groups. For example, Moorhead et al. (11) used a phylogenetic approach to identify a small number of diagnostic SNPs in the *sigB* gene that partitioned strains of *Listeria monocytogenes* into three previously described clonal lineages. Keim et al. (8) expanded upon this concept and proposed the idea of a “canonical SNP” (canSNP), an SNP that can be used to efficiently identify a point in the evolutionary history of a species. Mapping SNPs on a genetic population structure allows the identification of canSNPs that can be used diagnostically on a broad scale to define major genetic lineages in a species or, more narrowly, to define specific strains (8). A more recent publication used a similar approach to identify a streamlined set of SNPs that divide methicillin-resistant *Staphylococcus aureus* isolates into genetic groups that are consistent with its population structure (18).

\* Corresponding author. Mailing address: Department of Biological Sciences, Northern Arizona University, Flagstaff, AZ 86011-5640. Phone: (928) 523-1078. Fax: (928) 523-0639. E-mail: Paul.Keim@nau.edu.

† Supplemental material for this article may be found at <http://jcm.asm.org/>.

∇ Published ahead of print on 8 November 2006.

TABLE 1. Sequences of primers and TaqMan MGB probes

SNP	5' to 3' sequences for:		SNP change and genome position	GenBank accession no.
	Primers	TaqMan MGB probe <sup>a</sup>		
PS-1	F-TGATGGTTTTGATTTCTTAGGCTTT	FAM-AAGGGTCA <u>CA</u> CAACTC	G↔A, 7452	NC_003980
	R-CACTTTGGTTGGATGGTTAATGA	VIC-AAGGGTCG <u>CA</u> CAACTC		
PS-52	F-GTATCCTGAAATATAAAAAGTGTA AAGGTAATAAAATGGA	VIC-ATTAAGGACTCCCTCTTGTT	A↔C, 72924	NC_003981
	R-GATTCTTCAACGCAATATACCCTA CTAAAATTATACTAT	FAM-AAGGACTCCCTA <u>TT</u> GGTT		
Branch1-7	F-TCACCTCAATGACATCGCCA	FAM-CAAACCAATAC <u>CC</u> CCTTT	C↔A, 433277	NC_003997
	R-TTGTTGTGAAGACGGATAACTTT TATG	VIC-CAAACCAATA <u>AC</u> CCCTTT		
Branch1-26	F-GACGGGAGCCAACCAGAA	FAM-ATAGCTTTTT <u>TT</u> TCTATTCC	T↔C, 4624132	NC_003997
	R-CCGTTGAATAAGCAGTATGAAA TTTC	VIC-ATAGCTTTTT <u>CT</u> TCTATTCC		
Branch1-28	F-AATATCTTTCATACAAGGCGCAC TACT	FAM-CGTTGTAG <u>TT</u> TATTTTAC	T↔G, 4929186	NC_003997
	R-CCATAATCGTGCTTGTCCAAATC	VIC-CGTTGTAG <u>G</u> TATTTTAC		
Branch1-31	F-GAAGAACAAGCGAAAGACGT ACCT	VIC-CGTTACAT <u>G</u> GCAT	A↔G, 2749543	NC_003997
	R-GTAGTTCATAACGTTTGAAAAAGT AGGGATA	FAM-TCGGTTCACAT <u>A</u> GCAT		

<sup>a</sup> Note that all probes have a minor-groove binder, nonfluorescent quencher attached to the 3' end. FAM, 6-carboxyfluorescein. The Ames probes were labeled with FAM except in the case of PS-1, which was labeled with VIC. Underlined letters indicate SNP location.

<sup>b</sup> SNP changes are expressed as "Ames ↔ non-Ames."

In this study, we used a subset of about 1,000 previously published SNPs (15, 17) to identify canSNPs that effectively define the *B. anthracis* lineage that contains the Ames strain. To locate SNPs with the highest specificity for the Ames strain, we screened these SNPs across a panel of isolates that were identified as close genetic relatives of the Ames strain by using high-resolution MLVA subtyping (6, 17; M. N. Van Ert and P. Keim, unpublished data). Here we demonstrate the use of strain-specific canSNPs for subtyping *B. anthracis*, which can assist in epidemiological and forensic investigations.

#### MATERIALS AND METHODS

**Strains used in this study.** We selected a panel of 89 *B. anthracis* strains that represent 89 unique MLVA genotypes as described by Keim et al. (6) and includes an Ames strain isolate (genotype 62). In addition, we included two isolates that were previously identified as Ames ( $\Delta$ Ames [A0814; Ames cured of pX01 and pX02]), a clinical isolate from the 2001 letter attack ([A2012]), and an additional 10 isolates previously shown to be genetically related to the Ames strain by sequencing and MLVA analysis (17; Van Ert and Keim, unpublished; see Table S1 in the supplemental material).

**DNA isolation and quantitation.** DNA was isolated using one of three methods (see the supplemental material for details). DNA was quantified using a PicoGreen double-stranded DNA quantitation kit (Molecular Probes, Inc., Eugene, OR) according to the manufacturer's protocol. Sample fluorescence was measured using a TBS-300 minifluorometer (Turner Biosystems, Inc., Sunnyvale, CA).

**Identification and screening of Ames strain-specific SNPs.** About 3,500 SNPs located on the *B. anthracis* chromosome and plasmids were previously discovered by whole-genome comparative sequencing (17; J. Ravel, unpublished). Using a subset of about 1,000 SNPs, Pearson et al. (15) identified 32 chromosomal SNPs that were positioned on the phylogenetic branch contain-

ing the Ames strain (see Fig. S1 in the supplemental material). SNPs located on the *B. anthracis* plasmids were identified by Read et al. (17) by comparing the sequences of the Ames strain pX01 and pX02 plasmids to those of the previously published pX01 (*B. anthracis* Sterne [13]) and pX02 (*B. anthracis* Pasteur [14]) plasmids. The authors indicated that the SNPs exhibited specificity for the Ames strain in relation to the other strains involved in the study, including the closely related A0394 "Texas goat" isolate (see Table S1 in the supplemental material). Based upon these results, two SNPs located on the pX01 plasmid (PS-1 and PS-32) and one SNP located on the pX02 plasmid (PS-52) were selected as candidates.

**Initial screening of Ames strain-specific chromosomal SNPs.** We used DNA sequencing to screen the 32 chromosomal SNPs across a panel of DNA from the Ames strain, isolates that were genetically similar to the Ames strain and from a distantly related *B. anthracis* strain (Vollum; see Table S1 in the supplemental material). DNA sequencing was performed according to the manufacturer's protocol (Applied Biosystems, Foster City, CA). The SNPs located on the pX01 and pX02 plasmids were screened against these strains using allelic discrimination real-time PCR.

**TaqMan MGB allelic discrimination assays.** TaqMan minor-groove binding (MGB) allelic discrimination assays were designed around 4 of the 32 chromosomal SNPs which exhibited the highest specificity for the Ames strain. In addition, TaqMan MGB allelic discrimination assays were designed around three plasmid SNPs, PS-1, PS-32, and PS-52, reported by Read et al. (17).

TaqMan MGB probes and primers for the four chromosomal SNPs and the pX01 SNPs were designed by using ABI Primer Express software and guidelines, with the exception that allele-specific probe lengths were manually adjusted to match melting temperatures (12). Primers and probes to detect the pX02 PS-52 SNP were designed by Applied Biosystems. Probe and primer sequences are listed in Table 1. Each 10.0- $\mu$ l reaction mixture contained 1 $\times$  ABI Universal Master Mix, 250 nM of each probe, 600 nM each of forward and reverse primers, and 1.0  $\mu$ l of approximately 350 pg/ $\mu$ l template DNA. For all assays, thermal cycling parameters were 50°C for 2 min and 95°C for 10 min, followed by 40 to 50 cycles of 95°C for 15 s and 60°C for 1 min.

The TaqMan MGB assays were used to genotype a DNA collection representing a worldwide panel of 88 diverse *B. anthracis* strains, Ames relatives, and

TABLE 2. Profiles of chromosomal and plasmid-based single-nucleotide polymorphisms<sup>a,b</sup>

SNP	Profiles for indicated isolates:												
	A0462 Ames	A1115 Texas	A1117 Texas	A0394 Texas Goat	A0728 China	A0584 China	A0576 China	A0585 China	A0580 China	A0720 China	A0724 China	A0729 China	A0488 Vollum
PS-52	A	C	C	C	C	C	C	C	C	C	C	C	C
Branch1-7	C	A	A	A	A	A	A	A	A	A	A	A	A
Branch1-26	T	C	C	C	C	C	C	C	C	C	C	C	C
Branch1-28	T	G	G	G	G	G	G	G	G	G	G	G	G
Branch1-31	A	G	G	G	G	G	G	G	G	G	G	G	G
Branch1-8	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-15	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	T	T	T	T	T	T	T	T
Branch1-16	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-18	A	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	G	G	G	G	G	G	G	G
Branch1-22	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T	T	T	T	T	T	T
Branch1-23	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-27	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-30	A	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	C	C	C	C	C	C	C	C
PS-1	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-1	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-6	A	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	A	G	G	G	G	G	G	G
Branch1-9	A	<u>A</u>	<u>A</u>	<u>A</u>	<u>A</u>	T	T	T	T	T	T	T	T
Branch1-10	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-11	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T	T	T	T	T	T	T
Branch1-12	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-13	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-14	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-17	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T	T	T	T	T	T	T
Branch1-19	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T	T	T	T	T	T	T
Branch1-20	T	<u>T</u>	<u>T</u>	<u>T</u>	<u>T</u>	C	C	C	C	C	C	C	C
Branch1-24	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-25	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	A	A	A	A	A	A	A	A
Branch1-29	G	<u>G</u>	<u>G</u>	<u>G</u>	<u>G</u>	A	A	A	A	A	A	A	A
Branch1-21	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	C	C	C	T	T	T	T	T
Branch1-5	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T	T	T
PS-32	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T	T
Branch1-2	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T
Branch1-3	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T
Branch1-4	C	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	<u>C</u>	T

<sup>a</sup> Underlined letters indicate alleles shared with the Ames strain. One of the chromosomal SNPs was removed from the analysis since our sequence data indicated that the SNP was not specific for the phylogenetic branch containing the Ames strain (data not shown).

<sup>b</sup> SNP profiles for the chromosomal SNPs Branch1-1 to Branch1-31 were determined by DNA sequencing using flanking primers, whereas SNP profiles for the plasmid-based SNPs were determined by TaqMan MGB genotyping.

a panel of Ames strains. Real-time and endpoint fluorescence data were collected on the ABI 7900 to confirm robust allele-specific detection of the target sequences and clear endpoint allelic discrimination.

**Sensitivity and level of detection of TaqMan MGB assays.** To determine the limit of detection of the assays, serial dilutions of DNA from an Ames strain (A0462) and a non-Ames strain (A0488; Vollum) ranging from 100 pg to 10 fg were used as a template for the TaqMan MGB assays.

**RESULTS**

**Identification and screening of Ames strain-specific SNPs.**

Sequence analysis of the 32 chromosomal SNPs located along the *B. anthracis* Ames strain lineage (see Fig. S1 in the supplemental material; 15) indicated four SNPs (Branch1-7, Branch1-26, Branch1-28, and Branch1-31) separated the Ames MLVA type from all of the remaining genetic relatives (Table 2). Notably, this includes the separation of Ames from three closely related Texas isolates. These four SNPs were selected for further testing based upon their specificity for the Ames strain. Of the three TaqMan MGB assays designed around plasmid SNPs, the PS-52 SNP assay separated the Ames MLVA type from all of the remaining genetic relatives, whereas the PS-1 SNP assay exhibited less specificity and grouped the Ames strain with a single Chinese strain

and the three closely related Texas isolates (Table 2). The PS-32 SNP assay exhibited much lower specificity for the Ames strain (Table 2).

**TaqMan allelic discrimination genotyping assays.** To confirm the specificity of the Ames SNP assays (for SNPs, Branch1-7, Branch1-26, Branch1-28, Branch1-31, PS-1, and PS-52) for the Ames strain, we used TaqMan dual-probe allelic discrimination to genotype 88 globally and genetically diverse *B. anthracis* isolates. The cycle threshold ( $C_T$ ) values observed in the real-time data were similar among the assays designed around the chromosomal and pX02 SNPs (for Branch1-7, the mean was 21.1  $C_T$ , the standard deviation [SD] was  $\pm 0.96$ , and the range was 18.3 to 23.3; for Branch1-26, the mean was 23.6  $C_T$ , the SD was  $\pm 0.91$ , and the range was 20.5 to 26.0; for Branch1-28 the mean was 21.5  $C_T$ , the SD was  $\pm 0.93$ , and the range was 18.8 to 23.2; for Branch1-31 the mean was 22.6  $C_T$ , the SD was  $\pm 1.01$ , and the range was 19.5 to 24.7; and for PS-52, the mean was 22.9  $C_T$ , the SD was  $\pm 1.2$ , and the range was 19.6 to 27.7). The assay designed around the pX01 SNP PS-1 exhibited lower  $C_T$  values (the mean was 18.8  $C_T$ , the SD was  $\pm 0.90$ , and the range was 16.2 to 21.3).




FIG. 1. TaqMan endpoint allelic discrimination plots of the six Ames-specific SNP assays. Chromosomal SNPs for Branch1-7, Branch1-26, Branch1-28, and Branch1-31 loci are shown in panels A, B, C, and D, respectively. The dots along the y axis ( $n = 2$ ; blue) demonstrate the Ames-specific genotype; samples along the x axis (red) demonstrate genotyping of the non-Ames *B. anthracis* samples (diverse 88). Samples near the plot origin are negative, no-template controls ( $n = 3$ ). Panel E illustrates the PS-1 pX01 plasmid marker. The dots along the x axis (blue;  $n = 2$ ) demonstrate the Ames-specific genotype; samples along the y axis (red) demonstrate genotyping of the non-Ames *B. anthracis* samples (diverse 88). Samples near the plot origin are no-template controls ( $n = 2$ ) or pX01-negative *B. anthracis* strains ( $n = 2$ ). Panel F illustrates the pX02 plasmid marker PS-52. The dots along the y axis (blue) demonstrate the Ames-specific genotype; samples along the x axis (red) demonstrate genotyping of the “non-Ames” *B. anthracis* samples (diverse 88). Samples near the plot origin are negative controls ( $n = 2$ ) or pX02-negative *B. anthracis* strains ( $n = 1$ ). All PCRs were cycled 40 times.

The post-PCR allelic discrimination plot from these experiments illustrates the unambiguous separation of genotypes for the Ames strain and remaining “non-Ames” diverse 88 strains. All six SNP assays separated the Ames strains from the remaining 88 globally diverse *B. anthracis* genotypes (Fig. 1A to F). Although the PS-1 SNP assay separates Ames from the remaining 88 global strains, it will cluster Ames with three closely related Texas isolates and a single Chinese isolate (TaqMan data not shown; Table 2 presents SNP profile data).

**Level of detection of TaqMan MGB genotyping assays.** The TaqMan MGB allelic discrimination assays were used to reliably detect and genotype samples containing as little as 100 fg of template DNA (ca. 17 genome equivalents), whereas samples containing 10 fg of template DNA (ca. 1.7 genome equivalents) could be detected and genotyped sporadically. Fig. 2A illustrates the real-time amplification plots generated from analysis of a 10-fold serial dilution of template DNA from the Ames strain with the Branch1-7 SNP TaqMan assay. The post-PCR allelic discrimination plot from the same experiment (Fig.




FIG. 2. (A) Real-time plots of the Branch1-7 TaqMan MGB allelic-discrimination assay showing a dilution curve of Ames (A0462) DNA ranging from 100 pg to 10 fg (all quantities shown in triplicate; only fluorescence from the Ames allele probe is illustrated). (B) Endpoint allelic discrimination plots from the same Ames DNA dilution curve (blue cluster along the x axis) and from an equivalent non-Ames (A0488; Vollum) dilution curve (red cluster along the x axis) illustrate clear genotypic separation. Samples near the plot origin are negative controls (black squares;  $n = 6$ ). Note that the samples containing higher amounts of DNA (100 pg) clustered further from the origin, and samples with low DNA levels (10 fg) clustered nearer the origin. Note that all replicates at 100 and 10 fg amplified for both strains. The PCR was cycled 50 times.

2B) illustrates the unambiguous separation of genotypes for both the Ames strain and the non-Ames (Vollum) strain over a range of template DNA quantities from 100 pg to 10 fg.  $C_T$  values for all of the assays run across this serial dilution of template DNA, and numerous replicates of 10 fg input DNA are reported in Tables S2 and S3 in the supplemental material.

## DISCUSSION

The identification of strain-specific SNP markers in *B. anthracis* permits the development of rapid diagnostics to greatly assist in the investigation of biocrimes and natural outbreaks. Forensic and epidemiological investigations can require the analysis of hundreds or even thousands of specimens, including environmental and clinical samples. During the 2001 anthrax outbreak, the CDC used MLVA to subtype over 100 *B. anthracis* isolates and several dozen clinical samples (3). Significantly, these researchers used MLVA to include or exclude *B. anthracis* strains from the epidemiological investigation. Since MLVA requires PCR, post-PCR processing, electrophoresis, and fragment sizing, there is a practical limit to the number of samples that can be strain typed in a timely manner. In contrast, Ames-specific real-time assays allow thousands of DNA samples per day to be included or excluded as the Ames strain.

The combination of Ames strain-specific SNPs and real-time PCR allows for the development of assays that are both specific and sensitive. Five SNPs we identified were highly specific and were observed in only the Ames strains (A2012, A0462, and A0814 [chromosome only]), whereas a single SNP located on the pX01 plasmid (PS-1) partitioned the Ames strain with three closely related Texas isolates and a single Chinese strain. The pX01 SNP assay is still valuable because of its ability to monitor the plasmid composition of an isolate. Our data suggest that the SNPs which define the Ames genetic lineage are evolutionarily stable since SNPs found in the Ames strain, but not close genetic relatives, were strain specific when evaluated across a more comprehensive isolate collection (i.e., the diverse 88 *B. anthracis* strains).

The stability of these SNPs as diagnostic markers is likely a function of the low mutation rates of nucleotide substitutions in *B. anthracis* ( $10^{-10}$  per site per generation [21]), the recent evolutionary derivation of *B. anthracis* from *Bacillus cereus*, and the lack of recombination due to the highly clonal nature of *B. anthracis* (8, 15). The rarity of these mutational events within *B. anthracis* limits the likelihood of a SNP mutating again to a novel or ancestral state and lowers the probability of observing a false positive in any one of the SNP assays (i.e., a

non-Ames strain sharing the allele with Ames). The lack of recombination ensures that the few mutational events are restricted to a particular clade. The combinatorial use of the six Ames SNP assays vastly increases confidence in the inclusion/exclusion of a sample as the Ames strain since the likelihood that a different lineage of *B. anthracis* would convergently evolve to match this six-locus genotype is vanishingly small (e.g., the product of the six individual locus mutation rates is about  $\sim 10^{-60}$ ).

The Ames strain-specific PCR assays were sensitive, allowing reliable detection and genotyping of samples containing 100 fg of template DNA, which we estimate corresponds to approximately 17 genome equivalents. We routinely use one chromosomal SNP assay (Branch1-31) and two plasmid SNP assays (PS-1 and PS-52) on blinded proficiency samples and have achieved 100% accuracy in discriminating Ames versus non-Ames strain templates over a range of DNA levels (>260 samples [data not shown]). It is important to point out that the distinct genotypic clustering of samples in the endpoint analysis and the similar  $C_T$  values observed among *B. anthracis* strains argue against the presence of unexpected sequence variants (unexpected polymorphisms in probe/primer sites) in our diverse *B. anthracis* strain panel. If present, unexpected sequence variation among diverse strains can dramatically impact assay performance that ultimately results in less sensitivity and higher thresholds of detection.

Examining SNPs with various specificity for the Ames strain also provides insights on the phylogeny of the Ames evolutionary lineage. Based upon our SNP analyses, the nearest relatives to the Ames strain include two isolates cultivated from cattle in Texas in 2001 (Alex Hoffmaster, CDC) and a previously reported isolate cultivated from a goat in Texas in 1997 (17). These findings are supported by MLVA data and are consistent with the knowledge that the Ames strain was originally isolated from cattle in Jim Hogg County, Texas, in 1981 (Martin Hugh-Jones, personal communication). The isolation of two of these Ames relatives in 1997 and 2001 suggests the Ames lineage is endemic in this region. It is interesting that the remaining genetic near relatives to the Ames strain were isolated from human outbreaks in China (1957 and 1962), suggesting that the Chinese and Texas isolates share a common ancestral lineage. International trade of spore-contaminated products (bone meal, wool, etc.) can facilitate such long-distance dispersal of *B. anthracis* and may explain this geographically distant relationship.

Evaluating SNP data in relation to higher resolution genotyping data (MLVA genotypes) provides for fine-scale phylogenetic hypotheses that can be used to guide strain-specific SNP discovery. The applicability of this approach is not limited to *B. anthracis*, and we expect that this strategy will be used to identify strain-specific SNPs in other pathogens and biowarfare agents. For example, high-resolution variable-number tandem repeat markers have been identified in *Yersinia pestis* (9) and whole-genome sequence comparisons of this pathogen are currently underway. As whole-genome sequences and high-resolution molecular typing methods become available for biological threat agents, this strategy could assist efforts in the forensic and biodefense arena to develop assays coupling low-level detection with highly specific strain identification.

In conclusion, our strategy permits the identification of

SNPs that are diagnostic for narrowly defined *B. anthracis* genetic lineages. The Ames strain-specific SNP assays are based on six selected genomic differences between Ames and its genetic relatives and work at the single-molecule limit. Hence, we are approaching the theoretical limits for both specificity (single nucleotide) and sensitivity (single molecule) for strain subtyping. The theory of strain identification must be coupled with the practical limitation of laboratory analysis, necessitating extensive validation and quality control (7). With this caveat, the use of strain-specific SNPs for low-level, high-throughput genotyping may prove to be a valuable tool to assist in real-world forensic and public health studies.

#### ACKNOWLEDGMENTS

This work was supported by funding from the United States Department of Homeland Security and by the Federal Bureau of Investigation.

We thank Alex Hoffmaster and Bijang Wang for providing Ames-related strains and Marty Johnson and Lilly Wong (Applied Biosystems) for their technical assistance in assay design.

#### REFERENCES

1. Easterday, W. R., M. N. Van Ert, T. S. Simonson, D. M. Wagner, L. J. Kenefic, C. J. Allender, and P. Keim. 2005. Use of single nucleotide polymorphisms in the *plcR* gene for specific identification of *Bacillus anthracis*. *J. Clin. Microbiol.* **43**:1995–1997.
2. Easterday, W. R., M. N. Van Ert, S. Zanecki, and P. Keim. 2005. Specific detection of *Bacillus anthracis* using a TaqMan mismatch amplification mutation assay. *BioTechniques* **38**:731–735.
3. Hoffmaster, A. R., C. C. Fitzgerald, E. Ribot, L. W. Mayer, and T. Popovic. 2002. Molecular subtyping of *Bacillus anthracis* and the 2001 bioterrorism-associated anthrax outbreak, United States. *Emerg. Infect. Dis.* **8**:1111–1116.
4. Hurtle, W., E. Bode, D. A. Kulesh, R. S. Kaplan, J. Garrison, D. Bridge, M. House, M. S. Frye, B. Loveless, and D. Norwood. 2004. Detection of the *Bacillus anthracis gyrA* gene by using a minor groove binder probe. *J. Clin. Microbiol.* **42**:179–185.
5. Keim, P., A. Kalif, J. Schupp, K. Hill, S. E. Travis, K. Richmond, D. M. Adair, and M. Hugh-Jones. 1997. Molecular evolution and diversity in *Bacillus anthracis* as detected by amplified fragment length polymorphism markers. *J. Bacteriol.* **179**:818–824.
6. Keim, P., L. B. Price, A. M. Klevytska, K. L. Smith, J. M. Schupp, R. Okinaka, P. J. Jackson, and M. E. Hugh-Jones. 2000. Multiple-locus variable-number tandem repeat analysis reveals genetic relationships within *Bacillus anthracis*. *J. Bacteriol.* **182**:2928–2936.
7. Keim, P. 2003. Microbial forensics: a scientific assessment. American Academy of Microbiology, Washington, D.C.
8. Keim, P., M. N. Van Ert, T. Pearson, A. J. Vogler, L. Y. Huynh, and D. M. Wagner. 2004. Anthrax molecular epidemiology and forensics: using the appropriate marker for different evolutionary scales. *Infect. Genet. Evol.* **4**:205–213.
9. Klevytska, A. M., L. B. Price, J. M. Schupp, P. L. Worsham, J. Wong, and P. Keim. 2001. Identification and characterization of variable-number tandem repeats in the *Yersinia pestis* genome. *J. Clin. Microbiol.* **39**:3179–3185.
10. Lowe, C. A., M. A. Diggle, and S. C. Clarke. 2004. A single nucleotide polymorphism identification assay for the genotypic characterisation of *Neisseria meningitidis* using MALDI-TOF mass spectrometry. *Br. J. Biomed. Sci.* **61**:8–10.
11. Moorhead, S. M., G. A. Dykes, and R. T. Cursons. 2003. An SNP-based PCR assay to differentiate between *Listeria monocytogenes* lineages derived from phylogenetic analysis of the *sigB* gene. *J. Microbiol. Methods* **55**:425–432.
12. Morin, P. A., R. Saiz, and A. Monjaneb. 1999. High-throughput single nucleotide polymorphism genotyping by fluorescent 5' exonuclease activity. *BioTechniques* **27**:538–552.
13. Okinaka, R. T., K. Cloud, O. Hampton, A. R. Hoffmaster, K. K. Hill, P. Keim, T. M. Koehler, G. Lamke, S. Kumano, J. Mahillon, D. Manter, Y. Martinez, D. Ricke, R. Svensson, and P. J. Jackson. 1999. Sequence and organization of pXO1, the large *Bacillus anthracis* plasmid harboring the anthrax toxin genes. *J. Bacteriol.* **181**:6509–6515.
14. Okinaka, R., K. Cloud, O. Hampton, A. Hoffmaster, K. Hill, P. Keim, T. Koehler, G. Lamke, S. Kumano, D. Manter, Y. Martinez, D. Ricke, R. Svensson, and P. Jackson. 1999. Sequence, assembly and analysis of pXO1 and pXO2. *J. Appl. Microbiol.* **87**:261–262.
15. Pearson, T., J. D. Busch, J. Ravel, T. D. Read, S. D. Rhoton, J. M. U'Ren, T. S. Simonson, S. M. Kachur, R. R. Leadem, M. L. Cardon, M. N. Van Ert,

- L. Y. Huynh, C. M. Fraser, and P. Keim. 2004. Phylogenetic discovery bias in *Bacillus anthracis* using single-nucleotide polymorphisms from whole-genome sequencing. *Proc. Natl. Acad. Sci. USA* **101**:13536–13541.
16. Price, L. B., M. E. Hugh-Jones, P. J. Jackson, and P. Keim. 1999. Genetic diversity in the protective antigen gene of *Bacillus anthracis*. *J. Bacteriol.* **181**:2358–2362.
17. Read, T. D., S. L. Salzberg, M. Pop, M. Shumway, L. Umayam, L. Jiang, E. Holtzapple, J. D. Busch, K. L. Smith, J. M. Schupp, D. Solomon, P. Keim, and C. M. Fraser. 2002. Comparative genome sequencing for discovery of novel polymorphisms in *Bacillus anthracis*. *Science* **296**:2028–2033.
18. Stephens, A. J., F. Huygens, J. Inman-Bamber, E. P. Price, G. R. Nimmo, J. Schooneveld, W. Munckhof, and P. M. Gifford. 2006. Methicillin-resistant *Staphylococcus aureus* genotyping using a small number of polymorphisms. *J. Med. Microbiol.* **55**:43–51.
19. U'ren, J. M., M. N. Van Ert, J. M. Schupp, W. R. Easterday, T. S. Simonson, R. T. Okinaka, T. Pearson, and P. Keim. 2005. Use of a real-time PCR TaqMan assay for rapid identification and differentiation of *Burkholderia pseudomallei* and *Burkholderia mallei*. *J. Clin. Microbiol.* **43**:5771–5774.
20. Van Ert, M. N., S. A. Hofstadler, Y. Jiang, J. D. Busch, D. M. Wagner, J. J. Drader, D. J. Ecker, J. C. Hannis, L. Y. Huynh, J. M. Schupp, T. S. Simonson, and P. Keim. 2004. Mass spectrometry provides accurate characterization of two genetic marker types in *Bacillus anthracis*. *BioTechniques* **37**:642–651.
21. Vogler, A. J., J. D. Busch, S. Percy-Fine, C. Tipton-Hunton, K. L. Smith, and P. Keim. 2002. Molecular analysis of rifampin resistance in *Bacillus anthracis* and *Bacillus cereus*. *Antimicrob. Agents Chemother.* **46**:511–513.
22. Wahab, T., S. Hjalmarsson, R. Wollin, and L. Engstrand. 2005. Pyrosequencing *Bacillus anthracis*. *Emerg. Infect. Dis.* **11**:1527–1531.
23. Walters, J. J., W. Muhammad, K. F. Fox, A. Fox, D. Xie, K. E. Creek, and L. Pirisi. 2001. Genotyping single nucleotide polymorphisms using intact polymerase chain reaction products by electrospray quadrupole mass spectrometry. *Rapid Commun. Mass Spectrom.* **15**:1752–1759.