

addition to written reports is required to gain a comprehensive view of performance. In the United States this type of monitoring by a physician is recommended for all cases of cardiac arrest treated by emergency medical technicians using advanced skills.<sup>2</sup> No such recommendations exist in the United Kingdom. Monitoring of electrocardiograms and cassette tapes is time consuming and requires additional equipment.<sup>3</sup> Lack of resources or medical support, in addition to the lack of official directives, may result in ambulance services failing to monitor standards, and the full potential of advanced trained ambulance staff will not be achieved.

We thank Marks and Spencer for funding this research, Miss S Smith for preparing the manuscript, and Dr D D'Auria, Mr K G Smith, and the staff of the London Ambulance Service for allowing access to records and equipment.

- 1 Ambulance Staff Training Committee. *Extended training in ambulance aid*. Bristol: NHS Training Authority, 1987.
- 2 Cummins RO. EMT defibrillation: national guide-lines for implementation. *Am J Emerg Med* 1987;5:254-7.
- 3 Cummins RO, Austin D, Graves JR, et al. An innovative approach to medical control: semi-automatic defibrillators with solid-state memory modules for recording cardiac arrest events. *Ann Emerg Med* 1988;17:818-24.

(Accepted 23 December 1988)

## Screening for colonic cancer in patients with Barrett's oesophagus

D A F Robertson, R C S Ayres, C L Smith

Department of Medicine II,  
Southampton General  
Hospital, Southampton  
SO9 4XY

D A F Robertson, MD,  
lecturer in medicine  
R C S Ayres, MRCP, medical  
registrar  
C L Smith, MD, senior lecturer  
in medicine

Correspondence to: Dr  
Robertson.

*Br Med J* 1989;298:650

In 1985 Sontag *et al* reported an association between Barrett's oesophagus and colonic neoplasia, but their study did not include a control population.<sup>1</sup> In another uncontrolled series no colorectal cancers were seen in 36 patients with Barrett's oesophagus, although 33% of these patients had colorectal adenomas.<sup>2</sup> We performed colonoscopy prospectively in our patients with Barrett's oesophagus, none of whom had symptoms of colonic neoplasia, and compared the results with those obtained in a control group of patients thought to have the irritable bowel syndrome.<sup>3</sup>

### Subjects, methods, and results

Barrett's oesophagus was defined as columnar epithelium occurring in a tubular oesophagus at least 3 cm above the manometrically defined lower oesophageal sphincter.<sup>4</sup> We studied 32 patients (21 men and 11 women), whose mean age was 61 (range 29-81). None had a history of colorectal neoplasia. Two control patients thought on clinical grounds to have the irritable bowel syndrome, and matched for age and sex, were chosen for each patient with Barrett's oesophagus; their mean age was 55 (range 28-81). No additional screening tests, such as testing for faecal occult blood, were done to exclude colonic neoplasia.

Eight patients with Barrett's oesophagus (25%) had 12 colorectal tumours, three of which were malignant and nine benign (table). All were successfully treated, two by local resection, one by subtotal colectomy, and

five by endoscopic polypectomy. Three of the 64 controls (5%) had single adenomatous polyps, which were removed endoscopically. The relative risk of colonic tumour was 5.3 in the patients with Barrett's oesophagus (95% confidence interval 1.94 to 8.69).

### Comment

We found a high prevalence of colorectal tumours in our patients with Barrett's oesophagus, in keeping with the findings of Sontag *et al*.<sup>1</sup> Our study was entirely prospective, and the association between colonic neoplasia and Barrett's oesophagus seemed to be genuine and not due to some hidden bias; this may not have been the case in the uncontrolled study by Sontag *et al*, in which seven of 10 carcinomas were identified retrospectively.

Selecting an appropriate control group was difficult, and we considered that colonoscopic examination of asymptomatic people was unjustified because of the risks and discomfort entailed. We thus selected patients who had been referred for investigation and in whom the irritable bowel syndrome had been diagnosed on clinical grounds. The controls were on average six years younger than the study group, and this could have affected the prevalence of colonic tumours: the prevalence of colonic neoplasia related to age in asymptomatic subjects is unknown. The difference in age between the two groups, however, is unlikely to have accounted for all of the observed differences in prevalence.

The aetiology of both Barrett's oesophagus and carcinoma of the colon remains unknown. The symptoms of dyspepsia or chest pain are unlikely to have been caused by the colonic carcinoma or adenoma, and in each case the finding of a colonic lesion was incidental. An inherited or acquired common aetiological agent, such as a dietary factor or a tendency to neoplastic or metaplastic change throughout the gastrointestinal tract, may have been the cause. There was no evidence of dysplasia in biopsy specimens or resected colonic epithelium not affected by adenomatous or carcinomatous change.

It has been recommended that frequent oesophagoscopy should be undertaken in patients with Barrett's oesophagus to detect early oesophageal carcinoma or dysplasia. None of our patients had high grade dysplasia or oesophageal carcinoma but eight had colonic neoplasia. Colonoscopy may therefore be a more worthwhile screening procedure for cancer than oesophagoscopy in Barrett's oesophagus.

1 Sontag SJ, Schnell TG, Cheseb G, et al. Barrett's oesophagus and colonic tumours. *Lancet* 1985;ii:946-9.

2 Tripp MR, Sampliner RE, Kogan FJ, Morgan TR. Colorectal neoplasia and Barrett's oesophagus. *Am J Gastroenterol* 1986;81:1063-4.

3 Manning AP, Thompson WG, Heaton KW, Morris AF. Towards positive diagnosis of the irritable bowel. *Br Med J* 1978;ii:653-4.

4 Spechler SJ, Goyal RK. Barrett's oesophagus. *N Engl J Med* 1986;315:362-71.

(Accepted 23 December 1988)

Clinical details of patients and controls with colorectal tumours

Case No	Age (years) and sex	Site	Histological findings	Treatment
<i>Patients</i>				
1	79 M	Sigmoid colon Descending colon	Tubulovillous adenoma with severe dysplasia at both sites	Endoscopic polypectomy
2	66 M		Tubulovillous adenoma with severe dysplasia	Endoscopic polypectomy
3	53 M	Sigmoid colon	Tubular adenoma and mild dysplasia at two sites	Endoscopic polypectomy
4	72 F	Hepatic flexure	Adenomatous polyp	Endoscopic polypectomy
5	74 M	Sigmoid colon	Adenomatous polyp	Endoscopic polypectomy
6	52 M	Transverse colon	Tubular adenoma with mild dysplasia	Right hemicolectomy
7	44M	Hepatic flexure	Dukes's A carcinoma	Subtotal colectomy
		Splenic flexure	Dukes's B carcinoma	
8	71M	Sigmoid colon	Tubular adenoma	Sigmoid colectomy
		Sigmoid colon	Dukes's A carcinoma	
<i>Controls</i>				
9	52 M	Sigmoid colon	Adenomatous polyp	Endoscopic polypectomy
10	64 F	Descending colon	Adenomatous polyp	Endoscopic polypectomy
11	66 M	Rectum	Adenomatous polyp	Endoscopic polypectomy