

## Comparative Activities of Norfloxacin and Fifteen Other Antipseudomonal Agents Against Gentamicin-Susceptible and -Resistant *Pseudomonas aeruginosa* Strains

K. R. FORWARD, G. K. M. HARDING, G. J. GRAY, B. A. URIAS, AND A. R. RONALD

*Departments of Medical Microbiology and Medicine, University of Manitoba, and The Health Sciences Center, Winnipeg, Manitoba, Canada*

Volume 24, no. 4, p. 603, column 1, line 4: “. . . MIC of 64  $\mu\text{g/ml}$  . . .” should read “. . . MIC of 16  $\mu\text{g/ml}$ .”

---

## Synergism of Imipenem and Amikacin in Combination with Other Antibiotics Against *Nocardia asteroides*

MYLES E. GOMBERT AND TARYN M. AULICINO

*Division of Infectious Diseases, Department of Medicine, State University of New York Downstate Medical Center, Brooklyn, New York 11203*

Volume 24, no. 5, p. 811, column 1, lines 4–5: “The combinations of imipenem-cefotaxime and amikacin-cefotaxime . . .” should read “The combinations of imipenem-amikacin and amikacin-cefotaxime. . . .”

Page 811, column 1, lines 7–8: “The fractional inhibitory concentration was  $\leq 0.05$  . . .” should read “The fractional inhibitory concentration was  $\leq 0.5$ . . . .”