



Published in final edited form as:

Biochem Biophys Res Commun. 2007 March 2; 354(1): 127–132.

SEPS1 protects RAW264.7 cells from pharmacological ER stress agent-induced apoptosis

Kee-Hong Kim^{a,b,*}, Yuan Gao^c, Ken Walder^c, Greg R. Collier^c, Joseph Skelton^d, and Ahmed H. Kissebah^{a,b}

^aDivision of Endocrinology, Metabolism and Clinical Nutrition, the Department of Medicine, Medical College of Wisconsin, Milwaukee, WI 53226, USA

^bHuman and Molecular Genetics Center, Medical College of Wisconsin, Milwaukee, WI 53226, USA

^cMetabolic Research Unit, School of Health Sciences, Deakin University, Geelong, Victoria, Australia

^dDepartment of Pediatrics, Children's Research Institute, Medical College of Wisconsin, Milwaukee, WI, 53226, USA

Abstract

Selenoprotein S (SEPS1) is a novel endoplasmic reticulum (ER) resident protein and it is known to play an important role in production of inflammatory cytokines. Here, we show evidence that SEPS1 is stimulated by pharmacological ER stress agents in RAW264.7 macrophages as well as other cell types. Overexpression studies reveal a protective action of SEPS1 in macrophages against ER stress-induced cytotoxicity and apoptosis, resulting in promoting cell survival during ER stress. The protective action of SEPS1 is largely dependent on ER stress-mediated cell death signal with less effect on non-ER stress component cell death signals. Conversely, suppression of *SEPS1* in macrophages results in sensitization of cells to ER stress-induced cell death. These findings suggest that SEPS1 could be a new ER stress-dependent survival factor that protects macrophage against ER stress-induced cellular dysfunction.

Keywords

selenoprotein S; tanis; VIMP; endoplasmic reticulum stress; macrophages; apoptosis; tunicamycin; thapsigargin

Introduction

SEPS1 (Sels/Tanis/VIMP) has been previously identified as a putative ER stress response protein that is likely to be associated with an inflammatory response [1-4]. A genetic variation in the human *SEPS1* promoter region that is strongly associated with substantial increase in circulating levels of pro-inflammatory cytokine is located in the center of a putative ER stress-response element (ERSE) [5], suggesting a possibility of ER stress-dependent regulation of *SEPS1* transcription.

* correspondence author: Fax:+1 414 456 6516. E-mail address: kkim@mcw.edu (K. Kim)

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The ER is the major site for protein folding and maturation, cellular response to stress, and maintaining Ca^{2+} homeostasis [6]. Accumulation of misfolded proteins and alteration of Ca^{2+} homeostasis in the ER generate ER stress that triggers various cellular dysfunctions including apoptosis and inflammation [7-11]. ER stress is primarily sensed by three ER-bound proteins: PKR-like ER-associated kinase (PERK), a kinase and endonuclease inositol requiring enzyme1 (IRE1), and a basic leucine-zipper transcription factor activation of transcription factor 6 (ATF6) [8-12]. Activation of these proteins and their associated signaling pathways triggers attenuation of general protein synthesis and an increase in transcription of genes that are essential for molecular chaperones, protein folding and protein degradation in order to adapt to temporal ER stress [8,12]. Under the condition of prolonged and/or severe ER stress, the cell activates intracellular pathways that lead to programmed cell death [8,13,14] via several pathways, including caspase-12 and PERK-mediated activation of a transcriptional factor CHOP/GADD153 [8,13-18]. Elevated ER stress and its associated apoptosis are evidenced in many cell types including macrophages, pancreatic β -cells, neurons and endothelial cells with implication for various human diseases, including atherosclerosis, Parkinson's, Alzheimer's, and prion protein disease [15-17]. However, the protective mechanisms against ER stress-induced apoptosis have not yet been fully understood.

In this study, we have demonstrated an ER stress-dependent SEPS1 expression in macrophages as well as in various cell types. The overexpression and suppression studies of *SEPS1* suggest a survival role of SEPS1 in macrophages during ER stress and its potential role in controlling ER stress-associated signaling pathway.

Materials and Methods

Materials and Cell Culture

Tunicamycin, thapsigargin, dithiothreitol (DTT), cycloheximide and staurosporine were purchased from Fisher Scientific. Homocysteine was purchased from Sigma. Anti-Fas was purchased from eBioscience. RAW 264.7, HepG2 and HEK293 cells were obtained from American Tissue Culture Collection (ATCC) and all cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% (v/v) heat-inactivated fetal calf serum (FCS) and antibiotics (penicillin, 75 $\mu\text{g}/\text{ml}$; streptomycin, 50 $\mu\text{g}/\text{ml}$) in 5% CO_2 at 37°C.

Construction of Plasmid and Transient Transfection

A N-terminal flag-tagged open reading frame plus 3'-untranslated region of human *SEPS1* mRNA was amplified by PCR with the 5' primer, 5'-GCCACCATGGATTACAAGGATGACGACGATAAGGAACGCCAAGA-3' and 3' primer, 5'-GAAGTCCATAAATCTCCTTG-3'. The resultant PCR product was ligated into pTARGET vector (Promega) and then subcloned into pcDNA3.1 vector (Invitrogen). The pcDNA3.1 expression vector was transiently transfected into RAW 264.7 cells using Lipofectamine 2000 (Invitrogen) by following the manufacturer's instruction.

siRNA transfection

SEPS1 and scrambled siRNAs were synthesized *in vitro* using a kit from Ambion. Primer sequences used in this study and the condition of transient transfection of siRNA into RAW264.7 cells were described previously [5].

Western blot Analyses

Cells were lysed in a buffer containing 20 mM Tris-HCl, pH 7.5, 150 mM NaCl, 1% Nonidet P-40, 5 mM EDTA, and 1 mM phenylmethylsulfonyl fluoride for 10 min on ice. The protein

content was determined by Bradford assay (Bio-Rad). Ten μg of protein were subjected to SDS-polyacrylamide gel electrophoresis, electroblotted onto nitrocellulose membrane, and immunodetected using primary antibodies and goat anti-mouse IgG-horseradish peroxidase conjugate (Santa Cruz Biotechnology) by using an enhanced chemiluminescence detection kit (Pierce). The primary antibodies used in this study are as follow: anti-SEPS1 antibody [4], anti-GRP78 and anti-GAPDH (Santa Cruz Biotechnology), anti-CHOP, anti-phospho-PERK and anti-phospho-eIF2 α (Cell Signaling Technology).

DNA Fragmentation Assay

RAW 264.7 cells transfected with control or *SEPS1* expression vector were incubated with 10 $\mu\text{g}/\text{ml}$ tunicamycin or 10 μM thapsigargin for 24 h in the absence of serum. Cells were scraped, pelleted, washed in ice-cold PBS, and gently resuspended in lysis buffer (1 \times TE, 0.5 % SDS, 20 $\mu\text{g}/\text{ml}$ RNase) followed by incubation at 37°C for 1 h. After incubation, the lysates were treated with proteinase K (100 $\mu\text{g}/\text{ml}$) at 50°C for 1 h and DNA ladder formation was visualized by agarose gel electrophoresis.

Cell Cytotoxicity, Apoptosis and Viability Assays

RAW 264.7 cells transfected with control or *SEPS1* expression vector were challenged with indicated concentration of tunicamycin or thapsigargin for 24 h in the absence of serum. After 24 h, lactate dehydrogenase (LDH) activity in the media was determined using a cytotoxicity assay kit (Biovision). Caspase 3 activity in cell lysates was determined using a colorimetric Caspase-3 Assay System (Biovision). Cell viability was determined using either CellTiter Aqueous Assay (Promega) or the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) method [19].

Statistical Analysis

Student *t* test was used with $P < 0.05$ considered significant.

Results and Discussion

Tunicamycin- and thapsigargin-induced SEPS1 expression

We previously identified that human *SEPS1* promoter contains a putative ER stress response element (ERSE) [20]. This result suggests that SEPS1 expression and function can be regulated by ER stress. We first examined the effect of pharmacological ER stress agents on SEPS1 expression in macrophages. RAW264.7 cells were challenged with various concentration of pharmacological ER stress agents tunicamycin (Tm), an inhibitor of N-glycosylation, and thapsigargin (Tg), an inhibitor of ER Ca²⁺ ATPase activity for 24h [21-23]. RAW264.7 cells treated with Tm or Tg resulted in elevated SEPS1 protein levels in a dose-dependent manner with maximum induction at 10 $\mu\text{g}/\text{ml}$ or 5 μM , respectively (Fig. 1A). Tm- or Tg-induced ER stress in RAW264.7 cells was evidenced by a dose-dependent phosphorylation of ER stress response proteins PERK and eukaryotic initiation factor 2 α (eIF2 α) (Fig. 1A). SEPS1 expression in RAW264.7 cells was also dependent on the time of treatment with 10 $\mu\text{g}/\text{ml}$ Tm or 5 μM Tg with maximum induction after 8 h of treatment (Fig. 1B). Time-dependent induction of ER stress in Tm or Tg treated RAW264.7 cells was evidenced by induction of the two ER stress-response proteins GRP78 and CHOP (Fig. 1B). *SEPS1* mRNA levels in RAW264.7 cells were also observed to be markedly induced by Tm or Tg treatment (data not shown). ER stress-dependent induction of SEPS1 in RAW264.7 cells was further confirmed by challenging cells with other ER stress agents, including dithiothreitol (DTT) and homocysteine (Hcy) for 24 h. Both 2 mM DTT and 0.1 mM Hcy also induced SEPS1 expression in RAW264.7 cells (Fig. 1C). DTT- and Hcy-induced ER stress in RAW264.7 cells was evidenced by elevated level of GRP78 (Fig. 1C). These results clearly show that SEPS1 expression is up-regulated by ER

stress agents in macrophages. Since prolonged exposure of cells to ER stress is known to trigger apoptosis [8,13,14], we next questioned whether non-ER stress component apoptotic signals are able to induce SEPS1 expression. RAW264.7 cells were challenged with non-ER stress component apoptotic agents, including staurosporine (STS), a broad kinase inhibitor and potent apoptosis inducer, and anti-Fas, a mitochondrial-targeted apoptotic inducer, for 18 h. As expected, treatment of RAW264.7 cells with 5 μ M Tg resulted in an induction of SEPS1 and GRP78 (Fig. 1D). However, we found that non-ER stress component apoptotic signals tested in this study had little or no effect on SEPS1 expression in RAW264.7 cells. Treatment of RAW264.7 cells with STS resulted in slightly elevated levels of SEPS1 and GRP78 compared with those induced by Tg treatment. Moreover, anti-Fas treatment did not show any change in both SEPS1 and GRP78 expression in RAW264.7 cells (Fig. 1D). Tg-, STS- and Fas-induced apoptosis was evidenced by the generation of cleaved caspase 3 (Fig. 1D). Although the underlying basis for ER stress agent dependent induction of SEPS1 is not known at this time, this result indicates that SEPS1 expression is largely dependent upon ER stress with less significant association with STS- and Fas-induced general apoptotic signals in macrophages. To further support this notion, it will be of interest to examine whether intrinsic ER stress signal triggered by accumulated misfolded proteins [7-12] or free cholesterol [24] in the ER could be able to induce SEPS1 expression in macrophages.

Next, we tested whether induction of SEPS1 by ER stress agents can also be seen in other cell types. We employed MIN6 mouse pancreatic β -cells, HepG2 hepatoma cells and HEK 293 cells to test Tm- and Tg-dependent induction of SEPS1. Figure 1E shows the level of SEPS1 in response to a 24h treatment with concentrations of Tm or Tg through 10 μ g/ml or 5 μ M, respectively, revealing a dose-dependent effect in tested cell types. This result implies that induction of SEPS1 could be a common indicator of cells under the condition of ER stress.

SEPS1 protects macrophages from Tm- and Tg-induced cytotoxicity and apoptosis

To understand the physiological function of SEPS1 we determined the consequence of SEPS1 overexpression on pharmacological ER stress agent-induced cytotoxicity and apoptosis in RAW 264.7 cells. Cells were transiently transfected with either control or flag-SEPS1 expression vector, followed by challenging to serum-free medium containing various concentration of Tm (0-10 μ g/ml) or Tg (0-5 μ M) for 24 h. Figure 2A confirms transient expression of flag-tagged SEPS1 fusion protein in RAW 264.7 cells. We also observed that overexpression of SEPS1 in RAW264.7 cells is associated with reduced levels of Tm- and Tg-mediated GRP78 expression compared with those in control vector transfected cells (Fig. 2A). We then assessed the Tm- and Tg-induced cytotoxicity of these cells by measuring the activity of lactate dehydrogenase (LDH) in the medium released from the cells. Consistent with previous reports on the cytotoxic action of Tm and Tg [25, 26], treatment of control vector transfected RAW264.7 cells with Tm and Tg resulted in a dose-dependent increase in LDH activity in the cell culture media (Fig. 2B). However, Figure 2B illustrates that cells transfected with flag-SEPS1 expression vector were resistant to both Tm- and Tg-induced cytotoxicity indicating a cytoprotective action of SEPS1 during ER stress. Moreover, the basal cytotoxicity of cells transfected with flag-SEPS1 expression vector also was lower than that of cells transfected with control vector. This result implies that SEPS1 could also protect RAW264.7 cells against cytotoxicity caused by serum depletion.

Both pharmacological ER stress agents Tm and Tg have been known to promote apoptosis via activating ER stress signaling pathways [23]. Thus, we next further examined the consequence of SEPS1 overexpression on ER stress-induced apoptosis in RAW264.7 cells. We first determined the effect of SEPS1 overexpression on Tm- or Tg-induced DNA fragmentation in RAW264.7 cells. As described in "Materials and Methods", the genomic DNA was isolated from cells transfected with control or flag-SEPS1 expression vector followed by challenging

with 10 µg/ml Tm or 5 µM Tg for 24 h. The fragmented DNA was then visualized in an ethidium bromide-stained agarose gel. Consistent with previous report [27], both treatment of RAW264.7 cells with Tm- and Tg resulted in generation of fragmented DNA with size of 1 kb or less, an indicative of apoptosis (Fig. 2C). However, DNA fragmentation was barely detectable in Tm- or Tg-treated SEPS1 overexpressing cells (Fig. 2C). Effect of SEPS1 overexpression on Tm- or Tg-induced apoptosis was further determined by measuring caspase-3 activity in cell lysate isolated from RAW264.7 cells treated with 10 µg/ml Tm or 5 µM Tg for 24 h. As expected, the levels of caspase-3 activity in control cells challenged with Tm or Tg were approximately 2- or 4-fold greater, respectively, when compared with non-treated control cells (Fig. 2D). However, caspase-3 activity in RAW264.7 cells transfected with flag-SEPS1 expression vector was not influenced by Tm and Tg treatment (Fig. 2D). We also assessed the consequence of SEPS1 overexpression on RAW264.7 cell death induced by Tg treatment. Cells transfected with control vector exhibited a dose-dependent decrease in cell viability after 24 h of treatment with increasing concentration of Tg (0-10 µM) resulting in approximately 30% of cell viability at 10 µM Tg. However, RAW264.7 cells transfected with flag-SEPS1 expression vector were resistant to Tg-induced cell death with approximately 60% of cell viability at 10 µM Tg (Fig. 2E). Taken together, the results presented above suggest that SEPS1 protects macrophages from pharmacological ER stress agent-induced apoptosis thereby promoting cell survival during ER stress.

Protective action of SEPS1 against non-ER stress-induced macrophage death

We next attempted to test whether SEPS1 overexpression also improves RAW264.7 cell viability against non-ER stress-induced cell death signals. RAW264.7 cells transfected with control vector or flag-SEPS1 expression vector were challenged with STS (1 µM) or Fas-specific antibody (0.5 µg/ml) plus cycloheximide (CHX) (10 µg/ml), as well as an ER stress agent Tm (10 µg/ml) for 18 hr. As expected, RAW264.7 cells transfected with flag-SEPS1 expression vector exhibited improved cell viability compared with cells transfected with control vector against Tm-induced death (Fig. 3). SEPS1 overexpression, however, did not show significant protective effect against STS- and Fas-induced cell death (Fig. 3). Indeed, cells transfected with flag-SEPS1 expression vector exhibited increased sensitivity to STS-induced cell death. This result suggests that SEPS1 protects macrophages largely against pharmacological ER stress agent-induced apoptosis with less effect on non-ER stress-induced cell death signals.

Role of endogenous SEPS1 in ER stress-induced cell death

We next determined the role of endogenous *SEPS1* mRNA in pharmacological ER stress agent-induced cell death in macrophages. We tested the consequence of siRNA-mediated *SEPS1* suppression on Tm-induced RAW264.7 cell death. After 18h of *SEPS1* siRNA transfection into RAW264.7 cells, *SEPS1* mRNA level was suppressed approximately by 60 % (Fig. 4A). We then challenged these cells with various concentration of Tm (0-2 µg/ml) for 18 h and cell viability was determined using MTT assay. Control scrambled siRNA transfected RAW264.7 cells showed a dose-dependent cell death upon increasing concentration of Tm with approximately 60% of cell viability at 1 µg/ml Tm (Fig. 4B). However, siRNA-mediated suppression of endogenous *SEPS1* mRNA resulted in sensitization of cells to Tm-induced cell death with approximately 40 % of cell viability at 1 µg/ml Tm (Fig. 4B). This result suggests that endogenous *SEPS1* is required for macrophage survival against Tm-induced cell death. The remaining viability of *SEPS1* siRNA transfected cells exposed to Tm could be due to an incomplete *SEPS1* mRNA suppression or the presence of SEPS1-independent survival mechanisms. Nevertheless, this result together with the data shown in Figure 2 clearly demonstrate that SEPS1 plays an important role in promoting macrophage survival during the condition of pharmacological agent-induced ER stress.

Recently, SEPS1/VIMP has been identified as a novel ER membrane protein interacting with VCP/p97 and Derlin-1, key protein components of the retrotranslocation/ER-associated degradation (ERAD) machinery [1-3]. Given that retrotranslocation/ERAD is a critical step in counteracting ER stress and maintaining the ER homeostasis, the protective action of SEPS1 against ER stress-induced cell death presented in this study could be, at least in part, a potential mechanism by which retrotranslocation/ERAD alleviates ER stress-associated dysfunction. It will be of great interest to study the role of SEPS1 in VCP/p97-mediated retrotranslocation/ERAD.

In summary, our results provide evidence that SEPS1 is a novel ER stress-induced protein that attenuates pharmacological ER stress-induced cytotoxicity and apoptosis, resulting in promotion of cell survival. These results suggest that modulation of SEPS1 expression could be a novel mechanism to control ER stress-induced cell apoptosis.


Acknowledgement

We thank Drs. Donghee Kim (Rosalind Franklin University of Medicine and Science/The Chicago Medical School) and Susumu Seino (Kobe University, Japan) for the kind gift of MIN6 cell line. We also thank Mitja Pavlovic for assisting with the experiments.


References

- [1]. Ye Y, Shibata Y, Yun C, Ron D, Rapoport TA. A membrane protein complex mediates retrotranslocation from the ER lumen into the cytosol. *Nature* 2004;429:841–847. [PubMed: 15215856]
- [2]. Ye Y, Shibata Y, Kikkert M, van Voorden S, Wiertz E, Rapoport TA. Inaugural Article: Recruitment of the p97 ATPase and ubiquitin ligases to the site of retrotranslocation at the endoplasmic reticulum membrane. *Proc Natl Acad Sci U S A* 2005;102:14132–14138. [PubMed: 16186510]
- [3]. Lilley BN, Ploegh HL. Multiprotein complexes that link dislocation, ubiquitination, and extraction of misfolded proteins from the endoplasmic reticulum membrane. *Proc Natl Acad Sci U S A* 2005;102:14296–14301. [PubMed: 16186509]
- [4]. Walder K, Kantham L, McMillan JS, Trevaskis J, Kerr L, De Silva A, Sunderland T, Godde N, Gao Y, Bishara N, Windmill K, Tenne-Brown J, Augert G, Zimmet PZ, Collier GR. Tanis: a link between type 2 diabetes and inflammation? *Diabetes* 2002;51:1859–1866. [PubMed: 12031974]
- [5]. Curran JE, Jowett JB, Elliott KS, Gao Y, Gluschenko K, Wang J, Azim DM, Cai G, Mahaney MC, Comuzzie AG, Dyer TD, Walder KR, Zimmet P, Maccluer JW, Collier GR, Kissebah AH, Blangero J. Genetic variation in selenoprotein S influences inflammatory response. *Nat Genet* 2005;37:1234–1241. [PubMed: 16227999]
- [6]. Lehotsky J, Kaplan P, Babusikova E, Strapkova A, Murin R. Molecular pathways of endoplasmic reticulum dysfunctions: possible cause of cell death in the nervous system. *Physiol Res* 2003;52:269–274. [PubMed: 12790757]
- [7]. Zhang K, Shen X, Wu J, Sakaki K, Saunders T, Rutkowski DT, Back SH, Kaufman RJ. Endoplasmic reticulum stress activates cleavage of CREBH to induce a systemic inflammatory response. *Cell* 2006;124:587–599. [PubMed: 16469704]
- [8]. Harding HP, Ron D. Endoplasmic reticulum stress and the development of diabetes: a review. *Diabetes* 2002;51(Suppl 3):S455–461. [PubMed: 12475790]
- [9]. Ron D. Translational control in the endoplasmic reticulum stress response. *J Clin Invest* 2002;110:1383–1388. [PubMed: 12438433]
- [10]. Kaufman RJ. Orchestrating the unfolded protein response in health and disease. *J Clin Invest* 2002;110:1389–1398. [PubMed: 12438434]
- [11]. Mori K. Tripartite management of unfolded proteins in the endoplasmic reticulum. *Cell* 2000;101:451–454. [PubMed: 10850487]
- [12]. Rutkowski DT, Kaufman RJ. A trip to the ER: coping with stress. *Trends Cell Biol* 2004;14:20–28. [PubMed: 14729177]
- [13]. Szegezdi E, Fitzgerald U, Samali A. Caspase-12 and ER-stress-mediated apoptosis: the story so far. *Ann N Y Acad Sci* 2003;1010:186–194. [PubMed: 15033718]

- [14]. Nakagawa T, Zhu H, Morishima N, Li E, Xu J, Yankner BA, Yuan J. Caspase-12 mediates endoplasmic-reticulum-specific apoptosis and cytotoxicity by amyloid-beta. *Nature* 2000;403:98–103. [PubMed: 10638761]
- [15]. Barone MV, Crozat A, Tabae A, Philipson L, Ron D. CHOP (GADD153) and its oncogenic variant, TLS-CHOP, have opposing effects on the induction of G1/S arrest. *Genes Dev* 1994;8:453–464. [PubMed: 8125258]
- [16]. Tan Y, Dourdin N, Wu C, De Veyra T, Elce JS, Greer PA. Ubiquitous calpains promote caspase-12 and JNK activation during endoplasmic reticulum stress-induced apoptosis. *J Biol Chem* 2006;281:16016–16024. [PubMed: 16597616]
- [17]. Yang Q, Kim YS, Lin Y, Lewis J, Neckers L, Liu ZG. Tumour necrosis factor receptor 1 mediates endoplasmic reticulum stress-induced activation of the MAP kinase JNK. *EMBO Rep* 2006;7:622–627. [PubMed: 16680093]
- [18]. Hu P, Han Z, Couvillon AD, Kaufman RJ, Exton JH. Autocrine tumor necrosis factor alpha links endoplasmic reticulum stress to the membrane death receptor pathway through IRE1alpha-mediated NF-kappaB activation and down-regulation of TRAF2 expression. *Mol Cell Biol* 2006;26:3071–3084. [PubMed: 16581782]
- [19]. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods* 1983;65:55–63. [PubMed: 6606682]
- [20]. Gao Y, Hannan NR, Wanyonyi S, Konstantopolous N, Pagnon J, Feng HC, Jowett JB, Kim KH, Walder K, Collier GR. Activation of the selenoprotein SEPS1 gene expression by pro-inflammatory cytokines in HepG2 cells. *Cytokine* 2006;33:246–251. [PubMed: 16574427]
- [21]. Hetz C, Bernasconi P, Fisher J, Lee AH, Bassik MC, Antonsson B, Brandt GS, Iwakoshi NN, Schinzel A, Glimcher LH, Korsmeyer SJ. Proapoptotic BAX and BAK modulate the unfolded protein response by a direct interaction with IRE1alpha. *Science* 2006;312:572–576. [PubMed: 16645094]
- [22]. Marciniak SJ, Yun CY, Oyadomari S, Novoa I, Zhang Y, Jungreis R, Nagata K, Harding HP, Ron D. CHOP induces death by promoting protein synthesis and oxidation in the stressed endoplasmic reticulum. *Genes Dev* 2004;18:3066–3077. [PubMed: 15601821]
- [23]. Shiraishi H, Okamoto H, Yoshimura A, Yoshida H. ER stress-induced apoptosis and caspase-12 activation occurs downstream of mitochondrial apoptosis involving Apaf-1. *J Cell Sci* 2006;119:3958–3966. [PubMed: 16954146]
- [24]. Feng B, Yao PM, Li Y, Devlin CM, Zhang D, Harding HP, Sweeney M, Rong JX, Kuriakose G, Fisher EA, Marks AR, Ron D, Tabas I. The endoplasmic reticulum is the site of cholesterol-induced cytotoxicity in macrophages. *Nat Cell Biol* 2003;5:781–792. [PubMed: 12907943]
- [25]. Kim AJ, Shi Y, Austin RC, Werstuck GH. Valproate protects cells from ER stress-induced lipid accumulation and apoptosis by inhibiting glycogen synthase kinase-3. *J Cell Sci* 2005;118:89–99. [PubMed: 15585578]
- [26]. Lai JS, Zhao C, Warsh JJ, Li PP. Cytoprotection by lithium and valproate varies between cell types and cellular stresses. *Eur J Pharmacol* 2006;539:18–26. [PubMed: 16678157]
- [27]. Yang L, Carlson SG, McBurney D, Horton WE Jr. Multiple signals induce endoplasmic reticulum stress in both primary and immortalized chondrocytes resulting in loss of differentiation, impaired cell growth, and apoptosis. *J Biol Chem* 2005;280:31156–31165. [PubMed: 16000304]

**Fig. 1.**

Pharmacological ER stress agents-induced SEPS1 expression. RAW264.7 cells were exposed to the indicated concentrations of tunicamycin (Tm) and tunicamycin (Tm) for 24 h (A), Tm (10 $\mu\text{g/ml}$) or Tg (5 μM) for the indicated times (B), or Tm (10 $\mu\text{g/ml}$), Tg (5 μM), DTT (2.5 mM) or homocysteine (Hcy) (0.1 mM) for 24 hr (C). (D) RAW264.7 cells were also exposed to staurosporine (STS) (1 μM), Fas-antibody (0.5 $\mu\text{g/ml}$) or Tg (5 μM), for 18 hr. (E) MIN6, HepG2 and HEK293 cells were exposed to with indicated concentration of Tm or Tg for 24 h. Cell lysates isolated from these cells were probed with anti-SEPS1, anti-GRP78, anti-CHOP, anti-phospho-PERK, anti-phospho-eIF2 α or anti-GAPDH antibodies.

**Fig. 2.**

Effect of SEPS1 overexpression on pharmacological ER stress agents-induced cytotoxicity and apoptosis. (A) RAW264.7 cells were transfected with pcDNA3.1 control vector (control) or expression vector containing flag-tagged human *SEPS1* cDNA sequence (flag-SEPS1) as described under "Materials and Methods". Cell lysates isolated from these cells were probed with anti-SEPS1, anti-GRP78 or anti-GAPDH antibodies. (B) RAW264.7 cells transfected with pcDNA3.1 control vector (control) or expression vector for flag-SEPS1 were exposed to increased concentrations of Tm or Tg for 24 h and the cellular cytotoxicity was determined by measuring lactate dehydrogenase (LDH) activity in the media released from cells. (C) The generation of fragmented DNA in expression vector transfected RAW264.7 cells treated with Tm (10 µg/ml) or Tg (5 µM) for 24 hr was visualized by agarose gel electrophoresis. The first lane of the agarose gel contained 0.5 µg 1 kb plus DNA ladder (Invitrogen). (D) Caspase-3 activity was determined in the cell lysate as described under "Material and Methods". (E) Expression vector transfected cells were challenged with indicated concentration of Tg for 24 hr. Cell viability was determined in the cell lysates using CellTiter Aqueous Assay. Data are presented as the mean \pm S.E.M. (* $P < 0.05$; ** $P < 0.01$) and the experiment was repeated twice with similar results.




Fig. 3. Effect of SEPS1 on non-ER stress-induced cell death. RAW264.7 cells transfected with expression vectors for control pcDNA3.1 (control) or flag-SEPS1 were exposed to Tm (10 $\mu\text{g/ml}$), staurosporine (STS) (1 μM), or Fas-antibody (0.5 $\mu\text{g/ml}$) plus cycloheximide (CHX) (10 $\mu\text{g/ml}$) for 18 hr. Cell viability was determined using CellTiter Aqueous Assay. Data are presented as the mean \pm S.E.M. (* $P < 0.05$; ** $P < 0.01$) and the experiment was repeated twice with similar results.




Fig. 4. Effect of *SEPS1* mRNA suppression on pharmacological ER stress-induced cell death. RAW264.7 cells transfected with scrambled or *SEPS1* siRNA (20 nM) for 24 hr were exposed to indicated concentration of Tm for 24 hr. (A) *SEPS1* mRNA levels in RAW264.7 cells transfected with scrambled or *SEPS1* siRNA were determined by qRT-PCR and normalized against cyclophilin. (B) Cell viability of these cells was determined by MTT assay. Data are presented as the mean \pm S.E.M. (* $P < 0.05$; ** $P < 0.01$) and the experiment was repeated twice with similar results.