

GENOME ANNOUNCEMENT

The Complete Genome Sequence of *Bacillus thuringiensis* Al Hakam^V

Jean F. Challacombe,^{1,3*} Michael R. Altherr,³ Gary Xie,^{1,3} Smriti S. Bhotika,^{1,3§} Nancy Brown,³ David Bruce,^{1,3} Connie S. Campbell,^{1,3} Mary L. Campbell,^{1,3} Jin Chen,^{1,3†} Olga Chertkov,^{1,3} Cathy Cleland,⁶ Mira Dimitrijevic,^{1,3} Norman A. Doggett,³ John J. Fawcett,^{1,3} Tijana Glavina,^{2,4} Lynne A. Goodwin,^{1,3} Lance D. Green,^{1,3} Cliff S. Han,^{1,3} Karen K. Hill,³ Penny Hitchcock,^{6‡} Paul J. Jackson,^{3,5} Paul Keim,⁷ Avinash Ramesh Kewalramani,^{1,3} Jon Longmire,³ Susan Lucas,^{2,5} Stephanie Malfatti,^{2,5} Diego Martinez,^{1,3} Kim McMurry,^{1,3} Linda J. Meincke,^{1,3} Monica Misra,^{1,3} Bernice L. Moseman,^{1,3} Mark Mundt,⁸ A. Christine Munk,^{1,3} Richard T. Okinaka,³ B. Parson-Quintana,^{1,3} Lee Philip Reilly,^{1,3} Paul Richardson,^{2,4} Donna L. Robinson,^{1,3} Elizabeth Saunders,^{1,3} Roxanne Tapia,^{1,3} Judith G. Tesmer,^{1,3} Nina Thayer,^{1,3} Linda S. Thompson,^{1,3} Hope Tice,^{2,4} Lawrence O. Ticknor,⁶ Patti L. Wills,^{1,3} Paul Gilna,^{1,3¶} and Thomas S. Brettin^{1,3}

DOE Joint Genome Institute, Los Alamos National Laboratory, Los Alamos, New Mexico 87545¹; DOE Joint Genome Institute, Production Genome Facility, Walnut Creek, California 94598²; Los Alamos National Laboratory, Bioscience Division, Los Alamos, New Mexico 87545³; Lawrence Berkeley National Laboratory, Berkeley, California 94720⁴; Lawrence Livermore National Laboratory, Livermore, California 94550⁵; Los Alamos National Laboratory, Decision Applications Division, Los Alamos, New Mexico 87545⁶; Northern Arizona University, Department of Biological Sciences, Flagstaff, Arizona 86011-5640⁷; and Los Alamos National Laboratory, W Division, Los Alamos, New Mexico 87545⁸

Received 13 February 2007/Accepted 19 February 2007

Bacillus thuringiensis is an insect pathogen that is widely used as a biopesticide (E. Schnepf, N. Crickmore, J. Van Rie, D. Lereclus, J. Baum, J. Feitelson, D. R. Zeigler, and D. H. Dean, *Microbiol. Mol. Biol. Rev.* 62:775–806, 1998). Here we report the finished, annotated genome sequence of *B. thuringiensis* Al Hakam, which was collected in Iraq by the United Nations Special Commission (L. Radnedge, P. Agron, K. Hill, P. Jackson, L. Ticknor, P. Keim, and G. Andersen, *Appl. Environ. Microbiol.* 69:2755–2764, 2003).

The *Bacillus thuringiensis* Al Hakam genome was sequenced at the Joint Genome Institute using plasmid and fosmid DNA libraries. Draft assemblies were based on 246,217 total reads. All libraries provided 23-fold coverage of the genome. The Phred/Phrap/Consed software package was used for sequence assembly and quality assessment. After shotgun sequencing, reads were assembled with parallel phrap (High Performance Software, LLC). Possible misassemblies were corrected by transposon bombing (Epicenter Biotechnologies) of bridging clones. Gaps between contigs were closed by editing in Consed, by custom primer walks, or by PCR amplification. The complete genome of *B. thuringiensis* Al Hakam achieves an average of 24-fold se-

quence coverage per base with an error rate of less than 1 in 100,000. The sequences comprising the *B. thuringiensis* Al Hakam genome can be accessed using the GenBank accession numbers CP000485 and CP000486.

Gene predictions were obtained and annotation was performed as described previously (1). The 5.31-Mb genome of *B. thuringiensis* Al Hakam contains two replicons: a circular chromosome (5.26 Mb) encoding a predicted 4,969 open reading frames (ORFs), and the pALH1 circular phage, which contains 62 predicted ORFs. The G+C content of the chromosome is 35%, while that of the phage is 36%. The *B. thuringiensis* Al Hakam genome encodes 105 tRNAs and 13 rRNA operons and contains at least 21 pseudogenes. There were no additional plasmids identified in the assembly Blast searches against the *B. thuringiensis* Al Hakam genome using known insecticidal genes (*cry*, *cyt*, and *vip*) as queries revealed no chromosome-encoded (or phage-encoded) ORFs with significant similarity. Therefore, we conclude that this genome contains no homologues of the known *cry*, *cyt*, or *vip* genes. However, if they were present originally, it is possible that the plasmid(s) encoding these genes was lost during culture.

Previous AFLP analyses have shown that *B. thuringiensis* Al Hakam is phylogenetically more closely related to *B. anthracis*

* Corresponding author. Mailing address: Department of Energy Joint Genome Institute, Bioscience Division, MS M888, Los Alamos National Laboratory, Los Alamos, NM 87545. Phone: (505) 665-1485. Fax: (505) 665-3024. E-mail: jchalla@lanl.gov.

† Present address: Advanced Biomedical Computing Center, NCI-Frederick, Frederick, MD 21702.

‡ Present address: The Center for Biosecurity of UPMC, The Pier IV Building, 621 E. Pratt Street, Suite 210, Baltimore, MD 21202.

§ Present address: University of Florida, Gainesville, FL 32611.

¶ Present address: University of California, San Diego, La Jolla, CA 92093.

† Published ahead of print on 2 March 2007.

and other “Branch F” *Bacillus* isolates than to many of the commercially important *B. thuringiensis* isolates (2). The *B. thuringiensis* Al Hakam genome provides new sequence data that can be used to further study the evolutionary relationships among *B. cereus* group organisms.

We acknowledge the Intelligence Technology Innovation Center and the DOE Chemical and Biological Non-Proliferation Program for funding the sequencing and analysis of this genome.

REFERENCES

1. Han, C. S., G. Xie, J. F. Challacombe, M. R. Altherr, S. S. Bhotika, D. Bruce, C. S. Campbell, M. L. Campbell, J. Chen, O. Chertkov, C. Cleland, M. Dimitrijevic, N. A. Doggett, J. J. Fawcett, T. Glavina, L. A. Goodwin, K. K. Hill, P. Hitchcock, P. J. Jackson, P. Keim, A. R. Kewalramani, J. Longmire, S. Lucas, S. Malfatti, K. McMurry, L. J. Meincke, M. Misra, B. L. Moseman, M. Mundt, A. C. Munk, R. T. Okinaka, B. Parson-Quintana, L. P. Reilly, P. Richardson, D. L. Robinson, E. Rubin, E. Saunders, R. Tapia, J. G. Tesmer, N. Thayer, L. S. Thompson, H. Tice, L. O. Ticknor, P. L. Wills, T. S. Brettin, and P. Gilna. 2006. The pathogenomic sequence analysis of *Bacillus cereus* and *Bacillus thuringiensis* isolates closely related to *Bacillus anthracis*. *J. Bacteriol.* **188**:3382–3390.
2. Hoffmaster, A. R., K. K. Hill, J. E. Gee, C. K. Marston, B. K. De, T. Popovic, D. Sue, P. P. Wilkins, S. B. Avashia, R. Drumgoole, C. H. Helma, L. O. Ticknor, R. T. Okinaka, and P. J. Jackson. 2006. Characterization of *Bacillus cereus* isolates associated with fatal pneumonias: strains are closely related to *Bacillus anthracis* and harbor *B. anthracis* virulence genes. *J. Clin. Microbiol.* **44**:3352–3360.