

INFLAMMATORY BOWEL DISEASE

Bacterial and fungal microbiota in relation to probiotic therapy (VSL#3) in pouchitis

T Kühbacher, S J Ott, U Helwig, T Mimura, F Rizzello, B Kleessen, P Gionchetti, M Blaut, M Campieri, U R Fölsch, M A Kamm, S Schreiber



Gut 2006;55:833–841. doi: 10.1136/gut.2005.078303

See end of article for authors' affiliations

Correspondence to: Professor S Schreiber, Institute for Clinical Molecular Biology, and Department of General Internal Medicine, Christian Albrechts University Kiel, Schittenhelmstr. 12, 24105 Kiel, Germany; s.schreiber@mucosa.de

Revised version received 29 November 2005
Accepted for publication 7 December 2005
Published online first 9 January 2006

Background: The intestinal microbiota plays a critical role in the pathophysiology of pouchitis, a major complication after ileal pouch anal anastomosis in patients with ulcerative colitis. Recently, controlled trials have demonstrated that probiotics are effective in maintenance of remission in pouchitis patients. However, the mechanism by which therapy with probiotics works remains elusive. This study explores the role of the bacterial and fungal flora in a controlled trial for maintenance of remission in pouchitis patients with the probiotic VSL#3 compound.

Methods: The mucosa associated pouch microbiota was investigated before and after therapy with VSL#3 by analysis of endoscopic biopsies using ribosomal DNA/RNA based community fingerprint analysis, clone libraries, real time polymerase chain reaction (PCR), and fluorescence in situ hybridisation. Patients were recruited from a placebo controlled remission maintenance trial with VSL#3.

Results: Patients who developed pouchitis while treated with placebo had low bacterial and high fungal diversity. Bacterial diversity was increased and fungal diversity was reduced in patients in remission maintained with VSL#3 ($p=0.001$). Real time PCR experiments demonstrated that VSL#3 increased the total number of bacterial cells ($p=0.002$) and modified the spectrum of bacteria towards anaerobic species. Taxa specific clone libraries for *Lactobacilli* and *Bifidobacteria* showed that the richness and spectrum of these bacteria were altered under probiotic therapy.

Conclusions: Probiotic therapy with VSL#3 increases the total number of intestinal bacterial cells as well as the richness and diversity of the bacterial microbiota, especially the anaerobic flora. The diversity of the fungal flora is repressed. Restoration of the integrity of a "protective" intestinal mucosa related microbiota could therefore be a potential mechanism of probiotic bacteria in inflammatory barrier diseases of the lower gastrointestinal tract.

Probiotics are living microorganisms that, on ingestion in sufficient numbers, exert health benefits beyond the metabolic effect of nutritional components. It has been suggested that probiotic bacteria are effective and promising agents for the treatment of inflammatory gastrointestinal barrier disorders, including infectious colitis, antibiotic associated diarrhoea, and inflammatory bowel diseases (IBD).^{1–6} In recent years a series of placebo controlled randomised clinical trials have been conducted demonstrating the clinical efficacy of probiotics.^{7–11} However, understanding of the mechanism of action of probiotic bacteria is still incomplete.¹² In vitro and animal studies have shown that probiotics influence epithelial barrier function and gut permeability, restore the commensal bacterial microbiota, and reduce the production of inflammatory cytokines.^{13–16} Induction of anti-inflammatory cytokine expression, probably through modification of bacterial signalling in epithelial cells, has also been suggested.^{11, 17} This hypothesis is supported by the efficacy of local delivery of anti-inflammatory cytokines in animal models.¹⁸

Functional experiments using animal models suggest that probiotics may have profound effects on the composition of the faecal flora well beyond insertion of new bacterial strains.^{19, 20} However, the effects of probiotic bacteria on the resident mucosa related intestinal microbiota that is directly interacting with the host mucosal barrier organ is poorly understood. Culture based methods have identified a maximum of 10–30% of the bacterial spectrum (and probably less than 10% of fungal species) of the complex bacterial community in the human gut. Molecular techniques may

overcome these limitations.^{21–25} The 16S/18S rRNA marker gene system has been employed previously to analyse the bacterial communities of the intestine and to investigate the dominant species using oligonucleotide hybridisation, real time polymerase chain reaction (PCR), and fluorescent in situ hybridisation (FISH).^{22, 26–35} Community fingerprinting techniques have been shown to be a powerful tool for determining microbial diversity in complex samples.^{36–39}

Ulcerative colitis (UC) is a subform of IBD, a group of chronic recurrent gastrointestinal inflammatory barrier diseases. Restorative proctocolectomy with ileal pouch anal anastomosis is a well established procedure in the surgical treatment of UC and familial adenomatous polyposis.^{40–42} Approximately 15–46% of patients with UC develop pouchitis, a major complication after ileal pouch anal anastomosis, within five years of operation.^{43–46} A chronic relapsing form of pouchitis can be distinguished from a chronically active form.⁴⁷ The aetiology and pathophysiology of pouchitis are still unknown. Interaction between the mucosal immune system and the gut flora appears to play a crucial role in the initiation of pouchitis. Alterations of the bacterial microbiota may be an important factor triggering the disease process^{48, 49} and may explain the efficacy of antibiotic treatment in pouchitis.^{50, 51} However, remission periods are often short and the condition is complicated by frequent relapses. Recent

Abbreviations: IBD, inflammatory bowel disease; UC, ulcerative colitis; PDAI, pouchitis disease activity index; PCR, polymerase chain reaction; FISH, fluorescent in situ hybridisation; SSCP, single strand conformational polymorphism; DGGE, denaturing gradient gel electrophoresis

studies have shown that probiotic treatment with VSL#3, a mixture of eight different probiotic bacterial strains, is effective in maintaining remission in pouchitis.^{52–54}

A subset of patients (n = 15), who had participated in a double blind, randomised, placebo controlled clinical trial, were investigated.⁵² Patients with pouchitis in remission that had been induced by antibiotic therapy were recruited to receive either the VSL#3 probiotic compound or placebo for maintenance of remission. Biopsies were obtained before and two months after initiation of VSL#3 or placebo treatment. Probiotic therapy with VSL#3 increased the total number of intestinal bacterial cells as well as the richness and diversity of the bacterial microbiota, especially the anaerobic flora, whereas the fungal flora was repressed. In contrast, patients who relapsed while receiving placebo showed reduced diversity of the mucosal flora.

MATERIAL AND METHODS

Patients

Patients with recurrent or chronic active pouchitis, who were enrolled in a placebo controlled, randomised, remission maintenance trial with the probiotic preparation VSL#3⁵² were recruited for this substudy. The main inclusion criterion was prior successful induction of remission (defined as a pouchitis activity index of 0 or 1) by therapy with metronidazole and ciprofloxacin.⁵² Fifteen of 36 patients in the trial agreed to have additional biopsies for this study, taken for exploration of the mucosal microbiota. Pouchitis was defined by histological and endoscopic criteria using the pouchitis disease activity index (PDAI).⁵⁵ Age ranged from 22 to 64 years (median 34) in the 15 patients; there were six females and nine males. Total PDAI (expressed as medians) was 3 (1–7) at study entry in both groups.

Patients received VSL#3 (VSL pharmaceuticals Inc., Gaithersburg, Maryland, USA) 6 g once daily or identical placebo for 12 months. VSL#3 contains 300 billions viable lyophilised bacteria per gram, comprising four strains of lactobacilli (*Lactobacillus plantarum*, *Lactobacillus casei*, *Lactobacillus acidophilus*, and *Lactobacillus delbrueckii* subspecies *bulgaricus*), three strains of bifidobacteria (*Bifidobacterium infantis*, *Bifidobacterium longum*, and *Bifidobacterium breve*), and one strain of streptococcus (*Streptococcus salivarius* subspecies *thermophilus*). Placebo sachets contained maize without bacteria but did not differ in taste or physical appearance from VSL#3.

Mucosal biopsies were obtained during pouch endoscopy and immediately snap frozen in liquid nitrogen. Patients were studied before and after two months of therapy with study medication. Of the 15 patients recruited for the substudy, 10 received VSL#3 and five placebo. All VSL#3 patients in this substudy were still in remission after two months while all placebo patients showed signs of active pouchitis. All clinical and molecular examinations were carried out, and the results collated before the trial was unblinded.

Treatment of biopsy samples and DNA extraction

Biopsy and stool specimens were immediately snap frozen in liquid nitrogen after colonoscopy. DNA was extracted following a protocol adapted to the characteristics of the microorganisms⁵⁶: biopsies were incubated with 200 µl TL-buffer (200 mM HEPES, pH 7.5, 1 M KCl, 100 mM MgCl₂, 1 mM EDTA, 0.2% NaN₃) and 25 µl proteinase K (PeqLab, Erlangen, Germany) at 55°C for two hours. DNA was extracted using the FastDNA Spin Kit for soil after mechanical homogenisation (FastPrep FP 120 instrument) according to the manufacturer's instructions (both BIO 101, Carlsbad, California, USA). For sensitive and accurate quantification, DNA concentrations were determined using

PicoGreen according to the manufacturer's guidelines (dsDNA quantification kit; Molecular Probes, Leiden, the Netherlands).

Real time PCR

The real time PCR assay used in this study has been described previously.⁵⁷ Amplification and detection were carried out in 96 well optical plates on an ABI PRISM 7700 Sequence Detector with TaqMan Universal PCR 2× Master Mix, primer (0.4 µM), probe (0.2 µM), and 20 ng of sample DNA in a final volume of 50 µl per reaction. The universal probe UNI (5'-ACT GAG ACA CGG TCC A-3') binds to position 321-37 and is VIC labelled. In the present study, minor groove binder fluorescent probes with non-fluorescent quencher dyes (also called "dark" quencher) were used (Applied Biosystems, Foster City, California, USA). SDS software v1.7 or later was used to support non-fluorescent quencher probes. The universal primers used in this study hybridise to conserved regions on the 16S gene. The forward primer TPU1 (5'-AGA GTT TGA TCM TGG CTC AG-3') binds to position 8-27 and the reverse primer RTU8 (5'-AAG GAG GTG ATC CAN CCR CA-3') binds to position 1522-41 (*Escherichia coli* reference numbering). The absolute number of cells was normalised to the total amount of DNA extracted from each biopsy. The total number of cells was interpolated from the averaged standard curve as described.^{57 58}

Single strand conformational polymorphism (SSCP) analysis

For initial PCR, the two conserved primers COM-1 (5'-CAG CAG CCG CGG TAA TAC-3', position 519–536 on the reference *E coli* 16S gene) and Com2-Ph (5'-CCG TCA ATT CCT TTG AGT TT-3', position 907–926 on the reference *E coli* 16S gene) were used, as published by Schwieger and Tebbe.³⁶ PCR products were controlled for size and products by agarose gel electrophoresis. Preparation of single stranded DNA and silver staining of SSCP gels were performed according to Schwieger and Tebbe.³⁶ Image editing and normalisation were performed using GelCompare II-software (Applied Maths, Kortrijk, Belgium). The general diversity of bacterial species was calculated according to Shannon and Weaver, as described previously.⁵⁶

Denaturing gradient gel electrophoresis (DGGE) analysis

For PCR amplification of 18S rRNA fragments (1.650 bp), the two fungus specific primers NS1 (5'-GTA GTC ATA TGC TTG TCT C-3') and FR1-GC (5'-CCC CCG CCG CGC GCG GCG GGC GGG GCG GGG GCA CGG GCC GAI CCA TTC AAT CGG TAI T-3') were used, as published by Vainio and Hantula.⁵⁹ DGGE analysis and silver staining were performed according to Newton and colleagues.⁶⁰ Image editing, normalisation, and calculation of diversity indices were carried out as described above.

Sequencing of bands from SSCP/DGGE community profiles

To identify the bacteria correlated with the bands, selected products from the polyacrylamide gel after silver staining were excised with sterile scalpels. Gel slices were then mixed with elution buffer. After incubation, 5–20 µl were subjected to a PCR with COM primers (see above). After control with 2% agarose gel electrophoresis, 8 µl of the PCR products were digested with 0.3 U SAP (shrimp alkaline phosphatase) and 1.5 U ExoI. The sequencing reaction was performed on an ABI PRISM 3700 DNA Analyser using 1 µl of ABI PRISM BigDye (Applied Biosystems), 30 µM concentration of each sequencing primer (Eurogentec Seraing, Belgium), and 2 µl of digested PCR product. Forward and reverse sequences

Table 1 Oligonucleotides used for the group specific clone libraries

Bacterial group	Primer	Direction	Position	Sequence (5'-3')	Reference
Lactobacillus	Lac1	Forward	352-370*	AGCAGTAGGGAATCTCCA	Walter ³¹
	Lac2	Reverse	679-680*	ATTYCACCGCTACACATG	Walter ³¹
Bifidobacterium	Bif_164	Forward	164-181*	GGGTGGTAATGCCGGATG	Langendijk ²³
	Bif_662	Reverse	679-662*	TCCACCGTTACACCGGGAA	Langendijk ²³
Bacteroides/Prevotella	Bac_32	Forward	32-50*	AACGCTAGCTACAGGCTT	Bernhard ²⁷
	Bac_303	Reverse	303-321*	CCAATGTGGGGGACCTTC	Manz ⁶³
γ -Proteobacteria/Enterobacteriaceae	TPU5_F	Forward	906-26*	AAACTCAAATGAATTGACGG	von Wintzingerode ⁷⁸
	Enter1416	Reverse	1416-23*	CTTTGCAACCCACTCC	Sghir ⁷⁹
Sequencing	M13 (21)	Forward	389-404†	TGTA AACGACGGCCAGT	Guttman ⁸⁰
	M13 (24)	Reverse	208-25†	AACAGCTATGACCATG	Guttman ⁸⁰

*Position referring to the *Escherichia coli* reference sequence.

†M13 priming sites on pCCR 2.1 plasmid vector.

were aligned with Sequencher software package (Gene Codes Corp., Ann Arbor, Michigan, USA). Fragments were identified by NCBI BLAST database search.

Group specific clone libraries

Group specific clone libraries for the following bacterial taxa were generated: Lactobacillus, Bifidobacterium, Bacteroides/Prevotella, and γ -Proteobacteria/Enterobacteriaceae (for primer information see table 1). More than 600 clones were sequenced from 12 independent taxa specific libraries (Lactobacillus n = 135, Bifidobacterium n = 136, Bacteroides/Prevotella n = 124, and γ -Proteobacteria/Enterobacteriaceae n = 236) that were generated from pooled disease related (before study therapy, placebo, VSL#3) PCR products. A two step reconditioning PCR approach (35/10 cycles) was used as described elsewhere.^{61, 62} The specific fragment of the 16S rDNA was amplified and cloned into *E. coli* cells using the pCR 2.1 TOPO TA Cloning Kit for sequencing (Invitrogen, Karlsruhe, Germany), as described previously.⁵⁶

Cloning and sequencing of inserts was performed using 3.2 pmol of M13F and M13R sequencing primers (table 1), as described previously.⁵⁶ Alignment, assembly, and trimming of vector sequences were performed using the Sequencher software package (Gene Codes Corp.). Sequences were checked for vector contamination using the NCBI VecScreen tool. OTUs were identified by NCBI BLAST analysis using search results of at least 97% similarity. Sequences were examined for chimera using the Chimera Check tool of the Ribosomal Data Projects (RDP) of the Center for Microbial Ecology, Michigan State University, Michigan, USA.

Fluorescent in situ hybridisation (FISH) and oligonucleotide probes

Biopsies were fixed in 500 μ l of freshly prepared 4% buffered formalin and embedded in Histoplast (Sigma, St Louis, Missouri, USA) according to routine procedures. Cross sections (2 μ m) were cut and placed on coated microscope slides (Superfrost⁺/Plus). Sections were hybridised as described previously³⁵ using the following 16S/23S rRNA targeted oligonucleotide probes: (1) an equimolar mixture of five bacteria directed probes EUB 338, EUB 785, EUB 927, EUB 1055, and EUB 1088 (58, 59), referred to as EUB mix, to detect all bacteria; (2) Bac 303 to detect the *Bacteroides/Prevotella* cluster⁶³; (3) Erec 482 specific for most of the clostridia and eubacteria belonging to the *Clostridium coccoides-Eubacterium rectale* group⁶⁴; (4) bifidobacterial probe Bif 164²³; (5) Lab 158 for nearly all species of the genera *Lactobacillus* and *Enterococcus* in the gut⁶⁵; and Ec 1531 specific for a number of *Enterobacteriaceae* (for example, *Escherichia coli*, *Klebsiella pneumoniae*). The oligonucleotides were 5' labelled with the indocarbocyanine dye Cy3 (Thermo Hybaid GmbH, Waltham, Massachusetts, USA).




Figure 1 Bacterial diversity indicated as richness (No of bands) and diversity indices, obtained from single strand conformational polymorphism profiles of pouchitis patients in remission before randomisation (after treatment with antibiotics), after placebo (no therapy), and after VSL#3. For each group, the mean (SD) number of bands are shown. Bacterial diversity of the probiotic group was significantly higher compared with the starting value ($p=0.02$). Weighted diversity scores (SD) showed the same tendency.




Figure 2 Treatment with VSL#3 significantly increased the number of 16S rDNA copies compared with pretreatment samples and placebo ($p=0.002$, $p=0.013$, respectively).




Figure 3 Lactobacillus flora in remission (after antibiotic therapy), after placebo, and after VSL#3 therapy, as assessed by analysis of clone libraries. A total of 135 clones were randomly picked and sequenced. The graph shows the percentage of single species identified by BLAST analysis. Lactobacillus flora was more diverse under VSL#3 therapy in comparison with placebo. At inclusion (that is, before randomisation, after induction therapy with antibiotics), only two Lactobacillus species were detected, most of them *Lactobacillus mucosae* (>97%).

Expression of data

Normally distributed data are expressed as mean (SD), if not indicated otherwise. Statistical significance of the differences was examined using the Student’s *t* test for normally distributed data and the Mann Whitney U test or the Wilcoxon matched pairs test for non-normally distributed data. Distribution of data was evaluated by calculating Lilliefors probabilities based on the Komolgorov-Smirnov test.

RESULTS

Patients

Fifteen patients in remission induced by antibiotic therapy participated in this substudy (out of 36 participating in a double blind, randomised, placebo controlled trial of VSL#3 in pouchitis maintenance). No signs of morphological or histological inflammation were seen, as required by the inclusion criteria. The 10 patients who received VSL#3 were verified as being in remission at the time of the second biopsy (two months) while all five patients who were receiving placebo showed clinical and endoscopic signs of recurrent inflammation.

SSCP analysis and real time PCR of the bacterial microbiota

SSCP fingerprinting analysis of 16S rDNA fragments showed low bacterial richness and diversity in mucosal biopsies from patients at study entry after antibiotic induction of remission




Figure 4 Bifidobacterium flora in remission (after antibiotic therapy), after placebo, and after VSL#3 therapy, as assessed by analysis of clone libraries. A total of 136 clones were randomly picked and sequenced. The graph shows the percentage of single species identified by BLAST analysis. As seen for the Lactobacillus species (fig 3), the Bifidobacterium flora was more diverse in the VSL#3 group compared with both the placebo and samples before therapy. Before therapy (that is, before randomisation, after induction with antibiotics) *Bifidobacterium longum bv infantis* were seen almost exclusively.




Figure 5 Bacteroides/Prevotella flora in remission (after antibiotic therapy), after placebo, and after VSL#3 therapy, as assessed by analysis of clone libraries. A total of 135 clones were randomly picked and sequenced. The graph shows the percentage of single species identified by BLAST analysis. The diversity of Bacteroides/Prevotella bacteria was low (n=4), with the majority of sequences being assigned as uncultured *Bacteroides* species. There were no significant differences between the groups.

in comparison with normal colon. Probiotic therapy with VSL#3 increased bacterial richness and diversity of the pouch mucosal flora compared with both patients in remission before therapy and patients developing pouchitis while receiving placebo. The difference in bacterial diversity between patients before and after therapy with VSL#3 was statistically significant ($p = 0.02$) (fig 1). A universal real time PCR that amplified 16S rDNA was then used to compare the total number of bacterial cells. Treatment with VSL#3 caused a statistically significant increase in the number of 16S rDNA copies compared with pretreatment remission ($n = 10, p = 0.002$) and placebo treatment ($n = 5, p = 0.013$), as shown in fig 2. Separate statistical analysis showed no correlation between results of the SSCP analysis and real




Figure 6 γ -Proteobacteria/Enterobacteriaceae flora in remission (after antibiotic therapy), after placebo, and after VSL#3 therapy, as assessed by clone libraries. A total of 236 clones were randomly picked and sequenced because of the expected higher diversity. The graph shows the percentage of single species identified by BLAST analysis. The diversity of Enterobacteriaceae, especially *Enterobacter* species and *Escherichia coli*, was higher in the placebo and VSL#3 groups but there were only slight differences between these two groups. Remarkably, there was a high proportion of *Pseudomonas* species in the group before study therapy, which could be due to overgrowth induced by antibiotic selection pressure.

time PCR and PDAI (clinical, endoscopic, and histological subscores).

To investigate the nature of the species contributing to the increase in bacterial richness and diversity, which were consistently different between pouches in remission before and during VSL#3 treatment, bands were excised from the SCCP gel, reamplified, and sequenced. A BLAST suggested the presence of enterobacillus, *Enterococcus* sp, *Enterobacter* sp, and *Clostridium* sp in mucosal biopsies. Bacterial strains contained in VSL#3 were not detected as part of the mucosa adherent flora.

Taxa specific clone libraries

Community fingerprinting techniques are used for formal assessment of the richness and diversity of complex bacterial/fungal habitats, but for technical reasons often fail to provide sufficient taxonomic data. Clone libraries yield a realistic taxonomic representation of a complex microbial habitat that can be further narrowed and optimised by taxa specific amplification. To obtain data on the qualitative composition of the intestinal mucosa related microbiota after VSL#3 therapy, clone libraries with more than 600 sequenced clones were performed for the following taxa: *Lactobacillus*, *Bifidobacterium*, *Bacteroides/Prevotella*, and γ -Proteobacteria/Enterobacteriaceae. For each of the three experimental groups (before study treatment, but after antibiotic induction therapy, placebo treatment, and probiotic VSL#3 treatment) separate libraries were generated to assess differences in bacterial composition. As shown in figs 3 and 4, the spectrum of *Lactobacillus* and *Bifidobacterium* species was more diverse after VSL#3 treatment in comparison with placebo and samples before randomisation. With regard to lactobacilli and bacteroides species, patients at inclusion had only a narrow bacterial spectrum, with *Lactobacillus mucosae* and *Bifidobacterium longum* as the predominant species before treatment. The *Bacteroides/Prevotella* and γ -Proteobacteria/Enterobacteriaceae libraries showed only gradual differences between patients treated with placebo and VSL#3 (figs 5, 6). Most importantly, little differences were seen during treatment in general with regard to bacterial composition in the *Bacteroides/Prevotella* group, with the exception of the γ -Proteobacteria/Enterobacteriaceae flora. High levels of *Pseudomonas* species were observed after induction of remission by antibiotic therapy (probably as a result of overgrowth due to antibiotic selection pressure through preinclusion induction of remission).

Detection of mucosal bacteria by FISH

To complement the PCR based analysis, a second independent technique was used to identify and localise bacterial species. Colonisation was investigated in tissue sections of six patients in each group (before and during treatment) using the EUB mix of different fluorescence labelled 16S rRNA probes. With the different sets of probes described in the methods section it was found that nearly all bacteria detected by this technique represented *Enterobacteriaceae*. Bacteria were mainly detected within the epithelium and only rarely within the lamina propria. In uninfamed pouch (before treatment), intraepithelial *Enterobacteriaceae* (mainly *E coli*) were seen (fig 7). During treatment with VSL#3, a high mucosal count




Figure 7 The 16S rRNA targeted oligonucleotide mixture EUB that globally detects bacterial sequences was used in tissue sections from pouch mucosal biopsies. Subdifferentiation was carried out using a set of specific oligonucleotide probes. Almost all bacteria detected were classified as *Enterobacteriaceae* whereas the *Lactobacilli/Enterococcus* probes resulted only in occasional detection of signals in VSL#3 treated patients. The figure is a representation of six experiments per group (VSL#3 and placebo, before and after treatment). Mucosal biopsies in patients in remission prior to study drug therapy showed high numbers of intraepithelial bacteria (A). Colonisation was reduced after placebo treatment, which was accompanied by relapse of inflammation (B). Bacterial colonisation of the epithelium was maintained during VSL#3 treatment (C).

of these organisms was found. In contrast, recurrent inflammation during placebo treatment was associated with a reduction in the presence of *Enterobacteriaceae* (fig 4). No signals were detected with the *Bacteroides/Prevotella*, *Clostridium coccooides-Eubacterium rectale*, or bifidobacterial probes in any of the tissue sections. *Lactobacillus/Enterococcus* signals were seen only occasionally in tissue sections of VSL#3 treated patients.

DGGE analysis of the fungal microbiota

Because fungi are a potent, yet little explored, element of the intestinal microbiota, DGGE analysis was used to investigate fungal diversity. Due to the specific characteristics of the 18S rRNA gene, separation of 18S rDNA fragments by genetic fingerprinting requires longer amplicons. Therefore, DGGE was used because the length of the amplicons for SSCP analysis is restricted to 400–500 bp. Fungal diversity, indicated as number of bands and as weighted diversity index, was reciprocal compared with bacterial diversity. In remission before therapy (that is, following induction by antibiotics), fungal diversity was high ($n = 15$). During both placebo ($n = 5$) and VSL#3 ($n = 10$) treatment, fungal diversity decreased. In the VSL#3 group, a marked reduction in fungal diversity was seen both in comparison with pretreatment levels ($p = 0.001$) and with the placebo group ($p = 0.002$) (fig 8). Individual DGGE profiles showed wide interindividual variation in banding patterns. No consistent bands were identified using band matching analysis (fig 9). Separate statistical analysis of clinical, endoscopic, and histological PDAI scores revealed no significant correlation with fungal diversity.

DISCUSSION

The aetiology and pathogenesis of pouchitis is still unclear although ulcerative colitis and pouchitis share many similarities.^{66–69} This study investigated the influence of the probiotic agent VSL#3 on the bacterial microbiota in a placebo controlled clinical trial⁵² which was designed to assess the




Figure 8 Fungal diversity, indicated as richness (No of bands) and diversity indices, obtained from denaturing gradient gel electrophoresis profiles of pouchitis patients in remission (before study therapy, after induction with antibiotics), after placebo, and two months after treatment with VSL#3. For each group, the mean (SD) number of bands are shown. Fungal diversity in the VSL#3 group was significantly higher compared with both the starting value in remission ($p = 0.001$) and placebo ($p = 0.002$). Weighted diversity scores (SD) showed the same tendency.




Figure 9 Subset of denaturing gradient gel electrophoresis (DGGE) profiles showing wide interindividual variation in the composition of the fungal microflora. The gel demonstrates patients in remission at study start (R), and after two months of treatment with VSL#3 (V) or placebo (P). The fingerprints of fungal communities were generated by separation of 18S rDNA fragments. Defined fungal species were used as standards (lanes M), as described at Newton and colleagues⁶⁰. DGGE analysis was carried out for all samples and numeric calculations are shown in fig 5.

value of VSL#3 in maintaining antibiotic induced remission in patients with pouchitis.

In pouchitis, bacterial overgrowth and a decrease in beneficial bacteria, in particular lactobacilli and bifidobacteria, in comparison with conventional ileostomy flora, has been described.^{69–71} It appears likely that faecal stasis and immune stimulation by stool bacteria play an important role in the development of pouch inflammation. The mucosa associated flora is probably of immediate relevance to the disease process^{28–35} and differs significantly from the composition of the ileal and midstream/faecal flora.^{29–72}

Morphological data from FISH analysis indicate that normal *Enterobacteriaceae* species (mainly *E coli*) are found in uninfamed epithelium in ileoanal pouch biopsies (that is, from pouchitis patients who are in remission) in contrast with the normal ileum and colon of healthy volunteers in which no *E coli* was found within the epithelium.²⁸ When recurrent pouch inflammation occurred during placebo treatment, low bacterial diversity and a marked reduction in intramucosal *Enterobacteriaceae* species were found. The increase in *Enterobacteriaceae* within the mucosa during VSL#3 therapy indicates that remission maintenance under probiotics is associated with restoration of parts of the normal pouch flora.

For other relevant groups of intestinal bacteria, such as *Bacteroides/Prevotella* or the *Lactobacilli/Bifidobacteria* group, only sporadic signals were detected by FISH. The limited amount of mucosal biopsy material as well as the confined sensitivity of FISH analysis for single bacteria prompted us to complement the FISH results with PCR based molecular techniques.

The FISH results were supported by diversity analysis with SSCP and, at least in part, by taxa specific clone libraries. Successful maintenance of remission after VSL#3 treatment appeared to result in higher diversity of bacterial species that included members of the normal anaerobic enteric flora and not merely colonisation with strains contained only in VSL#3. With the exception of the *Bacteroides/Prevotella* group, the richness of bacteria was increased in the VSL#3 group compared with pretreatment levels. As diversity was increased during VSL#3 maintained remission, we suggest that this effect may be a component of the therapeutic mechanism of probiotics. The suggestion that low bacterial diversity could be an important mechanism for mucosal inflammation is supported by earlier investigations⁵⁶ in which

mucosal inflammation in IBD was associated with loss of normal anaerobic bacteria such as *Bacteroides* species, *Eubacterium* species, and *Lactobacillus* species.

Restoration of *Enterobacteriaceae* species in the mucosa and the increase in bacterial diversity are observed in the setting of the clinical trial, which demonstrated successful remission maintenance by VSL#3.⁵² The effects were observed on the specific background of remission induction with antibiotics. Antibiotics lead to significant alteration of the enteric bacterial balance. In neonates, antibiotic treatment induces complete eradication of *Lactobacillus* species together with a marked reduction in colonic total aerobic and anaerobic bacteria, in particular *Enterobacteriaceae* and *Enterococcus*.⁷³ The net effect of antimicrobial therapy therefore results in (i) a decrease in the total number of bacterial cells and (ii) a reduction in specific bacterial populations. Colonisation of the mucosa by *Enterobacteriaceae* species and diversification of *Lactobacillus* species and *Bifidobacterium* species in the VSL#3 group might be facilitated by eradication of complex populations by the preceding antibiotic therapy and the diminished competition by other dominant bacterial consortia, such as *Enterobacteriaceae* or *Enterococcus*. Therefore, expansion and diversification of the bacterial microbiota observed in the VSL#3 group could be specific to pretreatment with antibiotics on the background of the dysbiotic flora in IBD.⁵⁶⁻⁷⁴ In fact, this may also be the case for the clinical efficacy of the treatment in pouchitis.⁵²

It is well documented that pouchitis is associated with loss of beneficial intestinal bacteria, especially the subgroups lactobacilli and bifidobacteria.⁶⁹⁻⁷¹ We observed, as a net effect of VSL#3 associated remission, recolonisation and diversification of the potentially beneficial *Lactobacillus* and *Bifidobacterium* flora. As most patients in the placebo group relapsed and almost all patients in the VSL#3 group remained in remission, the specific effects of VSL#3 and the effect of an inflammation free mucosa interact and therefore cannot be separated in this study. Additional therapeutic mechanisms induced by VSL#3 treatment are possible that may be either primary or secondary to alteration of the constitution of the pouch microbiota. Specific induction of anti-inflammatory cytokines by probiotics could decrease the inflammatory activity in the mucosa and thereby provide better growth conditions for diversification of the flora.¹¹ Rachmilewitz *et al* demonstrated in Toll-like receptor and MyD88 deficient mice that the immune stimulatory effect of probiotic bacteria isolated from the VSL#3 compound was dependent on Toll-like receptor 9 signalling.⁷⁵ Therefore, direct interaction of probiotic bacteria with mucosal immunoregulation could either induce anti-inflammatory pathways¹¹ or directly interact with innate immune mechanisms⁷⁵ as the primary mechanisms altering the growth conditions of the mucosal microbiota, or could be secondary to changes in microbial constitution and diversity. The significant difference between the placebo and the VSL#3 groups indicates that VSL#3 induced remission is associated with reconstruction of the intestinal flora, but investigation of the hierarchy of mechanistic events has to await further exploration in model systems.

Fungi are important elements of the human flora. However, little is known about the composition of commensal fungal intestinal species. Fungal microbiotic diversity was inversely related to bacterial diversity. However, it has to be pointed out that fungal species represent only a small fraction of the total microbiota. Immunocompetent hosts have an efficient phagocytic capacity to prevent fungal invasion.⁷⁶ It can be hypothesised that alteration in the balance of bacterial and fungal species in the mucosal flora reflects a metabolic dysbalance of the complex microbial ecosystem with pathophysiological consequences for the

mucosal barrier. Most importantly, restored balance between bacterial and fungal diversity was seen in VSL#3 treated patients. The question of whether the increase in fungal diversity is a direct result of higher bacterial diversity or an independent parallel effect remains unclear. Although fungi are normal members of the intestinal microbiota to a certain extent, fungal overgrowth is a typical complication of any bacterial imbalance following antibiotic therapy or a specific dietary regimen, as for example in intensive care units. Most likely the fungal overgrowth seen here was a direct consequence of bacterial dysbalance supported by diminished control mechanisms and reduced competition through the total bacterial microbiota. Studies on the mutual dependencies of the different parts of the intestinal microbiota are urgently needed to clarify this point.

Our study demonstrated that VSL#3 maintained remission was accompanied by a higher bacterial and a reduced fungal diversity in comparison with placebo treatment. The increase in bacterial diversity was not caused by colonisation with bacterial strains contained in VSL#3 but represents an independent effect. It is not unclear at present whether the anti-inflammatory and immunoregulatory effects of probiotics are primary or secondary to induction of changes in the diversity of the mucosal microbiota. Increase in bacterial diversity may be a therapeutic mechanism for the probiotic mixture VSL#3 in maintenance of antibiotic induced remission in pouchitis. Methods for manipulating the complex microecology of the mucosal flora may be an important field for future therapeutic developments, especially addressing the secondary or primary prevention of pouchitis.

ELECTRONIC DATABASE AND WEBSITE INFORMATION

URLs used in this article are as follows:

- ARB software, ribosomal sequence database, and handling: <http://www.arb-home.de>
- NCBI BLAST homepage: <http://www.ncbi.nlm.nih.gov/BLAST/>
- Ribosomal Database Project II: <http://rdp.cme.msu.edu/>
- VecScreen database: <http://www.ncbi.nlm.nih.gov/VecScreen/VecScreen.html>

ACKNOWLEDGEMENTS

This paper describes a substudy to a randomised, controlled, clinical trial of VSL#3 versus placebo as maintenance therapy for pouchitis in remission, which has been reported separately.⁵² The work was supported by grants from the Deutsche Crohn und Colitis Vereinigung (DCCV), the Crohn's and Colitis Foundation of America (CCFA), the Deutsche Forschungsgemeinschaft (SFB 415), by Training and Mobility of Researchers (TMR-II) grant of the European commission, by the German Competence network IBD, and by the National Genome Research Network (both supported by the German Ministry of Education and Science (BMBF)). The authors thank the clinical investigators and the patients who provided research material for this study. The authors also thank Kornelia Smalla (Biologische Bundesanstalt für Land- und Forstwirtschaft, Institut für Pflanzenvirologie, Mikrobiologie und biologische Sicherheit, Braunschweig, Germany) for technical assistance with the DGGE analysis.

Authors' affiliations

T Kühbacher*, S J Ott*, S Schreiber, Department of General Internal Medicine, UKSH Campus Kiel, Kiel, Germany, and Institute for Clinical Molecular Biology, Christian-Albrechts-University Kiel, Germany
 U Helwig, U R Fölsch, Department of General Internal Medicine, UKSH Campus Kiel, Kiel, Germany
 T Mimura, M A Kamm, St Marks Hospital, London, UK
 F Rizzello, P Gionchetti, M Campieri, Department of Internal Medicine and Gastroenterology, University of Bologna, Italy

B Kleessen, M Blaut, German Institute of Human Nutrition (DIFE), Potsdam, Germany

*T Kühbacher and S J Ott contributed equally to this study.

Conflict of interest: None declared.

REFERENCES

- Madsen KL. The use of probiotics in gastrointestinal disease. *Can J Gastroenterol* 2001;**15**:817–22.
- Montaldo M, Arancio F, Izzi D, et al. Probiotics: history, definition, requirements and possible therapeutic applications. *Ann Ital Med Int* 2002;**17**:157–65.
- Shanahan F. Probiotics in inflammatory bowel disease. *Gut* 2001;**48**:609.
- Hart AL, Stagg AJ, Frame M, et al. The role of the gut flora in health and disease, and its modification as therapy. *Aliment Pharmacol Ther* 2002;**16**:1383–93.
- Cremonini E, Caro S, Santarelli L, et al. Probiotics in antibiotic associated diarrhoea. *Dig Liver Dis* 2002;**34**:S78–80.
- Isolauri E, Kirjavainen PV, Salminen S. Probiotic: a role in the treatment of intestinal infection and inflammation. *Gut* 2000;**50**:54–9.
- Gionchetti P, Rizzello F, Venturi A, et al. Oral bacteriotherapy as maintenance treatment in patients with chronic pouchitis: a double-blind, placebo-controlled trial. *Gastroenterology* 2000;**119**:305–9.
- Sen S, Mullan MM, Parker TJ, et al. Effect of *Lactobacillus plantarum* 299v on colonic fermentation and symptoms of irritable bowel syndrome. *Dig Dis Sci* 2002;**47**:2615–20.
- Pochapin M. The effect of probiotics on *Clostridium difficile* diarrhoea. *Am J Gastroenterol* 2000;**95**:S11–3.
- Kruis W, Schutz E, Frick P, et al. Double-blind comparison of an oral *Escherichia coli* preparation and mesalazine in maintaining remission of ulcerative colitis. *Aliment Pharmacol Ther* 1997;**11**:853–8.
- Furrie E, Macfarlane S, Kennedy A, et al. Synbiotic therapy (*Bifidobacterium longum*/Synergy 1) initiates resolution of inflammation in patients with active ulcerative colitis: a randomised controlled pilot trial. *Gut* 2005;**54**:242–9.
- Linskens RK, Huijsdens XW, Savelkoul PH, et al. The bacterial flora in inflammatory bowel disease: current insights in pathogenesis and the influence of antibiotics and probiotics. *Scand J Gastroenterol Suppl* 2001;**234**:29–40.
- Shibolet O, Karmeli F, Eliakim R, et al. Variable response to probiotics in two models of experimental colitis in rats. *Inflamm Bowel Dis* 2002;**8**:399–406.
- O'Mahony L, Freeney M, O'Halloran S, et al. Probiotic impact on microbial flora, inflammation and tumour development in IL-10 knock-out mice. *Aliment Pharmacol Ther* 2001;**15**:1219–25.
- Madsen K, Cornish A, Soper P, et al. Probiotic bacteria enhance murine and human intestinal epithelial barrier function. *Gastroenterology* 2001;**121**:580–91.
- Ulisse S, Gionchetti P, D'Álo S, et al. Expression of cytokines, inducible nitric oxide synthase, and matrix metalloproteinases in pouchitis: effects of probiotic treatment. *Am J Gastroenterol* 2001;**96**:2691–9.
- Borruel N, Carol M, Casellas F, et al. Increased mucosal tumour necrosis factor (alpha) production in Crohn's disease can be downregulated ex vivo by probiotic bacteria. *Gut* 2002;**51**:659–64.
- Steidler L, Hans W, Schotte L, et al. Treatment of murine colitis by *Lactococcus lactis* secreting interleukin-10. *Science* 2000;**289**:1352–5.
- Tannock GW, Munro K, Harmsen HJ, et al. Analysis of the fecal microflora of human subjects consuming a probiotic product containing *Lactobacillus rhamnosus* DR20. *Appl Environ Microbiol* 2000;**66**:2578–88.
- Bartosch S, Woodmansey EJ, Paterson JC, et al. Microbiological effects of consuming a synbiotic containing *Bifidobacterium bifidum*, *Bifidobacterium lactis*, and oligofructose in elderly persons, determined by real-time polymerase chain reaction and counting of viable bacteria. *Clin Infect Dis* 2005;**40**:28–37.
- Simon GL, Gorbach SL. Intestinal flora in health and disease. *Gastroenterology* 1984;**86**:174–93.
- Tannock GW. Molecular methods for exploring the intestinal ecosystem. *Br J Nutr* 2002;**87**:S199–201.
- Langendijk P, Schut F, Jansen G, et al. Quantitative fluorescence in situ hybridization of *Bifidobacterium* spp. with genus-specific 16S rRNA-targeted probes and its application in fecal samples. *Appl Environ Microbiol* 1995;**61**:3069–75.
- Wilson K, Blichington R. Human colonic biota studied by ribosomal DNA sequence analysis. *Appl Environ Microbiol* 1996;**62**:2273–8.
- McFarlane G, Gibson G. Metabolic activities of the normal colonic microflora. In: Gibson S, ed. *Human health: contribution of microorganisms*. Frankfurt: Springer, Frankfurt, Germany, 1994:17–38.
- Zheng D, Alm E, Stahl D, et al. Characterization of universal small-subunit rRNA hybridization probes for quantitative molecular microbial ecology studies. *Appl Environ Microbiol* 1996;**62**:4504–13.
- Zoetendal EG, Akkermans ADL, De Vos WM. Temperature gradient gel electrophoresis analysis of 16S rRNA from human fecal samples reveals stable and host-specific communities of active bacteria. *Appl Environ Microbiol* 1998;**64**:3854–59.
- Swidsinski A, Ladhoff A, Pernthaler A, et al. Mucosal flora in inflammatory bowel disease. *Gastroenterology* 2002;**122**:44–54.
- Zoetendal EG, von Wright A, Vilpponen-Salmela T, et al. Mucosa-associated bacteria in the human gastrointestinal tract are uniformly distributed along the colon and differ from the community recovered from feces. *Appl Environ Microbiol* 2002;**68**:3401–7.
- Seksik P, Rigottier-Gois L, Gramet G, et al. Alterations of the dominant faecal bacterial groups in patients with Crohn's disease of the colon. *Gut* 2003;**52**:237–42.
- Walter J, Hertel C, Tannock GW, et al. Detection of *Lactobacillus*, *Pediococcus*, *Leuconostoc*, and *Weissella* species in human feces by using group-specific PCR primers and denaturing gradient gel electrophoresis. *Appl Environ Microbiol* 2001;**67**:2578–85.
- Tannock GW. The bifidobacterial and *Lactobacillus* microflora of humans. *Clin Rev Allergy Immunol* 2002;**22**:231–53.
- Dore J, Sghir A, Hannequart-Gramet G, et al. Design and evaluation of a 16S rRNA-targeted oligonucleotide probe for specific detection and quantitation of human faecal *Bacteroides* populations. *Syst Appl Microbiol* 1998;**21**:65–71.
- Zoetendal EG, Ben-Amor K, Harmsen HJ, et al. Quantification of uncultured *Ruminococcus obeum*-like bacteria in human fecal samples by fluorescent in situ hybridization and flow cytometry using 16S rRNA-targeted probes. *Appl Environ Microbiol* 2002;**68**:4225–32.
- Kleessen B, Kroesen AJ, Buhr HJ, et al. Mucosal and invading bacteria in patients with inflammatory bowel disease compared with controls. *Scand J Gastroenterol* 2002;**37**:1034–41.
- Schwieger F, Tebbe CC. A new approach to utilize PCR-single-strand-conformation polymorphism for 16S rRNA gene-based microbial community analysis. *Appl Environ Microbiol* 1998;**64**:4870–6.
- Tebbe CC, Schmalenberger A, Peters S, et al. Single-strand conformation polymorphism (SSCP) for microbial community analysis. In: Rochelle PA, ed. *Environmental molecular microbiology: protocols and applications*. Wymondham, UK: Horizon Scientific Press, 2001:161–74.
- Valinsky L, Della Vedova G, Jiang T, et al. Oligonucleotide fingerprinting of rRNA genes for analysis of fungal community composition. *Appl Environ Microbiol* 2002;**68**:5999–6004.
- Valinsky L, Della Vedova G, Scupham AJ, et al. Analysis of bacterial community composition by oligonucleotide fingerprinting of rRNA genes. *Appl Environ Microbiol* 2002;**68**:3243–50.
- Stahlberg D, Gullberg K, Liljeqvist L, et al. Pouchitis following pelvic pouch operation for ulcerative colitis. Incidence, cumulative risk, and risk factors. *Dis Colon Rectum* 1996;**39**:1012–18.
- Kock NG, Darle N, Hultén L, et al. Ileostomy. *Curr Probl Surg* 1977;**14**:1–52.
- Parks AG, Nicholls RJ. Proctocolectomy without ileostomy. *BMJ* 1978;**2**:85–8.
- Sandborn WJ. Pouchitis following ileal pouch-anal anastomosis: definition, pathogenesis, and treatment. *Gastroenterology* 1994;**107**:1856–60.
- Svaninger G, Nordgren S, Oresland T, et al. Incidence and characteristics of pouchitis in the Kock continent ileostomy and the pelvic pouch. *Scand J Gastroenterol* 1993;**28**:695–700.
- Hurst RD, Molinari M, Chung TP, et al. Prospective study of the incidence, timing and treatment of pouchitis in 104 consecutive patients after restorative proctocolectomy. *Arch Surg* 1996;**131**:497–500.
- Madden MV, Farthing MJ, Nicholls RJ. Inflammation in the ileal reservoir: pouchitis. *Gut* 1990;**31**:247–9.
- Meagher AP, Farouk R, Dozois RR, et al. J ileal pouch-anal anastomosis for chronic ulcerative colitis: complication and long-term outcome in 1310 patients. *Br J Surg* 1998;**85**:800–3.
- Ruseler-van Embden JG, Schouten WR, van Lieshout LM. Pouchitis: result of microbial imbalance? *Gut* 1994;**35**:659–64.
- de-Silva HJ, Millard PR, Soper N, et al. Effects of the fecal stream and stasis on the ileal pouch mucosa. *Gut* 1991;**32**:1166–9.
- Madden M, McIntyre A, Nicholls RJ. Double-blind cross-over trial of metronidazole versus placebo in chronic unremitting pouchitis. *Dig Dis Sci* 1994;**39**:509–15.
- Mimura T, Rizzello F, Helwig U, et al. Randomised, placebo-controlled, double-blind trial of once daily, high dose probiotic therapy (VSL#3) for maintaining remission in patients with recurrent or refractory pouchitis. *Gut* 2004;**53**:108–14.
- Gionchetti P, Rizzello F, Venturi A, et al. Antibiotic combination therapy in patients with chronic, treatment-resistant pouchitis. *Aliment Pharmacol Ther* 1999;**13**:713–18.
- Campieri M, Gionchetti P. Probiotics in inflammatory bowel disease: new insight to pathogenesis or a possible therapeutic alternative? *Gastroenterology* 1999;**116**:1246–60.
- Gionchetti P, Rizzello F, Helwig U, et al. Prophylaxis of pouchitis onset with probiotic therapy: a double-blind, placebo-controlled trial. *Gastroenterology* 2003;**124**:1202–9.
- Sandborn WJ, Tremaine WJ, Batts KP, et al. Pouchitis after ileal pouch-anal anastomosis: a pouchitis disease activity index. *Mayo Clin Proc* 1994;**69**:409–15.
- Ott SJ, Musfeldt M, Wenderoth DF, et al. Reduction in diversity of the colonic mucosa associated bacterial microflora in patients with active inflammatory bowel disease. *Gut* 2004;**53**:685–93.
- Ott SJ, Musfeldt M, Ullmann U, et al. Quantification of intestinal bacterial populations by real-time PCR with a universal primer set and minor groove binder probes: a global approach to the enteric flora. *J Clin Microbiol* 2004;**42**:2566–72.
- Lyons SR, Griffen AL, Leys EJ. Quantitative real-time PCR for porphyromonas gingivalis and total bacteria. *J Clin Microbiol* 2000;**38**:2362–5.
- Vainio EJ, Hantula J. Direct analysis of wood-inhabiting fungi using denaturing gradient gel electrophoresis of amplified ribosomal DNA. *Mycol Res* 2000;**104**:927–36.
- Newton C, Gomes M, Fagbola O, et al. Dynamics of fungal communities in bulk and maize rhizosphere soil in the tropics. *Appl Environ Microbiol* 2003;**69**:3758–66.

- 61 **Thompson JR**, Marcelino LA, Polz MF. Heteroduplexes in mixed-template amplifications: formation, consequence and elimination by 'reconditioning PCR'. *Nucl Acids Res* 2002;**30**:2083-8.
- 62 **Acinas SG**, Klepac-Ceraj V, Hunt DE, *et al.* Fine-scale phylogenetic architecture of a complex bacterial community. *Nature* 2004;**430**:551-4.
- 63 **Manz W**, Amann R, Ludwig W, *et al.* Application of a suite of 16S rRNA-specific oligonucleotide probes designed to investigate bacteria of the phylum cytophaga-flavobacter-bacteroides in the natural environment. *Microbiology* 1996;**142**:1097-106.
- 64 **Franks AH**, Harmsen HJM, Raangs GC, *et al.* Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl Environ Microbiol* 1998;**64**:3336-45.
- 65 **Harmsen HJ**, Raangs GC, He T, *et al.* Extensive set of 16S rRNA-based probes for detection of bacteria in human feces. *Appl Environ Microbiol* 2002;**68**:2982-90.
- 66 **Luukkonen P**, Jarvinen H, Tanskanen M, *et al.* Pouchitis-recurrence of the inflammatory bowel disease? *Gut* 1994;**35**:243-6.
- 67 **de Silva HJ**, Jones M, Prince C, *et al.* Lymphocyte and macrophage subpopulations in pelvic ileal pouches. *Gut* 1991;**32**:1160-5.
- 68 **Mahida JR**, Patel S, Gionchetti P, *et al.* Macrophage subpopulations in the lamina propria of normal and inflamed colon and terminal ileum. *Gut* 1989;**30**:826-34.
- 69 **Onderdonk AB**, Dvorak AM, Cisneros RL, *et al.* Microbiologic assessment of tissue biopsy samples from ileal pouch patients. *J Clin Microbiol* 1992;**30**:312-17.
- 70 **Kmiot WA**, Youngs D, Tudor R, *et al.* Mucosal morphology, cell proliferation, and fecal bacteriology in acute pouchitis. *Br J Surg* 1993;**80**:1445-9.
- 71 **Sandborn WJ**, McLeod R, Jewell DP. Medical therapy for induction and maintenance of remission in pouchitis: a systemic review. *Inflamm Bowel Dis* 1999;**5**:33-9.
- 72 **Ott SJ**, Musfeldt M, Timmis KN, *et al.* In vitro alterations of intestinal bacterial microbiota in fecal samples during storage. *Diagn Microbiol Infect Dis* 2004;**50**:237-45.
- 73 **Schumann A**, Nutten S, Donnicola D, *et al.* Neonatal antibiotic treatment alters gastrointestinal tract developmental gene expression and intestinal barrier transcriptome. *Physiol Genomics*, 2005;**23**:235-45.
- 74 **Tamboli CP**, Neut C, Desreumaux P, *et al.* Dysbiosis in inflammatory bowel disease. *Gut* 2004;**53**:1-4.
- 75 **Rachmilewitz D**, Katakura K, Karmeli F, *et al.* Toll-like receptor 9 signaling mediates the anti-inflammatory effects of probiotics in murine experimental colitis. *Gastroenterology* 2004;**126**:520-8.
- 76 **Lionakis MS**, Kontoyiannis DP. Glucocorticoids and invasive fungal infections. *Lancet* 2003;**362**:1828-38.
- 77 **Bernhard AE**, Field KG. Identification of nonpoint sources of fecal pollution in coastal waters by using host-specific 16S ribosomal DNA genetic markers from fecal anaerobes. *Appl Environ Microbiol* 2000;**66**:1587-94.
- 78 **von Wintzingerode F**, Landt O, Ehrlich A, *et al.* Peptide nucleic acid-mediated PCR clamping as a useful supplement in the determination of microbial diversity. *Appl Environ Microbiol* 2000;**66**:549-57.
- 79 **Sghir A**, Gramet G, Suau A, *et al.* Quantification of bacterial groups within human fecal flora by oligonucleotide probe hybridization. *Appl Environ Microbiol* 2000;**66**:2263-6.
- 80 **Guttman DS**, Charlesworth D. An X-linked gene with a degenerate Y-linked homologue in a dioecious plant. *Nature* 1998;**393**:263-6.

EDITOR'S QUIZ: GI SNAPSHOT

Answer

From question on page 832

Absence of characteristic signs of Crohn's colitis, such as segmental involvement, deep ulcerations, and fistula, and lack of a therapeutic response, made the formerly suspected diagnosis appear unlikely. Therefore, parasitological and bacteriological investigations were performed.

Microscopy of a fresh stool sample revealed erythrocyclophagous "magna-form" trophozoites of *Entamoeba histolytica*. Differentiation from apathogenic but morphologically identical *Entamoeba dispar* or *Entamoeba moshkovskii* was confirmed by positive anti-*E histolytica* serum antibodies and a polymerase chain reaction specific for *E histolytica* DNA from a faecal sample. Antiparasitic treatment with metronidazole and subsequently with diloxanide furoate and paromomycin resulted in almost complete regression of colitis.^{1 2}

In the past, parasitological investigations had not been considered because there was no history of travel to countries where amoebiasis is endemic. Thus amoebic colitis was misinterpreted as Crohn's disease. Careful analysis of previous investigations revealed that the patient never had arthritis but polyarthrosis, probably due to his strenuous physical working activities. The source of amoeba infection was likely to be a contaminated waste water drain. Autochthonous *E histolytica* infections have been repeatedly reported in Western Europe.³⁻⁸ Amoeba infections may persist for months or even years prior to the development of invasive disease, but invasiveness may be facilitated by immunosuppressive therapy.²

doi: 10.1136/gut.2005.077487

REFERENCES

- 1 **Blessmann J**, Tannich E. Treatment of asymptomatic intestinal *Entamoeba histolytica* infection. *N Engl J Med* 2002;**347**:1384.
- 2 **Richter J**. Treatment of intestinal parasitic diseases. In: Wolfe T, Davis GA, Farroye FA, *et al.* *Therapy of digestive disorders*, 2nd edn. London: Elsevier, 2006:745-54.
- 3 **Doby JM**, Duval JM. Beauccournu. Amibiase, maladie professionnelle des egoutiers? *Nouv Presse Méd* 1980;**9**:532-3.
- 4 **Knobloch J**, Funke M, Bienzle U. Autochthonous liver abscess in Germany. *Tropenmed Parasit* 1980;**31**:414-16.
- 5 **Knobloch J**, Bialek R, Hagemann J. Intestinaler Protozoenbefall durch berufsbedingten Abwasserkontakt (in German). *Dtsch Med Wschr* 1983;**108**:57-60.
- 6 **Gatti S**, Cervini C, Bruno A, *et al.* Transmission of *Entamoeba histolytica* within a family complex. *Trans R Soc Trop Med Hyg* 1995;**89**:403-5.
- 7 **Vreden SGS**, Visser LG, Verweij JJ, *et al.* Outbreak of amoebiasis in a family in the Netherlands. *Clin Infect Dis* 2000;**31**:1101-4.
- 8 **Ambroise-Thomas P**, Goullier A, Grillot R, *et al.* Epidemic of autochthonous hepatic and intestinal amoebiasis in a place near Grenoble (in French). *Acta Trop* 1975;**32**:365-70.