

## WORLD VIEW

# Ethnicity-specific prevalences of refractive errors vary in Asian children in neighbouring Malaysia and Singapore

S-M Saw, P-P Goh, A Cheng, A Shankar, D T H Tan, L B Ellwein



*Br J Ophthalmol* 2006;90:1230–1235. doi: 10.1136/bjo.2006.093450

See end of article for authors' affiliations

Correspondence to:  
S-M Saw, Department of Community, Occupational and Family Medicine, National University of Singapore, 16 Medical Drive, Singapore 117597, Singapore;  
cofsawsm@nus.edu.sg

Accepted for publication  
1 June 2006  
Published Online First  
29 June 2006

**Aim:** To compare the prevalences of refractive errors in Malay, Chinese and Indian children in Malaysia and Singapore.

**Methods:** Children aged 7–9 years from three schools in the Singapore Cohort study of the Risk factors for Myopia (n = 1962) and similarly aged children from a random cluster sample in the metropolitan Kuala Lumpur area in the Malaysia Refractive Error Study in Children (n = 1752) were compared. Cycloplegic autorefractometry was conducted in both countries.

**Results:** The prevalence of myopia (spherical equivalent of at least –0.5 diopters (D) in either eye) was higher in Singapore Malays (22.1%) than in Malays in Malaysia (9.2%; 95% confidence interval (CI) 11.2 to 14.7;  $p < 0.001$ ). Similarly, Singapore Chinese (40.1%) had higher prevalences than Malaysian Chinese (30.9%; 95% CI 1.5 to 16.9). Singapore Indians had a higher prevalence (34.1%) than Malaysian Indians (12.5%; 95% CI 17.4 to 25.9). The multivariate odds ratio of astigmatism (cylinder at least 0.75 D in either eye) in Singapore Malays compared with Malaysian Malays was 3.47 (95% CI 2.79 to 4.32). Ethnicity-specific hyperopia rates did not differ in Singapore and Malaysia.

**Conclusion:** The ethnicity-specific prevalences of myopia in Singapore Malays, Chinese and Indians are higher than those in Malaysian Malays, Chinese and Indians. As Malays, Chinese and Indians in Malaysia have genetic make-up similar to that of Malays, Chinese and Indians in Singapore, environmental factors may contribute to the higher myopia rates.

The contributions of genetic and environmental factors towards the perceived “epidemic” of myopia in Asia need further evaluation.<sup>1</sup> In Asia, the prevalence of myopia is highest in urban Chinese populations such as Hong Kong, Taiwan, Singapore and southern China, and lowest in non-Chinese rural populations such as Nepal and India.<sup>2–7</sup> The differences in myopia rates may be attributed to variations in genetic susceptibility or environmental lifestyles in subpopulations. It has been purported that near work activity and a competitive education system may be the primary environmental factors related to myopia, whereas the genetic susceptibility to myopia may be polygenic in nature.<sup>8–9</sup>

Multiethnic populations allow the evaluation of genetic effects and inter-racial differences in culture and lifestyle. Comparative studies across different populations with similar genetic pools may provide insights into the importance of environmental influences in the development of myopia. Singapore and Malaysia are neighbouring countries, separated by a narrow waterway, and are examples of multiethnic societies in East Asia. The three major ethnic groups are the same in both countries: Chinese (77% in Singapore and 26% in Malaysia), Malays (14% in Singapore and 65% in Malaysia) and Indians (8% in Singapore and 8% in Malaysia), and other minorities (1% in Singapore and 1% in Malaysia).<sup>10–11</sup> Chinese in Singapore and Malaysia migrated from the same localities in south China (primarily Fujian and Guangdong Provinces), and Indians in Singapore and Malaysia migrated from the same parts of India, primarily from south India and Sri Lanka. Most Chinese and Indian families have lived for decades (primarily second, third or fourth generation), and the Malays are native to Singapore and Malaysia.

We aim to evaluate differences in the prevalences of refractive errors, including myopia, in 7–9-year-old children

of similar ethnicity—Malays, Chinese and Indians—from the Refractive Error Study in Children (RESC) and the Singapore COhort study of the Risk factors for Myopia (SCORM) conducted in two neighbouring countries, Malaysia and Singapore.<sup>12–16</sup>

## METHODS

### Singapore population

Children from grades 1 to 3 from the Northern and Western schools and grades 1 and 2 in the Eastern school were invited to participate. Ninety four children with serious medical or ocular disorders were excluded. The participation rate was 71.8% (693/965) in the Northern school, 80.0% (956/1195) in the Western school and 49.1% (313/638) in the Eastern school. The parents completed a baseline questionnaire, and the completed educational levels of fathers and mothers were assessed. Written informed consent was obtained from the parents.

Children were examined on the school premises in 1999 in the Northern and Eastern schools, and in 2001 in the Western school by a team of ophthalmologists and optometrists who were blinded to questionnaire data. After distance logarithm of the minimum angle of resolution visual acuity measurements, cycloplegia was induced with three drops of 1% cyclopentolate solution administered at 5-min intervals. At least 30 min after the last drop, refraction was obtained with one of two autokeratorefractometers (RK5, Canon, Tochigiken, Japan). A total of 1962 children were refracted: 851 (43.4%) were 7 years of age, 630 (32.1%) were 8 years of age and 481 (24.5%) were 9 years of age. There were 992 (50.6%) boys and 970 (49.4%) girls. Ethnicity was distributed

**Abbreviations:** RESC, Refractive Error Study in Children; SCORM, Singapore COhort study of the Risk factors for Myopia

as follows: in the Northern school, there were 415 (59.9%) Chinese, 227 (32.8%) Malays, 42 (6.1%) Indians and 9 (1.3%) of other ethnicity; in the Western school, there were 743 (77.7%) Chinese, 121 (12.7%) Malays, 83 (8.7%) Indians and 9 (0.9%) of other ethnicity; and in the Eastern school, there were 309 (98.7%) Chinese, 1 (0.3%) Indian and 3 (1.0%) of other ethnicity. Further details of the SCORM protocol have been described elsewhere, along with findings from the baseline examinations.<sup>13–16</sup>

### Malaysia population

The Malaysian study population was obtained by random cluster sampling of children aged 7–15 years in Gombak District in the metropolitan Kuala Lumpur area. Clusters were defined geographically, and eligible children were enumerated by name, sex and age using house-to-house visits within randomly selected clusters. All eligible children recruited attended primary school. The ethnicity of the father and years of schooling for each parent were recorded. Years of schooling was grouped as no formal or incomplete primary education (<6 years of schooling), completed primary education (6–9 years of schooling), completed secondary education (10–11 years of schooling) or completed tertiary education ( $\geq 12$  years of schooling). Written consent for each child was obtained from a parent or guardian.

Eye examinations were conducted in 140 schools by two clinical teams, mostly between March and July 2003. After distance logarithm of the minimum angle of resolution visual acuity measurements, cycloplegia was induced with two drops of 1% cyclopentolate administered 5 min apart by ophthalmic assistants, with a third drop after 20 min, if necessary. Cycloplegia and pupil dilation were evaluated after an additional 15 min—pupillary dilation of  $\geq 6$  mm with absence of light reflex was considered complete cycloplegia. Refraction was carried out by an optometrist with a handheld autorefractor (Retinomax K-Plus; Nikon, Tokyo, Japan).

Of the 2104 children between the ages of 7 and 9 years, 1781 (84.6%) were examined and 1752 (83.3%) had auto-refraction measurements after successful cycloplegic dilation: 581 (33.2%) were 7 years old, 601 (34.3%) were 8 years old and 570 (32.5%) were 9 years old. There were 924 (52.7%) boys and 828 (47.3%) girls. The ethnic composition was 1245 (71.1%) Malays, 285 (16.3%) Chinese, 152 (8.7%) Indians and 70 (4.0%) of other ethnicity.

The complete RESC protocol has been described elsewhere, as have further details regarding the specific sampling and examination methods used in Gombak District.<sup>12–17</sup> Examination participation rates in both Malaysia and Singapore were reasonably similar across age groups and sex. In Malaysia, the participation rate was highest in the other ethnicity category (92.1%), followed by Malays (84.1%), Chinese (81.0%) and Indians (77.6%). In Singapore, the participation rate was highest in Indians (82.1%), followed by Malays (78.9%), others (77.1%) and Chinese (67.4%).

The SCORM study was approved by the Institutional Review Board of the Singapore Eye Research Institute. The RESC study protocol was approved by the World Health Organization Secretariat Committee on Research Involving Human Subjects, Geneva, and the Ethical Committee of the Standing Committee for Medical Research, Malaysia Ministry of Health.

### Definitions and data analysis

Definitions of refraction from the RESC studies were used. Myopia was defined as spherical equivalent refractive error of at least  $-0.50$  diopter (D) and hyperopia as  $\geq +2.00$  D. Astigmatism was defined as cylinder of at least 0.75 D. Clustering effects associated with the cluster sampling design

in Malaysia were taken into account in all statistical analyses. Statistical analyses were carried out using Stata Statistical Software, Release 8.0 svy commands for analysing complex survey design data with clusters as primary sampling units.<sup>18</sup> The primary sampling unit was each geographical cluster in Malaysia, and the Singapore data were considered as one cluster. Differences in the prevalences of myopia, hyperopia and astigmatism between Malaysia and Singapore were considered significant if the 95% confidence intervals of the differences in the prevalences did not cross zero and  $p$  values were  $<0.05$ . The data were pooled to evaluate the effect of living in Singapore versus living in Malaysia on myopia, astigmatism and hyperopia in each ethnic group. Age, sex, education of the father or myopia-adjusted odds ratios (ORs) were presented.<sup>18</sup>

### RESULTS

The Singapore population had a greater proportion of Chinese, 7-year olds and fathers with tertiary educational level than the Malaysian population (table 1). Similarly, the Singapore site (10.7%) had a greater proportion of mothers with tertiary education than the Malaysian site (5.8%;  $p<0.001$ ). The mean (standard deviation (SD)) spherical equivalent refractive errors of the right eye in Malaysian Malay (+0.65 D (0.82)), Chinese ( $-0.14$  D (1.74)) and Indian children (+0.57 D (1.10)) indicated less myopia when compared with those of Singapore Malay (+0.15 D (1.42);  $p<0.001$ ), Chinese ( $-0.52$  D (1.69);  $p<0.001$ ) and Indian ( $-0.22$  D (1.74);  $p<0.001$ ) children (fig 1). Similarly, the mean (SD) refractive errors of both Malaysian boys and girls indicated less myopia compared with Singapore boys (+0.50 D (1.03)  $v$   $-0.49$  D (1.76);  $p<0.001$ ) and girls (+0.51 D (1.17)  $v$   $-0.25$  D (1.56);  $p<0.001$ ). The mean (SD) of refractive errors indicated less myopia in Malaysian children compared with Singapore children for 7 year olds (+0.64 D (0.92)  $v$   $-0.03$  D (1.38);  $p<0.001$ ), 8 year olds (+0.51 D (0.96)  $v$   $-0.29$  D (1.57);  $p<0.001$ ) and 9 year olds (+0.35 D (1.35)  $v$   $-1.08$  D (2.01);  $p<0.001$ ).

Overall, the prevalence of myopia was higher in Singapore (36.3%) than in Malaysia (13.4%; 95% CI of the difference in rates = 20.3 to 25.4;  $p<0.001$ ; table 2). The prevalence of high myopia (spherical equivalent in the worse eye of at least




Figure 1 Spherical equivalents in Malaysian Malay, Chinese and Indian children.

**Table 1** Characteristics of children in Malaysia and Singapore (n = 3714)

	All n (%)	Malaysia (n = 1752) n (%)	Singapore (n = 1962) n (%)	p Value ( $\chi^2$ )
Ethnicity				
Malay	1593 (42.9)	1245 (71.1)	348 (17.7)	<0.001
Chinese	1752 (47.2)	285 (16.3)	1467 (74.8)	
Indian	278 (7.5)	152 (8.7)	126 (6.4)	
Others	91 (2.5)	70 (4.0)	21 (1.1)	
Age (years)				
7	1432 (38.6)	581 (33.2)	851 (43.4)	<0.001
8	1231 (33.1)	601 (34.3)	630 (32.1)	
9	1051 (28.3)	570 (32.5)	481 (24.5)	
Sex				
Male	1916 (51.6)	924 (52.7)	992 (50.6)	0.19
Female	1798 (48.4)	828 (47.3)	970 (49.4)	
Completed education level of father				
No formal or incomplete education	117 (3.3)	46 (2.9)	71 (3.7)	<0.001
Primary education	857 (24.4)	393 (25.0)	464 (23.8)	
Secondary education	1518 (43.2)	771 (49.1)	747 (38.4)	
Pretertiary education	511 (14.5)	202 (12.9)	309 (15.9)	
Tertiary and above	513 (14.6)	158 (10.1)	355 (18.2)	
All	3714 (100)	1962 (100)	1752 (100)	

\*There are 16 missing values for education level of fathers in Singapore and 182 for education level of fathers in Malaysia.

-6 D) and moderate myopia (spherical equivalent in the worse eye < -6 D but at least -2 D) were 1.4% (95% CI 1.0 to 2.1) and 15.0% (95% CI 13.4 to 16.6), respectively, in Singapore, versus 0.5% (95% CI 0.1 to 0.8; p = 0.004) and 3.0% (95% CI 1.8 to 4.3; p < 0.001), respectively, in Malaysia. Myopia prevalences were significantly higher in Singapore than in Malaysia within specific strata defined by ethnicity alone (Chinese, Malays, Indians), age alone, sex alone, paternal education alone and maternal education alone.

The mean (SD) of astigmatic cylinder in the right eye for Malays in Malaysia versus Singapore was (-0.34 D (0.56)) v (-0.83 D (0.88); p < 0.001), that for Chinese (-0.51 D (0.68)) v -0.71 D (0.67); p < 0.001) and that for Indians (-0.46 D (0.72)) v -0.94 D (1.03); p < 0.001).

Overall, the prevalence of astigmatism was higher in Singapore (42.6%) than in Malaysia (22.2%; 95% CI 17.3 to 23.5; p < 0.001; table 3). The rates of astigmatism were higher in Singapore than in Malaysia within specific strata defined

by ethnicity, age, sex, paternal and maternal educational levels.

The overall prevalence of hyperopia was lower in Singapore children (1.7%) than in Malaysian children (2.9%; 95% CI -2.1 to -0.2; p = 0.005; table 4). The hyperopia rates were lower in Singapore than in Malaysia within the strata for 7-year-old children only (p < 0.001) and the strata of males only (p < 0.001), but similar within other strata defined by ethnicity, age, sex, paternal and maternal education levels.

The multivariate analyses show that the rates of myopia are higher in Singapore than in Malaysia for Malays only, Chinese only and Indians only (table 5). Singapore Malays and Indians had significantly different astigmatism rates compared with Malaysian Malays (p < 0.001) and Indians (p = 0.011) after adjusting for other factors including myopia. After adjusting for mother's education instead of father's education, the multivariate ORs for myopia in Singapore versus Malaysia for Malays only, Chinese only

**Table 2** Prevalences of myopia (spherical equivalent at least -0.5 D in either eye) in Malaysia and Singapore

	Malaysia (n = 1752)				Singapore (n = 1962)				Difference (Singapore - Malaysia)		p Value
	No at risk	Cases	Prevalence (%)	95% CI	No at risk	Cases	Prevalence (%)	95% CI	%	95% CI	
Ethnicity											
Malay	1245	114	9.2	7.4 to 10.9	348	77	22.1	17.9 to 26.9	13.0	11.2 to 14.7	<0.001
Chinese	285	88	30.9	23.4 to 38.3	1467	588	40.1	37.6 to 42.6	9.2	1.5 to 16.9	0.03
Indian	152	19	12.5	8.3 to 16.7	126	43	34.1	25.9 to 43.1	21.6	17.4 to 25.9	<0.001
Others	70	14	20.0	11.3 to 28.7	21	4	19.0	5.4 to 41.9	-1.0	-11.6 to 9.7	0.84
Age (years)											
7	581	58	10.0	6.8 to 13.1	851	243	28.6	25.5 to 31.7	18.6	15.4 to 21.7	<0.001
8	601	84	14.0	10.3 to 17.6	630	219	34.8	31.0 to 38.6	20.8	17.2 to 24.4	<0.001
9	570	93	16.3	11.7 to 20.9	481	250	52.0	47.4 to 56.5	35.7	31.0 to 40.3	<0.001
Gender											
Male	924	126	13.6	10.2 to 17.1	992	379	38.2	35.2 to 41.3	24.6	21.1 to 28.0	<0.001
Female	828	109	13.2	10.2 to 16.1	970	333	34.3	31.3 to 37.4	21.2	18.2 to 24.1	<0.001
Completed education level of father											
No formal education or incomplete education	46	4	8.7	1.5 to 15.9	71	18	25.4	15.8 to 37.1	16.7	9.2 to 24.1	0.008
Primary education	393	63	16.0	11.5 to 20.6	464	149	32.1	27.9 to 36.6	16.1	11.5 to 20.7	<0.001
Secondary education	771	91	11.8	8.8 to 14.8	747	268	35.9	32.4 to 39.4	24.1	21.0 to 27.1	<0.001
Pretertiary education	202	20	9.9	4.2 to 15.6	309	117	37.9	32.4 to 43.5	28.0	22.2 to 33.7	<0.001
Tertiary and above	158	30	19.0	13.4 to 24.6	355	153	43.1	37.9 to 48.4	24.1	18.3 to 29.9	<0.001
All	1752	235	13.4	10.8 to 16.0	1962	712	36.3	34.2 to 38.5	22.9	20.3 to 25.4	<0.001

**Table 3** Prevalence rates of astigmatism (cylinder  $\geq 0.75$  D in either eye) in Malaysia and Singapore

	Malaysia (n = 1752)				Singapore (n = 1962)				Difference (Singapore–Malaysia)		p Value
	No at risk	Cases	Prevalence (%)	95% CI	No at risk	Cases	Prevalence (%)	95% CI	(%)	95% CI	
Ethnicity											
Malay	1245	233	18.7	16.4 to 21.1	348	154	44.3	39.0 to 49.6	25.5	23.2 to 27.9	<0.001
Chinese	285	97	34.0	29.2 to 38.9	1467	623	42.5	39.9 to 45.0	8.4	3.4 to 13.4	0.003
Indian	152	34	22.4	15.1 to 29.6	126	52	41.3	32.6 to 50.4	18.9	11.5 to 26.3	<0.001
Others	70	25	35.7	20.7 to 50.7	21	7	33.3	14.6 to 57.0	-2.4	-20.8 to 16.0	0.76
Age (years)											
7	581	135	23.2	18.3 to 28.2	851	376	44.2	40.8 to 47.6	20.9	16.0 to 25.9	<0.001
8	601	144	24.0	20.4 to 27.5	630	267	42.4	38.5 to 46.3	18.4	14.9 to 21.9	<0.001
9	570	110	19.3	15.1 to 23.5	481	193	40.1	35.7 to 44.7	20.8	16.6 to 25.0	<0.001
Sex											
Male	924	206	22.3	19.1 to 25.5	992	441	44.5	41.3 to 47.6	22.2	18.9 to 25.4	<0.001
Female	828	183	22.1	17.7 to 26.5	970	395	40.7	37.6 to 43.9	18.6	14.2 to 23.0	<0.001
Completed education level of father											
No formal or incomplete education	46	8	17.4	4.6 to 30.2	71	39	54.9	42.7 to 66.8	37.5	24.3 to 50.8	<0.001
Primary education	393	107	27.2	23.0 to 31.5	464	191	41.2	36.6 to 45.8	13.9	9.7 to 18.2	<0.001
Secondary education	771	136	17.6	14.7 to 20.6	747	303	40.6	37.0 to 44.2	22.9	20.0 to 25.9	<0.001
Pretertiary education	202	44	21.8	16.3 to 27.3	309	137	44.3	38.7 to 50.1	22.6	17.0 to 28.1	<0.001
Tertiary and above	158	47	29.7	21.1 to 38.4	355	158	44.5	39.3 to 49.8	14.8	5.8 to 23.8	0.005
Myopia											
No	1517	262	17.3	14.6 to 20.0	1250	454	36.3	33.6 to 39.1	19.0	16.4 to 21.7	<0.001
Yes	235	127	54.0	47.3 to 60.8	712	382	53.7	49.9 to 57.4	-0.4	-7.1 to 6.3	0.91
All	1752	389	22.2	19.1 to 25.3	1962	836	42.6	40.4 to 44.8	20.4	17.3 to 23.5	<0.001

and Indians only were 3.57, 1.58 and 3.82, respectively. The multivariate ORs for astigmatism after controlling for mother’s education instead of father’s education for Malays only, Chinese only and Indians only were 3.30, 1.33 and 1.82, respectively.

**DISCUSSION**

The ethnicity-specific prevalences of myopia are higher in Singapore than in Malaysia. Malays are native to both Malaysia and Singapore, the Chinese to both countries migrated from the same parts of southern China, and the Indians to both countries migrated from similar localities in India (southern India) and Sri Lanka several decades ago.

Thus, the intercountry ethnicity-specific differences in myopia prevalences are unlikely to be due to genetic dissimilarities, and may be primarily environmental in nature. For example, most children in Singapore attend preschool (kindergarten or a childcare centre), and the syllabus maybe more structured and vigorous, with a greater use of information technology compared with the Malaysian preschool system, although the evidence is scarce and primarily anecdotal.<sup>19</sup> Differences in urban population density may also be relevant. Singapore is a small, urban city centre (population density of 6026/km<sup>2</sup>) compared with the Gombak District, with a population density of 851/km<sup>2</sup>, based on the 2000 census.<sup>11</sup> In Singapore, 82% of the people live in

**Table 4** Prevalences of hyperopia (spherical equivalent  $\geq +2$  D in either eye) in Malaysia and Singapore children

	Malaysia (n = 1752)				Singapore (n = 1962)				Difference (Singapore–Malaysia)		p Value
	No at risk	Cases	Prevalence (%)	95% CI	No at risk	Cases	Prevalence (%)	95% CI	(%)	95% CI	
Ethnicity											
Malay	1245	36	2.9	1.9 to 3.8	348	12	3.4	1.8 to 5.9	0.6	-0.4 to 1.5	0.29
Chinese	285	5	1.8	0.4 to 3.1	1467	18	1.2	0.7 to 1.9	-0.5	-2.0 to 0.9	0.37
Indian	152	6	3.9	0.4 to 7.5	126	3	2.4	0.5 to 6.8	-1.6	-5.2 to 2.1	0.27
Others	70	3	4.3	2.0 to 6.6	21	1	4.8	0.1 to 23.8	0.5	-2.3 to 3.3	0.71
Age (years)											
7	581	29	5.0	3.0 to 7.0	851	18	2.1	1.3 to 3.3	-2.9	-4.8 to -0.9	<0.001
8	601	12	2.0	0.7 to 3.3	630	12	1.9	1.0 to 3.3	-0.1	-1.4 to 1.2	0.88
9	570	9	1.6	0.4 to 2.8	481	4	0.8	0.2 to 2.1	-0.7	-1.9 to 0.5	0.09
Sex											
Male	924	30	3.2	1.9 to 4.6	992	13	1.3	0.7 to 2.2	-1.9	-3.3 to -0.6	<0.001
Female	828	20	2.4	1.2 to 3.6	970	21	2.2	1.3 to 3.3	-0.3	-1.5 to 1.0	0.66
Completed education level of father											
No formal or incomplete education	46	2	4.3	0.5 to 14.8	71	1	1.4	0.0 to 7.6	-2.9	-9.1 to 3.3	0.10
Primary education	393	13	3.3	1.4 to 5.2	464	11	2.4	1.2 to 4.2	-0.9	-2.8 to 1.0	0.25
Secondary education	771	20	2.6	1.4 to 3.8	747	11	1.5	0.7 to 2.6	-1.1	-2.3 to 0.1	0.02
Pretertiary education	202	3	1.5	0.3 to 4.3	309	5	1.6	0.5 to 3.7	0.1	-1.6 to 1.9	0.88
Tertiary and above	158	5	3.2	0.4 to 5.9	355	6	1.7	0.6 to 3.6	-1.5	-4.3 to 1.4	0.16
All	1752	50	2.9	1.9 to 3.8	1962	34	1.7	1.2 to 2.4	-1.1	-2.1 to -0.2	0.005



**Table 5** Odds ratio of myopia, astigmatism and hyperopia in Singapore and Malaysia, by ethnic group

	Crude OR (95% CI)	p Value	Age-sex-adjusted OR (95% CI)	p Value	Multivariate OR (95% CI)	p Value
<b>Myopia*</b>						
Malay site	2.82 (2.29 to 3.48)	<0.001	2.94 (2.33 to 3.69)	<0.001	3.32 (2.64 to 4.17)	<0.001
(Singapore v Malaysia)						
Chinese site	1.50 (1.04 to 2.15)	0.03	1.73 (1.17 to 2.58)	0.009	1.70 (1.13 to 2.56)	0.013
(Singapore v Malaysia)						
Indian site	3.63 (2.46 to 5.35)	<0.001	3.89 (2.52 to 6.01)	<0.001	5.12 (2.68 to 9.81)	<0.001
(Singapore v Malaysia)						
<b>Astigmatism†</b>						
Malay site	3.45 (2.95 to 4.03)	<0.001	3.45 (2.91 to 4.09)	<0.001	3.47 (2.79 to 4.32)	<0.001
(Singapore v Malaysia)						
Chinese site	1.43 (1.15 to 1.79)	0.003	1.42 (1.13 to 1.77)	0.004	1.26 (0.95 to 1.66)	0.10
(Singapore v Malaysia)						
Indian site	2.44 (1.60 to 3.73)	<0.001	2.41 (1.55 to 3.75)	<0.001	1.90 (1.18 to 3.08)	0.011
(Singapore v Malaysia)						
<b>Hyperopia*</b>						
Malay site	1.20 (0.85 to 1.69)	0.29	1.12 (0.79 to 1.58)	0.51	1.13 (0.76 to 1.69)	0.54
(Singapore v Malaysia)						
Chinese site	0.70 (0.30 to 1.60)	0.37	0.60 (0.26 to 1.41)	0.23	0.52 (0.23 to 1.16)	0.11
(Singapore v Malaysia)						
Indian site	0.59 (0.23 to 1.55)	0.27	0.53 (0.19 to 1.50)	0.22	0.73 (0.12 to 4.33)	0.71
(Singapore v Malaysia)						

\*For models with myopia and hyperopia as the dependent variable, adjusted for sex, age and father's completed education level.

†For models with astigmatism as the dependent variable, adjusted for sex, age, father's completed education level and myopia.

housing development board apartments, whereas most Malaysians live in private houses.<sup>10</sup> The per capita gross domestic product of Singapore is \$24 040 compared with \$9120 in Malaysia; hence, people with certain characteristics may tend to migrate to a relatively more prosperous Singapore.

The environmental hypothesis is also supported by comparisons of myopia prevalences among 7–9-year-old Chinese children in the two mainland China RESC studies: 18.4% in urban Guangzhou in southern China and 8.7% in semirural Shunyi District in northern China.<sup>5–20</sup> Prevalences of myopia in the RESC studies of 7–9-year-old Indian children in both urban and rural India were relatively low: 4.6% and 3.9%, respectively.<sup>7–21</sup>

Astigmatism is more common in Singapore Malays compared with Malaysian Malays, even after adjusting for age, sex, completed education level of fathers and myopia status. Evidence gathered from the RESC studies suggest that country-specific differences in astigmatism are not accounted for solely by differences in the prevalence of myopia. Other factors must come into play, although little is known about specific environmental risk factors for astigmatism. The prevalence of astigmatism in 7–9-year-old children in the RESC studies is as follows: Guangzhou, China (41.6%), Shunyi District, China (13.9%), urban India (14.6%) and rural India (8.4%).<sup>5–7–20–21</sup>

The prevalences of hyperopia are low in both Malaysia (2.9%) and Singapore (1.7%). Comparatively, the prevalence of hyperopia in 7–9-year olds is higher in Guangzhou, China (5.1%), Shunyi District, China (4.2%) and urban India (8.0%).<sup>5–20–21</sup> The prevalence of hyperopia is lowest in rural India (0.3%).<sup>7</sup>

Two features were particularly advantageous in this direct comparative evaluation of refractive errors from two separate studies in Malaysia and Singapore: similar school-going populations and essentially identical measurement methods. Firstly, the multi-ethnic similarity between neighbouring Malaysia and Singapore (Malays, Chinese and Indians), each with a common genetic origin, allowed us to better understand the relative influences of genes and environment. Secondly, the use of autorefraction for measurement of refractive error under cycloplegia induced with 1% cyclopentolate in both studies reduced the possibility of differences introduced by systematic measurement bias. Although a

handheld autorefractor was used for measurement in Malaysia and a table-mounted autorefractor in Singapore, the spherical equivalent readings using the handheld autorefractor were more negative compared with those using the table-mounted autorefractor.<sup>22–23</sup> In our study, because the handheld autorefractor readings in Malaysia may tend to deviate towards more negative readings, measurement biases may contribute to smaller perceived differences in the prevalences of myopia across countries.

It is recognised, however, that our comparisons may not necessarily apply to all children in Singapore or Malaysia. The three schools in Singapore may not be entirely representative of Singapore school children, nor is there evidence that children in Gombak District are representative of Malaysian children. Selection bias may occur because the response rate in the Eastern school (among the top 20 schools) in Singapore was only 49.1%: the overall rates of myopia in the Singapore sample could be underestimated. Perhaps parents of myopic children in Singapore may perceive that their child will not benefit from the study and may not enrol their children. The multivariate regression models show that the differences in myopia rates in Malaysia and Singapore cannot be completely explained by differences in parental educational level, and other environmental factors may contribute to the observed differences. The lack of near work activity, preschool activity, height or parental myopia from both countries may preclude definitive conclusions about the nature of country-specific differences. Future studies could investigate specific risk factors for differences in myopia rates across countries.

Nonetheless, we conclude that the prevalences of myopia in 7–9-year-old children are higher in Singapore Chinese than in Malaysian Chinese, higher in Singapore Indians than in Malaysian Indians and higher in Singapore Malays than in Malaysian Malays.

#### Authors' affiliations

S-M Saw, A Shankar, Department of Community, Occupational and Family Medicine, National University of Singapore, Singapore  
A Cheng, D T H Tan, Singapore Eye Research Institute, Singapore  
P-P Goh, Department of Ophthalmology, Hospital Selayang, Selangor, Malaysia

**L B Ellwein**, National Eye Institute, National Institutes of Health, Bethesda, Maryland, USA

Funding: This study was supported by the National Medical Research Council (NMRC), NMRC/0975/2005, Singapore; the World Health Organization under National Institutes of Health contract N01-EY-2103; and the Ministry of Health Malaysia under Major Research Grant 2003/13.

Competing interests: None.

## REFERENCES

- Muti DO**, Bullimore MA. Myopia: an epidemic of possibilities? *Optom Vis Sci* 1999;**76**:257–8.
- Fan DS**, Lam DS, Lam RF, *et al.* Prevalence, incidence, and progression of myopia of school children in Hong Kong. *Invest Ophthalmol Vis Sci* 2004;**45**:1071–5.
- Lin LL**, Shih YF, Tsai CB, *et al.* Epidemiologic study of ocular refraction among schoolchildren in Taiwan in 1995. *Optom Vis Sci* 1999;**76**:275–81.
- Wu HM**, Seet B, Yap EP, *et al.* Does education explain ethnic differences in myopia prevalence? A population-based study of young adult males in Singapore. *Optom Vis Sci* 2001;**78**:234–9.
- He M**, Zeng J, Liu Y, *et al.* Refractive error and visual impairment in urban children in Southern China. *Invest Ophthalmol Vis Sci* 2004;**45**:793–9.
- Pokharel GP**, Negrel AD, Munoz SR, *et al.* Refractive Error Study in Children: results from Mechi Zone, Nepal. *Am J Ophthalmol* 2000;**129**:436–44.
- Dandona R**, Dandona L, Srinivas M, *et al.* Refractive error in children in a rural population in India. *Invest Ophthalmol Vis Sci* 2002;**43**:615–22.
- Muti DO**, Mitchell GL, Moeschberger ML, *et al.* Parental myopia, near work, school achievement, and children's refractive error. *Invest Ophthalmol Vis Sci* 2002;**43**:3633–40.
- Klein AP**, Duggal P, Lee KE, *et al.* Support for polygenic influences on ocular refractive error. *Invest Ophthalmol Vis Sci* 2005;**46**:442–6.
- Singapore Population**. *Census 2000*. Singapore: Singapore Department of Statistics, 2001.
- Department of Statistics Malaysia**. *Preliminary count report. Population and housing census of Malaysia 2000*. Kuala Lumpur: Department of Statistics Publications, 2001.
- Goh PP**, Abgariyah Y, Pokharel GP, *et al.* Refractive error and visual impairment in school-age children in Gombak District, Malaysia. *Ophthalmology* 2005;**112**:678–85.
- Saw SM**, Hong CY, Chia KS, *et al.* Nearwork and myopia in young children. *Lancet* 2001;**357**:390.
- Saw SM**, Chua WH, Hong CY, *et al.* Nearwork in early-onset myopia. *Invest Ophthalmol Vis Sci* 2002;**43**:332–9.
- Saw SM**, Tong L, Chua WH, *et al.* Incidence and progression of myopia in Singaporean school children. *Invest Ophthalmol Vis Sci* 2005;**46**:51–7.
- Tong L**, Saw SM, Carkeet A, *et al.* Prevalence rates and epidemiological risk factors for astigmatism in Singapore school children. *Optom Vis Sci* 2002;**79**:606–13.
- Negrel AD**, Maul E, Pokharel GP, *et al.* Refractive Error Study in Children: sampling and measurement methods for a multi-country survey. *Am J Ophthalmol* 2000;**129**:421–6.
- StataCorp**. *Stata Statistical Software: Release 8.0*. College Station, TX: Stata Corporation, 2003.
- Nurturing early learners**. *A framework for a kindergarten curriculum in Singapore*. Singapore: Preschool Education Unit, Ministry of Education, [http://www.moe.gov.sg/preschooleducation/curriculum\\_framework.htm](http://www.moe.gov.sg/preschooleducation/curriculum_framework.htm) (accessed 1 Aug 2006).
- Zhao J**, Pan X, Sui R, *et al.* Refractive Error Study in Children: results from Shunyi District, China. *Am J Ophthalmol* 2000;**129**:427–35.
- Murthy GV**, Gupta SK, Ellwein LB, *et al.* Refractive error in children in an urban population in New Delhi. *Invest Ophthalmol Vis Sci* 2002;**43**:623–31.
- Kallay OP**, Cordonnier MJ, Dramaix MM. Cycloplegic refractive errors in children: comparison of a standard and a hand-held refractor. *Strabismus* 1998;**6**:3–7.
- Farook M**, Venkatramani J, Gazzard G, *et al.* Comparisons of the hand-held autorefractor, table-mounted autorefractor and subjective refraction in Singapore adults. *Optom Vis Sci* 2005;**82**:1066–70.