

Endocrine Disrupting Contaminants—Beyond the Dogma

Louis J. Guillette Jr.

Department of Zoology, University of Florida, Gainesville, Florida, USA

Descriptions of endocrine disruption have largely been associated with wildlife and driven by observations documenting estrogenic, androgenic, antiandrogenic, and antithyroid actions. These actions, in response to exposure to ecologically relevant concentrations of various environmental contaminants, have now been established in numerous vertebrate species. However, many potential mechanisms and endocrine actions have not been studied. For example, the DDT [1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)ethane] metabolite, *p,p'*-DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)ethylene] is known to disrupt prostaglandin synthesis in the uterus of birds, providing part of the explanation for DDT-induced egg shell thinning. Few studies have examined prostaglandin synthesis as a target for endocrine disruption, yet these hormones are active in reproduction, immune responses, and cardiovascular physiology. Future studies must broaden the basic science approach to endocrine disruption, thereby expanding the mechanisms and endocrine end points examined. This goal should be accomplished even if the primary influence and funding continue to emphasize a narrower approach based on regulatory needs. Without this broader approach, research into endocrine disruption will become dominated by a narrow dogma, focusing on a few end points and mechanisms. **Key words:** adrenal steroidogenesis, endocrine disruption, estrogens, hormone metabolism, nitrate, progesterone, prostaglandins. *Environ Health Perspect* 114(suppl 1):9–12 (2006). doi:10.1289/ehp.8045 available via <http://dx.doi.org/> [Online 21 October 2005]

In 1991 a hypothesis was formulated to suggest that numerous xenobiotic chemicals used in everyday commerce or natural chemicals released into the environment by human activity had the potential to disrupt the endocrine system of wildlife and humans at ecologically relevant concentrations. This hypothesis has become known as the endocrine-disrupting contaminants (EDCs) hypothesis (Colborn and Clement 1992). Since the presentation of this original hypothesis more than a decade ago, significant scientific and public debates have raged concerning its validity. Much of the early work was driven by wildlife observations that documented *a*) estrogenic, androgenic, antiandrogenic, and antithyroid actions in fish found below outfalls of sewage or paper pulp mills; *b*) exposure of alligators to agricultural chemicals; and *c*) exposure of birds, fish, and mammals to complex mixtures of chemicals in the Laurentian Great Lakes of North America or the Baltic Sea of Northern Europe [for representative reviews, see Crain et al. (2000); Fox (1992); Guillette and Gunderson (2001); Tyler et al. (1998)]. A growing literature supports these early observations and has extended them, thus supporting the hypothesis that various chemicals alter endocrine function in a wide array of vertebrates and invertebrates (Gray et al. 2002; Iguchi et al. 2001; Markey et al. 2002; Oberdörster and Cheek 2000). The questions being addressed today are not whether endocrine disruption occurs because of contaminant exposure but rather at what concentrations does it occur? Or, what is the mechanism of action driving the response? Or, is the response observed adverse at the population level?

Major debates in the scientific and public arena continue to arise around the issues of EDC-induced human health effects, especially breast cancer, human semen quality, and birth defects of the genitalia and reproductive system (McLachlan 2001; National Academy of Sciences 1999). As with wildlife, a number of studies report an association between contaminant exposure and alterations in the development and functioning of the male reproductive system (Sharpe and Irvine 2004; Toppari et al. 1996). However, numerous studies also document the difficulty of establishing the link between exposure and health outcomes in human populations [for discussion, see Birnbaum and Fenton (2003); Carpenter et al. (2002); Harvey and Johnson (2002); Safe et al. (2002)]. Thus, many studies resort to examining mechanisms in model species or tissues and then relating this information to human populations (e.g., Mori et al. 2003). This approach was dramatically supported recently, when a study of baby boys revealed that end points used in rodent studies were useful in examining the developmental biology of humans. Swan et al. (2005) demonstrated that anogenital distance, a frequently used end point in rodent developmental toxicology studies, decreased significantly with increasing environmental exposure to phthalates, as reported in prenatally exposed rodents. In short, as we have learned from drug exposure studies, many end points in rodents can predict adverse outcomes in humans.

Estrogens and estrogenic actions have dominated the EDC literature since the early 1990s. The scientific literature associated with understanding the legacy of diethylstilbestrol (DES)

exposure during pregnancy has been a prominent model (Bern 1992). That is, the scientific community in collaboration with DES advocacy groups composed of mothers and exposed children has studied the effects of embryonic and neonatal exposure to a potent synthetic estrogen. These data, from affected human populations and laboratory rodent model systems, have provided much of the basic scientific literature for arguing adverse health effects in human populations exposed to EDCs (Markey et al. 2003; McLachlan 2001). This dependence on documented adverse human health effects and laboratory rodent models has led to a vast majority of EDC research focused on estrogenic actions, especially those associated with estrogen receptor-mediated actions. This work has been reinforced by wildlife studies using estrogenic end points such as yolk production in male fish (Sumpter and Jobling 1995). Additional studies that demonstrate the antiandrogenic actions of *p,p'*-DDE [1,1-dichloro-2,2-bis(*p*-chlorophenyl)ethylene], the major bioaccumulated metabolite of DDT [1,1,1-trichloro-2,2-bis(*p*-chlorophenyl)ethane], and various fungicides that act on the androgen receptor (Gray et al. 2001) have supported the perception that EDC action primarily produces an “estrogenic” feminizing or demasculinizing response. Discussion of the antithyroidal actions of polychlorinated biphenyls (PCBs) and polybrominated diphenyl ethers (PBDEs) has broadened the research in EDCs, but current approaches focus almost exclusively on the neurological or neuroendocrine actions of these compounds in developing embryos (Rolland 2000; Zoeller et al. 2002). All these actions are worth study, but the focus on estrogenic, antiestrogenic,

This article is part of the monograph “The Ecological Relevance of Chemically Induced Endocrine Disruption in Wildlife.”

Address correspondence to L.J. Guillette Jr., Department of Zoology, 223 Bartram Hall – Box 118525, University of Florida, Gainesville, FL 32611 USA. Telephone: 352-392-1098. Fax: 352-392-3704. E-mail: ljjg@zoo.ufl.edu

I thank T. Iguchi, J. McLachlan, E. Orlando, D. Pickford, and my graduate students T. Edwards, B. Moore, M. Milnes, and M. Gunderson for stimulating discussions that led to this commentary.

The research described from my laboratory was supported by grants from the U.S. Environmental Protection Agency, the National Institute of Environmental Health Sciences, and the Marisla Foundation

The authors declare they have no competing financial interests.

Received 31 January 2005; accepted 22 August 2005.

and antithyroidal outcomes has produced a dogma that EDC research is largely about receptor binding or mechanisms that are largely based on sex steroid or thyroid action. From the perspective of regulatory science, a focus on one or two mechanisms is cost- and time-effective, and given the economic and regulatory issues surrounding EDC research, a “realistic” approach (Ankley et al. 1997; Kavlock et al. 1996). But from the perspective of advancing the science, this limited view has had a restrictive effect. Widening this view is essential if further advances in our understanding are to occur.

In this article, I suggest some areas where future research is needed to broaden our understanding of endocrine disruption by environmental contaminants. This article is by no means a review but rather a short outline of areas where recent data suggest that future work could produce insight into mechanisms of endocrine disruption.

Estrogens: Steroid Actions beyond Receptor Binding

Although many contaminant-induced estrogenic actions described to date appear to be caused by receptor-mediated interactions, other responses are less clear (McLachlan 2001). Several studies have provided support for the hypothesis that estrogenic actions could be due to increased aromatase activity. For example, the feminizing effect of atrazine on some species has been hypothesized to be caused by its ability to alter aromatase-induced conversion of androgens to estrogens (Crain et al. 1997; Hayes et al. 2003). Several studies with mammalian or fish cell lines suggest that such a mechanism is possible (Sanderson et al. 2000, 2001; Sanderson and van den Berg 2003). Future studies must consider a mechanism by which contaminants alter steroid action not by receptor binding but rather via alterations in the synthesis of these hormones by modification in the enzymatic pathways, either by direct changes in enzyme synthesis or in enzyme activity levels (i.e., alterations in available cofactors) [for discussion see Harvey and Johnson (2002); Sanderson and van den Berg (2003)].

Glucocorticoids and Progestins: Steroid Actions beyond Estrogens and Androgens

Before the presentation of the EDC hypothesis, a metabolite of DDT, *o,p'*-DDD [1,1-dichloro-2,2-bis(4-chlorophenyl)ethane], was known to disrupt adrenal steroidogenesis in some mammalian species, including humans (Benecke et al. 1991; Ruppert and Kraft 1999). In fact, this compound has been marketed as a drug to treat adrenal cancer and Cushing's disease because of its suppressive action on adrenal steroidogenesis (Benecke et al. 1991). Even with this knowledge, few investigators have examined wildlife species or humans for alterations in

adrenal function in areas where DDT use is still extensive, as in many tropical regions with a high risk of malaria. Not all species will respond to this compound with altered adrenal steroidogenesis (Breuner et al. 2000), but future studies on a wide array of species are needed.

Furthermore, because of the action of *o,p'*-DDD in humans, that it is a major metabolite of DDT, and that DDT is still used extensively in tropical regions, more studies are needed that examine human populations for adrenal-based end points, especially pregnant women. At least one recent study has suggested that coincident with the use of DDT and elevated DDE plasma levels, a significant increase in human preterm parturition occurred (Longnecker et al. 2001). It is now commonly accepted that the hypothalamo-pituitary-adrenal axis is central to the control of gestation length and the timing of birth in humans, and probably in eutherian mammals in general (Chrousos et al. 1998). Alterations in adrenal steroidogenesis and in glucocorticoid feedback to the hypothalamo-pituitary axis are associated with disrupted gestation and birth, which suggests that chemicals capable of acting on this system are potentially important in human reproductive health.

Progesterone and related progestins are another group of steroid hormones with a central role in gestation maintenance in every vertebrate species studied to date (Jones and Baxter 1991). These hormones are essential for oocyte maturation as well. Recent studies identified a receptor for these steroids in the cell membrane (Zhu et al. 2003a, 2003b), in addition to the traditional perinuclear receptor, which acts as a transcription factor (Carson-Jurica et al. 1990). Some species can have more than one form of the perinuclear receptor with unique tissue distributions (Custodia-Lora and Callard 2002). Very few studies have examined the action of environmental contaminants on progestin-based phenomena. Various contaminants can bind the nuclear progesterone receptor (aPR) from the American alligator (Vonier et al. 1996). Although many of the chemicals tested, including most of the DDT metabolites, did not reduce [³H]R5020 binding to aPR, 2,2-bis(*p*-chlorophenyl)ethanol (DDOH, a DDT metabolite), endosulfan, alachlor, dicofol, atrazine, and kepone did (Vonier et al. 1996). Pickford and Morris (1999) have demonstrated that methoxychlor disrupts progesterone-induced oocyte maturation in the frog *Xenopus leavis*. This action appears to be independent of the nuclear receptor. Progesterone plays a major role in gestation maintenance; thus, a disruption in progesterone signaling could also affect the frequency of spontaneous miscarriage, as described above. In short, we need to broaden our concepts of the mechanisms and end points used to assess steroid mimicry by environmental contaminants.

Prostaglandins and Uterine Functioning: the Past and the Future?

Dramatic declines in the populations of numerous bird species occurred during the decades following the wide-scale use of DDT for insect control in the United States and other countries around the world. Upon examination, DDE, the major metabolite of DDT was associated with altering the physiological process of eggshell formation that subsequently led to eggshell thinning and population declines of numerous avian species, particularly raptors and shorebirds [see review by Lundholm (1997)]. These studies indicate that the supply of calcium to the eggshell gland is not impeded by DDE, but rather this organochlorine contaminant disrupts calcium transport within the eggshell gland (Lundholm 1997). In addition, prostaglandins (PGs) have been implicated in eggshell thinning because DDE disrupts PG synthesis, which reduces bicarbonate transport in the duck shell gland lumen, thereby reducing calcium transport.

Given the important roles of various PGs during ovulation, pregnancy, and parturition in a wide array of vertebrate species (Guillette et al. 1991; Jenkin and Young 2004), it is surprising that few investigators have examined the disruption of this important class of hormones. PG synthesis, like steroidogenesis, is controlled by a pathway of enzymes (Helliwell et al. 2004a). Steroids, particularly estrogens, regulate the expression of a number of these enzymes. Thus, with a wide range of chemicals capable of altering the “estrogenic milieu,” it could be hypothesized that PG synthesis is altered in the reproductive system of any vertebrate, as described previously for the avian reproductive tract. Interestingly, recent studies have suggested that PGs could work through perinuclear receptors as well as through the established membrane-G protein receptors (Helliwell et al. 2004b). If this is the case, disruption at the level of transcription could occur via several mechanisms leading to different outcomes. Finally, it is important to recognize that PGs are not just reproductive hormones. Rather, they play essential roles in the immune response, including inflammation and arthritis, cardiovascular regulation encompassing hypertension, and respiration including asthma. These are just some of the roles that ubiquitous PG hormones play in vertebrates. Moreover, PGs are evolutionarily ancient and are important in reproduction and development in invertebrates as well (Stanley and Howard 1998). In brief, a major research effort is needed to examine the possible linkages between the wide-ranging alterations seen in organisms living in contaminated environments and alterations in prostaglandin synthesis.

Hepatic Metabolism of Hormones: Changing the Hormonal Milieu

Many vertebrates exhibit species-specific, sexually dimorphic patterns of hepatic enzyme activity that appear to be regulated by sex steroids and/or growth hormone (Gustafsson 1994). In humans and rodents, 50% or more of the drugs and pesticides currently used induce the expression of the hepatic enzyme cytochrome P450 (CYP)3A (Gibson et al. 2002). Numerous steroids, including testosterone, 17 β -estradiol, progesterone, and androstenedione, are metabolized by CYP3A (Gibson et al. 2002). It is therefore conceivable that wildlife or humans exposed to contaminated environments would exhibit elevated CYP3A activity that leads to increased clearance of such steroids as testosterone and 17 β -estradiol from the plasma, as reported in chickens (Chen et al. 1993, 1994) and humans (Bammel et al. 1992).

Blumberg et al. (1998) first proposed the "steroid sensor hypothesis" by suggesting that the induction of hepatic enzymes responsible for the metabolism of endogenous and exogenous compounds was regulated by a nuclear receptor with broad specificity. A group of nuclear receptors (e.g., steroid xenobiotic receptor, SXR) was discovered, and this hypothesis has been further supported by studies that have observed a positive relationship between SXR expression and CYP3A expression (Blumberg and Evans 1998). Transcription of CYP3A, as well as of other hepatic enzymes involved in the biotransformation of testosterone, pharmaceutical agents, and xenobiotics, appears to be regulated, at least in part, by SXR and related nuclear transcription factors (Blumberg et al. 1998; Xie et al. 2000). Differences in the activity of the hepatic enzymes responsible for the biotransformation of sex steroids exist among alligators collected from contaminated and relatively uncontaminated sites in Florida (Gunderson et al. 2001, 2004). We have hypothesized that differences in plasma sex steroid concentrations could be caused, in part, by differences in hepatic clearance of these sex steroids, as has been demonstrated in other species (Wilson and LeBlanc 1998, 2000). Future studies need to consider the complex interactions among contaminant exposure, hepatic enzyme induction, and regulation of circulating hormone concentrations (e.g., Tabb et al. 2004) because these issues are not separate but are, in fact, tightly integrated.

It has been observed relatively recently that the promoter region of the human gene for inducible nitric oxide (NO) synthase (iNOS) has response elements for the constitutive androstane receptor (CAR) and the pregnane X receptor (PXR). These orphan receptors and transcription factors are related to SXR [see above and Toell et al. (2002)]. Specifically,

PXR activation by xenobiotics appears to up-regulate iNOS and subsequently increases NO synthesis. Inflammation and apoptosis are phenomena regulated by NO, as are various actions of the hypothalamo-pituitary-gonad axis. Interestingly, xenobiotic effects along this axis could be explained by modulations in iNOS that lead to altered hormonal action of NO. However, this explanation is not the only possible source of elevated NO because nitrate pollutants could also alter homeostasis in NO synthesis (Guillette and Edwards 2005).

Novel Mechanisms of Endocrine Disruption

Nitrates: an endocrine disruptor? It appears that nitrate, a global pollutant of most aquatic systems, has the potential to be an endocrine-disrupting contaminant (Guillette and Edwards 2005). Nitrate (NO₃) and nitrite (NO₂) have known physiological influences (Jensen 2003; Levallois and Phaneuf 1994; Zrally et al. 1997). Nitrate and nitrite have been reported for decades to be toxic in humans and animals (Avery 1999). As early as 1945, methemoglobinemia was associated with drinking nitrate-contaminated well water on farms from the Midwest United States. Those concerns as well as new ones continue today with increasing contamination of groundwater with nitrates (Avery 1999; Porter et al. 1999). Nitrates have been implicated as a disruptor of gonadal steroidogenesis (Panesar 1999; Panesar and Chan 2000; Zrally et al. 1997) and thyroid function (Kursa et al. 2000; Lahti et al. 1985; Wyngaarden et al. 1952, 1953). A recent retrospective study of our work on alligators from various lakes in Florida suggests that nitrate could contribute to some of the altered endocrine parameters previously reported in juvenile animals (Guillette and Edwards 2005). Nitrate could alter steroidogenesis by *a*) conversion to nitrite and nitric oxide in the mitochondria (Meyer 1995; Nohl et al. 2001, 2000; Zweier et al. 1999), which is the site of initial steroid synthesis (Stocco and Clark 1996); *b*) altering Cl⁻ ion concentrations in the cell by substituting for Cl⁻ in the membrane transport pump (Alrefai et al. 2001; Panesar 1999); or *c*) binding to the heme region of various cytochrome P450 enzymes associated with steroidogenesis and altering enzymatic action (Danielson 2002; White et al. 1987). More research is needed to examine the possible endocrine disruptive action of this ubiquitous pollutant.

Novel research systems in endocrine disruption. Several exciting studies have expanded our understanding of systems capable of endocrine disruption, such as the studies documenting contaminant-induced disruption of the interspecies signaling system associated with the nitrogen fixation pathway (Fox et al. 2001), thyroid hormone-dependent gene expression during metamorphosis in amphibians (Crump

et al. 2002) or pheromonal communication among salamanders (Park et al. 2001; Park and Propper 2002). These examples are just a few of the natural systems that require further research efforts. Although regulatory biology and most research funding will continue to focus on a few model systems and mechanisms, a broad basic research agenda is needed if we are to move the field forward. From my perspective, research in endocrine disruption represents less a new field, but rather more an integration of many fields, including biology, chemistry, epidemiology, and atmospheric and earth sciences. It is the new synthesis of science by which we will begin to understand the complexity of the world around us and our impact on it. New understanding of the effect of contamination has been gained because comparative endocrinologists, developmental biologists, ecologists and evolutionary biologists have exchanged ideas, knowledge, and hypotheses with physicians, epidemiologists, toxicologists and biochemists. Understanding the complex interactions between the plethora of environmental contaminants and the organisms living in contaminated environments will require many approaches. The most important is a broad, creative approach to the science and its interpretation.

REFERENCES

- Alrefai WA, Ramaswamy K, Dudeja PK. 2001. Mechanism(s) of chloride transport in human distal colonic apical membrane vesicles. *Dig Dis Sci* 46:2209–2218.
- Ankley GT, Johnson RD, Detenbeck NE, Bradbury SP, Toth G, Folmar L. 1997. Development of a research strategy for assessing the ecological risk of endocrine disruptors. *Rev Toxicol* 1:71–106.
- Avery AA. 1999. Infantile methemoglobinemia: reexamining the role of drinking water nitrates. *Environ Health Perspect* 107:583–586.
- Bammel A, Vandermeer K, Ohnhaas EE, Kirch W. 1992. Divergent effects of different enzyme-inducing agents on endogenous and exogenous testosterone. *Eur J Clin Pharmacol* 42:641–644.
- Benecke R, Keller E, Vetter B, De Zeeuw RA. 1991. Plasma level monitoring of mitotane (*o,p*-DDD) and its metabolite (*o,p*-DDE) during long-term treatment of Cushing's disease with low doses. *Eur J Clin Pharmacol* 14:421–426.
- Bern H. 1992. The fragile fetus. In: *Chemically-Induced Alterations in Sexual and Functional Development: The Wildlife/Human Connection* (Colborn T, Clement C, eds). Princeton, NJ:Princeton Scientific Publications, 9–15.
- Birnbaum LS, Fenton SE. 2003. Cancer and developmental exposure to endocrine disruptors. *Environ Health Perspect* 111:389–394.
- Blumberg B, Evans RM. 1998. Orphan nuclear receptors—new ligands and new possibilities. *Gene Develop* 12:3149–3155.
- Blumberg B, Sabbagh W, Juguilon H, Bolado J, van Meter CM, Ono ES, et al. 1998. SXR, a novel steroid and xenobiotic-sensing nuclear receptor. *Genes Dev* 12:3195–3205.
- Breuner CW, Jennings DH, Moore MC, Orchinik M. 2000. Pharmacological adrenalectomy with mitotane. *Gen Comp Endocrinol* 120:27–34.
- Carpenter, DO, Arcaro, K, Spink, DC. 2002. Understanding the human health effects of chemical mixtures. *Environ Health Perspect* 110(suppl 1): 25–42.
- Carson-Jurica MA, Schrader WT, O'Malley BW. 1990. Steroid receptor family. *Endocr Rev* 11:201–220.
- Chen S-W, Dzuik PJ, Francis BM. 1994. Effect of four environmental toxicants on plasma Ca and estradiol 17 beta and hepatic P450 in laying hens. *Environ Toxicol Chem* 13:789–796.
- Chen S-W, Francis BM, Dzuik PJ. 1993. Effect of concentration of mixed-function oxidase on concentration of estrogen, rate of egg lay, eggshell thickness, and plasma calcium in laying hens. *J Anim Sci* 71:2700–2707.

- Chrousos GP, Torpy DJ, Gold PW. 1998. Interactions between the hypothalamic-pituitary-adrenal axis and the female reproductive system: clinical implications. *Ann Intern Med* 129:229–240.
- Colborn T, Clement C, eds. 1992. Chemically-Induced Alterations in Sexual and Functional Development: The Wildlife/Human Connection. Princeton, NJ:Princeton Scientific Publications
- Crain DA, Guillette LJ Jr, Rooney AA, Pickford DB. 1997. Alteration in steroidogenesis in alligators (*Alligator mississippiensis*) exposed naturally and experimentally to environmental contaminants. *Environ Health Perspect* 105:528–533.
- Crain DA, Rooney AA, Orlando EF, Guillette LJ Jr. 2000. Endocrine-disrupting contaminants and hormone dynamics: lessons from wildlife. In: *Endocrine-Disrupting Contaminants: An Evolutionary Perspective* (Guillette LJ Jr, Crain DA, eds). Philadelphia:Francis and Taylor, 1–21.
- Crump D, Werry K, Veldhoen N, Van Aggelen G, Helbing CC. 2002. Exposure to the herbicide acetochlor alters thyroid hormone-dependent gene expression and metamorphosis in *Xenopus laevis*. *Environ Health Perspect* 110:1199–1205.
- Custodia-Lora N, Callard IP. 2002. Progesterone and progesterone receptors in reptiles. *Gen Comp Endocrinol* 127:1–7.
- Danielson PB. 2002. The cytochrome P450 superfamily: biochemistry, evolution, and drug metabolism in humans. *Curr Drug Metab* 3:561–597.
- Fox GA. 1992. Epidemiological and pathological evidence of contaminant-induced alterations in sexual development in free-living wildlife. In: *Chemically-Induced Alterations in Sexual and Functional Development: The Wildlife/Human Connection* (Colborn T, Clement C, eds). Princeton, NJ:Princeton Scientific Publications, 147–158.
- Fox JE, Starcevic M, Kow KY, Burow ME, McLachlan JA. 2001. Nitrogen fixation: endocrine disruptors and flavonoid signaling. *Nature* 413:128–129.
- Gibson GG, Plant NJ, Swales KE, Ayrton A, El-Sankary W. 2002. Receptor-dependent transcriptional activation of cytochrome P4503A genes: induction mechanisms, species differences, and interindividual variation in man. *Xenobiotica* 32:165–206.
- Gray LE Jr, Ostby J, Furr J, Wolf C, Lambright C, Parks L, et al. 2001. Effects of environmental antiandrogens on reproductive development in experimental animals. *Hum Reprod Update* 7:248–264.
- Gray LE Jr, Parks L, Lambright C, Orlando EF, Guillette LJ Jr. 2002. The effects of endocrine-disrupting chemicals on the reproduction and development of wildlife. In: *Endocrine Disruptors in the Environment* (Metzler M, ed). Heidelberg:Springer-Verlag, 209–248.
- Guillette LJ Jr, Dubois DH, Cree A. 1991. Prostaglandins, oviducal function, and parturient behavior in nonmammalian vertebrates. *Am J Physiol* 260:R854–R861.
- Guillette LJ Jr, Edwards T. 2005. Is nitrate an ecologically relevant endocrine disruptor in vertebrates? *Integr Comp Biol* 45:19–27.
- Guillette LJ Jr, Gunderson MP. 2001. Alterations in the development of the reproductive and endocrine systems of wildlife exposed to endocrine disrupting contaminants. *Reproduction* 122:857–864.
- Gunderson MP, LeBlanc GA, Guillette LJ Jr. 2001. Alterations in sexually dimorphic biotransformation of testosterone in juvenile American alligators (*Alligator mississippiensis*) from contaminated lakes. *Environ Health Perspect* 109:1257–1264.
- Gunderson MP, Oberdörster E, Guillette LJ Jr. 2004. EROD, MROD, and GST activities in juvenile alligators collected from 3 sites in the Kissimmee-Everglades drainage, Florida (USA). *Comp Biochem Physiol C Toxicol Pharmacol* 139:39–46.
- Gustafsson J-A. 1994. Regulation of sexual dimorphism in rat liver. In: *The Differences Between the Sexes* (Short RV, Balaban E, eds). Cambridge, UK:Cambridge University Press, 231–241.
- Harvey PW, Johnson I. 2002. Approaches to the assessment of toxicity data with endpoints related to endocrine disruption. *J Appl Toxicol* 22:241–247.
- Hayes TB, Haston K, Tsui M, Hoang A, Haeffele C, Vonk A. 2003. Atrazine-induced hermaphroditism at 0.1 ppb in American Leopard frogs (*Rana pipiens*): laboratory and field evidence. *Environ Health Perspect* 111:568–575.
- Helliwell RJA, Adams LF, Mitchell MD. 2004a. Prostaglandin synthases: recent developments and a novel hypothesis. *Prostaglandins Leukot Essent Fatty Acids* 70:101–113.
- Helliwell RJA, Berry EBE, O'Carroll SJ, Mitchell MD. 2004b. Nuclear prostaglandin receptors: role in pregnancy and parturition? *Prostaglandins Leukot Essent Fatty Acids* 70:149–165.
- Iguchi T, Watanabe H, Katsu Y. 2001. Developmental effects of estrogenic agents on mice, fish, and frogs: a mini-review. *Horm Behav* 40:248–251.
- Jenkin G, Young IR. 2004. Mechanisms responsible for parturition; the use of experimental models. *Anim Repro Sci* 82–83:567–581.
- Jensen F. 2003. Nitrite disrupts multiple physiological functions in aquatic animals. *Comp Biochem Physiol A Mol Integr Physiol* 135:9–24.
- Jones RE, Baxter DC. 1991. Gestation, with emphasis on corpus luteum biology, placentation and parturition. In: *Vertebrate Endocrinology: Fundamental and Biomedical Implications* (Pang PKT, Schreiber MP, Jones RE, eds). New York:Academic Press, 205–302.
- Kavlock RJ, Daston GP, DeRosa D, Fenner-Crisp P, Gray LE, Kaattari S, et al. 1996. Research needs for the risk assessment of health and environmental effects of endocrine disruptors: a report of the U.S. EPA-sponsored workshop. *Environ Health Perspect* 104(suppl 4):715–740.
- Kursa J, Travnicek J, Rambeck VA, Kroupova V, Vitovec J. 2000. Goiterogenic effects of extracted rapeseed meal and nitrates in sheep and their progeny. *Vet Med Czech* 45:129–140.
- Lahti E, Gharri M, Lindqvist OV. 1985. Uptake and distribution of radioiodine, and the effect of ambient nitrate, in some fish species. *Comp Biochem Physiol* 80A:337–342.
- Levallois P, Phaneuf D. 1994. Nitrate contamination of drinking water—an evaluation of health risks. *Can J Public Health* 85:192–196.
- Longnecker MP, Klebanoff MA, H Z, Brock JW. 2001. Association between maternal serum concentrations of the DDT metabolite DDE and preterm and small-for-gestational-age babies at birth. *Lancet* 358:110–114.
- Lundholm CE. 1997. DDE-induced eggshell thinning in birds: Effects of *p,p'*-DDE on calcium and prostaglandin metabolism of the eggshell gland. *Comp Biochem Physiol C Pharmacol Toxicol Endocrinol* 118:113–128.
- Markey CM, Coombs MA, Sonnenschein C, Soto AM. 2003. Mammalian development in a changing environment: exposure to endocrine disruptors reveals the developmental plasticity of steroid-hormone target organs. *Evol Dev* 5:67–75.
- Markey CM, Rubin BS, Soto AM, Sonnenschein C. 2002. Endocrine disruptors: from wingspread to environmental developmental biology. *J Steroid Biochem Molec Biol* 83:235–244.
- McLachlan JA. 2001. Environmental signaling: what embryos and evolution teach us about endocrine disrupting chemicals. *Endocr Rev* 22:319–341.
- Meyer D. 1995. Enzymatic/non-enzymatic formation of nitric oxide. *Nat Med* 1:1103–1104.
- Mori C, Komiyama M, Adachi T, Sakurai K, Nishimura D, Takashima K, et al. 2003. Application of toxicogenomic analysis to risk assessment of delayed long-term effects of multiple chemicals, including endocrine disruptors in human fetuses. *Environ Health Perspect* 111:803–809.
- National Academy of Sciences. 1999. *Hormonally Active Agents in the Environment*. Washington, DC:National Academy Press.
- Nohl H, Kozlov AV, Staniek K, Gille L. 2001. The multiple functions of coenzyme Q. *Bioorg Chem* 29:1–13.
- Nohl H, Staniek K, Sobhian S, Bahrami S, Redl H, Kozlov AV. 2000. Mitochondria recycle nitrite back to the bioregulator nitric monoxide. *Acta Biochemica Polonica* 47:913–921.
- Oberdörster E, Cheek AO. 2000. Gender benders at the beach: endocrine disruption in marine and estuarine organisms. *Environ Toxicol Chem* 20:23–36.
- Panesar NS. 1999. Role of chloride and inhibitory action of inorganic nitrate on gonadotropin-stimulated steroidogenesis in mouse Leydig tumor cells. *Metabolism* 48:693–700.
- Panesar NS, Chan KW. 2000. Decreased steroid hormone synthesis from inorganic nitrite and nitrate: studies *in vitro* and *in vivo*. *Toxicol Appl Pharmacol* 169:222–230.
- Park D, Hempleman SC, Propper CR. 2001. Endosulfan exposure disrupts pheromonal systems in the red-spotted newt: a mechanism for subtle effects of environmental chemicals. *Environ Health Perspect* 109:669–673.
- Park D, Propper CR. 2002. Endosulfan affects pheromonal detection and glands in the male red-spotted newt, *Notophthalmus viridescens*. *Bull Environ Contam Toxicol* 69:609–616.
- Pickford DB, Morris ID. 1999. Effects of endocrine-disrupting contaminants on amphibian oogenesis: methoxychlor inhibits progesterone-induced maturation of *Xenopus laevis* oocytes *in vitro*. *Environ Health Perspect* 107:285–292.
- Porter WP, Jaeger JW, Carlson H. 1999. Endocrine, immune, and behavioral effects of aldicarb (carbamate), atrazine (triazine), and nitrate (fertilizer) mixtures at groundwater concentrations. *Toxicol Ind Health* 15:133–150.
- Rolland R. 2000. A review of chemically-induced alterations in thyroid and vitamin A status from field studies of wildlife and fish. *J Wildl Dis* 36:615–635.
- Ruppert C, Kraft W. 1999. Examination in dogs with spontaneous hyperadrenocorticism. Part 3. Thyroid hormone concentrations before and after treatment with mitotane (Lysodren®). *Tier Prax aus klein Heimtiere* 27:167–171.
- Safe SH, Pallaroni L, Yoon K, Gaido KW, Ross S, McDonnell DP. 2002. Problems for risk assessment of endocrine-active estrogenic compounds. *Environ Health Perspect* 110(suppl 6):925–929.
- Sanderson JT, Letcher RJ, Heneweer M, Giesy JP, van den Berg M. 2001. Effects of chloro-S-triazine herbicides and metabolites on aromatase activity in various human cell lines and on vitellogenin production in male carp hepatocytes. *Environ Health Perspect* 109:1027–1031.
- Sanderson JT, Seinen W, Giesy JP, van den Berg M. 2000. 2-Chloro-S-triazine herbicides induce aromatase (CYP19) activity in H295R human adrenocortical carcinoma cells: a novel mechanism for estrogenicity? *Toxicol Sci* 54:121–127.
- Sanderson T, van den Berg M. 2003. Interactions of xenobiotics with the steroid hormone biosynthesis pathway. *Pure Appl Chem* 75:1957–1971.
- Sharpe RM, Irvine DS. 2004. How strong is the evidence of a link between environmental chemicals and adverse effects on human reproductive health. *BMJ* 328:447–451.
- Stanley DW, Howard RW. 1998. The biology of prostaglandins and related eicosanoids in invertebrates: cellular, organismal, and ecological actions. *Am Zool* 38:369–381.
- Stocco DM, Clark B. 1996. Regulation of the acute production of steroids in steroidogenic cells. *Endocr Rev* 17:221–244.
- Sumpter JP, Jobling S. 1995. Vitellogenesis as a biomarker for estrogenic contamination of the aquatic environment. *Environ Health Perspect* 103(suppl 7):173–178.
- Swan, SH, Main KM, Liu F, Stewart SL, Kruse RL, Calafat AM, et al., and the Study for Future Families Research Team. 2005. Decrease in anogenital distance among male infants with prenatal phthalate exposure. *Environ Health Perspect* 113:1056–1061.
- Tabb MM, Kholodovych V, Grun F, Zhou C, Welsh WJ, Blumberg B. 2004. Highly chlorinated PCBs inhibit the human xenobiotic response mediated by the steroid and xenobiotic receptor (XSR). *Environ Health Perspect* 112:163–169.
- Toell A, Kroncke K-D, Kleinert H, Carlberg C. 2002. Orphan nuclear receptor binding site in the human inducible nitric oxide synthase promoter mediates responsiveness to steroid and xenobiotic ligands. *J Cell Biochem* 85:72–82.
- Toppari J, Larsen JC, Christiansen P, Giwercman A, Grandjean P, Guillette LJ Jr, et al. 1996. Male reproductive health and environmental xenoestrogens. *Environ Health Perspect* 104 (suppl 4):741–803.
- Tyler CR, Jobling S, Sumpter JP. 1998. Endocrine disruption in wildlife: a critical review of the evidence. *Crit Rev Toxicol* 28:319–361.
- Vonier PM, Crain DA, McLachlan JA, Guillette LJ Jr, Arnold SF. 1996. Interaction of environmental chemicals with the estrogen and progesterone receptors from the oviduct of the American alligator. *Environ Health Perspect* 104:1318–1322.
- White PC, New MI, Dupont B. 1987. Congenital adrenal hyperplasia. *N Engl J Med* 316:1519–1524.
- Wilson VS, LeBlanc GA. 1998. Endosulfan elevates testosterone biotransformation and clearance in CD-1 mice. *Toxicol Appl Pharmacol* 148:158–168.
- Wilson VS, LeBlanc GA. 2000. The contribution of hepatic inactivation of testosterone to the lowering of serum testosterone levels by ketoconazole. *Toxicol Sci* 54:128–137.
- Wynngaarden JB, Stanburg JB, Rapp B. 1953. The effects of iodine, perchlorate, tricyanate, and nitrate administration upon the iodine concentration mechanism of the rat thyroid. *Endocrinology* 52:568–574.
- Wynngaarden JB, Wright BM, Ways P. 1952. The effect of certain anions upon the accumulation and retention of iodine by the thyroid gland. *Endocrinology* 50:537–549.
- Xie W, Barwick JL, Downes M, Blumberg B, Simon CM, Nelson MC, et al. 2000. Humanized xenobiotic response in mice expressing nuclear receptor XSR. *Nature* 206:435–439.
- Zhu Y, Bond J, Thomas P. 2003a. Identification, classification, and partial characterization of genes in humans and other vertebrates homologous to a fish membrane progesterin receptor. *Proc Natl Acad Sci USA* 100:2237–2242.
- Zhu Y, Rice CD, Pang Y, Pace M, Thomas P. 2003b. Cloning, expression, and characterization of a membrane progesterin receptor and evidence it is an intermediary in meiotic maturation of fish oocytes. *Proc Natl Acad Sci USA* 100:2231–2236.
- Zoeller RT, Dowling ALS, Herzog CTA, Iannacone EA, Gauger KJ, Bansal R. 2002. Thyroid hormone, brain development, and the environment. *Environ Health Perspect* 110 (suppl 3):355–361.
- Zraly Z, Bendova J, Svecova D, Faldikova L, Veznik Z, Zajicova A. 1997. Effects of oral intake of nitrates on reproductive functions of bulls. *Vet Med Czech* 42:345–354.
- Zweier JL, Samouilov A, Kuppusamy P. 1999. Non-enzymatic nitric oxide synthesis in biological systems. *Biochim Biophys Acta* 1411:250–262.