

- Symptoms of bronchial hyperreactivity and asthma in relation to environmental factors. *Arch Dis Child* 1988;63:473-8.
- 18 Lunn JE, Knowelden J, Roe JW. Patterns of respiratory illness in Sheffield junior schoolchildren—a follow up study. *Br J Prev Soc Med* 1970;24:223-8.
 - 19 Ware JH, Ferris BG Jr, Dockery DW, Spengler JD, Stram DO, Speizer FE. Effects of ambient sulfur oxides and suspended particles on respiratory health of pre-adolescent children. *Am Rev Respir Dis* 1986;133:834-42.
 - 20 Braun-Fahrlander C, Ackermann-Lieblich U, Schwartz J, Rutishauser M, Gnehm HP, Wonne HU. Air pollution and respiratory symptoms in preschool children. *Am Rev Respir Dis* 1992;148:42-7.
 - 21 Molino NA, Wright SC, Katz I, Tarlo S, Silverman F, McClean PA, et al. Effect of low concentrations of ozone on inhaled allergen responses in asthmatic subjects. *Lancet* 1991;338:199-203.
 - 22 Ferris BG. Epidemiology standardization project. *Am Rev Respir Dis* 1978;118 (suppl):7-53.
 - 23 Nicolai T, Mutius E von, Reitmeir P, Wjst M. Reactivity to cold air hyperventilation in normal and asthmatic children in a survey of 5697 school children in southern Bavaria. *Am Rev Respir Dis* (in press).
 - 24 Fleiss JL. *Statistical methods for rates and proportions*. New York: John Wiley, 1980.
 - 25 SAS. *Statistical analysis system. Version 6.06*. Cary, North Carolina: SAS, 1990.
 - 26 Steinbrück P, Herrmann H, Landmann H. Empfehlungen der Problemkommission für Lungenkrankheiten und Tuberkulose zur Definition von chronischer Bronchitis, Asthma bronchiale und Lungenemphysem. *Monatsschrift für Lungenkrankheiten und Tuberkulose Bekämpfung* 1969;12: 289-98.
 - 27 Logan WPD. Mortality in London fog incident. *Lancet* 1953;i:336-8.
 - 28 Gore AT, Shaddick CW. Atmospheric pollution and mortality in the county of London. *Br J Prev Soc Med* 1968;12:104-13.
 - 29 Lunn JE, Knowelden J, Handyside AJ. Patterns of respiratory illness in Sheffield infant school children. *Br J Prev Soc Med* 1967;21:7-16.
 - 30 Dodge R, Solomon P, Moyers J, Hayes C. A longitudinal study of children exposed to sulfur oxides. *Am J Epidemiol* 1985;121:720-36.
 - 31 Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG. Effects of inhalable particles on respiratory health of children. *Am Rev Respir Dis* 1989;139:587-94.
 - 32 Koenig JQ, Morgan MS, Horike M, Pierson WE. The effects of sulfur dioxide on nasal and lung function in adolescents with extrinsic asthma. *J Allergy Clin Immunol* 1985;76:813-8.
 - 33 Sheppard D, Wong WS, Uehara JA, Nadel JA, Boushey HA. Lower threshold and greater bronchomotor responsiveness of asthmatic subjects to sulphur dioxide. *Am Rev Respir Dis* 1980;122:873.
 - 34 Jorres R, Magnussen H. Airways response of asthmatics after a 30 min exposure, at resting ventilation, to 0.25 ppm NO₂ or 0.5 ppm SO₂. *Eur Respir J* 1990;3:132-7.
 - 35 Pope CA. Respiratory disease associated with community air pollution and steel mill, Utah Valley. *Am J Public Health* 1989;79:623-8.
 - 36 Takafuji S, Suzuki S, Koizumi K, Tadokoro K, Miyamoto T, Ikemori R, et al. Diesel-exhaust particulates inoculated by the intranasal route have an adjuvant activity for IgE production in mice. *J Allergy Clin Immunol* 1987;79:639-45.

(Accepted 6 October 1992)

Medical management of missed abortion and anembryonic pregnancy

H El-Refaey, K Hinshaw, R Henshaw, N Smith, A Templeton

Department of Obstetrics and Gynaecology, Aberdeen Maternity Hospital, Aberdeen AB9 2ZD

H El-Refaey, research fellow
K Hinshaw, research fellow
R Henshaw, lecturer
N Smith, consultant
A Templeton, professor

Correspondence to: Professor Templeton.

BMJ 1992;305:1399

Miscarriage is the most common complication of pregnancy and probably accounts for around 50 000 inpatient admissions in the United Kingdom each year. Generally, its management has changed little with time. Recently developments in non-surgical treatment, refined in induced abortion, offer an opportunity to improve management and to remove the need for surgery and anaesthesia. We report our initial experience of the use of mifepristone (an antiprogesterone) and misoprostol (a synthetic analogue of prostaglandin E₁) in the management of missed abortion and anembryonic pregnancy (gestation sac present but no developing embryo).

Patients, methods, and results

Sixty women with a diagnosis of missed abortion or anembryonic pregnancy equivalent to 13 weeks' gestation or less were recruited after counselling. Twenty five of the women had been referred for ultrasound scanning because of bleeding in early pregnancy, while in the remainder the diagnosis had been made on routine scanning when they booked. Patients in whom an incomplete abortion was diagnosed were not included.

Each patient was given a single oral dose of mifepristone 600 mg and was admitted to the gynaecological ward as an outpatient 36-48 hours later, when misoprostol 600 µg was given orally in a divided dose (400 µg and, two hours later, 200 µg). The patient's pulse, blood pressure, and temperature were recorded hourly, as were any side effects and requests for analgesia. If the products of conception were not expelled and verified within four hours vaginal ultrasonography was performed. If the gestation sac was not seen and bleeding had occurred the procedure was considered to have been successful, but if not the patient was offered evacuation of the uterus under general anaesthesia. All patients were reviewed 10-14 days later.

The median age of the 60 women was 27 (range 15-44), and the median duration of amenorrhoea was 71 (42-110) days. Twenty nine patients were diagnosed as having anembryonic pregnancies and the remaining 31 as having had a missed abortion. One of the patients thought to have an anembryonic pregnancy was

eventually found to have an ectopic pregnancy when she failed to abort. She has been excluded from further consideration.

Eight patients aborted with mifepristone alone. Of the 51 remaining patients, 43 aborted after taking misoprostol 600 µg and five more aborted after receiving a second divided dose of 600 µg misoprostol. In three patients the treatment failed, and they underwent evacuation of the uterus under general anaesthesia. Exploratory curettage was performed in two other patients at 14 and 22 days after treatment with misoprostol, but no products of conception were obtained.

The median time from administration of misoprostol to abortion was 4 (range 1-11) hours. The median duration of bleeding after abortion was 10 (2-22) days. There were no cases of infection, and no patient required antibiotic treatment. Side effects from misoprostol treatment (nausea, vomiting, and diarrhoea) were few: antiemetic drugs were given to five patients, and diarrhoea was reported by seven. Thirty nine women did not ask for any pain relief, 13 requested oral analgesia, and seven required parenteral analgesia.

Comment

The medical management of induced abortion is now established practice, and several centres have developed considerable skill in using mifepristone with a variety of prostaglandin analogues.¹⁻⁴ We have shown the clinical feasibility of managing missed abortion and anembryonic pregnancy medically without resort to surgery or anaesthesia. Patients' perceptions at this early stage of development of the method are positive but need to be assessed more thoroughly, particularly in relation to the alternatives. The implications in terms of resources are considerable since if this method became standard practice a major part of emergency work in gynaecology would be removed from the operating room and could be managed at times more convenient to patients and staff.

- 1 UK Multicentre Trial. The efficacy and tolerance of mifepristone and prostaglandin in first trimester termination of pregnancy. *Br J Obstet Gynaecol* 1990;97:480-6.
- 2 World Health Organisation Task Force on Postovulatory Methods for Fertility Regulation. Termination of early human pregnancy with RU486 (mifepristone) and the prostaglandin analogue sulprostone: a multicentre randomized comparison between two treatment regimens. *Hum Reprod* 1989;4:718.
- 3 World Health Organisation. Pregnancy termination with mifepristone and gemeprost: a multicentre comparison between repeated doses and a single dose of mifepristone. *Fertil Steril* 1990;56:132-40.
- 4 Norman JE, Thong KJ, Baird DT. Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifepristone. *Lancet* 1991;338:1233-6.

(Accepted 23 October 1990)