

Heterologous Expression and Identification of the Genes Involved in Anaerobic Degradation of 1,3-Dihydroxybenzene (Resorcinol) in *Azoarcus anaerobius*[∇]

Paula I. Darley,^{1†} Jutta A. Hellstern,^{1†‡} Javier I. Medina-Bellver,² Silvia Marqués,² Bernhard Schink,¹ and Bodo Philipp^{1*}

Fachbereich Biologie, Universität Konstanz, D-78457 Constance, Germany,¹ and Estación Experimental del Zaidín, C/ Profesor Albareda 1, E-18008 Granada, Spain²

Received 9 November 2006/Accepted 9 March 2007

Azoarcus anaerobius, a strictly anaerobic, gram-negative bacterium, utilizes resorcinol as a sole carbon and energy source with nitrate as an electron acceptor. Previously, we showed that resorcinol degradation by this bacterium is initiated by two oxidative steps, both catalyzed by membrane-associated enzymes that lead to the formation of hydroxyhydroquinone (HHQ; 1,2,4-benzenetriol) and 2-hydroxy-1,4-benzoquinone (HBQ). This study presents evidence for the further degradation of HBQ in cell extracts to form acetic and malic acids. To identify the *A. anaerobius* genes required for anaerobic resorcinol catabolism, a cosmid library with genomic DNA was constructed and transformed into the phylogenetically related species *Thauera aromatica*, which cannot grow with resorcinol. By heterologous complementation, a transconjugant was identified that gained the ability to metabolize resorcinol. Its cosmid, designated R⁺, carries a 29.88-kb chromosomal DNA fragment containing 22 putative genes. In cell extracts of *T. aromatica* transconjugants, resorcinol was degraded to HHQ, HBQ, and acetate, suggesting that cosmid R⁺ carried all of the genes necessary for resorcinol degradation. On the basis of the physiological characterization of *T. aromatica* transconjugants carrying transposon insertions in different genes of cosmid R⁺, eight open reading frames were found to be essential for resorcinol mineralization. Resorcinol hydroxylase-encoding genes were assigned on the basis of sequence analysis and enzyme assays with two mutants. Putative genes for hydroxyhydroquinone dehydrogenase and enzymes involved in ring fission have also been proposed. This work provides the first example of the identification of genes involved in the anaerobic degradation of aromatic compounds by heterologous expression of a cosmid library in a phylogenetically related organism.

Resorcinol (1,3-dihydroxybenzene) is produced and utilized in large amounts by the timber, adhesives, and oil industries and enters freshwater environments through the release of effluents. Also, roots of aquatic plants such as *Nuphar lutea* exude resorcinol in considerable amounts into the aquatic environment (42). Resorcinol is photochemically transformed upon exposure to sunlight. However, these chemical modifications are, in general, slow processes (45) and do not degrade resorcinol to carbon dioxide. Therefore, most of the detoxification and reintroduction of resorcinol constituents back into the carbon cycle must be catalyzed by microbial degradation.

There have been several reports of bacteria able to utilize resorcinol as carbon and energy sources in the presence or absence of oxygen by different biochemical strategies. While aerobes dearomatize the benzene ring by using molecular oxygen as a cosubstrate for monooxygenases and dioxygenases, anaerobes have to replace all of the oxygen-dependent steps with alternative sets of reactions (6, 16, 21, 36). Notably, in anaerobic pathways, the aromatic ring is reduced rather than oxidized and benzoyl coenzyme A (CoA) emerges as the most

common intermediate in the degradation of a large diversity of aromatic substrates (4, 5, 7, 21, 36, 46).

Since the late 1960s, several microorganisms such as *Pseudomonas putida* strain ORC (10), *Trichosporon cutaneum* (12), and *Azotobacter vinelandii* (18) were reported to be able to degrade resorcinol aerobically. In all of these cases, resorcinol is oxidized in reactions catalyzed by oxygenases to hydroxyhydroquinone (HHQ; 1,2,4-benzenetriol) or pyrogallol (1,2,3-benzenetriol). HHQ undergoes further diol ring cleavage with subsequent formation of maleylacetate and later β -keto adipate (10, 12), while pyrogallol is cleaved to form oxalocrotonate and later pyruvate plus acetate (18). Very recently, a new *Rhodococcus opacus* strain named RW was isolated and shown to degrade resorcinol (29). So far, nothing is known about the biochemistry that *R. opacus* RW uses in the mineralization of resorcinol.

Anaerobic resorcinol degradation has been documented for various microorganisms, such as sulfate-reducing bacteria (38), fermenting bacteria (44), and denitrifiers (17, 26, 40). Notably, none of these bacteria use the benzoyl-CoA pathway in resorcinol catabolism. This is due to the fact that resorcinol as a dihydroxybenzene is less stabilized by resonance of the benzene ring and can be attacked reductively or oxidatively without prior activation. Both strategies have been described for resorcinol destabilization under anoxic conditions. While a fermenting *Clostridium* sp. uses reductive biochemistry to convert resorcinol to cyclohexanedione (26, 44), which is further

* Corresponding author. Mailing address: Fach M654, Mikrobielle Ökologie, Fachbereich Biologie, Universität Konstanz, D-78457 Constance, Germany. Phone: 49 7531 88 4541. Fax: 49 7531 88 4047. E-mail: bodo.philipp@uni-konstanz.de.

† P.I.D. and J.A.H. contributed equally to this work.

‡ Present address: RCC Ltd., 4452 Itingen, Switzerland.

[∇] Published ahead of print on 16 March 2007.

hydrolyzed to 5-oxohexanoate (36), *Anaerobius anaerobius*, a nitrate-reducing bacterium, uses two oxidative reactions (32). Resorcinol is hydroxylated at position 4 of the aromatic ring with $K_3Fe(CN)_6$ or nitrate as an electron acceptor to form hydroxyhydroquinone (HHQ) in a reaction catalyzed by resorcinol hydroxylase. HHQ is subsequently oxidized with nitrate to HBQ (2-hydroxy-1,4-benzoquinone) in a reaction catalyzed by hydroxyhydroquinone dehydrogenase (32). Both reactions were measured in the membrane fraction of resorcinol-grown cells (32). The further fate of HBQ has not been studied yet. Hydroxylation of resorcinol in *A. anaerobius* provides a less energy-consuming alternative to the benzoyl-CoA pathway, in which benzoyl-CoA formation requires stoichiometric hydrolysis of ATP (46). Moreover, the electrons from resorcinol and HHQ oxidation can directly enter the denitrification process and may allow energy conservation by proton translocation (32).

The aims of our study were to further elucidate the resorcinol degradation pathway in *A. anaerobius* and to identify genes that code for resorcinol metabolism. Analysis of the sequences of 29.88-kb chromosomal DNA from a cosmid library should provide new insight into the function of the resorcinol-degrading enzymes.

MATERIALS AND METHODS

Chemicals. All of the chemicals and biochemicals used in this study were purchased from Fluka (Neu Ulm, Germany), Merck (Darmstadt, Germany), Serva (Heidelberg, Germany), Sigma-Aldrich (Deisenhofen, Germany), or Biozym (Oldendorf, Germany) and were of the highest quality available. Enzymes used for cloning and other materials used for molecular biology techniques were obtained from MBI Fermentas (St. Leon-Rot, Germany), Roche Diagnostics (Mannheim, Germany), Eppendorf (Hamburg, Germany), Stratagene (Heidelberg, Germany), or QIAGEN (Hilden, Germany). 2-Hydroxy-1,4-benzoquinone (HBQ) was prepared nonenzymatically by auto-oxidation of hydroxyhydroquinone (10). Custom sequencing was performed at GATC (Constance, Germany).

Bacterial strains, vectors, and culture conditions. The bacterial strains and vectors used in this work are summarized in Table 1. *Azoarcus anaerobius* strain LuFRes1, *T. aromatica* strains AR1 and K172, and their transconjugants were cultured anaerobically at 30°C in 100-ml infusion bottles containing 50 ml of nonreduced mineral medium (32) under nitrogen gas. The medium was buffered with 30 mM 3-(*N*-morpholino)propanesulfonic acid instead of bicarbonate and supplemented with 8 mM nitrate. Carbon sources were stored as 0.5 M stock solutions in sterile infusion bottles under nitrogen gas and added to cultures with syringes to final concentrations of 2 mM resorcinol (*A. anaerobius*), 2 mM 3,5-dihydroxybenzoate, 2 mM benzoate (*T. aromatica* AR1), and 5 mM succinate (*T. aromatica* K172). For biochemical studies, all strains were cultured in 1-liter infusion bottles and harvested at an optical density at 578 nm (OD_{578}) of 0.3 to 0.4. To study resorcinol degradation and nitrate reduction by *T. aromatica* transconjugants containing pLAFR3 recombinant cosmid or EZ::TN<KAN-2> transposon insertional derivatives of cosmid R⁺, 100-ml serum bottles containing 50 ml of anaerobic mineral medium supplemented with resorcinol (2 mM) and nitrate (8 mM) were inoculated with 0.5% preculture. *Escherichia coli* HB101(pRK600) was used as a helper strain, and strains HB101 and DH5 α (pLAFR3 derivative cosmids) were used as the donor strains in triparental matings. All *E. coli* strains were grown aerobically at 37°C in Luria-Bertani (LB) medium (35). Antibiotics were used at the following concentrations: tetracycline, 20 μ g ml⁻¹; chloramphenicol, 30 μ g ml⁻¹; kanamycin, 50 μ g ml⁻¹.

Growth and substrate depletion analysis. Samples of the culture were withdrawn anaerobically with a sterile plastic syringe flushed with N₂ at the designated time points. Growth was monitored spectrophotometrically by measuring OD_{578} in a Hitachi U-100 spectrophotometer. For substrate depletion analysis, the cells from the culture samples were pelleted (for 5 min by microcentrifuge) and the supernatant was stored at -20°C until analyzed by high-pressure liquid chromatography (HPLC) for resorcinol and/or 3,5-dihydroxybenzoate degradation, nitrate consumption, and nitrite formation as described below.

Preparation of cell extracts. All of the steps in the preparation of cell extracts were performed anaerobically. Cultures of *A. anaerobius* and *T. aromatica* transconjugants were harvested at an OD_{578} of 0.3 and washed once under anoxic conditions with 100 ml of 50 mM potassium phosphate buffer (pH 7.0) as described previously (28). Unless used immediately, cell pellets were frozen in liquid N₂ and stored at -20°C. To prepare cell extracts, cells were suspended in 50 mM potassium phosphate buffer (pH 7.0), passed through a French press at 138 MPa, and then centrifuged at 20,000 \times g for 20 min at 4°C to remove cell debris. The membranes were collected by centrifugation at 100,000 \times g for 1 h at 4°C and resuspended in anoxic 50 mM potassium phosphate buffer (pH 7.0) (half the volume of the cytosolic fraction). Both fractions were either used immediately or frozen in liquid N₂ and stored at -20°C.

Protein determination. Protein content was quantified by the Bradford method (8) with bovine serum albumin as the standard.

Enzymatic tests. Resorcinol hydroxylase and hydroxyhydroquinone dehydrogenase activities were measured at 30°C under anoxic conditions as previously described (32), except that potassium phosphate buffer (50 mM, pH 7.0) was used as the buffer in all of the assays. A standard reaction mixture used to assay resorcinol hydroxylase contained 0.2 to 0.3 mg of protein, 1 mM $K_3Fe(CN)_6$, and 1 mM resorcinol. A standard reaction mixture used to assay hydroxyhydroquinone dehydrogenase contained 0.2 to 0.3 mg of protein, 1 mM NaNO₃, and 1 mM resorcinol. NAD(P)H-dependent degradation of resorcinol beyond HBQ was coupled to nitrate reduction and monitored by measuring the formation of acetate, malate, and succinate by ion-exchange HPLC as described below. A standard reaction (1 ml) was started by addition of 1 mM resorcinol to the assay mixture containing the membrane fraction (1 mg protein ml⁻¹), supplemented or not with the cytosolic fraction (1 mg protein ml⁻¹), and incubated at 30°C in 50 mM potassium phosphate (pH 7.0) with 4 to 8 mM nitrate and 1 mM NADH. At defined time points, samples (25 μ l) were withdrawn, mixed with ice-cold potassium phosphate buffer (25 μ l), and immediately analyzed for resorcinol degradation by HPLC. Additional samples (150 μ l) were withdrawn anoxically from the assay mixture, and the enzymatic reaction was stopped by adding 50 μ l of 1 M sulfuric acid. Denatured proteins were removed by centrifugation, and the sample was analyzed immediately by HPLC for organic acid formation or stored at -20°C.

HPLC analysis. Resorcinol, HHQ, and HBQ were analyzed with a C₁₈ reversed-phase column (Grom-Sil 120 ODS-5ST, 5 μ m, 150 by 4.6 mm; Grom, Herrenberg, Germany) as described elsewhere (32). Resorcinol degradation in cultures by *T. aromatica* transconjugants and resorcinol hydroxylase enzyme activities were measured discontinuously by HPLC with detection at 278 and 206 nm, respectively. The mobile phase, comprising a mixture of 100 mM ammonium phosphate buffer (pH 2.6) and methanol, was used at a flow rate of 1 ml min⁻¹. The solvent phase (5% [vol/vol] methanol) was initially held for 1 min, and then the concentration was increased to 45% over a period of 6 min and then lowered within 0.5 min to 5% and held for an additional 7 min. Acetic and succinic acids were separated on an ion-exchange column (Aminex HPX-87H; Bio-Rad, Munich, Germany) that was operated at a flow rate of 0.6 ml min⁻¹ at 40°C with 5 mM sulfuric acid solution as the mobile phase. Acetic acid eluted after 15.7 min, succinic acid eluted after 13.0 min, and malic acid eluted after 10.5 min. All three acids were quantified against external standards. Nitrate and nitrite were determined with an A06 anion-exchange column (3 by 120 mm; Sykam, Freising, Germany). A 40 mM NaCl solution was used as the eluent at a flow rate of 1 ml min⁻¹. Nitrate eluted after 5.6 min, and nitrite eluted after 3.1 min. Both compounds were detected at 210 nm and quantified against external standards.

DNA manipulation. Standard methods were used for genomic DNA preparation, DNA digestion with restriction endonucleases, ligation, agarose gel electrophoresis, and transformation of *E. coli* (35). Cosmids were prepared for sequencing and in vitro transposon mutagenesis experiments with the QIAGEN Large construct kit (QIAGEN, Hilden, Germany).

Cosmid library construction. Genomic DNA of *A. anaerobius* was isolated by an established protocol (35). Partial digestion of the DNA with PstI was carried out, and fragments of 20 and 30 kb were isolated and ligated with the PstI-digested and dephosphorylated pLAFR3 (41) vector. The resulting ligation products were then packaged into lambda phage heads with a Gigapack III packaging extract (Stratagene). The phage particles were transduced into *E. coli* HB101, and colonies were grown on LB agar containing tetracycline (27). The resulting colonies were harvested in liquid LB medium, grown overnight, and used as the donors in triparental mating as described below, by using *T. aromatica* AR1 as the acceptor.

Triparental mating. *T. aromatica* strains AR1 and K172 were used as recipients and grown for 60 h anaerobically in minimal medium supplemented with

TABLE 1. Bacterial strains and cosmids used in this study

Strain or vectors	Relevant characteristics ^a	Source or reference
Strains		
<i>A. anaerobius</i> LuFRes1 (DSM12081)	Wild type; Res ⁺	40
<i>T. aromatica</i>		
AR1 (DSM11528)	Wild type; Res ⁻ DHB ⁺ HHQ pathway ⁺	13
K172 (DSM6984 ¹)	Wild type; Res ⁻ DHB ⁻ HHQ pathway ⁻	2, 43
<i>E. coli</i>		
DH5 α	F ⁻ ϕ 80 <i>dlacZ</i> Δ M15 Δ (<i>lacZYA-argF</i>)U169 <i>deoR recA1 endA1 hsdR17</i> (r_K^- m_K^+) <i>phoA supE44</i> λ^- <i>thi-1 gyrA96 relA1</i>	35
HB101	<i>supE44 hsdS20</i> (r_B^- m_B^-) <i>recA13 ara-14 proA2 lacY1 galK2 rpsL20 xyl-5 mtl-1 leuB6 thi-1</i>	35
Plasmid pRK600		
	ColE1 Tra ⁺ Cm ^r	22
Cosmids		
pLAFR3	Tc ^r Tra ⁻ Mob ⁺ <i>cos</i> ; RK2 replicon	41
R ⁺	Tc ^r , pLAFR3 containing a 29.9-kb resorcinol gene cluster of <i>A. anaerobius</i>	This study
Mu_4	Tc ^r Km ^r R ⁺ <i>orf14::TN<KAN-2></i> ; knockout mutant	This study
Mu_13	Tc ^r Km ^r R ⁺ <i>btdhS::TN<KAN-2></i> ; knockout mutant	This study
Mu_14	Tc ^r Km ^r R ⁺ <i>orf12::TN<KAN-2></i> ; knockout mutant	This study
Mu_16	Tc ^r Km ^r R ⁺ <i>orf10::TN<KAN-2></i> ; knockout mutant	This study
Mu_21	Tc ^r Km ^r R ⁺ <i>rhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_22	Tc ^r Km ^r R ⁺ <i>bqdhS::TN<KAN-2></i> ; knockout mutant	This study
Mu_31	Tc ^r Km ^r R ⁺ <i>orf3::TN<KAN-2></i> ; knockout mutant	This study
Mu_36	Tc ^r Km ^r R ⁺ <i>orf14::TN<KAN-2></i> ; knockout mutant	This study
Mu_40	Tc ^r Km ^r R ⁺ <i>rhS::TN<KAN-2></i> ; knockout mutant	This study
Mu_41	Tc ^r Km ^r R ⁺ <i>orf5::TN<KAN-2></i> ; knockout mutant	This study
Mu_42	Tc ^r Km ^r R ⁺ <i>orf7::TN<KAN-2></i> ; knockout mutant	This study
Mu_44	Tc ^r Km ^r R ⁺ <i>rhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_45	Tc ^r Km ^r R ⁺ <i>orf10::TN<KAN-2></i> ; knockout mutant	This study
Mu_46	Tc ^r Km ^r R ⁺ <i>orf11::TN<KAN-2></i> ; knockout mutant	This study
Mu_51	Tc ^r Km ^r R ⁺ <i>bqdhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_53	Tc ^r Km ^r R ⁺ <i>bqdhS::TN<KAN-2></i> ; knockout mutant	This study
Mu_54	Tc ^r Km ^r R ⁺ <i>orf2::TN<KAN-2></i> ; knockout mutant	This study
Mu_56	Tc ^r Km ^r R ⁺ <i>orf1::TN<KAN-2></i> ; knockout mutant	This study
Mu_63	Tc ^r Km ^r R ⁺ <i>orf9::TN<KAN-2></i> ; knockout mutant	This study
Mu_64	Tc ^r Km ^r R ⁺ <i>orf2::TN<KAN-2></i> ; knockout mutant	This study
Mu_68	Tc ^r Km ^r R ⁺ <i>orf2::TN<KAN-2></i> ; knockout mutant	This study
Mu_73	Tc ^r Km ^r R ⁺ <i>rhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_76	Tc ^r Km ^r R ⁺ <i>btdhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_80	Tc ^r Km ^r R ⁺ <i>bqdhL::TN<KAN-2></i> ; knockout mutant	This study
Mu_82	Tc ^r Km ^r R ⁺ <i>orf1::TN<KAN-2></i> ; knockout mutant	This study
Mu_93	Tc ^r Km ^r R ⁺ <i>orf13::TN<KAN-2></i> ; knockout mutant	This study

^a Res⁺, growth on resorcinol; Res⁻, no growth on resorcinol; DHB⁺, growth on 3,5-dihydroxybenzoate; DHB⁻, no growth on 3,5-dihydroxybenzoate; HHQ pathway⁺, has HHQ pathway; HHQ pathway⁻, lacks HHQ pathway; Km^r, kanamycin resistance; Cm^r, chloramphenicol resistance; Tc^r, tetracycline resistance.

3,5-dihydroxybenzoate (2 mM) and succinate (5 mM), respectively. The following *E. coli* strains were grown in LB medium overnight: strain HB101 containing helper plasmid pRK600 with chloramphenicol and strain HB101 and DH5 α donors containing pLAFR3 recombinant cosmids with tetracycline or EZ::TN<KAN-2> transposon insertional derivatives of cosmid R⁺ with kanamycin. All steps in the triparental mating were done aerobically as follows. A 0.5-ml volume of *E. coli* culture and 3 ml of the recipient culture in the exponential growth phase were centrifuged at 13,000 \times g for 10 min at 10°C. Cell pellets were washed twice with 1 ml LB medium. The resulting cell pellets of all three types of cells were resuspended together in 30 μ l LB and distributed on a sterile 47-mm-diameter, 0.22- μ m-pore-size filter (Schleicher & Schuell, Dassel, Germany) that was placed on an LB plate and incubated overnight at 30°C. The filter was transferred into 1 ml of mineral medium in a sterile tube. Cells were washed off by vigorous vortexing and transferred to 100-ml serum bottles containing 50 ml anoxic mineral medium supplemented with resorcinol (2 mM) and nitrate (8 mM). The cultures were incubated at 30°C for 1 to 3 weeks in the dark

without shaking. The ability of these transconjugants to grow on resorcinol was confirmed by substrate depletion analysis.

Isolation of a *T. aromatica* AR1 transconjugant clone that degrades resorcinol. Heterologous expression of an *A. anaerobius* cosmid library was achieved in *T. aromatica* strain AR1 at 30°C under anoxic conditions with resorcinol (2 mM) and nitrate (8 mM) as substrates in the presence of tetracycline. After 2 weeks of incubation, 2% of the grown cells were transferred into fresh anoxic medium supplemented with resorcinol (2 mM) and nitrate (8 mM) and allowed to continue to grow. After 2 days, the medium was turbid, indicating that resorcinol degradation took place. In order to isolate single transconjugant colonies, agar shake dilutions were performed as described elsewhere (31). Single transconjugant colonies were analyzed for colony shape and motility, and 16S rRNA gene analysis confirmed that the transconjugant was indeed *T. aromatica* AR1. The cosmid was isolated, sequenced, and designated R⁺. Cosmid R⁺ was propagated in *E. coli* strain HB101 and used as the donor in the second triparental mating with *T. aromatica* K172 as the acceptor.

Construction, isolation, and sequencing of transposon insertion mutants. Gene knockouts were created by the in vitro transposon insertion mutagenesis technique with an EZ::TN<KAN-2> insertion kit (Epicenter, Oldendorf, Germany) and cosmid R⁺ as the template according to the manufacturer's instructions. *E. coli* DH5 α competent cells were electroporated with the resulting transposon insertion products with a GenePulser (Bio-Rad, Munich, Germany). Transposon insertion clones were selected on kanamycin-containing LB plates since the transposon confers resistance to this antibiotic. To identify clones having a single transposon insertion at different positions in the coding region of cosmid R⁺, the cosmids of about 100 of the 3,000 colonies obtained were isolated and analyzed by restriction digestion with a selected set of restriction enzymes. Those cosmids considered interesting were introduced into *T. aromatica* AR1 and K172 via triparental mating, and the resulting transconjugants were grown in mineral medium supplemented with kanamycin, resorcinol, and nitrate as described above. Samples were withdrawn anaerobically with N₂-flushed, sterile plastic syringes at the designated time points and analyzed for growth and resorcinol degradation. The precise sites of EZ::TN<KAN-2> transposon insertion were determined by sequencing the derivative cosmid R⁺ with transposon-specific primers provided in the in vitro transposon mutagenesis kit (KAN-2 FP-1 and KAN-2 RP-1).

Sequence analysis. Nucleotide and amino acid sequences were analyzed with tools provided by the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST/>) and the ExPASy molecular biology server (<http://www.expasy.ch/>). Transmembrane helices in proteins were predicted with TMHMM server 2.0 (<http://www.cbs.dtu.dk/services/TMHMM/>).

Nucleotide sequence accession number. The nucleotide sequence of the coding region of cosmid R⁺ was deposited in the NCBI (GenBank) nucleotide sequence database under accession no. EF078692.

RESULTS AND DISCUSSION

HBQ degradation in cell suspension. Various attempts were made to establish culture conditions for in vivo degradation of the resorcinol degradation intermediates HHQ and HBQ in *A. anaerobius*. No growth was observed with HHQ or HBQ prepared by auto-oxidation of HHQ at concentrations between 0.5 and 2 mM. No degradation and no product formation were observed in experiments with dense suspensions of intact cells or with cells permeabilized with cetyltrimethylammonium bromide. These results suggest two things. Either (i) *A. anaerobius* has no uptake system for such polar compounds that are readily degraded once formed in the cell, or (ii) neither intermediate is an inducer in the resorcinol degradation pathway. Nevertheless, *A. anaerobius* was able to take up and metabolize 5 mM acetate, succinate, or malate, each of which is a carboxylic acid formed during resorcinol degradation (see below).

Identification of HBQ degradation products in cell extracts of *A. anaerobius*. HBQ, the first nonaromatic intermediate in resorcinol degradation, is highly reactive (32, 47) and should be prone to ring fission. HBQ is not commercially available but can be produced by auto-oxidation of HHQ (10). In air, HBQ spontaneously oxidizes and forms polymers, which renders its detection by HPLC difficult. It also reacts with chemicals and components of cell extracts carrying thiol groups, giving rise to addition products. Therefore, HBQ depletion and product formation were studied with extracts prepared from resorcinol-grown cells which were incubated anoxically with resorcinol (1 to 2 mM) and nitrate (8 mM). In these experiments, nitrate was used as the electron acceptor for two reasons. In a previous study, it was shown that (i) oxidation of resorcinol and HHQ was dependent on electron acceptors with a positive redox potential such as K₃Fe(CN)₆ and nitrate (32) and (ii) nitrate was the only electron acceptor that did not react chemically with HHQ (32). Nitrate was used in excess to provide an electron acceptor for the first two reactions, namely, resorcinol hydroxylation and HHQ dehydrogenation, and to provide an




FIG. 1. Utilization of resorcinol (○) and formation of succinate (■), acetate (●), and malate (▲) by a mixture of the membrane fraction and cytosolic fraction of resorcinol-grown *A. anaerobius*.

electron acceptor for potential oxidations involved in the further degradation of HBQ. In assays with nitrate and membrane fractions prepared from resorcinol-grown cells, resorcinol was converted at an average rate of 2.1 mU mg⁻¹. During the reaction, the assay mixture turned from colorless to brick red, which is indicative of HBQ accumulation (24, 30, 32), which was confirmed by HPLC analysis (32). No substrate degradation and no HBQ formation were detected in assay mixtures in which resorcinol or nitrate was omitted or in those containing membranes prepared from acetate-grown cells. These data are in agreement with the results previously reported (32). Addition of a cytosolic fraction of resorcinol-grown cells to assay mixtures containing membrane fractions of resorcinol-grown cells increased the rate of resorcinol degradation twofold (5.8 mU mg⁻¹ protein). Nevertheless, HBQ accumulation was still revealed by red color development. Various coenzymes and cosubstrates, such as ascorbic acid, ATP, biotin, CoA, cysteine, dithiothreitol, glutathione, lipoic acid, naphthoquinone, thiamine pyrophosphate (TPP), NAD(P)⁺, NADPH, and NADH, were supplied in the assay mixtures containing resorcinol, nitrate, and both membrane and cytosolic fractions of resorcinol-induced cells. Except for NAD(P)H, none of them prevented HBQ accumulation. The resorcinol degradation rate increased to 9.9 mU mg⁻¹ with either NADH or NADPH. HPLC chromatograms obtained by separation on an ion-exchange column displayed two products within the first 6 h of incubation, which increased proportionally with resorcinol consumption. One of the two metabolic products formed during the first 6-h reaction almost disappeared, and a third one appeared (Fig. 1). By comparison of their retention times with standards of different carboxylic acids and by coelution studies, the unknown metabolites were identified. The primary products were identified as acetic acid and malic acid, and the later one was identified as succinic acid. As shown in Fig. 1, almost stoichiometric amounts of succinic and acetic acids were formed from resorcinol. Acetate and malate were formed concomitantly as resorcinol was metabolized, and succinate was formed later while malate was consumed, indicating that malate was the primary reaction product. Control assay mix-




FIG. 2. Degradation of resorcinol by transconjugants *T. aromatica* AR1 (A) and K172 (B) harboring cosmid R⁺. Resorcinol (○) is metabolized with concomitant consumption of nitrate (△), release of nitrite (▲), and an increase in OD₅₇₈ (●).

tures without resorcinol but with NADH showed that only negligible nitrate reduction took place. These results clearly indicated (i) that acetate, malate, and later succinate were formed in the resorcinol degradation pathway; (ii) that their formation was dependent on a soluble and resorcinol-induced enzyme(s) that required NAD(P)H; and (iii) that resorcinol could act as an inducer of all catabolic pathway genes (39). Similar findings were reported also for the anaerobic metabolism of other aromatic compounds by *T. aromatica* (9, 20, 37) and *Azoarcus* sp. strain EbN1 (34).

Cosmid R⁺ contains all of the genes involved in resorcinol degradation by *A. anaerobius*. *A. anaerobius* and *T. aromatica* belong to the class *Betaproteobacteria* and are capable of growing at the expense of aromatic compounds under denitrifying conditions (32, 33, 40). A cosmid library with *A. anaerobius* chromosomal DNA was constructed and heterologously expressed in two *T. aromatica* strains, AR1 and K172, that are unable to metabolize resorcinol. Nevertheless, *T. aromatica* AR1 degrades α -resorcyate (3,5-dihydroxybenzoate) via the HHQ pathway (14, 33) and can therefore be expected to harbor genes similar to those of *A. anaerobius*. *T. aromatica* K172 is known to degrade neither resorcinol nor α -resorcyate. Transconjugants of both strains containing cosmid R⁺ were able to grow with resorcinol even after seven successive passages without resorcinol and the selective marker tetracycline (data not shown), demonstrating that cosmid R⁺ was stable in both hosts. *T. aromatica* transconjugants degraded resorcinol

at a rate similar to that of *A. anaerobius* (about 1 mM resorcinol day⁻¹) (data not shown). The increase in OD₅₇₈ was coupled to resorcinol degradation and to a simultaneous reduction of nitrate to nitrite by both transconjugants (Fig. 2A and B), indicating that at least the resorcinol hydroxylase gene(s) should be contained in cosmid R⁺, and this was confirmed by enzymatic tests. Resorcinol hydroxylase was measured in membrane fractions of both transconjugants prepared from cells grown with resorcinol but not from cells grown with an alternative substrate (Table 2). These results are in good agreement with previously reported data (32). As expected, no resorcinol hydroxylase was found in either wild-type strain of *T. aromatica*. To determine if the HHQ dehydrogenase-encoding gene(s) is on the cosmid R⁺, enzyme activities were measured in transconjugants and compared with those found in the wild-type strains. HHQ dehydrogenase activity was not detected in *T. aromatica* K172 (wild type), as expected, but was found in all other cell extracts (nonfractionated by ultracentrifugation) of both transconjugants grown on resorcinol or on a different aromatic substrate. In both transconjugants, the activities measured were in the same range as those measured in *A. anaerobius*. A twofold higher HHQ dehydrogenase activity than in *A. anaerobius* extracts was detected in cell extracts prepared from an AR1 transconjugant grown with α -resorcyate. Although *A. anaerobius* is unable to grow with α -resorcyate, this finding can be explained only by its structural similarity to resorcinol, which may make it act as a good gratuitous inducer of HHQ dehydrogenase. In this context, benzoate, which carries no hydroxyl groups, may act as a bad gratuitous inducer. It will be interesting to determine if other dihydroxybenzoates, trihydroxybenzenes, and trihydroxybenzoate isomers can act as inducers of both HHQ dehydrogenases of *A. anaerobius* and *T. aromatica* AR1. To determine if also the ring cleavage enzyme(s) is encoded by cosmid R⁺, membrane fractions of transconjugants grown with resorcinol were mixed with cytosolic fractions of the respective transconjugants grown with either resorcinol or an alternative carbon source and tested for acetate, malate, and succinate formation. These assay mixtures contained HHQ (1 mM), nitrate (4 mM), and NADH (1 mM).

TABLE 2. Specific activities of resorcinol hydroxylase in membrane fractions and HHQ dehydrogenase in unfractionated cell extracts of *A. anaerobius* and *T. aromatica* strains

Strain	Growth substrate	Sp act (mU mg ⁻¹ protein)	
		Resorcinol hydroxylase	HHQ dehydrogenase
<i>A. anaerobius</i>	Resorcinol	98	33
	Benzoate	0	8
<i>T. aromatica</i> AR1/R ⁺	Resorcinol	80	37
	α -Resorcyate	0	80
<i>T. aromatica</i> K172/R ⁺	Resorcinol	150	20
	Benzoate	0	3
<i>T. aromatica</i> AR1 wild type	α -Resorcyate	0	27
<i>T. aromatica</i> K172 wild type	Benzoate	0	0

TABLE 3. Properties of genes and gene products encoded by cosmid R⁺

Gene product	Gene properties		Protein properties			Protein correspondence				Protein with highest similarity
	Range	% GC	Length (amino acids)	Mol mass (kDa)	pI	% Identity	% Similarity	E value	Reference	
ORF1	1–1150	68	384	41.8	8.9	59	70	9e-104	gi 121608751 ref YP_996558.1	Transposase (<i>Verminephrobacter eiseniae</i>)
ORF2	1218–2147	63	309	33.3	6.0	61	77	1e-94	gi 118592793 ref ZP_01550182.1	ABC transporter substrate-binding protein (<i>Stappia aggregata</i>)
ORF3	2509–3618	67	369	38.9	6.3	85	90	1e-95	gi 56479427 ref YP_161016.1	Possibly flavin-dependent dehydrogenase (<i>Azoarcus</i> sp. strain EbN1)
ORF4	3757–4656	57	299	32.9	6.2	72	84	8e-121	gi 56478704 ref YP_160293.1	Possibly involved in regulation of phenolic degradation (<i>Azoarcus</i> sp. strain EbN1)
ORF5	5096–5701	62	201	22.7	5.3	70	78	1e-68	gi 89358401 ref ZP_01196223.1	Conserved hypothetical protein (<i>Xanthobacter autotrophicus</i>)
ORF6	5712–6680	62	322	36.7	6.7	61	72	1e-109	gi 110634534 ref YP_674742.1	Putative glutathione S-transferase (<i>Mesorhizobium</i> sp.)
RHS	6763–7659	62	298	33.2	7.5	55	70	5e-92	gi 118698587 ref ZP_01556663.1	Pyrogallol hydroxytransferase (<i>Burkholderia ambifaria</i>)
RHL	7708–10455	63	915	102.7	6.6	67	66	0.0	gi 118698588 ref ZP_01556664.1	Pyrogallol hydroxytransferase (<i>Burkholderia ambifaria</i>)
BQDHL	11332–13554	65	740	81.1	5.6	57	73	0.0	gi 119946408 ref YP_944088.1	Pyruvate dehydrogenase (<i>Psychromonas ingrahamii</i>)
BQDHS	13589–14854	69	421	43.3	5.9	55	70	2e-102	gi 83593214 ref YP_426966.1	Dihydrolipoamide acetyltransferase (<i>Rhodospirillum rubrum</i>)
BQDHM	14866–16263	65	465	49.6	6.2	68	83	3e-180	gi 23013388 ref ZP_00053288.1	Pyruvate/2-oxoglutarate dehydrogenase (<i>Magnetospirillum magnetotacticum</i>)
ORF7	16611–18260	68	549	60.3	7.7	38	59	6e-110	gi 56419244 ref YP_146562.1	Acetoin operon expression regulatory protein (<i>Geobacillus kaustophilus</i>)
ORF8	18310–18933	61	207	23.8	5.4	38	53	2e-38	gi 88940881 ref ZP_01146311.1	Putative protein-disulfide isomerase (<i>Acidiphilium cryptum</i>)
ORF9	19004–20239	65	411	42.6	9.8	54	69	3e-76	gi 71907514 ref YP_285101.1	Major facilitator superfamily MFS_1 (<i>Dechloromonas aromatica</i>)
ORF10	20236–21306	63	356	38.8	5.5	44	58	2e-66	gi 83308699 emb CAJ01609.1	Cobalamin synthesis protein cobW/p47k family protein (<i>Methylocapsa acidiphila</i>)
ORF11	21339–21920	65	193	21.8	7.1	54	67	2e-38	gi 91790635 ref YP_551587.1	Antibiotic biosynthesis monooxygenase (<i>Polaromonas</i> sp.)
ORF12	21954–23414	65	486	51.9	5.6	73	86	0.0	gi 118744335 ref ZP_01592329.1	Succinic semialdehyde dehydrogenase (<i>Geobacter lovleyi</i>)
BTDHL	23426–24336	68	303	31.3	6.0	45	63	4e-46	gi 27377791 ref NP_769320.1	Dehydrogenase (<i>Bradyrhizobium japonicum</i>)
BTDHS	24352–24957	60	201	22.2	10.4	39	53	1e-14	gi 78693872 ref ZP_00858386.1	Hypothetical protein (<i>Bradyrhizobium</i> sp.)
ORF13	25135–26388	65	417	46.6	5.5	48	68	1e-104	gi 119946410 ref YP_944090.1	Peptidase M24 (<i>Psychromonas ingrahamii</i>)
ORF14	26433–27548	61	371	40.8	6.4	51	67	3e-102	gi 74317595 ref YP_315335.1	NADH-dependent flavin oxidoreductase (<i>Thiobacillus denitrificans</i>)
ORF15	27752–29737	68	661	72.8	6.9	37	57	3e-117	gi 114567214 ref YP_754368.1	Acetoin operon expression regulatory protein (<i>Syntrophomonas wolfei</i>)

Substoichiometric amounts of acetate were formed in all assay mixtures containing cytosolic fractions of resorcinol-grown transconjugants. After 6 h, 0.21 mM acetate was formed with the cytosolic fraction of resorcinol-grown cells of strain AR1/R⁺ and 0.11 mM acetate was formed with the cytosolic fraction of strain K1721/R⁺. No malate or succinate was detected in these assay mixtures. No acetate was formed in the assay mixtures containing cytosolic fractions prepared from cells grown without resorcinol. These results demonstrate that

the genes for the whole resorcinol degradation pathway are encoded by cosmid R⁺.

Genes and gene organization on cosmid R⁺. Sequencing of both strands of the coding region of cosmid R⁺ resulted in sequence information for 29.88 kb, which comprised 22 open reading frames (ORFs). Table 3 and Figure 3 summarize the locations of all 22 genes, the number of amino acids encoded by each protein, and the percent similarity and percent identity of the amino acid sequences of all 22 proteins with known




FIG. 3. Organization of the 29,880-bp gene cluster of cosmid R⁺ involved in anaerobic resorcinol metabolism in *A. anaerobius*. Arrows indicate the direction of translation of each ORF. Genes encoding enzymes required for resorcinol degradation are shaded dark gray. Genes shown to be nonessential for resorcinol degradation in growth experiments with transposon insertion mutants are in light gray, and the genes encoding proteins so far unknown to be essential or nonessential for resorcinol degradation are unshaded. Sizes of gaps between genes are indicated in base pairs above the schematically drawn ORFs.

sequences in the databases. No stop codon was detected for *orf1*, indicating that this gene was incomplete. Except for *btdH1*, all of the other ORFs were preceded by putative, typical Shine-Dalgarno ribosomal binding sites, which are very similar to the consensus sequence AGGAGG located 4 to 13 nucleotides upstream of the starting codon. On the basis of a BLAST similarity search, an alternative start codon, GTG, was identified for *orf3*. Except for *orf3*, in all of the genes translation is initiated with ATG. Taking into account the ORF's direction of transcription and the size of the intergenic spaces, it appears that the coding region of cosmid R⁺ is organized into three putative operons that may form three transcriptional units. In the products encoded by *orf2*, *orf9*, *orf11*, and *btdH2*, 1, 12, 2, and 4 transmembrane helices were predicted, respectively. The analysis of the 29.88-kb coding region of cosmid R⁺ allowed us to tentatively assign functions to 20 out of 22 gene products—except that it showed that cosmid R⁺ contains all of the elements required for high-affinity nutrient acquisition systems (three transporters), detoxification (one glutathione-S-transferase), catabolism (10 oxidoreductases), and components with regulatory functions (σ factors, regulators, and a stress response system).

Isolation of mutants deficient in resorcinol utilization. To analyze the resorcinol degradation pathway of *A. anaerobius* in greater detail, in vitro transposon mutagenesis was carried out with cosmid R⁺ by using the EZ::TN<KAN2> transposon. This particular transposon is a very useful tool for studies aiming to identify essential genes since no polar effects of transposon insertions on downstream genes can take place. This is due to the fact that the transposon contains multiple *E. coli*-type promoters oriented outward in both directions (15). This procedure generated a reasonable number (3,000) of stable mutated cosmids in *E. coli*, 100 of which were isolated and sequenced. Sequencing of the transposon insertion sites revealed a duplication of 9 bp at each point of insertion, which is consistent with the properties of the EZ::TN<KAN2> transposon. Only 26 out of the 100 transposon insertional cosmids initially selected were found to contain one transposon inserted in the *A. anaerobius* chromosomal DNA of cosmid R⁺. Both strains of *T. aromatica*, AR1 and K172, were transformed by mating with the selected 26 cosmid R⁺ derivatives and screened for the ability to grow on resorcinol. In repeated experiments with at least three independent matings, we confirmed that 12 mutants, *T. aromatica* AR1 and K172 transconjugants, were unable to grow with resorcinol (2 mM) but were able to grow in the presence of kanamycin on benzoate (2 mM) and succinate (5 mM), respectively. These results strongly indicate that the failure to metabolize resorcinol was not due to

an unsuccessful transconjugation but rather was caused by the disruption of key enzymes involved in resorcinol degradation. Interestingly, cultures of AR1 transconjugants containing Mu_13 and Mu_36 cosmids developed a red-orange color during incubation with resorcinol which indicated that HHQ or HBQ was formed. In addition to the 12 mutants of *T. aromatica* K172, 1 more mutant, Mu_76, was also impaired in resorcinol degradation and growth with resorcinol. Therefore, we assume that this mutant had a defect in the gene encoding HHQ dehydrogenase, which in strain AR1 could be complemented by its chromosomal gene. Since it was shown in this study that α -resorcylate could induce *A. anaerobius* HHQ dehydrogenase from cosmid R⁺, we believe that resorcinol can act as an inducer of the *T. aromatica* AR1 chromosomal gene. Table 4 summarizes the transposon insertion sites in the coding region of cosmid R⁺ and the phenotypes of *T. aromatica*

TABLE 4. Transposon insertion sites in the genes of cosmid R⁺ and mutant phenotypes in recombinant *T. aromatica* strains AR-1 and K172

Mutant	Transposon insertion site	Growth ^a of strain K172		Growth ^a of strain AR1	
		Resorcinol (2 mM)	Succinate (2 mM)	Resorcinol (2 mM)	Benzoate (2 mM)
Mu_4	27028	–	+	–	+
Mu_13	24529	–	+	–	+
Mu_14	22541	+	ND	+	ND
Mu_16	21134	+	+	+	ND
Mu_21	10148	–	+	–	+
Mu_22	14692	–	+	–	+
Mu_31	3212	+	ND	+	ND
Mu_36	27319	–	+	–	+
Mu_40	7196	–	+	–	+
Mu_41	5656	+	ND	+	ND
Mu_42	17257	+	ND	+	ND
Mu_44	9933	–	+	–	+
Mu_45	21277	+	ND	+	ND
Mu_46	21371	+	ND	+	ND
Mu_51	12756	–	+	–	+
Mu_53	13965	–	+	–	+
Mu_54	1438	+	ND	+	ND
Mu_56	924	+	ND	+	ND
Mu_63	19259	+	ND	+	ND
Mu_64	1466	+	ND	+	ND
Mu_68	1527	+	ND	+	ND
Mu_73	8306	–	+	–	+
Mu_76	24230	–	+	+	ND
Mu_80	11364	–	+	–	+
Mu_82	925	+	ND	+	ND
Mu_93	26058	–	+	–	+

^a +, growth; –, no growth; ND, not determined.




FIG. 4. Percentage of carbon sources utilized by *T. aromatica* AR1 mutants at the end of 4 days of incubation with a mixture of resorcinol (black bars) and α -resorcylyate (gray bars).

AR1 and K172 transconjugants carrying mutated cosmids. To determine if host chromosomal genes could complement any of the defects carried by the 12 *T. aromatica* AR1 mutants unable to grow on resorcinol, we monitored the resorcinol consumption of mutants grown with nitrate (8 mM) on mixtures of resorcinol (2 mM) and α -resorcylyate (2 mM) over a 1-week period. These experiments were carried out with only one mutant for each key gene that had the transposon inserted nearest to the N terminus of the gene. In Fig. 4, the results obtained at the end of 4 days of growth are shown since after that the medium was almost depleted of the electron acceptor. The mutants were grouped into three classes. Mu_93 was able to metabolize both substrates at the same time but at different rates (data not shown). The Mu_21, Mu_40, Mu_53, and Mu_80 mutants did not degrade resorcinol at all but did degrade α -resorcylyate. The Mu_13 and Mu_36 mutants grew poorly and degraded both substrates partially at the same time. In addition, these cultures accumulated a red-orange color which was observed as well in the cultures incubated with resorcinol only. However, no intermediate was detected by HPLC, which might mean that either it was not stable and decayed during sample preparation, analysis, or storage or that it was present in undetectable amounts in the culture samples analyzed. Studies concerning these two mutants are in progress.

Functional assignment of the identified key gene products in resorcinol metabolism. In the present study, we have obtained biochemical evidence of NAD(P)H- and nitrate-dependent reactions involved in further degradation of HBQ to acetate, malate, and succinate. HHQ, the precursor of HBQ, is a common intermediate in the aerobic degradation of various aromatic compounds, such as resorcinol by *T. cutaneum* (12), *P. putida* (10), or *Corynebacterium glutamicum* (23); 4-nitrophenol by *Arthrobacter* sp. (24); 4-nitrocatechol by *Burkholderia cepacia* (11); and chlorohydroxyquinol by *B. cepacia* AC1100 (47). For all of the above pathways, except 4-nitrocatechol, it has been shown or proposed that the hydroxyhydroquinone intermediate is converted to maleylacetate, which is subsequently transformed to β -keto adipate, which is cleaved to succinate and acetate, which enter the trichloroacetic acid cycle. In the aerobic degradation of 4-nitrocatechol, hydroxyhydroquinone is converted to 1,4-benzenediol (hydroquinone) and γ -hydroxymuconic semialdehyde. Maleylacetate reductase and

CoA transferase, two enzymes characteristic of aerobic metabolism, were found to be insignificantly low and unstable in *A. anaerobius* (32). Moreover, none of the proteins encoded by cosmid R⁺ showed similarity to the corresponding proteins involved in the aerobic degradation of resorcinol or HHQ. All of these findings strongly indicate that *A. anaerobius* uses for resorcinol degradation a set of enzymes that differ from that which aerobes use when dealing with either resorcinol or HHQ. Mapping of transposon insertions in cosmid R⁺ derivatives of those mutants impaired in resorcinol utilization revealed that eight ORFs code for enzymes specific for resorcinol degradation in *A. anaerobius*. These products are encoded by *rhLS*, *bqdhLS*, *bidhLS*, *orf13*, and *orf14*.

Genes of resorcinol hydroxylase. The product of *rhLS* was identified as resorcinol hydroxylase by monitoring the growth and resorcinol degradation of *T. aromatica* transconjugants containing the cosmid R⁺ derivatives Mu_21, Mu_40, Mu_44, and Mu_73. In addition, mutants Mu_21 and Mu_40 were grown on resorcinol and α -resorcylyate and the membrane fractions were prepared and confirmed for resorcinol hydroxylase activity. The membrane fraction of a transconjugant containing cosmid R⁺ that was grown and prepared in the same way was used as a positive control. Neither activity nor color development was detected in assay mixtures with membranes prepared from mutants grown with resorcinol and α -resorcylyate (Table 5). According to BLAST analysis, the products of *rhL* and *rhS* are similar to anaerobic molybdopterin oxidoreductases and anaerobic dimethyl sulfoxide reductase-like enzymes. The best match for the *rhL* product to a biochemically studied enzyme was 52% identity to the α subunit of *Pelobacter acidigallici* pyrogallol-phloroglucinol transhydroxylase, and for the *rhS* product it was 49% identity to the β subunit of the same *P. acidigallici* transhydroxylase. Pyrogallol-phloroglucinol transhydroxylase is a soluble protein that catalyzes the conversion of pyrogallol to phloroglucinol (1,3,5-trihydroxybenzene). Its sequence is available (3), and its crystal structure was resolved as well (1, 28). The holoenzyme contains a molybdenum ion coordinated to two molybdopterin guanidine dinucleotide cofactors in the large subunit and three four-iron, four-sulfur clusters in the small subunit (3, 28). Binding sites for molybdenum and two molybdopterin guanidine dinucleotide molecules were identified in the *rhL* product, and eight conserved cysteine residues that could coordinate two iron-sulfur clusters were found in the *rhS* product. The cellular localization of this enzyme is still an unsolved problem. Resorcinol hydroxylase is predicted from the amino acid sequence to be soluble; however, it was measured almost exclusively in the membranes of *A. anaerobius* (32) and *T. aromatica* transconjugants. One may speculate that it could interact with an as-yet-unknown membrane anchor. On the basis of all of these findings, we propose

TABLE 5. Specific activities of resorcinol hydroxylase in the membrane fraction of *T. aromatica* AR1 transconjugants

Strain	Growth substrate	Resorcinol hydroxylase sp act (mU mg ⁻¹ protein)
<i>T. aromatica</i> AR1/R ⁺	Resorcinol- α -resorcylyate	31
<i>T. aromatica</i> AR1/Mu_21	Resorcinol- α -resorcylyate	0
<i>T. aromatica</i> AR1/Mu_40	Resorcinol- α -resorcylyate	0

produce *A. anaerobius* enzymes in *E. coli* to facilitate a detailed study of each protein and to finally elucidate the degradation pathway. Such studies are under way in our laboratory.

ACKNOWLEDGMENTS

This work was supported by the Deutsche Forschungsgemeinschaft, Bonn, Germany, and by research funds of the University of Constance. Work in Spain was financed by grant VEM2003-20075-CO2-01 from the Spanish Ministry of Science and Education. JIM-B work in Germany was partially supported by a Spanish-German exchange program (Acciones Integradas) of the Ministry of Science and Education.

REFERENCES

- Abt, D. J., O. Einsle, H. Niessen, R. Krieger, A. Messerschmidt, B. Schink, and P. M. Kroneck. 2002. Crystallization and preliminary X-ray analysis of the molybdenum-dependent pyrogallol-phloroglucinol transhydroxylase of *Pelobacter acidigallici*. *Acta Crystallogr. D Biol. Crystallogr.* **58**:343–345.
- Anders, H.-J., A. Kaetzke, P. Kämpfer, W. Ludwig, and G. Fuchs. 1995. Taxonomic position of aromatic-degrading denitrifying pseudomonad strains K 172 and KB 740 and their description as new members of the genera *Thauera*, as *Thauera aromatica* sp. nov., and *Azoarcus*, as *Azoarcus evansii* sp. nov., respectively, members of the beta subclass of the *Proteobacteria*. *Int. J. Syst. Bacteriol.* **45**:327–333.
- Baas, D., and J. Rety. 1999. Cloning, sequencing and heterologous expression of pyrogallol-phloroglucinol transhydroxylase from *Pelobacter acidigallici*. *Eur. J. Biochem.* **265**:896–901.
- Boll, M. 2005. Dearomatizing benzene ring reductases. *J. Mol. Microbiol. Biotechnol.* **10**:132–142.
- Boll, M. 2005. Key enzymes in the anaerobic aromatic metabolism catalysing Birch-like reductions. *Biochim. Biophys. Acta* **1707**:34–50.
- Boll, M., and G. Fuchs. 2005. Unusual reactions involved in anaerobic metabolism of phenolic compounds. *Biol. Chem.* **386**:989–997.
- Boll, M., G. Fuchs, and J. Heider. 2002. Anaerobic oxidation of aromatic compounds and hydrocarbons. *Curr. Opin. Chem. Biol.* **6**:604–611.
- Bradford, M. M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- Breinig, S., E. Schiltz, and G. Fuchs. 2000. Genes involved in anaerobic metabolism of phenol in the bacterium *Thauera aromatica*. *J. Bacteriol.* **182**:5849–5863.
- Chapman, P. J., and D. W. Ribbons. 1976. Metabolism of resorcinolic compounds by bacteria: alternative pathways for resorcinol catabolism in *Pseudomonas putida*. *J. Bacteriol.* **125**:985–998.
- Chauhan, A., S. K. Samanta, and R. K. Jain. 2000. Degradation of 4-nitro-catechol by *Burkholderia cepacia*: a plasmid-encoded novel pathway. *J. Appl. Microbiol.* **88**:764–772.
- Gaal, A., and H. Y. Neujahr. 1979. Metabolism of phenol and resorcinol in *Trichosporon cutaneum*. *J. Bacteriol.* **137**:13–21.
- Gallus, C., N. Gorny, W. Ludwig, and B. Schink. 1997. Anaerobic degradation of α -resorcylyate by a nitrate-reducing bacterium, *Thauera aromatica* strain AR-1. *Syst. Appl. Microbiol.* **20**:540–544.
- Gallus, C., and B. Schink. 1998. Anaerobic degradation of α -resorcylyate by *Thauera aromatica* strain AR-1 proceeds via oxidation and decarboxylation to hydroxyhydroquinone. *Arch. Microbiol.* **169**:333–338.
- Gerdes, S. Y., M. D. Scholle, M. D'Souza, A. Bernal, M. V. Baev, M. Farrell, O. V. Kurnasov, M. D. Daugherty, F. Mseeh, B. M. Polanuyer, J. W. Campbell, S. Anantha, K. Y. Shatalin, S. A. Chowdhury, M. Y. Fonstein, and A. L. Osterman. 2002. From genetic footprinting to antimicrobial drug targets: examples in cofactor biosynthetic pathways. *J. Bacteriol.* **184**:4555–4572.
- Gibson, J., and C. S. Harwood. 2002. Metabolic diversity in aromatic compound utilization by anaerobic microbes. *Annu. Rev. Microbiol.* **56**:345–369.
- Gorny, N., G. Wahl, A. Brune, and B. Schink. 1992. A strictly anaerobic nitrate-reducing bacterium growing with resorcinol and other aromatic compounds. *Arch. Microbiol.* **158**:48–53.
- Groseclose, E. E., and D. W. Ribbons. 1981. Metabolism of resorcinolic compounds by bacteria: new pathway for resorcinol catabolism in *Azotobacter vinelandii*. *J. Bacteriol.* **146**:460–466.
- Hawkins, C. F., A. Borges, and R. N. Perham. 1989. A common structural motif in thiamin pyrophosphate-binding enzymes. *FEBS Lett.* **255**:77–82.
- Heider, J., M. Boll, K. Breese, S. Breinig, C. Ebenau-Jehle, U. Feil, N. Gad'on, D. Laempe, B. Leuthner, M. E. Mohamed, S. Schneider, G. Burchhardt, and G. Fuchs. 1998. Differential induction of enzymes involved in anaerobic metabolism of aromatic compounds in the denitrifying bacterium *Thauera aromatica*. *Arch. Microbiol.* **170**:120–131.
- Heider, J., and G. Fuchs. 1997. Anaerobic metabolism of aromatic compounds. *Eur. J. Biochem.* **243**:577–596.
- Herrero, M., V. de Lorenzo, and K. N. Timmis. 1990. Transposon vectors containing non-antibiotic resistance selection markers for cloning and stable chromosomal insertion of foreign genes in gram-negative bacteria. *J. Bacteriol.* **172**:6557–6567.
- Huang, Y., K. X. Zhao, X. H. Shen, M. T. Chaudhry, C. Y. Jiang, and S. J. Liu. 2006. Genetic characterization of the resorcinol catabolic pathway in *Corynebacterium glutamicum*. *Appl. Environ. Microbiol.* **72**:7238–7245.
- Jain, R. K., J. H. Dreisbach, and J. C. Spain. 1994. Biodegradation of *p*-nitrophenol via 1,2,4-benzenetriol by an *Arthrobacter* sp. *Appl. Environ. Microbiol.* **60**:3030–3032.
- Kleiger, G., and D. Eisenberg. 2002. GXXXG and GXXXA motifs stabilize FAD and NAD(P)-binding Rossmann folds through C^α-H...O hydrogen bonds and van der Waals interactions. *J. Mol. Biol.* **323**:69–76.
- Kluge, C., A. Tschech, and G. Fuchs. 1990. Anaerobic metabolism of resorcylic acids (m-dihydroxybenzoic acids) and resorcinol (1,3-benzenediol) in a fermenting and in a denitrifying bacterium. *Arch. Microbiol.* **155**:68–74.
- Kretz, P. L., S. W. Kohler, and J. M. Short. 1991. Identification and characterization of a gene responsible for inhibiting propagation of methylated DNA sequences in *mcrA mcrBI Escherichia coli* strains. *J. Bacteriol.* **173**:4707–4716.
- Messerschmidt, A., H. Niessen, D. Abt, O. Einsle, B. Schink, and P. M. Kroneck. 2004. Crystal structure of pyrogallol-phloroglucinol transhydroxylase, an Mo enzyme capable of intermolecular hydroxyl transfer between phenols. *Proc. Natl. Acad. Sci. USA* **101**:11571–11576.
- Ngugi, D. K., M. K. Tsanuo, and H. I. Boga. 2005. *Rhodococcus opacus* strain RW, a resorcinol-degrading bacterium from the gut of *Macrotermes michaelseni*. *Afr. J. Biotechnol.* **4**:639–645.
- Nordin, K., M. Unell, and J. K. Jansson. 2005. Novel 4-chlorophenol degradation gene cluster and degradation route via hydroxyquinol in *Arthrobacter chlorophenolicus* A6. *Appl. Environ. Microbiol.* **71**:6538–6544.
- Pfennig, N. 1978. *Rhodocyclus purpureus* gen. nov. and sp. nov., a ring-shaped, vitamin B₁₂-requiring member of the family *Rhodospirillaceae*. *Int. J. Syst. Bacteriol.* **28**:283–288.
- Philipp, B., and B. Schink. 1998. Evidence of two oxidative reaction steps initiating anaerobic degradation of resorcinol (1,3-dihydroxybenzene) by the denitrifying bacterium *Azoarcus anaerobius*. *J. Bacteriol.* **180**:3644–3649.
- Philipp, B., and B. Schink. 2000. Two distinct pathways for anaerobic degradation of aromatic compounds in the denitrifying bacterium *Thauera aromatica* strain AR-1. *Arch. Microbiol.* **173**:91–96.
- Rabus, R., M. Kube, A. Beck, F. Widdel, and R. Reinhardt. 2002. Genes involved in the anaerobic degradation of ethylbenzene in a denitrifying bacterium, strain EbN1. *Arch. Microbiol.* **178**:506–516.
- Sambrook, J., and D. W. Russell. 1989. Molecular cloning: a laboratory manual, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schink, B., B. Philipp, and J. Müller. 2000. Anaerobic degradation of phenolic compounds. *Naturwissenschaften* **87**:12–23.
- Schmeling, S., A. Narmadakh, O. Schmitt, N. Gad'on, K. Schuhle, and G. Fuchs. 2004. Phenylphosphate synthase: a new phosphotransferase catalyzing the first step in anaerobic phenol metabolism in *Thauera aromatica*. *J. Bacteriol.* **186**:8044–8057.
- Schnell, S., F. Bak, and N. Pfennig. 1989. Anaerobic degradation of aniline and dihydroxybenzenes by newly isolated sulfate-reducing bacteria and description of *Desulfobacterium anilini*. *Arch. Microbiol.* **152**:556–563.
- Shingler, V. 2003. Integrated regulation in response to aromatic compounds: from signal sensing to attractive behaviour. *Environ. Microbiol.* **5**:1226–1241.
- Springer, N., W. Ludwig, B. Philipp, and B. Schink. 1998. *Azoarcus anaerobius* sp. nov., a resorcinol-degrading, strictly anaerobic, denitrifying bacterium. *Int. J. Syst. Bacteriol.* **48**:953–956.
- Staskawicz, B., D. Dahlbeck, N. Keen, and C. Napoli. 1987. Molecular characterization of cloned avirulence genes from race 0 and race 1 of *Pseudomonas syringae* pv. *glycinea*. *J. Bacteriol.* **169**:5789–5794.
- Sütfeld, R., F. Petereit, and A. Nahrstedt. 1996. Resorcinol in exudates of *Nuphar lutea*. *J. Chem. Ecol.* **22**:2221–2231.
- Tschech, A., and G. Fuchs. 1987. Anaerobic degradation of phenol by pure cultures of newly isolated denitrifying pseudomonads. *Arch. Microbiol.* **148**:213–217.
- Tschech, A., and B. Schink. 1985. Fermentative degradation of resorcinol and resorcylic acids. *Arch. Microbiol.* **143**:52–59.
- Vialaton, D., and C. Richard. 2002. Phototransformation of aromatic pollutants in solar light: photolysis versus photosensitized reactions under natural water conditions. *Aquat. Sci.* **64**:207–215.
- Wischgoll, S., D. Heintz, F. Peters, A. Erxleben, E. Sarnighausen, R. Reski, A. Van Dorsselaer, and M. Boll. 2005. Gene clusters involved in anaerobic benzoate degradation of *Geobacter metallireducens*. *Mol. Microbiol.* **58**:1238–1252.
- Zaborina, O., D. L. Daubaras, A. Zago, L. Xun, K. Saido, T. Klem, D. Nikolic, and A. M. Chakrabarty. 1998. Novel pathway for conversion of chlorohydroxyquinol to maleylacetate in *Burkholderia cepacia* AC1100. *J. Bacteriol.* **180**:4667–4675.