

Upregulation of Signalase Processing and Induction of prM-E Secretion by the Flavivirus NS2B-NS3 Protease: Roles of Protease Components

VLADIMIR F. YAMSHCHIKOV,^{1*} DENNIS W. TRENT,¹ AND RICHARD W. COMPANS²

Division of Viral Products, Center for Biologics Evaluation and Research, Food and Drug Administration, Bethesda, Maryland 20892,¹ and Department of Microbiology and Immunology, Emory University School of Medicine, Atlanta, Georgia 30322²

Received 21 November 1996/Accepted 4 March 1997

Recently, we have shown that the ability of the flavivirus NS2B-NS3 protease complex to promote efficient signalase processing of the C-prM precursor, as well as secretion of prM and E, does not appear to depend strictly on cleavage of the precursor at its Lys-Arg-Gly dibasic site by the protease. We suggested that the association of the protease with the precursor via NS2B may be sufficient by itself for the above effects. To study the proposed association in more detail, we have developed an assay in which processing at the C-prM dibasic cleavage site is abolished by Lys → Gly conversion. We constructed deletion mutants and chimeras of the West Nile (WN) flavivirus NS2B protein and expressed them in the context of {5'-C→NS3₂₄₃} containing either wild-type C-prM or its cleavage site mutant. All NS2B variants were able to form active protease complexes. Deletion of the carboxy-terminal cluster of hydrophobic amino acids in NS2B had no apparent effect on the formation of prM and prM-E secretion for the cassettes containing either wild-type or mutated C-prM precursor. Deletion of the amino-terminal hydrophobic cluster in NS2B did not affect prM-E secretion for the cassettes with wild-type C-prM but abrogated prM-E secretion for the cassettes with the mutated dibasic cleavage site in C-prM. Similarly, the NS2B-NS3₁₇₈ protease of Japanese encephalitis (JE) virus, when substituted for the WN virus NS2B-NS3₂₄₃ protease, was able to promote prM-E secretion for the cassette with the wild-type C-prM precursor but not with the mutated one. Replacement of the deleted amino-terminal hydrophobic cluster in the WN virus NS2B protein with an analogous JE virus sequence restored the ability of the protease to promote prM-E secretion. On the basis of these observations, roles of individual protease components in upregulation of C-prM signalase processing are discussed.

The flavivirus genome consists of an infectious single-stranded RNA molecule of approximately 11,000 nucleotides. The genome is translated into a large polyprotein, which is processed to individual flavivirus proteins by host cell and viral proteases. In the flavivirus polyprotein precursor, structural proteins (C, prM, and E) occupy the amino-terminal one-third of the polyprotein, with the rest of the precursor consisting of nonstructural (NS) proteins. The proteins are arranged in the order C-prM-E-NS1-NS2A-NS2B-NS3-NS4A-NS4B-NS5 (reviewed in reference 8). The majority of the nonstructural region in the precursor is processed by the viral protease (3, 15, 28), which functions as an NS2B-NS3 complex (6, 10, 12, 32) and cleaves at NS2A-NS2B-NS3-NS4A and NS4B-NS5 junctions.

In contrast, the majority of the structural region is processed by host cell signalases. While the prM-E and E-NS1 junctions appear to be processed cotranslationally (4, 25, 26, 33) both in vitro and in vivo, signalase cleavage of the C-prM precursor occurs less efficiently when assayed in vitro (22, 26). However, no such difference was observed in flavivirus-infected cells (9). It has been shown (19, 34) that signalase processing of the flavivirus C-prM precursor occurs efficiently only in the presence of the viral NS2B-NS3 protease. We have demonstrated (36) that the NS2B-NS3 protease specifically cleaves at the dibasic site in the C-prM precursor, with formation of the virion form of the capsid protein (C_{vir}) and presumably a prM precursor containing the signal sequence at its amino terminus.

Cleavage at this site was found to facilitate prM formation, apparently due to an increase in the efficiency of signalase processing of the latter precursor (1, 19, 30). However, it has been reported (37) that upregulation of C-prM signalase cleavage and secretion of prM and E could be observed even when the dibasic site in the C-prM was disrupted by Lys → Gly mutation, which abolished *trans* cleavage at this site by the protease in vitro (36). As one possible explanation, it was suggested that NS2B plays a specific role in upregulation of C-prM processing (37). An alternative explanation is that while disruption of the dibasic site indeed abolished cleavage by the protease in *trans*, processing at the mutated site may occur when the protease is presented in *cis* to the precursor. However, cleavage at some cryptic site or nonspecific degradation of C in the mutated precursor by the protease also cannot be ruled out.

In the present study, we investigated the ability of the protease complex to upregulate C-prM signalase processing and induction of prM and E secretion in *trans*. We also constructed deletion variants of the West Nile (WN) flavivirus NS2B protein and examined the effects of these deletions on C-prM signalase processing and secretion of prM and E. To address the specificity of these processes in more detail, chimeric cassettes containing homologous sequences from Japanese encephalitis (JE) virus were also studied in the above assays.

MATERIALS AND METHODS

Cells, viruses, and antisera. HeLa T4 cells (20) were used throughout this study as described previously (34). Anti-WN virus hyperimmune mouse ascites fluid was obtained from the American Type Culture Collection (Rockville, Md.),

* Corresponding author.




FIG. 1. Schematic representation of proteins encoded by expression cassettes used in this study. The 5'-terminal half of the WN virus genome, with the amino-terminal half of the encoded polyprotein drawn as an open bar, is presented in the middle. Individual flavivirus proteins are marked, with the vertical bars denoting their boundaries. Hydrophobic regions serving as signal and stop-transfer sequences are represented by the thick vertical bars. In the upper part of the figure, the thick horizontal bars represent the expression cassettes used for expression in the vaccinia virus-T7 polymerase system; open bars represent JE virus-specific sequences in chimeric cassettes. The lower part of the figure demonstrates mutated regions of the expression cassettes above. On the left side, the single-letter amino acid representation of the wild-type WN virus polyprotein sequence at the C-prM junction is shown. The carboxy terminus of C, the amino terminus of prM, and the viral protease and signalase cleavage sites are denoted below the sequence. The dibasic site KRG at the carboxy terminus of C which was disrupted by the Lys → Gly mutation is boxed. On the right side, an exploded view of the NS2B protein is shown. The amino- and carboxy-terminal hydrophobic regions are represented by the solid boxes and labeled N and C, respectively, and the 40-aa conserved domain (see the text for details) is represented by the open box. Dashed lines represent deletions which were introduced into NS2B, and the open bar in the NJH cassette represents the JE virus-specific sequence.

and JE virus-specific NS2B and NS3 mouse antisera (17) were supplied by C.-J. Lai (National Institute of Allergy and Infectious Diseases, Bethesda, Md.).

Construction of gene cassettes. DNA manipulations, PCR, and cloning were performed as described previously (21, 37). The gene cassettes used in this study are depicted in Fig. 1.

Construction of $W\{5'-C \rightarrow NS1\}$ and $W\{5'-C \rightarrow NS3_{243}\}$ expression cassettes was described previously (34, 35). The $W\{5'-C \rightarrow NS2A\}$ cassette was prepared by PCR amplification of the WN virus NS2A gene and building up the $W\{5'-C \rightarrow NS1\}$ cassette using the *NsiI* (at the beginning of NS2A) and *SalI* (in the polylinker) sites. All cassettes are controlled by the T7 promoter. Previously (37), we had constructed variants of these cassettes containing a mutant form of C, referred to below as $C(G_2)$, in which the dibasic cleavage site KR/G located at the C carboxy terminus in the C-prM precursor was disrupted by a Lys → Gly substitution. This mutation abolished the processing of the C-prM precursor by the viral NS2B-NS3₂₄₃ protease in vitro (36). Construction of the $W\{NS2A \rightarrow NS3_{243}\}$ cassette which encodes wild-type components of the viral protease was also described (34).

PCR mutagenesis was used to construct derivatives of the $W\{5'-C \rightarrow NS3_{243}\}$ and $W\{5'-C(G_2) \rightarrow NS3_{243}\}$ cassettes carrying variants of NS2B, which are depicted in Fig. 1. For construction of deletion variants, two DNA fragments corresponding to NS2A-NS2B(part) and NS2B(part)-NS3 regions upstream and downstream from the intended deletion in NS2B were obtained by PCR amplification. After purification, the fragments were ligated, the ligation products were digested with *NsiI* (situated at the beginning of the NS2A gene) and *BamHI* (situated in the amino-terminal part of the NS3 gene), and desired NS2B deletion variants were recovered after cloning into *NsiI* and *BamHI* sites of a plasmid carrying the $\{NS2A \rightarrow NS3_{243}\}$ cassette. In the (Δ H) variant, amino-terminal residues 2 to 52 of NS2B were removed; the (Δ AH) variant lacks carboxy-terminal amino acids 94 to 129 of NS2B. The integrity of the open reading frames (ORFs) in the resulting constructs was verified by sequencing. Subsequently, *NsiI*-*BamHI* fragments from selected plasmids were used to construct appropriate deletion derivatives of the $W\{5'-C \rightarrow NS3_{243}\}$ and $W\{5'-C(G_2) \rightarrow NS3_{243}\}$ cassettes.

The NS2B-NS3₂₄₃-encoding fragment was prepared by PCR amplification with the start and stop codons supplied by primers; it was cloned into the *SmaI* site of pGEM3 under control of the T7 promoter.

Both WN and JE virus cDNAs feature an *Eco47III* site at the same position in the end of NS2A gene, which was used for construction of chimeric WN/JE virus cassettes. This site is a part of the Leu-Ala-Leu encoding sequence, which is found in both WN (5) and JE (31) virus polyproteins 27 amino acids (aa) upstream from the NS2A/NS2B cleavage site. A $W/J\{5'-C \rightarrow NS3_{178}\}$ cassette (Fig. 1), and its $C(G_2)$ counterpart were prepared by joining of the *EcoRI*-

Eco47III fragment encoding the WN virus part of the cassette with the *Eco47III*-*HindIII* fragment encoding the JE virus part of the cassette. In both of these cassettes, C, prM, E, NS1, and most of NS2A are derived from the WN virus, while the carboxy-terminal 24 aa of NS2A, all of NS2B, and the NS3₁₇₈ protease domain are of JE virus origin; the vector supplied three unrelated amino acids and a stop codon. In the second pair of chimeric $W/J\{5'-C \rightarrow NS3_{243}\}$ cassettes (Fig. 1), only the amino-terminal hydrophobic cluster in WN NS2B (positions 2 to 52 of NS2B) was replaced with that from JE virus NS2B.

Expression, labeling, immunoprecipitation, and gel electrophoresis. Throughout this study, the vaccinia virus T7 polymerase system (14) was used for expression as described previously (37). In vivo labeling and immunoprecipitation were described previously (34), except that in cotransfection experiments, 5 μ g of a plasmid encoding the $\{5'-C \rightarrow NS2A\}$ reporter cassette and 2.5 μ g of a plasmid encoding the NS2B-NS3₂₄₃ protease cassette were used. Unless otherwise indicated, cells were labeled for 1 h with 100 μ Ci of a [³⁵S]Met-Cys mixture (Amersham, Arlington Heights, Ill.), washed twice with phosphate-buffered saline, and chased in the growth medium for 5 h. In general, immunoprecipitated samples were deglycosylated with endo- β -N-acetylglucosaminidase F and resolved by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) as described previously (37). Dried gels were quantitated with PhosphorImager and ImageQuant software (Molecular Dynamics), and band intensities were normalized to NS1 as an internal standard (37).

RESULTS

Construction of expression cassettes. The flavivirus protease is a complex of two proteins, NS2B and NS3 (6, 10, 12, 32), which both may contribute to the observed upregulation of C-prM processing by signalases. The function of the NS3 protease domain in this upregulation was studied previously (1, 19, 37). To investigate the role of NS2B in this process, we have constructed several expression cassettes encoding modified NS2B proteins. A hydrophobicity plot of WN virus NS2B reveals two clusters of hydrophobic amino acids (Fig. 2) surrounding the region which is critical for activity of the NS2B-NS3 protease complex (13). Using PCR deletion mutagenesis, we constructed NS2B variants in which either the amino-ter-




FIG. 2. Hydrophobicity profile of WN virus NS2B. The NS2B sequence (5) hydrophobicity was plotted by the method of Kyte and Doolittle (18). Above the plot, amino- and carboxy-terminal hydrophobic regions are shown as solid boxes and marked N and C, respectively. The conserved 40-aa domain (13) is shown as an open box.

terminal or the carboxy-terminal hydrophobic cluster was removed (Fig. 1). We also constructed two chimeric WN/JE virus cassettes. In the first chimeric cassette, designated W/J{5'-C→NS3₁₇₈}, the entire protease of JE virus was used instead of that of WN virus. To ensure proper formation of the amino terminus of JE virus NS2B, the carboxy-terminal 24 aa of JE virus NS2A (approximately 10% of its length), together with the native JE virus NS2A/NS2B cleavage site, were also included in this cassette. In the second chimeric cassette, only the amino-terminal hydrophobic cluster in WN virus NS2B was replaced with that from JE virus; the rest of the WN virus protease components were retained.

First, we cloned these NS2B variants as {NS2A→NS3_x} cassettes to verify whether ORFs remained intact and resulting protease complexes (upon transient expression from plasmids) retained protease activity. After such verification, genes encoding the protease components were transferred in the context of {5'-C→NS3_x} cassettes.

In our previous study (37), we used the W{5'-C(G₂)→NS3₂₄₃} cassette in which the dibasic site in the C-prM precursor was

disrupted by a Lys → Gly mutation. In vitro, this mutation abolished cleavage of the C-prM precursor by the viral protease supplied in *trans* (36). However, both upregulation of C-prM signalase cleavage and secretion of prM and E were observed upon in vivo expression of the above cassette. As one possible explanation, it was suggested (37) that association of the protease with the C-prM precursor by itself (without cleavage at the dibasic site) upregulates signalase cleavage and induces prM-E secretion. To verify this hypothesis and to identify the roles of the protease components, we prepared appropriate C(G₂) counterparts of all the above-mentioned cassettes (Fig. 1).

On the other hand, prM-E secretion observed for the cassette containing the mutated C-prM precursor could be simply explained by the ability of the protease to process the precursor either at the disrupted dibasic site or at the upstream RRS site (36) when the protease is present in *cis*. We have prepared an NS2B-NS3₂₄₃ cassette to determine whether the induction of secretion for the cassette with mutated C-prM could be observed when the protease is supplied in *trans*. Since NS2A has been shown to be required for proper processing of NS1 (11), this protein was also included in the reporter cassette, {C→NS2A}.

Effects of NS2B deletions on the activity of the NS2B-NS3₂₄₃ protease complex. Upon expression of an {NS2A→NS3₂₄₃} encoding cassette, two WN virus-specific proteins, NS2B and NS3₂₄₃, are readily detected in cell lysates by immunoprecipitation (32, 34, 36). Both proteins result from self-cleavage of the above precursor, thus indicating formation of the active protease complex (6, 10, 12, 32). A similar assay was successfully used (13) to show that deletions in hydrophobic regions of NS2B upstream or downstream from the 40-aa hydrophilic domain do not abolish protease activity. Using this assay, we first verified that our modifications in NS2B did not result in inactivation of the protease. As demonstrated in Fig. 3A (lane 2), deletion of the amino-terminal hydrophobic domain had no apparent effect on the activity of the (NΔH)NS2B-NS3₂₄₃ pro-




FIG. 3. Effect of modifications in NS2B on the formation of the active protease complexes. {NS2A→NS3₂₄₃} (A and B) or {5'-C→NS3_x} (C and D) cassettes were expressed with the vaccinia virus-T7 polymerase system, and cells were labeled for 1 h and lysed. Proteins were recovered by immunoprecipitation with WN virus-specific antisera (unless stated otherwise) and resolved by PAGE. Identified proteins are depicted beside the gels; positions of prestained molecular weight (in thousands) markers are also shown. (A) Lanes: 1, wild-type NS2B; 2, (NΔH)NS2B-containing cassette. (B) Lanes: 1, wild-type NS2B; 2, vaccinia virus-T7 polymerase-infected and mock-transfected cells; 3, (CΔH)NS2B-containing cassette. (C) Lanes: 1 and 4, wild-type W{5'-C→NS3₂₄₃}; 2 and 5, W/J{5'-C→NS3₁₇₈}; 3 and 6, W/J{5'-C(G₂)NS3₁₇₈} cassette. Proteins were immunoprecipitated with JE virus NS2B-specific (lanes 1 to 3) or NS3-specific (lanes 4 to 6) antisera. (D) Lanes: 1, wild-type W{5'-C→NS3₂₄₃}; 2, W/J{5'-C→NS3₂₄₃}; 3, W/J{5'-C(G₂)NS3₂₄₃} cassette.

tease. For comparison, expression of the wild-type protease is shown in lane 1. The only observed difference was an increase in the apparent mobility of the (NΔH)NS2B protein (calculated 10 kDa versus 14.4 kDa of wild-type NS2B) and of either the NS2A-(NΔH)NS2B or (NΔH)NS2B-NS3₂₄₃ precursor(s), which have similar calculated molecular masses. Both cassettes produce apparently the same NS3₂₄₃ protein (calculated 26.7 kDa), as expected.

In contrast, a shift toward processing at the alternative site(s) in NS2A and/or NS3₂₄₃ is profoundly exhibited by the cassette containing (CΔH)NS2B with the carboxy-terminal hydrophobic domain deleted (Fig. 3B, lane 3). The deletion removes 36 aa from NS2B; two possible dibasic cleavage sites in the carboxy-terminal region of NS2A are situated 33 aa (KKG) and 40 aa (KRS) upstream from the NS2A/NS2B junction. Alternative cleavage in this region of NS2A has been observed for JE and yellow fever (YF) viruses (17, 24). Utilization of either site would result in a protein with approximately the same calculated molecular mass as that for wild-type NS2B. Indeed, among other products, a protein with slightly lower mobility than that of wild-type NS2B can be observed in Fig. 3B (lane 3). Since a product with the same mobility as that of wild-type NS3₂₄₃ was also observed in the immunoprecipitates, we concluded that despite aberrant self-processing, expression of the above cassette results in formation of the active protease.

The correct processing of the JE virus NS2B-NS3₁₇₈ protease precursor was observed upon expression of the W/J{5'-C→NS3₁₇₈} chimeric cassette after immunoprecipitation of NS2B and NS3₁₇₈ with JE virus NS2B- and NS3-specific mouse antisera. As demonstrated in Fig. 3C, both JE virus protease-containing cassettes produced correctly processed NS3₁₇₈, since a polypeptide of the expected size (calculated molecular mass, 19.4 kDa) was observed (lanes 5 and 6) after immunoprecipitation with the JE virus NS3-specific antiserum. Apparently, the same polypeptide was coprecipitated with NS2B (calculated molecular mass, 14.3 kDa) by the JE virus NS2B-specific antiserum (lanes 2 and 3), indicating that both proteins form a complex (2, 7, 17). Neither WN virus NS2B (lane 1) nor NS3₂₄₃ (lane 4) was immunoprecipitated by the JE virus-specific antiserum.

Formation of the active protease complex containing WN/JE virus chimeric NS2B is demonstrated in Fig. 3D, where NS2B and NS3₂₄₃ proteins of the same size were immunoprecipitated from cell lysates upon expression of either the wild-type W{5'-C→NS3₂₄₃} cassette (lane 1) or both its NS2B chimeric derivatives (lanes 2 and 3).

From these data, we concluded that expression of all the above constructs results in the formation of active protease complexes. As compared with the wild-type {NS2A→NS3₂₄₃} cassette, both (NΔH)NS2B- and (CΔH)NS2B-containing cassettes produced correctly processed NS3₂₄₃, which contains the active center of the protease (3, 15). The JE virus protease precursor also appeared to function properly in the context of WN virus polyprotein-encoding cassettes, producing correctly processed components of the JE virus protease complex, and replacement of the amino-terminal hydrophobic domain of NS2B with the homologous sequence from JE virus NS2B also had no apparent effect on formation of the active protease.

Effects of deletions in NS2B on C-prM processing and secretion of prM-E heterodimers. Recently, Chambers et al. (7) have reported that mutations in hydrophobic regions of NS2B, while having little or no effect on the activity of the resulting protease, appeared to be deleterious for the formation of infectious virus. The precise stage at which such mutations block virus replication remains unknown. It has been suggested (37) that NS2B plays a certain role in the upregulation of C-prM

processing by the viral protease. In this study, we were interested in investigating whether these hydrophobic regions might be involved in such upregulation.

To address this question, we modified the W{5'-C→NS3₂₄₃} and W{5'-C(G₂)→NS3₂₄₃} cassettes (37) by replacing the wild-type NS2B gene with its (NΔH) variant carrying a deletion of the amino-terminal hydrophobic cluster (Fig. 1), and all four cassettes were expressed in the vaccinia virus-T7 polymerase system. As seen in Fig. 4A, all cassettes produced significant amounts of prM, E, and NS1 proteins, which were readily detectable in cell lysates by immunoprecipitation. In all cases, expression resulted in the formation of the active protease complexes, as indicated by the presence of NS2B (lanes 2 and 4) or (NΔH)NS2B (lanes 6 and 8), and NS3₂₄₃ proteins in immunoprecipitates. No apparent difference in the formation of intracellular prM, E, and NS1 was observed for all four cassettes. In agreement with the previous report (37), both cassettes with the wild-type NS2B produced secreted prM and E (Fig. 4A, lanes 1 and 3). However, while the (NΔH)NS2B-containing cassette with wild-type C-prM also produced secreted prM and E (lane 5), secretion of these proteins was almost abolished for the cassette containing the mutated C(G₂)-prM precursor (lane 7).

Mutations introduced into the carboxy-terminal hydrophobic cluster of NS2B were also shown (7) to be deleterious for virus replication, since no infectious progeny was recovered. To investigate whether it is also important for upregulation of C-prM processing by the protease, we modified the W{5'-C→NS3₂₄₃} and W{5'-C(G₂)→NS3₂₄₃} cassettes (37) by replacing the wild-type NS2B gene with its (CΔH) variant in which this hydrophobic cluster was deleted (Fig. 1). This deletion did not affect the secretion of prM and E for cassettes containing either wild-type C-prM (Fig. 4B, lanes 1 and 3) or mutated C(G₂)-prM (lanes 2 and 4).

In the above experiments, a 5-h chase period was required to detect secreted proteins in the media, and comparable amounts of intracellular prM were detected for all four cassettes at the end of the chase. However, as shown in Fig. 4C, efficient formation of prM could be observed even after 5-min labeling and chase periods in the presence of either wild-type NS2B-NS3₂₄₃ (lanes 1 and 2) or (CΔH)NS2B-NS3₂₄₃ (lanes 5 and 6) proteases, whereas significantly less prM was detected for both wild-type C-prM- and C(G₂)-prM-containing cassettes with the (NΔH)NS2B-NS3₂₄₃ protease (lanes 3 and 4).

From the above data, we concluded that the amino-terminal hydrophobic cluster of NS2B, but not its carboxy-terminal one, appears to be important for the upregulation of C-prM signalase processing and for the induction of prM-E secretion by the NS2B-NS3₂₄₃ protease.

trans induction of prM and E secretion by the NS2B-NS3₂₄₃ protease. Previously, it has been shown (36) that disruption of the dibasic cleavage site KR/G at the C-prM junction (25) by Lys → Gly mutation completely abolished the processing of this precursor by the viral protease in cell-free assays when the protease was supplied in *trans*. Disruption of the dibasic cleavage site, however, did not affect the secretion of prM and E upon expression of the W{5'-C(G₂)→NS3₂₄₃} cassette (37). On the one hand, this would suggest that the ability of the protease to upregulate the signalase processing of C-prM and to induce the secretion of prM and E does not depend on its ability to process the C-prM precursor at the upstream dibasic site. However, on the other hand, the observed secretion might still result from the cleavage at the disrupted dibasic site due to altered specificity of the protease acting in *cis*. For the YF virus protease, some tolerance to modifications in the dibasic motif has been reported (24). To address this question, we attempted




FIG. 4. Effects of deletions in NS2B on upregulation of C-prM signalase processing and secretion of prM-E heterodimers. The NS2B variants were expressed by using the vaccinia virus-T7 polymerase system in the context of the $\{5'-C \rightarrow NS3_{243}\}$ cassettes containing either wild-type C-prM or C(G₂)-prM as described in Materials and Methods. Proteins were recovered by immunoprecipitation with the WN virus-specific antiserum, deglycosylated with PNGase F, and resolved by PAGE. Identified proteins are depicted. (A) Lanes: 1 and 2, W{5'-C→NS3₂₄₃} ; 3 and 4, W{5'-C(G₂)→NS3₂₄₃} ; 5 to 8, the same as in lanes 1 to 4, respectively, but for (ΔH)NS2B-containing cassettes. Odd-numbered lanes, medium samples; even-numbered lanes, cell lysate samples. On the graph below, band intensities of prM, E, and NS1 were normalized to amounts of intracellular NS1 for each cassette, and numbers under the graph correspond to the lane numbers on the gel. (B) Lanes: 1, W{5'-C→NS3₂₄₃} ; 2, W{5'-C(G₂)→NS3₂₄₃} ; 3 and 4, the same as in lanes 1 and 2, respectively, but for (ΔH)NS2B-containing cassettes. Only medium samples are shown. On the graph below, band intensities of prM, E, and NS1 were normalized to amounts of secreted NS1 for each cassette. (C) Cell lysate samples obtained after 5 min of labeling and 5 min of chase. Lanes: 1 and 2, wild-type NS2B; 3 and 4, (ΔH)NS2B; 5 and 6, (CΔH)NS2B-containing cassettes. Odd-numbered lanes, wild-type C-prM; even-numbered lanes, C(G₂)-prM-containing cassettes. Samples were not deglycosylated; E and NS1 are not resolved (34). On the graph below, band intensities of prM and E plus NS1 were normalized to the total of E plus NS1 for each cassette.

to determine whether induction of prM and E secretion depends specifically on the presence of the protease in *cis*. As demonstrated in Fig. 5, secretion of prM and E may be observed in cotransfection experiments upon coexpression of the NS2B-NS3₂₄₃ cassette with {C→NS2A} cassettes containing either the wild-type C-prM (lane 1) or that with the mutated dibasic site (lane 3). In agreement with previous reports (1, 19, 23, 34), secretion of prM and E was not observed upon expression of either {C→NS2A} cassette without the protease (results not shown). In contrast, proper processing and secretion of NS1 do not require the presence of the protease (4, 7–9, 19, 23, 24, 27, 33, 34) but depend on the presence of downstream NS2A protein (11).

From the data presented above, we conclude that induction of prM and E secretion from the cassette with an uncleavable C-prM precursor at the dibasic site cannot be explained by the altered specificity of the *cis*-acting protease.

Effects of heterologous and chimeric proteases on C-prM processing and secretion of prM-E heterodimers. The above findings indicate that the ability of the NS2B-NS3₂₄₃ protease to upregulate the signalase processing of C-prM and induce the secretion of prM and E does not appear to depend on its ability to process the C-prM precursor at the upstream dibasic site. In this study, we were interested in addressing the specificity and roles of individual protease components in this process.

For this purpose, we used the W/J{5'-C→NS3₁₇₈} cassette in which the entire WN virus protease was replaced with that of JE virus. Such replacement had no apparent effect on the secretion of prM and E or on the formation of intracellular prM (Fig. 6A, lanes 3 and 4; compare with lanes 1 and 2) for the cassette containing wild-type C-prM. However, the secretion of prM and E was greatly reduced for the chimeric cassette with the mutated C-prM precursor (lanes 7 and 8), compared to the same cassette encoding the homologous protease (lanes 5 and 6), thus apparently resembling the behavior of the (ΔH)NS2B-containing cassette. The important role of the amino-terminal hydrophobic cluster of NS2B in upregulation and induction of prM-E secretion was demonstrated above, thus providing one possible explanation. If this domain is involved in specific interaction of the protease with the C-prM precursor, heterologous NS2B may fail to provide the necessary association. To address this question, we replaced only the amino-terminal domain in WN virus NS2B with that of JE virus NS2B, but the rest of the WN virus protease was retained. While both these viruses belong to the same group and their polyprotein sequences display a high degree of sequence identity (5, 31), there are 14 amino acid differences in the 54-aa amino-terminal hydrophobic domains of their NS2B proteins. As seen in Fig. 6B, such replacement did not affect the ability of the protease to induce the secretion of prM and E for both wild-type and mutated C-prM-containing cassettes, indicating




FIG. 5. Upregulation of C-prM signalase cleavage and induction of prM-E secretion by the NS2B-NS3₂₄₃ protease supplied *in trans*. W{5'-C→NS2A} cassettes containing either wild-type C-prM (lanes 1 and 2) or C(G₂)-prM (lanes 3 and 4) were coexpressed with the NS2B-NS3₂₄₃ protease-encoding cassette, and proteins were recovered by immunoprecipitation with WN virus-specific antiserum, deglycosylated with PNGase F, and resolved by PAGE. Odd-numbered lanes, medium samples; even-numbered lanes, cell lysate samples. Identified proteins and positions of prestained markers are depicted. The band corresponding to NS3₂₄₃ is more clearly visible upon longer exposure; however, this resulted in a lack of resolution of E and NS1 in lanes 2 and 4. On the graph below, band intensities of prM, E, and NS1 were normalized to amounts of intracellular NS1 for each cassette, and numbers under the graph correspond to the lane numbers on the gel.

that involvement of the amino-terminal hydrophobic cluster of NS2B is probably not highly sequence specific.

From the present experiments, we conclude that the JE virus protease is able to process the WN virus C-prM precursor at its dibasic site. With the intact site, such processing results in upregulation of signalase cleavage and induction of prM and E secretion. However, when cleavage is blocked due to disruption of the dibasic motif, the JE virus protease fails to induce prM and E secretion. Thus, the chimeric cassettes containing the entire JE virus protease appear to resemble the (NΔH) NS2B-containing cassettes. However, from the latter experiments, we conclude that the NS2B amino-terminal hydrophobic domain alone cannot provide for the observed specificity of the protease.

DISCUSSION

Previously, it has been shown (19, 34) that both efficient processing of the C-prM precursor by host cell signalases and secretion of the prM and E require the presence of the active viral NS2B-NS3₂₄₃ protease. Subsequently, specific cleavage by the viral protease at the dibasic site situated at the carboxy terminus of the C protein in the C-prM precursor has been demonstrated *in vitro* (36). It was observed, however, that disruption of the dibasic site in the C-prM precursor by a Lys → Gly mutation had no effect on upregulation of prM formation and did not abolish the secretion of prM and E (37),

even though mutant C-prM was not processed *in vitro* by the protease supplied *in trans* (36). This might indicate that both prM upregulation and induction of prM-E secretion constitute a distinct function of the protease and do not appear to be simply the consequence of prior cleavage of C-prM at the dibasic site. However, an alternative explanation cannot be ruled out. It suggests that in *cis*, in contrary to the *trans* assays mentioned above, the protease may be less specific and, when expressed as part of the same polyprotein, is able to process C-prM at the modified dibasic site. Indeed, some tolerance for modifications in the dibasic motif was observed for YF virus protease (24). It was of interest to determine if the same effects of upregulation and secretion could be observed for the mutated C-prM-containing cassettes with the protease supplied *in trans*, i.e., by coexpression of the protease components from a separate cassette. In this study, we found that both upregulation of prM formation and induction of prM-E secretion can be successfully achieved upon cotransfection of {C→NS2A} reporter and {NS2B-NS3₂₄₃} cassettes, and the protease complex appeared to be equally effective for both wild-type and mutated C-prM-containing cassettes. Since the protease failed to process mutated C-prM *in trans* (36) but retained its ability to upregulate prM formation and induce prM-E secretion when present either *in cis* (37) or *in trans* (this study), neither process seems to depend solely on the above cleavage event. It has been shown, however, that signalase cleavage occurs efficiently after treatment of membrane-associated C-prM with trypsin (30), which may result in rather nonspecific degradation of C. Utilization of the cryptic site(s) or nonspecific degradation of C in the presence of the viral protease was not observed *in vitro* (36). Strictly, this still cannot be ruled out *in vivo*. However, we consider such nonspecific degradation by the viral protease to be a very unlikely scenario. We suggest that association of the protease complex with the C-prM precursor by itself might be sufficient for upregulation of prM formation and induction of prM-E secretion. Such association may be responsible for the partial escape of prM and E antigens in nonvirion form (23) from flavivirus-infected cells (27, 29).

Since the flavivirus protease consists of two proteins (6, 10, 12, 32), we were interested in investigating the role that each component protein plays in the above processes. Earlier (34, 37), we had suggested that NS2B may be responsible for the association of the protease complex with the C-prM precursor. If so, this protein might contain a specific region which is crucial for such an association, and deletion of this region would affect the upregulation and induction of secretion for the cassette containing the uncleavable C-prM precursor. In this regard, the amino- and carboxy-terminal hydrophobic clusters in NS2B were of particular interest. It has been reported that while deletions in the amino- or carboxy-terminal hydrophobic clusters of NS2B did not impair the proteolytic activity of the protease (13), modifications in these regions appeared to be deleterious for the formation of an infectious virus progeny (7). We found that the presence or absence of the carboxy-terminal hydrophobic cluster had no effect on the ability of the protease to upregulate prM formation and induce prM-E secretion for cassettes containing either wild-type or mutated C-prM. In contrast, deletion of the amino-terminal hydrophobic cluster nearly abolished prM-E secretion for the cassette with mutated but not wild-type C-prM. These results imply that for the wild-type cassette, as also might occur in flavivirus infection, the effect of the protease on processing of the C-prM region may indeed be a composite result of two distinct mechanisms: (i) association of the protease complex with the C-prM precursor and (ii) protease cleavage at the dibasic site in the




FIG. 6. Expression of WN/JE virus chimeric $\{5'-C \rightarrow NS3_v\}$ cassettes. The cassettes containing either wild-type C-prM or C(G₂)-prM were expressed in the vaccinia virus-T7 polymerase system, and proteins were recovered by immunoprecipitation with WN virus-specific antiserum, deglycosylated with PNGase F, and resolved by PAGE. Identified proteins and positions of prestained markers are depicted. On the graphs below the gel, band intensities of prM, E, and NS1 were normalized to amounts of intracellular NS1 for each cassette, and numbers under the graphs correspond to the lane numbers on the gels. (A) Lanes: 1 and 2, $W\{5'-C \rightarrow NS3_{243}\}$; 3 and 4, $W/J\{5'-C \rightarrow NS3_{178}\}$; 5 and 6, $W\{5'-C(G_2) \rightarrow NS3_{243}\}$; 7 and 8, $W/J\{5'-C(G_2) \rightarrow NS3_{178}\}$. Odd-numbered lanes, medium samples; even-numbered lanes, cell lysate samples. (B) Lanes: 1 and 2, $W\{5'-C \rightarrow NS3_{243}\}$; 3 and 4, $W/J\{5'-C \rightarrow NS3_{243}\}$; 5 and 6, $W\{5'-C(G_2) \rightarrow NS3_{243}\}$; 7 and 8, $W/J\{5'-C(G_2) \rightarrow NS3_{243}\}$. Odd-numbered lanes, medium samples; even-numbered lanes, cell lysate samples.

precursor. Both of these mechanisms, in the case of wild-type C-prM, may contribute to upregulation of signalase cleavage and induction of prM-E secretion. In the cassettes with the modified dibasic sites, however, cleavage of C-prM by the protease becomes impossible, but association of the protease with the precursor still can effect upregulation of its signalase processing and induction of prM-E secretion. The present data indicate that NS2B indeed plays an important role in both these processes. In contrast to the wild-type and its (ΔH) variant, the (NΔH)NS2B-containing protease has lost its ability to facilitate prM formation and prM-E secretion for the cassette with an uncleavable dibasic site in C-prM. Apparently, either deletion of the amino-terminal hydrophobic domain impaired the association of the protease with C-prM precursor, or such an association has lost its ability to upregulate signalase cleavage and induce secretion. Thus, in the doubly mutated cassette containing uncleavable C-prM and (NΔH)NS2B, both effects are blocked and secretion of prM and E was not observed.

Subsequently, we investigated the specificity of such association of the protease with the C-prM precursor. To address this question, first we replaced the entire WN virus protease with that of JE virus. Both viruses belong to the same serologic group and share extensive amino acid sequence identity, especially for NS3 and NS5 proteins. We have found that while expression of the chimeric $W/J\{5'-C \rightarrow NS3_{178}\}$ cassette resulted in efficient prM formation and prM-E secretion, the C(G₂)-prM counterpart of this cassette failed to produce secreted prM and E. Since secretion was observed with the wild-

type dibasic site in C-prM, the JE virus protease was probably capable of cleaving the WN virus C-prM precursor at its dibasic site. However, since no secretion was observed when the dibasic site was rendered uncleavable, the JE virus protease evidently was not capable of proper association with the C-prM precursor. Thus, the JE virus protease expressed in the context of the WN virus reporter cassette had properties resembling the (NΔH)NS2B-containing WN virus protease. On the basis of the above results demonstrating the importance of the N-terminal hydrophobic domain in NS2B, we speculated that this domain provides for the observed specificity of association of the protease complex with the C-prM precursor. To verify this hypothesis, we prepared a pair of $W/J\{5'-C \rightarrow NS3_{243}\}$ chimeric cassettes [containing either wild-type C-prM or C(G₂)-prM] in which only the N-terminal hydrophobic domain in WN virus NS2B was replaced with that from JE virus NS2B. The chimeric cassettes, however, appeared to be almost indistinguishable from the parental $W\{5'-C \rightarrow NS3_{243}\}$ and $W\{5'-C(G_2) \rightarrow NS3_{243}\}$ WN virus cassettes, since efficient formation of prM and prM-E secretion was again observed for both wild-type and mutated C-prM. Thus, the N-terminal hydrophobic domain does not seem to be solely responsible for the observed specificity of protease association with C-prM. The JE virus protease used in the above experiments included only 178 aa of NS3, which is just enough to incorporate only the protease active center (3, 15). The WN virus protease, however, included 243 amino acids of NS3, thus also incorporating the amino-terminal part

of the NS3 helicase domain (16), whose contribution to the observed specificity cannot be ruled out.

The results of this study support and further elaborate the model of late events in flavivirus polyprotein processing and virion formation that we proposed previously (34). The apparent complexity of C-prM processing suggests that this region might constitute a central point of flavivirus replication. Elucidation of its molecular mechanisms may uncover approaches to new generations of safe vaccines.

ACKNOWLEDGMENTS

We thank C.-J. Lai (NIAID, Bethesda, Md.) for JE NS2B- and NS3-specific antisera and B. Falgout (FDA/CBER, Bethesda, Md.) for critical comments.

This study was supported in part by research award VRD/V22/181/72 from the World Health Organization.

REFERENCES

- Amberg, S. M., A. Nestorowicz, D. W. McCourt, and C. M. Rice. 1994. NS2B-3 proteinase-mediated processing in the yellow fever virus structural region: in vitro and in vivo studies. *J. Virol.* **68**:3794-3802.
- Arias, C. F., F. Preugschat, and J. M. Strauss. 1993. Dengue 2 virus NS2B and NS3 form a stable complex that can cleave NS3 within the helicase domain. *Virology* **193**:888-899.
- Bazan, J. F., and R. J. Fletterick. 1989. Detection of a trypsin-like serine protease domain in flaviviruses and pestiviruses. *Virology* **171**:637-639.
- Biedrzycka, A., M. R. Cauchi, A. Bartholomusz, J. J. Gorman, and P. J. Wright. 1987. Characterization of protease cleavage sites involved in the formation of envelope glycoprotein and three nonstructural proteins of dengue virus type 2, New Guinea C strain. *J. Gen. Virol.* **68**:1317-1326.
- Castle, E., U. Leidner, T. Nowak, G. Wengler, and G. Wengler. 1986. Primary structure of the West Nile flavivirus genome region coding for all nonstructural proteins. *Virology* **149**:10-26.
- Chambers, T. J., A. Grakoui, and C. M. Rice. 1991. Processing of yellow fever virus nonstructural polyprotein: a catalytically active NS3 proteinase domain and NS2B are required for cleavages at dibasic sites. *J. Virol.* **65**:6042-6050.
- Chambers, T. J., A. Nestorowicz, S. M. Amberg, and C. M. Rice. 1993. Mutagenesis of the yellow fever virus NS2B protein: effects on proteolytic processing, NS2B-NS3 complex formation, and viral replication. *J. Virol.* **67**:6797-6807.
- Chambers, T. J., C. S. Hahn, R. Galler, and C. M. Rice. 1990. Flavivirus genome organization, expression and replication. *Annu. Rev. Microbiol.* **44**:649-688.
- Chambers, T. J., D. W. McCourt, and C. M. Rice. 1990. Production of yellow fever virus proteins in infected cells: identification of discrete polyprotein species and analysis of cleavage kinetics using region-specific polyclonal antisera. *Virology* **177**:159-174.
- Chambers, T. J., R. C. Weir, A. Grakoui, D. W. McCourt, J. F. Bazan, R. J. Fletterick, and C. M. Rice. 1990. Evidence that the N-terminal domain of nonstructural protein NS3 from yellow fever virus is a serine protease responsible for site-specific cleavages in the viral polyprotein. *Proc. Natl. Acad. Sci. USA* **87**:8898-8902.
- Falgout, B., R. Chanock, and C.-J. Lai. 1989. Proper processing of dengue virus nonstructural glycoprotein NS1 requires the N-terminal hydrophobic signal sequence and the downstream nonstructural protein NS2a. *J. Virol.* **63**:1852-1860.
- Falgout, B., M. Pethel, Y.-M. Zhang, and C.-J. Lai. 1991. Both nonstructural protein NS2B and NS3 are required for the proteolytic processing of dengue virus nonstructural proteins. *J. Virol.* **65**:2467-2475.
- Falgout, B., R. H. Miller, and C.-J. Lai. 1993. Deletion analysis of dengue virus type 4 nonstructural protein NS2B: identification of a domain required for NS2B-NS3 proteinase activity. *J. Virol.* **67**:2034-2042.
- Fuerst, T. R., E. G. Niles, F. W. Studier, and B. Moss. 1986. Eukaryotic transient-expression system based on recombinant vaccinia virus that synthesizes bacteriophage T7 RNA polymerase. *Proc. Natl. Acad. Sci. USA* **83**:8122-8126.
- Gorbalenya, A. E., A. P. Donchenko, E. V. Koonin, and V. M. Blinov. 1989. N-terminal domains of putative helicases of flavi- and pestiviruses may be serine proteases. *Nucleic Acids Res.* **17**:3889-3897.
- Gorbalenya, A. E., E. V. Koonin, A. P. Donchenko, and V. M. Blinov. 1989. Two related superfamilies of putative helicases involved in replication, recombination, repair and expression of DNA and RNA genomes. *Nucleic Acids Res.* **17**:4713-4729.
- Jan, L. R., C. S. Yang, D. W. Trent, B. Falgout, and C. J. Lai. 1995. Processing of Japanese encephalitis virus non-structural proteins: NS2B-NS3 complex and heterologous proteases. *J. Gen. Virol.* **76**:573-580.
- Kyte, J., and R. F. Doolittle. 1982. A method for displaying the hydropathic character of a protein. *J. Mol. Biol.* **157**:105-132.
- Lobigs, M. 1993. Flavivirus pre-membrane protein cleavage and spike heterodimer secretion require the function of the viral proteinase NS3. *Proc. Natl. Acad. Sci. USA* **90**:6218-6222.
- Maddon, P. J., A. G. Dalgleish, J. S. McDougal, P. R. Clapham, R. A. Weiss, and R. Axel. 1986. The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. *Cell* **47**:333-348.
- Maniatis, T., E. F. Fritsch, and J. Sambrook. 1982. *Molecular cloning: a laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Markoff, L. 1989. In vitro processing of dengue virus structural proteins: cleavage of the pre-membrane protein. *J. Virol.* **63**:3345-3352.
- Mason, P. W., S. Pincus, M. J. Fournier, T. L. Mason, R. E. Shope, and E. Paoletti. 1991. Japanese encephalitis virus-vaccinia recombinants produce particulate forms of the structural membrane proteins and induce high levels of protection against lethal JEV infection. *Virology* **180**:294-305.
- Nestorowicz, A., T. J. Chambers, and C. M. Rice. 1994. Mutagenesis of the yellow fever virus NS2A/2B cleavage site: effects on proteolytic processing, viral replication and evidence for alternative processing of the NS2A protein. *Virology* **199**:114-123.
- Nowak, T., P. M. Farber, G. Wengler, and G. Wengler. 1989. Analyses of the terminal sequences of West Nile virus structural proteins and of the in vitro translation of these proteins allow the proposal of a complete scheme of the proteolytic cleavages involved in their synthesis. *Virology* **169**:365-376.
- Ruiz-Linares, A., A. Cahour, P. Despres, M. Girard, and M. Bouloy. 1989. Processing of yellow fever virus polyprotein: role of cellular proteases in maturation of the structural proteins. *J. Virol.* **63**:4199-4209.
- Russell, P. K., W. E. Brandt, and J. M. Dalrymple. 1980. Chemical and antigenic structure of flaviviruses, p. 503-529. *In* R. W. Schlesinger (ed.), *The togaviruses*. Academic Press, Inc., New York, N.Y.
- Speight, G., G. Coia, M. D. Parker, and E. G. Westaway. 1988. Gene mapping and positive identification of the nonstructural proteins NS2A, NS2B, NS3, NS4B, and NS5 of the flavivirus Kunjin and their cleavage sites. *J. Gen. Virol.* **69**:23-34.
- Stevens, T. M., and R. W. Schlesinger. 1965. Studies on nature of dengue viruses. 1. Correlation of particle density, infectivity, and RNA content of type 2 virus. *Virology* **27**:103-112.
- Stocks, C. E., and M. Lobigs. 1995. Posttranslational signal peptidase cleavage at the flavivirus C-prM junction in vitro. *J. Virol.* **69**:8123-8126.
- Sumiyoshi, H., C. Mori, I. Fuke, K. Morita, S. Kuhara, J. Kondou, Y. Kukushi, H. Nagamatu, and A. Igarashi. 1987. Complete nucleotide sequence of the Japanese encephalitis virus genome RNA. *Virology* **161**:497-510.
- Wengler, G., G. Czaya, P. M. Farber, and J. H. Hegemann. 1991. In vitro synthesis of West Nile virus proteins indicates that the amino-terminal segment of the NS3 protein contains the active center of the protease which cleaves the viral polyprotein after multiple basic amino acids. *J. Gen. Virol.* **72**:851-858.
- Wright, P. S., M. R. Cauchi, and M. L. Ng. 1989. Definition of the carboxy termini of the three glycoproteins specified by dengue virus type 2. *Virology* **171**:61-67.
- Yamshchikov, V. F., and R. W. Compans. 1993. Regulation of late events in flavivirus polyprotein processing and maturation. *Virology* **192**:38-51.
- Yamshchikov, V. F., and R. W. Compans. 1994. Generation of long flavivirus expression cassettes by in vivo recombination and transient dominant selection. *Gene* **149**:193-201.
- Yamshchikov, V. F., and R. W. Compans. 1994. Processing of the intracellular form of the West Nile virus capsid protein by the viral NS2B-NS3 protease: an in vitro study. *J. Virol.* **68**:5765-5771.
- Yamshchikov, V. F., and R. W. Compans. 1995. Formation of the flavivirus envelope: role of the viral NS2B-NS3 protease. *J. Virol.* **69**:1995-2003.