

A Small Region of the Ecotropic Murine Leukemia Virus (MuLV) *gag* Gene Profoundly Influences the Types of Polytypic MuLVs Generated in Mice

MARC LAVIGNON,¹† JENNIFER RICHARDSON,² AND LEONARD H. EVANS^{1*}

Laboratory of Persistent Viral Diseases, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, Hamilton, Montana 59840,¹ and Institut Cochin de Genetique Moleculaire, INSERM U363, 75014 Paris, France²

Received 27 May 1997/Accepted 7 August 1997

The vast majority of recombinant polytypic murine leukemia viruses (MuLVs) generated in mice after infection by ecotropic MuLVs can be classified into two major antigenic groups based on their reactivities to two monoclonal antibodies (MAbs) termed Hy 7 and 516. These groups very likely correspond to viruses formed by recombination of the ecotropic MuLV with two distinct sets of polytypic *env* genes present in the genomes of inbred mouse strains. We have found that nearly all polytypic MuLVs identified in mice infected with a substrain of Friend MuLV (F-MuLV₅₇) are reactive with Hy 7, whereas mice infected with Moloney MuLV (Mo-MuLV) generate major populations of both Hy 7- and 516-reactive polytypic MuLVs. We examined polytypic MuLVs generated in NFS/N mice after inoculation with Mo-MuLV-F-MuLV₅₇ chimeras to determine which regions of the viral genome influence this difference between the two ecotropic MuLVs. These studies identified a region of the MuLV genome which encodes the nucleocapsid protein and a portion of the viral protease as the only region that influenced the difference in polytypic-MuLV generation by Mo-MuLV and F-MuLV₅₇.

Mice inoculated with exogenous ecotropic murine leukemia viruses (MuLVs) or mice expressing endogenous ecotropic MuLVs frequently generate polytypic viruses which have been implicated in the induction of a number of proliferative diseases in mice (1, 3, 15, 16, 18, 21, 22, 25, 30). These viruses, also termed dualtropic or mink cell focus-forming MuLVs, utilize a cellular receptor different from that used by ecotropic MuLVs and exhibit a distinct host range (22, 38). Ecotropic MuLVs infect only murine cell lines, while polytypic MuLVs infect murine cell lines as well as lines from several other species. Polytypic MuLVs are generated by recombination between the inoculated ecotropic MuLV and endogenous retroviral sequences resulting in a replacement of the 5' region of the ecotropic *env* gene (4, 6, 7, 10, 12–14, 41, 42), which determines the specificity for different MuLV cellular receptors (2, 33, 38).

We have previously reported that nearly all polytypic MuLV isolates can be classified as members of two major antigenic subclasses defined by their reactivity with two monoclonal antibodies (MAbs) termed Hy 7 and 516 (26). These two MAbs react with mutually exclusive epitopes, both of which have been mapped to the same amino acid in the first third of the SU protein and very likely reflect the existence of two discrete families of endogenous polytypic sequences. In this study we have found that the populations of polytypic MuLVs generated after infection of NFS/N or IRW (inbred Rocky Mountain White) mice with Moloney MuLV (Mo-MuLV) (44) are strikingly different than those generated after inoculation with a substrain of F-MuLV (F-MuLV₅₇) (31) and that this

difference is determined by a small region encompassing the sequences encoding the viral nucleocapsid (NC) protein and a portion of the viral protease (PR).

Different antigenic subclasses of polytypic MuLVs are generated after neonatal inoculation with Mo-MuLV or F-MuLV₅₇. Our initial analyses examined polytypic MuLVs generated in mice after inoculation with Mo-MuLV or F-MuLV₅₇. Quantification of the two antigenic subclasses of polytypic MuLVs in mixed populations was accomplished by a focal immunofluorescence assay (46) with MAbs Hy 7 (11) and 516 (8), specifically reactive with each of the two polytypic MuLV subclasses. Viruses were quantified as infectious centers (ICs) of splenocytes and thymocytes from infected NFS/N mice as well as from mice of a second inbred strain, IRW (9). Endogenous ecotropic or polytypic MuLVs are not expressed in uninfected mice of either strain.

Following infection of NFS/N or IRW mice by Mo-MuLV, most animals exhibited substantial proportions of both antigenic polytypic subclasses, although a large range in the relative levels of Hy 7- and 516-reactive MuLVs was observed among individual mice (Fig. 1A and B). In contrast, mice infected with F-MuLV₅₇ consistently exhibited predominantly Hy 7-reactive polytypic viruses, with several mice expressing this antigenic subclass exclusively (Fig. 1C and D).

The 5' region of the ecotropic MuLV genome influences the types of polytypic MuLVs generated in infected mice. Mo-MuLV and F-MuLV₅₇ infect the spleen with approximately equal efficiencies starting about 3 to 4 days after inoculation. In contrast, infection of the thymus by F-MuLV₅₇ or by chimeric viruses containing the F-MuLV₅₇ transcriptional regulatory sequences resident in the long terminal repeat occurs only after a substantial delay and is variable among different mice (17). Several of the chimeras examined in this study failed to consistently express a sufficiently high level of viruses in the thymus to accurately assess. Accordingly, we have evaluated polytypic MuLVs generated after inoculation of the chimeric

* Corresponding author. Mailing address: Laboratory of Persistent Viral Diseases, Rocky Mountain Laboratories, National Institute of Allergy and Infectious Diseases, Hamilton, MT 59840.

† Present address: Institut Gustave Roussy, URA 147 CNRS 39, 94800 Villejuif, France.




FIG. 1. Expression of polytropic MuLV antigenic subclasses in NFS/N and IRW mice after inoculation with F-MuLV₅₇ or Mo-MuLV. Mice were inoculated neonatally and sacrificed at 4 to 8 weeks of age. IC assays were conducted on thymocytes and splenocytes as previously described (26). For determination of the percentage of polytropic MuLVs reactive with MAb Hy 7, the total polytropic MuLV population was taken to be the sum of Hy 7-reactive and 516-reactive polytropic MuLVs. Each data point represents the result obtained for a single animal.

MuLVs by assays of polytropic viruses from splenocytes 4 to 8 weeks after neonatal inoculation.

Analyses of polytropic MuLVs generated after infection by chimeric viruses in which the U3 region of the long terminal repeat, the *env* gene, or the *pol* gene of F-MuLV₅₇ had been replaced by the analogous gene sequences from Mo-MuLV revealed no significant differences from those generated in mice inoculated with F-MuLV₅₇ (Fig. 2B to 2D). These results indicated that these regions, which encode a number of functions, such as transcriptional regulation, tissue tropism, viral genome replication, integration, and receptor binding, do not determine the difference in the polytropic MuLV subclasses generated by F-MuLV₅₇ and Mo-MuLV. In contrast, mice infected with the chimeric virus FMU5G, in which the R, U5, and *gag* regions of F-MuLV₅₇ were replaced by analogous Mo-MuLV sequences, exhibited a profile of polytropic MuLVs clearly different from that of F-MuLV₅₇-infected mice (Fig. 2E). Mice infected with this chimera generated substantial proportions of both Hy 7- and 516-reactive polytropic MuLVs, similar to that observed after Mo-MuLV infection. As was the case with mice infected with Mo-MuLV, large variations in the percentages of the two antigenic subclasses of polytropic MuLVs were observed among individual mice.

Determinants which influence the relative levels of polytropic MuLV subclasses in infected mice map to the nucleocapsid/protease-encoding sequences of the inoculated ecotropic MuLV. FMU5G contains, in addition to the structural *gag* gene, sequences which are involved in genomic RNA packaging, dimer formation, mRNA splicing, and transcriptional initiation (20, 27, 29, 35–37, 39). To define more closely the sequences of FMU5G which influence the generation of the two subclasses of polytropic MuLVs, chimeric viruses containing smaller substitutions within the U5G region were examined (Fig. 3). Mice infected with chimeric MuLVs corresponding to F-MuLV₅₇ containing M-MuLV packaging sequences (FM Ψ) or Mo-MuLV packaging sequences along with the Mo-MuLV sequences encoding the matrix and capsid proteins (FM Ψ MACA) exhibited predominantly Hy 7-reactive polytropic MuLVs similar to F-MuLV₅₇ (Fig. 3B and C). In contrast, polytropic MuLVs generated in mice infected with the chimera containing Mo-MuLV sequences encoding the nucleocapsid protein and the viral protease (FMNCPR) (Fig. 3D) differed markedly from those observed in mice infected with F-MuLV₅₇, indicating the presence of determinants in this region which strongly influence the generation of the polytropic MuLV antigenic subclasses.

The NCPR region of F-MuLV₅₇ may facilitate the generation of Hy 7-reactive polytropic MuLVs in infected mice. The difference in the relative levels of polytropic MuLV subclasses observed in mice inoculated with MuLVs containing Mo-MuLV NCPR sequences from those in mice infected with MuLVs containing F-MuLV₅₇ NCPR region could be the result of a difference in the levels of expression of one or of both subclasses of polytropic MuLVs. A comparison of the IC titers in splenocytes from mice infected with viruses containing the F-MuLV₅₇ NCPR region with the titers in splenocytes from mice infected with viruses in which this region was derived from Mo-MuLV indicated that mice infected with the former viruses exhibit a significantly higher level of Hy 7-reactive polytropic MuLVs ($P < 0.001$) (Fig. 4). These results suggest that the NCPR sequences of F-MuLV₅₇ facilitate the generation of an Hy 7-reactive recombinant(s). The emergence of a new population of Hy 7-reactive MuLVs could result from amplification of a single species of polytropic MuLV or, alternatively, from coordinate amplification of a family of Hy 7-reactive MuLVs. The depressed levels of MAb 516-reactive




FIG. 2. Expression of polytropic MuLV antigenic subclasses in NFS/N mice inoculated with chimeric viruses corresponding to F-MuLV₅₇ in which the transcriptional regulatory sequences (FMI-2), the *env* gene (FME), the *pol* gene (FMP), or the *gag* gene (FMU5G) was replaced by Mo-MuLV sequences. NFS/N mice were inoculated neonatally, and polytropic-MuLV expression in the spleens was evaluated as described for Fig. 1. (A) Schematic depicting the ecotropic viral RNA genomes of F-MuLV₅₇ and Mo-MuLV (M-MuLV). The locations of the viral gene boundaries in the ecotropic viral genomes are indicated above the F-MuLV₅₇ diagram. The restriction endonuclease cleavage sites are indicated at the bottom of the F-MuLV₅₇ diagram and closely approximate the sites on the analogous proviral DNA used to construct the F-MuLV₅₇-Mo-MuLV chimeras. (B to E) Schematics of chimeric MuLVs and the percentages of Hy 7-reactive polytropic MuLVs detected in splenocytes. The chimeric MuLV FMI-2 was originally described by Chatis et al. (5). All other chimeras have been previously reported (40). Each data point in the graphs represents a determination for an individual mouse.

polytropic MuLVs in mice infected with viruses containing the F-MuLV₅₇ NCPR region compared to the levels in mice infected with ecotropic MuLVs possessing the Mo-MuLV NCPR sequences (Fig. 4) may be the result of interference by an emerging Hy 7-reactive population. In this regard we have observed that coinoculation of F-MuLV₅₇ or Mo-MuLV with

polytropic MuLV isolates results in nearly complete suppression of de novo polytropic MuLV expression in mice, presumably through viral interference mediated by the exogenously administered polytropic MuLV (24).




FIG. 3. Expression of polytropic MuLV antigenic subclasses in NFS/N mice inoculated with chimeric viruses corresponding to F-MuLV₅₇ in which sequences encoding the RNA packaging region (FMΨ), the packaging region plus the MA and CA protein (FMΨMACA), or the NC protein and PR (FMNCPR) have been replaced by Mo-MuLV elements. NFS/N mice were inoculated neonatally, and polytropic-MuLV expression in the spleens was evaluated as described for Fig. 1. (A) Schematic indicating the region of the viral genome encompassing the substitutions in the chimeras. The locations of the genetic elements and the restriction sites used in the construction of the chimeras are indicated in the enlarged view. (B to D) Schematics of the chimeric MuLVs and the percentages of Hy 7-reactive polytropic MuLVs detected in splenocytes. The region of the viral genome in corresponding to F-MuLV₅₇ or Mo-MuLV is indicated by white bars or black bars, respectively. The chimeras have been previously reported (40). Each data point in the graphs represents a determination for an individual mouse.




FIG. 4. Polytypic-MuLV titers expressed in NFS/N mice inoculated with ecotropic MuLVs. Inoculated MuLVs contained the F-MuLV₅₇ NCPR sequences (F-MuLV₅₇, FME, FMP, FMU5G, FMΨ, and FMΨMACA) (55 mice) or the Mo-MuLV NCPR sequences (M-MuLV, FMU5G, and FMNCPR) (51 mice). Average titers of the polytypic subclasses are indicated. The error bars indicate the standard errors of the means.

Sequence difference that may influence the difference in polytypic MuLVs generated by F-MuLV₅₇ and Mo-MuLV. From the analyses of FMU5G (Fig. 2E) and FMNCPR (Fig. 3D) it appears that critical sequences which influence the types of polytypic MuLVs generated after inoculation of the viruses are between the *Bgl*II restriction endonuclease site near the 3' end of the CA gene sequences and the *Sac*I site near the 3' end of the PR gene sequences. A comparison of the nucleotide sequences of F-MuLV₅₇ (accession no. J02192) and Mo-MuLV (accession no. J02255) in this region reveals differences that result in nine different amino acids. These include two differences in the carboxyl-terminal region of the CA protein, five differences in the NC protein, and two differences in the portion of the PR protein encoded by these sequences.

The carboxyl-terminal region of the CA protein of Mo-MuLV differs from that of the CA protein of F-MuLV₅₇ at residue 252 (threonine in Mo-MuLV versus alanine in F-MuLV₅₇) and residue 257 (lysine in Mo-MuLV versus arginine in F-MuLV₅₇). Although the threonine/alanine difference alters the hydrophobicity of this region, these differences in the CA protein do not suggest a mechanism which might influence the types of polytypic MuLVs generated by recombination.

Most of the viral PR is encoded by the sequences between the *Bgl*II and *Sac*I sites. The PR catalyzes cleavage of the polyprotein precursor to the proteins encoded by the *gag* gene as well as cleavage of a 16-amino-acid carboxyl-terminal peptide from the TM protein (32, 45). Since all polytypic MuLVs possess an altered *env* gene compared to that of their ecotropic MuLV parents, changes in PR function could conceivably affect processing of the TM protein and perhaps influence the propagation of recombinant viruses. The two differences in the PR between Mo-MuLV and F-MuLV₅₇ are at residue 71 (lysine in Mo-MuLV versus arginine in F-MuLV₅₇) and residue 96 (asparagine in Mo-MuLV versus histidine in F-MuLV₅₇). Neither of these residues is in the catalytic site or the substrate binding pocket of the PR (28, 32); however, this does not exclude the possibility that these differences affect PR function.

The NC protein is encoded entirely by Mo-MuLV sequences in the two chimeras (FMU5G and FMNCPR) that differed from F-MuLV₅₇ with regard to the types of polytypic MuLVs generated in infected mice. If the effect of the NCPR region is at the level of generation of polytypic MuLV recombinants, the NC protein seems an attractive candidate. This protein is involved in the dimerization of viral genomic RNA, annealing of the tRNA^{Pro} primer, and in vivo proviral synthesis and has recently been reported to influence the specificity as well as the efficiency of viral genomic RNA packaging (20, 29, 35–37, 39, 47, 48). The Mo-MuLV NC differs from that of F-MuLV₅₇ at residue 4 (valine in Mo-MuLV versus isoleucine in F-MuLV₅₇), residue 8 (lysine in Mo-MuLV versus arginine in F-MuLV₅₇), residue 19 (serine in Mo-MuLV versus proline in F-MuLV₅₇), residue 23 (arginine in Mo-MuLV versus histidine in F-MuLV₅₇), and residue 37 (lysine in Mo-MuLV versus arginine in F-MuLV₅₇). A region critical for the functions of the NC protein is a centrally located zinc finger and flanking regions containing several basic amino acids (20, 23, 29, 39, 47). Residue 37 lies within the zinc finger, and residue 23 is in the amino terminus-flanking region (19, 43). The lysine/arginine difference at residue 37 is conservative with regard to steric and charge considerations. However, the histidine/arginine difference at residue 23 in the flanking region may be notable. At physiological pH virtually all arginine residues bear a positive charge, in contrast to less than 5% of histidine residues. Although the effects of mutations at this precise position have not been reported, replacement of other basic amino acids with neutral residues in the flanking regions have marked effects on infectivity and packaging of viral genomic RNAs (23, 39). Recombination may require the formation and reverse transcription of heterodimeric RNAs. It is conceivable that the differences in the generation of polytypic MuLV by F-MuLV₅₇ and Mo-MuLV are a result of selective formation and/or packaging of heterodimers mediated by the NC protein.

The NCPR region of a second F-MuLV, FB29, is nearly identical to that of F-MuLV₅₇ (34). Infection of NFS/N mice with FB29 frequently results in a higher proportion of Hy 7-reactive than 516-reactive recombinants in the spleen; however, more MAb 516-reactive polytypic MuLVs are detected in FB29-infected mice than in mice infected with F-MuLV₅₇ (26). It is possible that regions of the genome other than the NCPR region in other ecotropic MuLVs may influence the types of polytypic MuLVs generated. Although our discussion has focused on differences in the proteins encoded by the retroviral sequences which affect the generation of polytypic MuLVs, it is also possible that the effect is a result of differences in the nucleic acid sequences rather than the proteins encoded by these sequences. Further studies which include site-directed mutagenesis of the NCPR region, determination of the heterogeneity of Hy 7-reactive polytypic SU sequences in F-MuLV-induced recombinants, and identification and analyses of the polytypic progenitor sequences present in the mouse genome may help elucidate the mechanism(s) by which the NCPR sequences influence the generation of polytypic MuLVs.

REFERENCES

- Adachi, A., K. Sakai, N. Kitamura, S. Nakanishi, O. Niwa, M. Matsuyama, and A. Ishimoto. 1984. Characterization of the *env* gene and long terminal repeat of molecularly cloned Friend mink cell focus-inducing virus DNA. *J. Virol.* **50**:813–821.
- Battini, J.-L., J.-M. Heard, and O. Danos. 1992. Receptor choice determinants in the envelope glycoproteins of amphotropic, xenotropic, and polytropic murine leukemia viruses. *J. Virol.* **66**:1468–1475.
- Blatt, C., K. Mileham, M. Haas, M. N. Nesbitt, M. E. Harper, and M. I. Simon. 1983. Chromosomal mapping of the mink cell focus-inducing and

- xenotropic *env* gene family in the mouse. Proc. Natl. Acad. Sci. USA **80**:6298–6302.
4. **Bosselman, R. A., L. J. L. D. Van Griensven, M. Vogt, and I. M. Verma.** 1979. Genome organization of retroviruses. VI. Heteroduplex analysis of ecotropic and xenotropic sequences of Moloney mink cell focus-inducing viral RNA obtained from either a cloned isolate or a thymoma cell line. J. Virol. **32**:968–978.
 5. **Chatis, P. A., C. A. Holland, J. W. Hartley, W. P. Rowe, and N. Hopkins.** 1983. Role for the 3' end of the genome in determining disease specificity of Friend and Moloney murine leukemia viruses. Proc. Natl. Acad. Sci. USA **80**:4408–4401.
 6. **Chattopadhyay, S. K., M. W. Cloyd, D. L. Linemeyer, M. R. Lander, E. Rands, and D. R. Lowy.** 1982. Cellular origin and role of mink cell focus-forming viruses in murine thymic lymphoma. Nature (London) **295**:25–31.
 7. **Chattopadhyay, S. K., M. R. Lander, S. Gupta, E. Rands, and D. R. Lowy.** 1981. Origin of mink cytopathic focus-forming (MCF) viruses: comparison with ecotropic and xenotropic murine leukemia virus genomes. Virology **113**:465–483.
 8. **Chesebro, B., W. Britt, L. Evans, K. Wehrly, J. Nishio, and M. Cloyd.** 1983. Characterization of monoclonal antibodies reactive with murine leukemia viruses: use in analysis of strains of Friend MCF and Friend ecotropic murine leukemia virus. Virology **127**:134–148.
 9. **Chesebro, B., J. L. Portis, K. Wehrly, and J. Nishio.** 1983. Effect of murine host genotype on MCF virus expression, latency, and leukemia cell type of leukemias induced by Friend murine leukemia helper virus. Virology **128**:221–233.
 10. **Chien, Y.-H., I. M. Verma, T. Y. Shih, E. M. Scolnick, and N. Davidson.** 1978. Heteroduplex analysis of the sequence relations between the RNAs of mink cell focus-inducing and murine leukemia viruses. J. Virol. **28**:352–360.
 11. **Cloyd, M. W., B. Chesebro, J. L. Portis, and M. Weir.** 1982. MCF-specific murine monoclonal antibodies made against AKR-247 MCF virus recognize a unique determinant associated with the gp70-p15(E) complex. J. Virol. **41**:1112–1117.
 12. **Donoghue, D. J., E. Rothenberg, N. Hopkins, D. Baltimore, and P. A. Sharp.** 1978. Heteroduplex analysis of nonhomology region between Moloney MuLV and the dual host range derivative HIX virus. Cell **14**:959–970.
 13. **Elder, J. H., J. W. Gautsch, F. C. Jensen, R. A. Lerner, J. W. Hartley, and W. P. Rowe.** 1977. Biochemical evidence that MCF murine leukemia viruses are envelope (*env*) gene recombinants. Proc. Natl. Acad. Sci. USA **74**:4676–4680.
 14. **Evans, L., M. Nunn, P. H. Duesberg, D. Troxler, and E. Scolnick.** 1980. RNAs of defective and nondefective components of Friend anemia and polycythemia virus strains identified and compared. Cold Spring Harbor Symp. Quant. Biol. **44**:823–835.
 15. **Evans, L. H., and M. W. Cloyd.** 1984. Generation of mink cell focus-forming viruses by Friend murine leukemia virus: recombination with specific endogenous proviral sequences. J. Virol. **49**:772–781.
 16. **Evans, L. H., and M. W. Cloyd.** 1985. Friend and Moloney murine leukemia viruses specifically recombine with different endogenous retroviral sequences to generate mink cell focus-forming viruses. Proc. Natl. Acad. Sci. USA **82**:459–463.
 17. **Evans, L. H., and J. D. Morrey.** 1987. Tissue-specific replication of Friend and Moloney murine leukemia viruses in infected mice. J. Virol. **61**:1350–1357.
 18. **Fischinger, P. J., S. Nomura, and D. P. Bolognesi.** 1975. A novel murine oncornavirus with dual eco- and xenotropic properties. Proc. Natl. Acad. Sci. USA **72**:5150–5155.
 19. **Friedrich, R. W., W. Koch, U. von Maydell-Livonius, H. Schrewe, and W. Zimmermann.** 1990. Direct submission to the EMBL Data Library.
 20. **Gorelick, R., L. E. Anderson, J. P. Hanser, and A. Rein.** 1988. Point mutants of Moloney murine leukemia virus that fail to package viral RNA: evidence for specific RNA recognition by a zinc finger-like protein sequence. Proc. Natl. Acad. Sci. USA **85**:8420–8424.
 21. **Green, N., H. Hiai, J. H. Elder, R. A. Schwartz, R. H. Khirya, C. Y. Thomas, P. N. Tschlis, and J. M. Coffin.** 1980. Expression of leukemogenic recombinant viruses associated with a recessive gene in HRS/J mice. J. Exp. Med. **152**:249–264.
 22. **Hartley, J. W., N. K. Wolford, L. J. Old, and W. P. Rowe.** 1977. A new class of murine leukemia virus associated with development of spontaneous lymphomas. Proc. Natl. Acad. Sci. USA **74**:789–792.
 23. **Housset, V., H. De Rocquigny, B. P. Roques, and J.-L. Darlix.** 1993. Basic amino acids flanking the zinc finger of Moloney murine leukemia virus nucleocapsid protein NCp10 are critical for virus infectivity. J. Virol. **67**:2537–2545.
 24. **Lavignon, M., and L. Evans.** Unpublished observations.
 25. **Lavignon, M., and L. Evans.** 1996. A multistep process of leukemogenesis in Moloney murine leukemia virus-infected mice that is modulated by retroviral pseudotyping and interference. J. Virol. **70**:3852–3862.
 26. **Lavignon, M., J. L. Walker, S. M. Perryman, F. G. Malik, A. S. Khan, T. S. Theodore, and L. H. Evans.** 1994. Characterization of epitopes defining two major subclasses of polytropic murine leukemia viruses (MuLVs) which are differentially expressed in mice infected with different ecotropic MuLVs. J. Virol. **68**:5194–5203.
 27. **Mann, R., F. C. Mulligan, and D. Baltimore.** 1983. Construction of a retrovirus packaging mutant and its use to produce helper-free defective retrovirus. Cell **33**:153–159.
 28. **Menendez-Arias, L., I. Weber, and S. Oroszlan.** 1995. Mutational analyses of the substrate binding pocket of murine leukemia virus protease and comparison with human immunodeficiency virus proteases. J. Biol. Chem. **270**:29162–29168.
 29. **Méric, C., and S. P. Goff.** 1989. Characterization of Moloney murine leukemia virus mutants with single-amino-acid substitutions in the Cys-His box of the nucleocapsid protein. J. Virol. **63**:1558–1568.
 30. **O'Donnell, P. V., E. Stockert, Y. Obata, and L. J. Old.** 1981. Leukemogenic properties of AKR dualtropic (MCF) viruses: amplification of murine leukemia virus-related antigens on thymocytes and acceleration of leukemia development in AKR mice. Virology **112**:548–563.
 31. **Oliff, A. I., G. L. Hager, E. H. Chang, E. M. Scolnick, H. W. Chan, and D. R. Lowy.** 1980. Transfection of molecularly cloned Friend murine leukemia virus DNA yields a highly leukemogenic helper-independent type C virus. J. Virol. **33**:475–486.
 32. **Oroszlan, S., and R. B. Luftig.** 1990. Retroviral proteinases. Curr. Top. Microbiol. Immunol. **157**:153–185.
 33. **Ott, D., and A. Rein.** 1992. Basis for receptor specificity of noncancerous murine leukemia virus surface glycoprotein gp70^{su}. J. Virol. **66**:4632–4638.
 34. **Perryman, S., J. Nishio, and B. Chesebro.** 1991. Complete nucleotide sequence of Friend murine leukemia virus, strain FB29. Nucleic Acids Res. **19**:6950.
 35. **Prats, A. C., V. Housset, G. de Billy, F. Cornille, B. Roques, and J. L. Darlix.** 1991. Viral annealing activities of the nucleocapsid protein of Moloney murine leukemia virus are zinc independent. Nucleic Acids Res. **19**:3533–3541.
 36. **Prats, A.-C., C. Roy, P. Wang, M. Erard, V. Housset, C. Gabus, C. Paoletti, and J.-L. Darlix.** 1990. *cis* elements and *trans*-acting factors involved in dimer formation of murine leukemia virus RNA. J. Virol. **64**:774–783.
 37. **Prats, A. C., L. Sarih, C. Gabus, S. Litvak, G. Keith, and J. L. Darlix.** 1988. Small finger protein of avian and murine retroviruses has nucleic acid annealing activity and positions the replication primer tRNA onto genomic RNA. EMBO J. **7**:1777–1783.
 38. **Rein, A.** 1982. Interference grouping of murine leukemia viruses: a distinct receptor for the MCF-recombinant viruses on mouse cells. Virology **120**:251–257.
 39. **Rein, A., D. P. Harvin, J. Mirro, S. M. Ernst, and R. J. Gorelick.** 1994. Evidence that a central domain of nucleocapsid protein is required for RNA packaging in murine leukemia virus. J. Virol. **68**:6124–6129.
 40. **Richardson, J., A. Corbin, F. Pozo, S. Orsoni, and M. Sitbon.** 1993. Sequences responsible for the distinctive hemolytic potentials of Friend and Moloney murine leukemia viruses are dispersed but confined to the Ψ -*gag-PR* region. J. Virol. **67**:5478–5486.
 41. **Rommelaere, J., D. V. Faller, and N. Hopkins.** 1978. Characterization and mapping of RNase T1-resistant oligonucleotides derived from the genomes of Akv and MCF murine leukemia viruses. Proc. Natl. Acad. Sci. USA **75**:495–499.
 42. **Shih, T. Y., M. O. Weeks, D. H. Troxler, J. M. Coffin, and E. M. Scolnick.** 1978. Mapping host range-specific oligonucleotides within genomes of the ecotropic and mink cell focus-inducing strains of Moloney murine leukemia virus. J. Virol. **26**:71–83.
 43. **Shinnick, T. M., R. A. Lerner, and J. G. Sutcliffe.** 1981. Nucleotide sequence of Moloney murine leukaemia virus. Nature (London) **293**:543–548.
 44. **Shoemaker, C., S. Goff, E. Gilboa, M. Paskind, S. W. Mitra, and D. Baltimore.** 1980. Structure of a cloned circular Moloney murine leukemia virus DNA molecule containing an inverted segment: implications for retrovirus integration. Proc. Natl. Acad. Sci. USA **77**:3932–3936.
 45. **Shultz, A., and A. Rein.** 1985. Maturation of murine leukemia virus *env* proteins in the absence of other viral proteins. Virology **145**:335–339.
 46. **Sitbon, M., J. Nishio, K. Wehrly, D. Lodmell, and B. Chesebro.** 1985. Use of a focal immunofluorescence assay on live cells for quantitation of retroviruses: distinction of host range classes in virus mixtures and biological cloning of dual-tropic murine leukemia viruses. Virology **141**:110–118.
 47. **Yu, Q., and J. L. Darlix.** 1996. The zinc finger of nucleocapsid protein of Friend murine leukemia virus is critical for proviral DNA synthesis in vivo. J. Virol. **70**:5791–5798.
 48. **Zhang, Y., and E. Barklis.** 1995. Nucleocapsid protein effects on the specificity of retrovirus RNA encapsidation. J. Virol. **69**:5716–5722.