



Published in final edited form as:

Neurochem Res. 2005 January ; 30(1): 15–23.

Cytidine 5'-Diphosphocholine (CDP-Choline) in Stroke and Other CNS Disorders

Rao Muralikrishna Adibhatla^{1,2,3,4} and J. F. Hatcher¹

¹ Department of Neurological Surgery, University of Wisconsin, Madison, WI

² Cardiovascular Research Center University of Wisconsin, Madison, WI

³ Veterans Administration Hospital, Madison, WI

Abstract

Brain phosphatidylcholine (PC) levels are regulated by a balance between synthesis and hydrolysis. Pro-inflammatory cytokines such as tumor necrosis factor- α (TNF- α) and interleukin-1 (IL-1 α/β) activate phospholipase A₂ (PLA₂) and PC-phospholipase C (PC-PLC) to hydrolyze PC. PC hydrolysis by PLA₂ releases free fatty acids including arachidonic acid, and lyso-PC, an inhibitor of CTP-phosphocholine cytidyltransferase (CCT). Arachidonic acid metabolism by cyclooxygenases/lipoxygenases is a significant source of reactive oxygen species. CDP-choline might increase the PC levels by attenuating PLA₂ stimulation and loss of CCT activity. TNF- α also stimulates proteolysis of CCT. TNF- α and IL-1 β are induced in brain ischemia and may disrupt PC homeostasis by increasing its hydrolysis (increase PLA₂ and PC-PLC activities) and inhibiting its synthesis (decrease CCT activity). The beneficial effects of CDP-choline may result by counteracting TNF- α and/or IL-1 mediated events, integrating cytokine biology and lipid metabolism. Re-evaluation of CDP-choline phase III stroke clinical trial data is encouraging and future trials are warranted. CDP-choline is non-xenobiotic, safe, well tolerated, and can be considered as one of the agents in multi-drug treatment of stroke.

Keywords

Cerebral ischemia; citicoline; clinical trials; interleukin-1 β ; phospholipases; phospholipids; reactive oxygen species; tumor necrosis factor; lipidomics

INTRODUCTION

Stroke or “brain attack” is the first leading cause of long-lasting disability, third leading cause of death and continues to be a problem of vast clinical significance. Approximately 3.9 million Americans are stroke survivors, and the after-effects of stroke require more than \$51 billion in healthcare costs annually. Presently, tissue plasminogen activator (tPA) is the only FDA approved drug for the treatment of acute ischemic stroke but needs to be administered within 3 h (1). However, there are some concerns that tPA has neurotoxic side effects in addition to its beneficial (thrombolytic) actions (2). Many neuroprotective agents have undergone phase III clinical trials for stroke; most of the trials were abandoned due to ineffectiveness or toxicity of the drug.

⁴Address reprint requests to: R. M. Adibhatla Department of Neurological Surgery, H4-330, Clinical Science Center, 600 Highland Avenue, University of Wisconsin-Madison, Madison, WI 53792-3232. Tel: +608-263-1791; Fax: +608-263-1409; E-mail: adibhatl@neurosurg.wisc.edu

Cytidine-5'-diphosphocholine (CDP-choline or Citicoline) is composed of cytidine and choline linked by a diphosphate bridge and is an essential intermediate in the synthesis of phosphatidylcholine (PC), a major brain phospholipid, via Kennedy pathway. Exogenous CDP-choline is hydrolyzed and absorbed as cytidine and choline (3), and CDP-choline is re-synthesized from cytidine triphosphate (CTP) and phosphocholine by CTP-phosphocholine cytidyltransferase (CCT), the rate-limiting enzyme in PC synthesis (see [4] and references therein). CDP-choline also serves as a choline donor in the biosynthesis of the neurotransmitter acetylcholine (5). As the intermediate in PC biosynthesis, it was believed that CDP-choline would rectify membrane damage and provide benefit in CNS disorders and injury (including stroke).

CDP-choline has been studied in >11,000 volunteers and patients and showed beneficial effects in cerebral ischemia, traumatic brain injury, hypoxia, Alzheimer's and Parkinson's diseases, learning and memory disorders, alcoholism, drug addiction, amblyopia and glaucoma (Table I.) Citicoline, an international non-proprietary name of CDP-choline, is marketed as a prescription drug (in Japan, Spain, France and Italy) or as an over-the-counter dietary supplement in USA. CDP-choline was originally developed for stroke treatment by Ferrer Internacional, S. A. (Barcelona, Spain) and is being tested for treatment of Alzheimer's and Parkinson's disorders (7). CDP-choline is the only agent that is non-xenobiotic and has virtually no side effects. In 1983, 22 articles were published that described the physi-co-chemical properties, pharmacokinetics, toxicity and bioavailability of this agent (54). CDP-choline (600 or 1000 mg/day) or placebo to healthy volunteers did not show any abnormal side effects in terms of hematological or clinical analysis (3). No clinically significant ECG and EEG abnormalities were noticed. Neurological tests, tendon reflexes, blood pressure and heart rate were not affected by any dose of the drug or placebo. The tolerance of CDP-choline is excellent and side effects were rare, never severe and consisted mainly of digestive intolerance, gastrointestinal discomfort and restlessness. In no case was it necessary to interrupt the treatment for side effects attributed to CDP-choline use (55). Recent re-evaluation of USA phase III stroke clinical trial data is encouraging (49) and this agent still holds promise for treatment of acute ischemic stroke. Indevus Inc. licensed exclusive North American rights from Ferrer Internacional, S. A. for the manufacture, use and sale of CDP-choline for the treatment of stroke.

CDP-CHOLINE IN CLINICAL TRIALS

Stroke

There have been 13 stroke clinical trials of CDP-choline since 1980 (nine in Europe and Japan and four in the USA) (5). The European clinical trials showed that CDP-choline improved global and neurological function and promoted earlier motor and cognitive recovery. A large multi-center study in Japan found that CDP-choline showed improvement in a global outcome rating scale. Four major clinical trials in the USA have provided ambiguous results, and thus the beneficial effects of CDP-choline have not been established (49 and references therein). In the first study, CDP-choline improved functional outcome and reduced neurologic deficit. However, two subsequent studies failed to demonstrate improvement in the outcome. On *post-hoc* analysis, CDP-choline was shown to provide beneficial effects in a subgroup of moderate-to-severe stroke cases. Subsequent pooling of individual patient data from four US trials showed that CDP-choline treatment for 6 weeks improved overall recovery at 12 weeks in acute ischemic stroke patients (49 and references therein). Pooled diffusion-weighted magnetic resonance imaging (DW-MRI) data from two clinical trials showed a significant dose-dependent reduction on percent change in lesion volume (56). We have summarized recent experimental data on the effects of CDP-choline in cerebral ischemia and evaluated several factors which might have hindered efficacy of CDP-choline in stroke clinical trials in the USA. One of the factors is the brain uptake of CDP-choline. The European and Japanese trials used

i.v. administration in contrast to oral route used in the USA trials. In animal studies, brain uptake of CDP-choline or its metabolites was 0.5% of the oral dose, whereas i.v. administration elevated brain uptake to 2%. Liposome encapsulation of the drug can further increase brain uptake up to 23% (8,21) and also circumvent CCT, the key rate-limiting enzyme in PC synthesis. Liposome encapsulation suggests a possible strategy to increase the CDP-choline levels in the CNS and enhance its clinical effectiveness (8). In the light of recent clinical evaluations, experimental data in cerebral ischemia, and realization that oral administration was not appropriate, new phase III stroke clinical trials are warranted. Benefit from CDP-choline in humans is far from proven, however future trials are essential before making any conclusions that CDP-choline is ineffective for stroke treatment.

Alzheimer's Disease and Other Memory Disorders

Clinical studies have demonstrated that CDP-choline improves cognitive performance in elderly subjects (36). Blocking synthesis of PC is sufficient in itself to cause cell death (57, 58), and a 10% loss of cellular membrane is a threatening situation for neuronal viability (59). Alzheimer's brains have shown loss of PC and phosphatidylethanolamine (60) and CDP-choline may rectify the membrane damage in these brains (36). Additionally, the cholinergic system is dysfunctional in Alzheimer's brain (50), and CDP-choline may provide benefit by enhancing acetylcholine synthesis (Fig. 1A).

Parkinson's Disease

CDP-choline stimulates tyrosine hydroxylase activity and dopamine release (3), which may be due to increases in brain acetylcholine since choline administration produced the same effects (61). Parkinson's disease is characterized by a selective degeneration of the dopaminergic neurons of the substantia nigra (62), however the phospholipid abnormalities present in Alzheimer's brains were not observed (60). Levodopa is the main therapeutic option for treatment of Parkinson's disease; its main disadvantage is progressive loss of efficacy (63). CDP-choline has been tested in treatment of Parkinson's disease because of its ability to increase the availability of dopamine (3). Combination treatment of Parkinson's patients with CDP-choline and levodopa allowed significant reduction of the levodopa dose, thus minimizing side effects of levodopa therapy (3).

CEREBRAL ISCHEMIA

The energy needs of the brain are supplied by metabolism of glucose and oxygen for the phosphorylation of ADP to ATP. Rapid loss of ATP occurs following cerebral ischemia, resulting in uncontrolled leakage of ions across the cell membrane, membrane depolarization and release of neurotransmitters glutamate and dopamine (64). Excess glutamate release and stimulation of its receptors results in phospholipases activation, phospholipid hydrolysis and arachidonic acid release (5), ultimately leading to apoptotic or necrotic cell death (65). Apoptotic cell death is mediated by activation of caspases. CDP-choline attenuated expression of pro-caspases, cleaved caspase-3 and nuclear DNA fragmentation after focal cerebral ischemia (27). CDP-choline in combination with nimodipine reduced infarction and increased expression of anti-apoptotic Bcl-2 after focal cerebral ischemia (26). CDP-choline pre-treatment prevented excitotoxic death caused by excessive glutamate exposure in cerebellar granule neurons (45) and in *in vivo* focal cerebral ischemia model (101). These findings have been summarized in Fig. 1A.

Phospholipases

In addition to being an essential membrane structural component, PC is the source of bioactive lipids such as phosphatidates, 1,2-diacylglycerol, and arachidonic acid, among others (58). Thus, PC hydrolysis serves important roles in signal transduction mediated by various stimuli

including cytokines (59,84). PC can be hydrolyzed (66,67) by PC-phospholipase C (PLC) (68), PC-phospholipase D (PLD) (69), or PLA₂ (70). There is substantial evidence that PLA₂ is activated in transient ischemia and contributes to neuronal damage (12,24).

PLA₂ isozymes occur in multiple forms (71,72) in the mammalian cell and are classified as calcium independent, cytosolic (cPLA₂) and secretory (sPLA₂). sPLA₂ isoforms are low molecular weight (~14 kDa) and require millimolar Ca²⁺ concentrations for activity. Our studies demonstrated significant increases in PLA₂ activity in membrane and mitochondrial fractions following transient cerebral ischemia (13,24). The majority of PLA₂ activity required mM Ca²⁺ for optimal activity, characteristic of sPLA₂. CDP-choline treatment significantly attenuated PLA₂ activity in both membrane and mitochondrial fractions. *In vitro*, CDP-choline and its components cytidine and choline had no effect on PLA₂ activity, and thus CDP-choline is not as such a PLA₂ inhibitor (13). Since CDP-choline does not directly inhibit PLA₂, our data suggests that CDP-choline prevents activation of sPLA₂ (Fig. 1B). These findings are consistent with our data that CDP-choline attenuated loss of phospholipids and increase in free fatty acids including arachidonic acid after both focal and global transient cerebral ischemia (14–16,24).

TNF- α and IL-1 β Disrupt PC Homeostasis

Brain PC homeostasis is regulated by a balance between synthesis and hydrolysis by phospholipases. Two forms of IL-1 are present in brain tissue, IL-1 α and IL-1 β , and act on the IL-1 receptor (73). TNF- α and IL-1 β are induced following brain ischemia (74–77). TNF- α (78–81) and IL-1 (82–88) stimulate hydrolysis of PC through induction of PLA₂ and PC-phospholipase C (PC-PLC), which mediate their cytotoxicity (84,86,87). Activation of PLA₂ by TNF- α /IL-1 may be mediated by increases in PLA₂ activating protein (PLAP) (88–90). Treatment with TNF- α antibody attenuated infarction (24,77), loss of phospholipids and increase in ceramide and free fatty acids including arachidonic acid after focal cerebral ischemia (24).

CCT and Lyso-PC (4)

With the exception of liver, the CDP-choline pathway is the main source of *de novo* PC synthesis in mammalian tissues. Thus, CCT is a key enzyme in regulation of PC synthesis in non-hepatic tissues including the brain. CCT is an amphitropic enzyme in mammalian tissue and is distributed between cytosol and membrane. The catalytic activity of CCT has an absolute requirement for a lipid environment, which is provided by the phospholipids of the cellular membrane. Lyso-PC, a PLA₂ hydrolysis product of PC, inhibits CCT (Fig. 2A) (91).

TNF- α can also stimulate proteolytic degradation of CCT, resulting in a decrease in CCT protein (Fig. 2A) (93). Thus, TNF- α /IL-1 disrupt PC homeostasis by increasing PC hydrolysis (increase PLA₂ and PC-PLC activities) and inhibiting its synthesis (decrease CCT activity) (Fig. 2B).

We have shown stimulation of PLA₂, loss of phospholipids including PC (13–15,24), decrease in CCT activity and increase in lyso-PC (4) following transient brain ischemia. CDP-choline significantly increased CCT activity (4) and attenuated lyso-PC. Thus, many of the effects of CDP-choline are in a direction opposite to those of TNF- α /IL-1; one hypothesis is that the beneficial effects of CDP-choline result by partly counteracting TNF- α /IL-1 mediated events (Fig. 2B). However, further studies are required to validate this hypothesis.

These findings may have important implications for treatment of ischemic brain injury with CDP-choline. Loss of CCT was an early event in cerebral ischemia that persisted throughout 1 day reperfusion. While CDP-choline increased CCT activity in the membrane fraction, this

did not occur until 6-h reperfusion. Thus, even with CDP-choline treatment, PC synthesis may be impaired for several hours after cerebral ischemia.

CDP-choline treatment delayed by 3 h did not offer any neuroprotection (8). This later treatment could further delay the recovery of CCT activity during the critical early reperfusion time. Identification of the factor(s) responsible for CCT loss may lead to new therapeutic interventions in stroke, or combination treatments that could enhance CDP-choline's efficacy. Cholinephosphotransferase (CDP-choline-1,2-diacylglycerol cholinephospho-transferase (CPT), EC 2.7.8.2) catalyzes the final step of the Kennedy (CDP-choline) pathway for the *de novo* synthesis of PC. Though CPT is non-rate limiting under normal conditions, it may act as regulatory switch under some pathological conditions and set in motion the apoptotic cell death program (102,103). The role of this enzyme in CNS injury and the effect of CDP-choline need to be examined. Emerging lipidomics (104,105) may offer some solutions on elucidating CDP-choline mechanisms through integration of cytokine biology and lipid metabolism (106,107) in the near future.

Reactive Oxygen Species (ROS) Lipid Peroxidation and Glutathione

Formation of ROS and the resultant oxidation of biological molecules is a well-recognized mechanism of tissue damage in ischemia/reperfusion (65,94). ROS induce lipid peroxidation, resulting in formation of malondialdehyde (MDA), 4-hydroxy-nonenal (HNE) and acrolein (95). HNE and acrolein induce neuronal apoptosis by covalently cross-linking with proteins (95,96). CDP-choline significantly attenuated ischemia-induced hydroxyl radicals (OH^{\bullet}) and MDA formation (12,13). PLA_2 releases free fatty acids including arachidonic acid from membrane phospholipids. Arachidonic acid metabolism by COX/LOX is considered to be a significant source of ROS (12,17,97). The other important free fatty acid, docosahexaenoic acid (DHA) is a major source of ROS, lipid peroxidation and neuronal injury. Bazan's group (106,107) showed some elegant relationship between pro-inflammatory gene expression and lipid metabolism by tandem liquid chromatography-photodiode array-electrospray ionization-mass spectrometric-mass spectrometry (LC-PDA-ESI-MS-MS) using lipidomic (104,105) approaches. The decrease in OH^{\bullet} production in transient brain ischemia following CDP-choline treatment may be due to attenuation of PLA_2 activation. Glutathione is the primary endogenous antioxidant defense system and removes H_2O_2 and lipid peroxides in the brain. CDP-choline increased total glutathione levels, glutathione reductase activity, decreased GSSG and glutathione oxidation ratio (an indicator of the redox status of glutathione) after transient cerebral ischemia (15).

SINGLE DRUG INTERVENTIONS MAY NOT BE EFFECTIVE

Due to the multiple pathways involved in ischemic injury, no single agent is likely to provide complete neuroprotection (98,99). CDP-choline in combination with NMDA receptor antagonist MK801 (100), thrombolytic agent (recombinant tPA) (29), urokinase (28), or basic fibroblast growth factor (23) showed synergistic benefit in experimental ischemia models. There seems to be a growing consensus to adopt a multi-pronged approach using drug cocktails since the nature of stroke injury is complex and multi-dimensional (98,99,108). CDP-choline is non-xenobiotic, safe and well tolerated, which makes it a viable choice to be used in combinational therapy for the treatment of stroke and Parkinson's disease (with levodopa).

Acknowledgements

This study was supported in part by grants to RMA from NIH/NINDS, (NS42008); University of Wisconsin Medical School Research Grants (161-9904), and laboratory resources provided by William S. Middleton VA Hospital.

References

1. Fisher M, Brott TG. Emerging therapies for acute ischemic stroke: New therapies on trial. *Stroke* 2003;34:359–361. [PubMed: 12574536]
2. Siao CJ, Fernandez SR, Tsirka SE. Cell type-specific roles for tissue plasminogen activator released by neurons or microglia after excitotoxic injury. *J Neurosci* 2003;23:3234–3242. [PubMed: 12716930]
3. Secades JJ. CDP-choline: Update and review of its pharmacology and clinical use. *Methods Find Exp Clin Pharmacol* 2002;24(Supp B):1–53.
4. Adibhatla RM, Hatcher JF, Dempsey RJ. Cytidine-5'-diphosphocholine (CDP-choline) affects CTP:phosphocholine cytidyltransferase and lyso-phosphatidylcholine after transient brain ischemia. *J Neurosci Res* 2004;76:390–396. [PubMed: 15079868]
5. Adibhatla RM, Hatcher JF, Dempsey RJ. Citicoline: Neuroprotective mechanisms in cerebral ischemia. *J Neurochem* 2002;80:12–23. [PubMed: 11796739]
6. Conant R, Schauss AG. Therapeutic applications of citicoline for stroke and cognitive dysfunction in the elderly: A review of the literature. *Altern Med Rev* 2004;9:17–31. [PubMed: 15005642]
7. Alexandrov AV. Citicoline: Ferrer Internacional. *Curr Opin Invest Drugs* 2001;2:1757–1762.
8. Adibhatla RM, Hatcher JF. Citicoline mechanisms and clinical efficacy in cerebral ischemia. *J Neurosci Res* 2002;70:133–139. [PubMed: 12271462]
9. D'Orlando KJ, Sandage BW Jr. Citicoline (CDP-choline): Mechanisms of action and effects in ischemic brain injury. *Neurol Res* 1995;17:281–284. [PubMed: 7477743]
10. Weiss GB. Metabolism and actions of CDP-choline as an endogenous compound and administered exogenously as citicoline. *Life Sci* 1995;56:637–660. [PubMed: 7869846]
11. Grieb P, Rejdak R. Pharmacodynamics of citicoline relevant to the treatment of glaucoma. *J Neurosci Res* 2002;67:143–148. [PubMed: 11782957]
12. Adibhatla RM, Hatcher JF, Dempsey RJ. Phospholipase A₂, hydroxyl radicals and lipid peroxidation in transient cerebral ischemia. *Antioxid Redox Signal* 2003;5:647–654. [PubMed: 14580322]
13. Adibhatla RM, Hatcher JF. Citicoline decreases phospholipase A₂ stimulation and hydroxyl radical generation in transient cerebral ischemia. *J Neurosci Res* 2003;73:308–315. [PubMed: 12868064]
14. Rao AM, Hatcher JF, Dempsey RJ. Lipid alterations in transient forebrain ischemia: Possible new mechanisms of CDP-choline neuroprotection. *J Neurochem* 2000;75:2528–2535. [PubMed: 11080206]
15. Adibhatla RM, Hatcher JF, Dempsey RJ. Effects of citicoline on phospholipid and glutathione levels in transient cerebral ischemia. *Stroke* 2001;32:2376–2381. [PubMed: 11588329]
16. Rao AM, Hatcher JF, Dempsey RJ. Does CDP-choline modulate phospholipase activities after transient forebrain ischemia? *Brain Res* 2001;893:268–272. [PubMed: 11223016]
17. Rao AM, Hatcher JF, Dempsey RJ. Lipid metabolism in ischemic neuronal death. *Recent Res Develop Neurochem* 1999;2:533–549.
18. Rao AM, Hatcher JF, Dempsey RJ. CDP-choline: Neuroprotection in transient forebrain ischemia of gerbils. *J Neurosci Res* 1999;58:697–705. [PubMed: 10561698]
19. Grieb P, Gadamski R, Wojda R, Janisz M. CDP-choline, but not cytidine, protects hippocampal CA₁ neurones in the gerbil following transient forebrain ischaemia. *Folia Neuropathol* 2001;39:175–180. [PubMed: 11770128]
20. Trovarelli G, de Medio GE, Dorman RV, Piccinin GL, Horrocks LA, Porcellati G. Effect of CDP-choline on ischemia-induced alterations of brain lipid in the gerbil. *Neurochem Res* 1981;6:821–833. [PubMed: 6796897]
21. Fresta M, Puglisi G. Reduction of maturation phenomenon in cerebral ischemia with CDP-choline-loaded liposomes. *Pharm Res* 1999;16:1843–1849. [PubMed: 10644072]
22. Kakihana M, Fukuda N, Suno M, Nagaoka A. Effects of CDP-choline on neurologic deficits and cerebral glucose metabolism in a rat model of cerebral ischemia. *Stroke* 1988;19:217–222. [PubMed: 3344538]
23. Schabitz WR, Li F, Katsumi I, Sandage BW, Locke KW, Fischer M. Synergistic effects of a combination of low-dose basic fibroblast growth factor and citicoline after temporary experimental focal ischemia. *Stroke* 1999;30:427–432. [PubMed: 9933283]

24. Adibhatla RM, Hatcher JF, Tureyen K, Tsao F, Dempsey RJ. Does CDP-choline partly counteract TNF- α mediated events after stroke? Soc Neurosci Abstr 2004;#455.10.
25. Goldberg WJ, Dorman RV, Dabrowiecki Z, Horrocks LA. The effects of ischemia and CDPamines on Na⁺, K⁺-ATPase and acetylcholinesterase activities in rat brain. Neurochem Pathol 1985;3:237–248. [PubMed: 3008046]
26. Sobrado M, Lopez MG, Carceller F, Garcia AG, Roda JM. Combined nimodipine and citicoline reduce infarct size, attenuate apoptosis and increase bcl-2 expression after focal cerebral ischemia. Neuroscience 2003;118:107–113. [PubMed: 12676142]
27. Krupinski J, Ferrer I, Barrachina M, Secades JJ, Mercadal J, Lozano R. CDP-choline reduces pro-caspase and cleaved caspase-3 expression, nuclear DNA fragmentation, and specific PARP-cleaved products of caspase activation following middle cerebral artery occlusion in the rat. Neuropharmacology 2002;42:846–854. [PubMed: 12015211]
28. Shuaib A, Yang Y, Li Q. Evaluating the efficacy of citicoline in embolic ischemic stroke in rats: Neuroprotective effects when used alone or in combination with urokinase. Exp Neurol 2000;161:733–739. [PubMed: 10686092]
29. Andersen M, Overgaard K, Meden P, Boysen G. Effects of citicoline combined with thrombolytic therapy in a rat embolic stroke model. Stroke 1999;30:1464–1470. [PubMed: 10390324]
30. Alkan T, Kahveci N, Goren B, Korfali E, Ozluk K. Ischemic brain injury caused by interrupted versus uninterrupted occlusion in hypotensive rats with subarachnoid hemorrhage: Neuroprotective effects of citicoline. Arch Physiol Biochem 2001;109:161–167. [PubMed: 11780777]
31. Savci V, Cavun S, Goktalay G, Ulus IH. Cardiovascular effects of intracerebroventricularly injected CDP-choline in normotensive and hypotensive animals: The involvement of cholinergic system. Arch Pharmacol 2002;365:388–398.
32. Baskaya MK, Dogan A, Rao AM, Dempsey RJ. Neuroprotective effects of citicoline on brain edema and blood–brain barrier breakdown after traumatic brain injury. J Neurosurg 2000;92:448–452. [PubMed: 10701532]
33. Dempsey RJ. Cytidinediphosphocholine treatment to decrease traumatic brain injury-induced hippocampal neuronal death, cortical contusion volume, and neurological dysfunction in rats. J Neurosurg 2003;98:867–873. [PubMed: 12691414]
34. Dixon CE, Ma XC, Marion DW. Effects of CDP-choline treatment on neurobehavioral deficits after TBI and on hippocampal and neocortical acetylcholine release. J Neurotrauma 1997;14:161–169. [PubMed: 9104933]
35. Barrachina M, Dominguez I, Ambrosio S, Secades J, Lozano R, Ferrer I. Neuroprotective effect of citicoline in 6-hydroxydopamine-lesioned rats and in 6-hydroxydopamine-treated SH-SY5Y human neuroblastoma cells. J Neurol Sci 2003;215:105–110. [PubMed: 14568136]
36. Alvarez XA, Sampedro C, Lozano R, Cacabelos R. Citicoline protects hippocampal neurons against apoptosis induced by brain β -amyloid deposits plus cerebral hypoperfusion in rats. Methods Find Exp Clin Pharmacol 1999;21:535–540. [PubMed: 10599052]
37. Alvarez XA, Vecino B, Perea JE, Daniele D, Cacabelos R. Citicoline antagonizes bromazepam-induced amnesia in rats. Human Psychopharmacol 1997;12:547–556.
38. Gimenez R, Aguilar J. Effects of cytidine 5'-diphosphocholine on plasma homocysteine levels in rat. Comp Biochem Physiol 2003;134:271–276.
39. Teather LA, Wurtman RJ. Dietary cytidine (5')-diphosphocholine supplementation protects against development of memory deficits in aging rats. Prog Neuro-psychopharmacol Biol Psychiatry 2003;27:711–717.
40. Lopez-Coviella I, Agut J, Savci V, Ortiz JA, Wurtman RJ. Evidence that 5'-cytidinediphosphocholine can affect brain phospholipid composition by increasing choline and cytidine plasma levels. J Neurochem 1995;65:889–894. [PubMed: 7616250]
41. Clark W, Gunionrinker L, Lessov N, Hazel K. Citicoline treatment for experimental intracerebral hemorrhage in mice. Stroke 1998;29:2136–2139. [PubMed: 9756595]
42. Alberghina M, Viola M, Serra I, Mistretta A, Giuffrida AM. Effect of CDP-choline on the biosynthesis of phospholipids in brain regions during hypoxic treatment. J Neurosci Res 1981;6:421–433. [PubMed: 7299849]

43. Bruhwylter J, Liegeois JF, Geczy J. Facilitatory effects of chronically administered citicoline on learning and memory processes in the dog. *Prog Neuro-Psychopharm Biol Psych* 1998;22:115–128.
44. Wang CS, Lee RKK. Choline plus cytidine stimulate phospholipid production, and the expression and secretion of amyloid precursor protein in rat PC12 cells. *Neurosci Lett* 2000;283:25–28. [PubMed: 10729625]
45. Mir C, Clotet J, Aledo R, Durany N, Argemí J, Lozano R, Cervós-Navarro J, Casals N. CDP-choline prevents glutamate-mediated cell death in cerebellar granule neurons. *J Mol Neurosci* 2003;20:53–56. [PubMed: 12663935]
46. Barrachina M, Secades J, Lozano R, Gómez-Santos C, Ambrosio S, Ferrer I. Citicoline increases glutathione redox ratio and reduces caspase-3 activation and cell death in staurosporine-treated SH-SY5Y human neuroblastoma cells. *Brain Res* 2002;957:84–90. [PubMed: 12443983]
47. Oshitari T, Fujimoto N, Adachi-Usami E. Citicoline has a protective effect on damaged retinal ganglion cells in mouse culture retina. *Neuroreport* 2002;13:2109–2111. [PubMed: 12438935]
48. Plataras C, Angelogianni P, Tsakiris S. Effect of CDP-choline on hippocampal acetylcholinesterase and Na⁺, K⁽⁺⁾-ATPase in adult and aged rats. *Z Natur-forsch [C]* 2003;58:277–281.
49. Davalos A, Castillo J, Alvarez-Sabin J, Secades JJ, Mercadal J, Lopez S, Cobo E, Warach S, Sherman D, Clark WM, Lozano R. Oral citicoline in acute ischemic stroke: An individual patient data pooling analysis of clinical trials. *Stroke* 2002;33:2850–2857. [PubMed: 12468781]
50. Cacabelos R, Caamano J, Gomez MJ, Fernandez-Novoa L, Franco-Maside A, Alvarez XA. Therapeutic effects of CDP-choline in Alzheimer's disease. Cognition, brain mapping, cerebrovascular hemodynamics, and immune factors. *Ann NY Acad Sci* 1996;777:399–403. [PubMed: 8624120]
51. Wurtman RJ, Regan M, Ulus I, Yu L. Effect of oral CDP-choline on plasma choline and uridine levels in humans. *Biochem Pharmacol* 2000;60:989–992. [PubMed: 10974208]
52. Alvarez XA, Laredo M, Corzo D, Fernandez-Novoa L, Mouzo R, Perea JE, Daniele D, Cacabelos R. Citicoline improves memory performance in elderly subjects. *Methods Find Exp Clin Pharmacol* 1997;19:201–210. [PubMed: 9203170]
53. Renshaw PF, Daniels S, Lundahl LH, Rogers V, Lukas SE. Short-term treatment with citicoline (CDP-choline) attenuates some measures of craving in cocaine-dependent subjects: A preliminary report. *Psycho-pharmacology* 1999;142:132–138.
54. Anonymous. Cytidine diphosphate choline (citicoline, Somazina). *Arzneim-Forsch* 1983;33:1011–1080.
55. Lozano Fernandez R. Efficacy and safety of oral CDP-choline. Drug surveillance study in 2817 cases *Arzneim-Forsch* 1983;33:1073–1080.
56. Warach S. Dose dependent reduction in infarct growth with citicoline treatment: evidence of neuroprotection in human stroke? *Stroke* 2002;33:354.
57. Cui Z, Houweling M, Chen MH, Record M, Chap H, Vance DE, Terce F. A genetic defect in phosphatidylcholine biosynthesis triggers apoptosis in Chinese hamster ovary cells. *J Biol Chem* 1996;271:14668–14671. [PubMed: 8663247]
58. Cui Z, Houweling M. Phosphatidylcholine and cell death. *Biochim Biophys Acta* 2002;1585:87–96. [PubMed: 12531541]
59. Klein J. Membrane breakdown in acute and chronic neurodegeneration: Focus on choline-containing phospholipids. *J Neur Transm* 2000;107:1027–1063.
60. Nitsch RM, Blusztajn JK, Pittas AG, Slack BE, Growdon JH, Wurtman RW. Evidence for a membrane defect in Alzheimer disease brain. *Proc Natl Acad Sci USA* 1992;89:1671–1675. [PubMed: 1311847]
61. Blusztajn JK, Wurtman RJ. Choline and cholinergic neurons. *Science* 1983;221:614–620. [PubMed: 6867732]
62. Rego AC, Oliveira CR. Mitochondrial dysfunction and reactive oxygen species in excitotoxicity and apoptosis: Implications for the pathogenesis of neurodegenerative diseases. *Neurochem Res* 2003;28:1563–1574. [PubMed: 14570402]
63. Romrell J, Fernandez HH, Okun MS. Rationale for current therapies in Parkinson's disease. *Expert Opin Pharmacother* 2003;4:1747–1761. [PubMed: 14521485]
64. Ter Horst, GJ.; Korf, J., editors. *Clinical pharmacology of cerebral ischemia*. NJ: Humana; 1997.

65. Mattson MP, Culmsee C, Yu ZF. Apoptotic and antiapoptotic mechanisms in stroke. *Cell Tissue Res* 2000;301:173–187. [PubMed: 10928290]
66. Exton JH. Phosphatidylcholine breakdown and signal transduction. *Biochim Biophys Acta* 1994;1212:26–42. [PubMed: 8155724]
67. Tronchere H, Record M, Terce F, Chap H. Phosphatidylcholine cycle and regulation of phosphatidylcholine biosynthesis by enzyme translocation. *Biochim Biophys Acta* 1994;1212:137–151. [PubMed: 8180240]
68. Li YH, Maher P, Schubert D. Phosphatidylcholine-specific phospholipase C regulates glutamate-induced nerve cell death. *Proc Natl Acad Sci USA* 1998;95:7748–7753. [PubMed: 9636222]
69. Klein J, Chalifa V, Liscovitch M, Loffelholz K. Role of phospholipase D activation in nervous system physiology and pathophysiology. *J Neurochem* 1995;65:1445–1455. [PubMed: 7561837]
70. Six DA, Dennis EA. The expanding superfamily of phospholipase A₂ enzymes: Classification and characterization. *Biochim Biophys Acta* 2000;1488:1–19. [PubMed: 11080672]
71. Balsinde J, Winstead MV, Dennis EA. Phospholipase A₂ regulation of arachidonic acid mobilization. *FEBS Lett* 2002;531:2–6. [PubMed: 12401193]
72. Sun GY, Xu J, Jensen MD, Simonyi A. PLA₂ in the central nervous system: Implications for neurodegenerative diseases. *J Lipid Res* 2004;45:205–213. [PubMed: 14657205]
73. Allan SM, Rothwell NJ. Cytokines and acute neurodegeneration. *Nature Neurosci* 2001;2:734–744.
74. Allan SM, Rothwell NJ. Inflammation in central nervous system injury. *Philos Trans R Soc Lond B Biol Sci* 2003;358:1669–1677. [PubMed: 14561325]
75. Hallenbeck JM. The many faces of TNF in stroke. *Nature Med* 2002;8:1363–1368. [PubMed: 12457181]
76. Saito K, Suyama K, Nishida K, Sei Y, Basile AS. Early increases in TNF- α , IL-6 and IL-1- β levels following transient cerebral ischemia in gerbil brain. *Neurosci Lett* 1996;206:149–152. [PubMed: 8710173]
77. Wang CX, Shuaib A. Involvement of inflammatory cytokines in central nervous system injury. *Prog Neurobiol* 2002;67:161–172. [PubMed: 12126659]
78. Anthonsen MW, Solhaug A, Johansen B. Functional coupling between secretory and cytosolic PLA₂ modulates TNF- α - and IL-1 β -induced NF- κ B activation. *J Biol Chem* 2001;276:30527–30536. [PubMed: 11390371]
79. Heller RA, Kronke M. TNF receptor-mediated signaling pathways. *J Cell Biol* 1994;126:5–9. [PubMed: 8027185]
80. Schutze S, Berkovic D, Tomsing O, Unger C, Kronke M. TNF- α induces rapid production of 1',2'-diacylglycerol by a phosphatidylcholine-specific PLC. *J Exp Med* 1991;174:975–988. [PubMed: 1658188]
81. Woo CH, Eom YW, Yoo MH, You HJ, Han HJ, Song WK, Yoo YJ, Chun JS, Kim JH. TNF- α generates reactive oxygen species via a cytosolic PLA₂-linked cascade. *J Biol Chem* 2000;275:32357–32362. [PubMed: 10934206]
82. Balboa MA, Varela-Nieto I, Killermann Lucas K, Dennis EA. Expression and function of phospholipase A₂ in brain. *FEBS Lett* 2002;531:12–17. [PubMed: 12401195]
83. Galella G, Medini L, Stragliotto E, Stefanini P, Rise P, Tremoli E, Galli C. In human monocytes IL-1 stimulates a PLC active on phosphatidylcholine and inactive on phosphatidylinositol. *Biochem Pharmacol* 1992;44:715–720. [PubMed: 1510717]
84. Schutze S, Machleidt T, Kronke M. The role of diacylglycerol and ceramide in TNF- α and IL-1 signal transduction. *J Leukocyte Biol* 1994;56:533–541. [PubMed: 7964160]
85. Sun GY, Hu ZY. Stimulation of PLA₂ expression in rat cultured astrocytes by LPS, TNF- α and IL-1 β . *Prog Brain Res* 1995;105:231–238. [PubMed: 7568882]
86. Hayakawa M, Ishida N, Takeuchi K, Shibamoto S, Hori T, Oku N, Ito F, Tsujimoto M. Arachidonic acid-selective cytosolic PLA₂ is crucial in the cytotoxic action of TNF. *J Biol Chem* 1993;268:11290–11295. [PubMed: 7684376]
87. Machleidt T, Kramer B, Adam D, Neumann B, Schutze S, Wiegmann K, Kronke M. Function of the p55 TNF receptor “death domain” mediated by phosphatidylcholine-specific PLC. *J Exp Med* 1996;184:725–733. [PubMed: 8760826]

88. Bomalaski JS, Clark MA. PLA₂ and arthritis. *Arthrit Rheum* 1993;36:190–198.
89. Bomalaski JS, Steiner MR, Simon PL, Clark MA. IL-1 increases PLA₂ activity, expression of PLA₂-activating protein, and release of linoleic acid from the murine T helper cell line EL-4. *J Immunology* 1992;148:155–160. [PubMed: 1530793]
90. Clark MA, Chen MJ, Crooke ST, Bomalaski JS. TNF (cachectin) induces PLA₂ activity and synthesis of a PLA₂-activating protein in endothelial cells. *Biochem J* 1988;250:125–132. [PubMed: 3128274]
91. Boggs KP, Rock CO, Jackowski S. Lysophosphatidylcholine and 1-O-octadecyl-2-O-methyl-*rac*-gly-cero-3-phosphocholine inhibit the CDP-choline pathway of phosphatidylcholine synthesis at the CTP-phosphocholine cytidyltransferase step. *J Biol Chem* 1995;270:7757–7764. [PubMed: 7706325]
92. Shohami E, Ginis I, Hallenbeck JM. Dual role of TNF- α in brain injury. *Cytokine Growth Fact or Rev* 1999;10:119–130.
93. Mallampalli RK, Ryan AJ, Salome RG, Jackowski S. TNF- α inhibits expression of CTP : Phosphocholine cytidyltransferase. *J Biol Chem* 2000;275:9699–9708. [PubMed: 10734122]
94. Kontos HA. Oxygen radicals in cerebral ischemia. *Stroke* 2001;32:2712–2716. [PubMed: 11692043]
95. Uchida K. Current status of acrolein as a lipid peroxidation product. *Trends Cardiovasc Med* 1999;9:109–113. [PubMed: 10639724]
96. McCracken E, Graham DI, Nilsen M, Stewart J, Nicoll JA, Horsburgh K. 4-Hydroxynonenal immunoreactivity is increased in human hippocampus after global ischemia. *Brain Pathol* 2001;11:414–421. [PubMed: 11556686]
97. Katsuki H, Okuda S. Arachidonic acid as a neurotoxic and neurotrophic substance. *Prog Neurobiol* 1995;46:607–636. [PubMed: 8545546]
98. De Keyser J, Sulter G, Luiten PG. Clinical trials with neuroprotective drugs in acute ischaemic stroke: Are we doing the right thing? *Trends Neurosci* 1999;22:535–540. [PubMed: 10542428]
99. White BC, Sullivan JM, DeGracia DJ, O'Neil BJ, Neumar RW, Grossman LI, Rafols JA, Krause GS. Brain ischemia and reperfusion: Molecular mechanisms of neuronal injury. *J Neurol Sci* 2000;179:1–33. [PubMed: 11054482]
100. Onal MZ, Li FH, Tatlisumak T, Locke KW, Sandage BW, Fisher M. Synergistic effects of citicoline and MK-801 in temporary experimental focal ischemia in rats. *Stroke* 1997;28:1060–1065. [PubMed: 9158650]
101. Hurtado O, Moro MA, Cardenas A, Sanchez V, Fernandez-Tome P, Leza JC, Lorenzo P, Secades JJ, Lozano R, Davalos A. Neuroprotection afforded by prior citicoline administration in experimental brain ischemia: effects on glutamate transport. *Neurobiol Disease*. 2005in press
102. Wright MM, Howe AG, Zarembeg V. Cell membranes and apoptosis: role of cardiolipin, phosphatidylcholine, and anticancer lipid analogues. *Biochem Cell Biol* 2004;82:18–26. [PubMed: 15052325]
103. Wright MM, Henneberry AL, Lagace TA, Ridgway ND, McMaster CR. Uncoupling farnesol-induced apoptosis from its inhibition of phosphatidylcholine synthesis. *J Biol Chem* 2001;276:25254–25261. [PubMed: 11306571]
104. Han X, Gross RW. Global analyses of cellular lipidomes directly from crude extracts of biological samples by ESI mass spectrometry: a bridge to lipidomics. *J Lipid Res* 2003;44:1071–1079. [PubMed: 12671038]
105. Forrester JS, Milne SB, Ivanova PT, Brown HA. Computational lipidomics: a multiplexed analysis of dynamic changes in membrane lipid composition during signal transduction. *Mol Pharmacol* 2004;65:813–821. [PubMed: 15044609]
106. Marcheselli VL, Hong S, Lukiw WJ, Tian XH, Gronert K, Musto A, Hardy M, Gimenez JM, Chiang N, Serhan CN, Bazan NG. Novel docosanoids inhibit brain ischemia-reperfusion-mediated leukocyte infiltration and pro-inflammatory gene expression. *J Biol Chem* 2003;278:43807–43817. [PubMed: 12923200]
107. Bazan NG. Synaptic lipid signaling: significance of polyunsaturated fatty acids and platelet-activating factor. *J Lipid Res* 2003;44:2221–2233. [PubMed: 13130128]
108. Labiche LA, Grotta J. Clinical trials for cytoprotection in stroke. *NeuroRx* 2004;1:46–70. [PubMed: 15717007]

Abbreviations

CDP-choline (Citicoline)	cytidine-5'-diphosphocholine
CMP	cytidine 5'-monophosphate
CCT, CTP	phosphocholine cytidyltransferase
CTP	cytidine triphosphate
GSH	glutathione
IL-1	interleukin 1
PC	phosphatidylcholine
PLA₂	phospholipase A ₂
PLC	phospholipase C
ROS	reactive oxygen species
PLAP	PLA ₂ activating protein
TNF-α	tumor necrosis factor- α





Fig 1. CDP-choline: (A) possible neuroprotective pathways based on the published reports, (B) effects mediated by attenuating PLA₂ stimulation (based on authors' work). ↑ indicates increase; ↓ indicates decrease.

**Fig 2.**

TNF- α , IL-1 β , CDP-choline and PC homeostasis. (A) Potential pathways of CCT inhibition mediated by TNF- α . TNF- α and IL-1 β are induced after brain ischemia (74,77,92) and stimulate PLA₂ (78) and releases lyso-PC. TNF- α can inhibit CCT activity by two pathways: (1) proteolysis of CCT (93), and/ or (2) inhibition by lyso-PC (91). (B) TNF- α and IL-1 β may be responsible for the loss of PC through modulation of PLA₂ (78), PC-PLC (80) and CCT (84, 93) following transient cerebral ischemia. Hypothesis: CDP-choline may counteract the TNF- α /IL-1 β mediated disruption of PC homeostasis by attenuating PLA₂ activation and increasing CCT activity (Fig. 1B). \uparrow indicates increase; \downarrow indicates decrease. PLAP, PLA₂ activating protein.

Table I
CDP-Choline Studies in *in vivo* (Normal and Pathological) and *in vitro* Conditions

	Effect/Outcome by CDP-choline	References
<i>Reviews</i>		
Pharmacology and clinical use	Review of pharmacology and clinical use Development for stroke trials by drug companies (Ferrer, Takeda, and Indevus, formerly Interneuron)	(3,6) (7)
CNS injury	Mechanisms in cerebral ischemia & clinical efficacy Action in CNS injury and stroke trials	(5,8) (9,10)
Glaucoma	Application for glaucoma treatment	(11)
<i>In vivo studies</i>		
Transient forebrain ischemia in gerbil	Affected CCT activity, lyso-PC, OH• radical, lipid peroxidation, lipids, phospholipases A ₂ and C, arachidonic acid release, leukotriene C ₄ , GSH, and hippocampal neuronal death	(4,12–18)
Permanent forebrain ischemia in gerbil	CDP-choline but not cytidine alone provided hippocampal neuroprotection	(19)
Transient global ischemia in rat	Intracerebroventricular pre-treatment affected brain lipids Liposome encapsulation decreased lipid peroxidation and increased survival	(20) (21)
Transient focal cerebral ischemia in rat	Restored glucose metabolism and increased acetylcholine synthesis Combination with MK-801 or basic fibroblast growth factor synergistically reduced infarction Attenuated PLA ₂ activity, phospholipid loss and increase in FFA and ceramide	(22) (23) and references therein (24)
Decapitation ischemia in rat	Effects on lipids, Na ⁺ -K ⁺ -ATPase and acetylcholinesterase activities	(25)
Permanent focal cerebral ischemia in SD rat	In combination with nimodipine reduced apoptosis, infarct size, and increased Bcl-2 expression	(26)
Embolitic ischemic stroke in rat	Attenuated caspase cleaved products of PARP and DNA fragmentation Combination with urokinase improved neurobehavioral score and reduced infarction	(27) (28)
Subarachnoid hemorrhage	Combination with rtPA promoted functional recovery and reduced infarction Neuroprotection could be mediated by recovery of arterial pressure in hypotensive rats	(29) (30)
Cardiovascular effects on rat	Increased blood pressure and reversed hypotension in hemorrhagic shock	(31)
Traumatic brain injury in rat	Attenuated blood-brain barrier dysfunction, edema, cortical contusion volume, CA ₃ neuronal death, and neurological dysfunction Improved cognitive deficits	(32,33) (34)
Parkinson's disease model in rat	Attenuated substantia nigra dopaminergic cell dropout and tyrosine hydroxylase immunoreactivity	(35)
β-amyloid deposit + hypoperfusion in rat	Attenuated hippocampal neuronal apoptosis, and reduced microglia activation	(36)
Rat	Reduced benzodiazepine-induced cognitive impairments	(37)
Female rats	Increased CCT activity and decreased platelet activating factor Protected selective impairment in hippocampal deficits PC levels increased only after 42 days of CDP-choline administration	(38) (39) (40)
Intracerebral hemorrhage in Swiss albino mice	Improved functional outcome, reduced infarct volume	(41)
Hypoxia in guinea pig	Affected phospholipid biosynthesis in brain mitochondria.	(42)
Dog	Increased memory and learning	(43)
<i>In vitro studies</i>		
PC12 cells	Cytidine + choline stimulated phospholipid production, expression and secretion of amyloid precursor protein	(44)
Cerebellar granule neurons/MCAO	Prevented glutamate mediated cell death	(45,101)
Human neuroblastoma SH-SY5Y cells	Increased glutathione redox ratio and attenuated caspase-3 activation in staurosporine-treated cells	(46)
Mouse retinal ganglion cells	Showed protective effects on damaged retinal ganglion.	(47)
Rat brain homogenates	Stimulated acetylcholinesterase and Na ⁺ -K ⁺ -ATPase activities in tissue homogenates	(48)
<i>Clinical studies</i>		
Stroke	Review of US clinical trials. Oral treatment of patients within the first 24 h after stroke increased probability of recovery at 3 months	(49 and references cited therein)
Alzheimer's disease	Improved mental performance in Alzheimer's disease	(50)
Parkinson's disease	In combination with levodopa showed functional improvement and decreased side effects of levodopa	(3)
Normal subjects	Increased choline and uridine but not cytidine levels	(51)
Memory	Improved memory deficits in elderly subjects	(52)
Amblyopia/glaucoma	Improved the visual activity, function in patients	(11)
Drug addiction	Reduced craving for cocaine	(53)
Alcoholism	Improved concentration in patients with alcoholic abstinence syndrome	(3)