

Complete Genome Sequence of the Oral Pathogenic Bacterium *Porphyromonas gingivalis* Strain W83

Karen E. Nelson,^{1*} Robert D. Fleischmann,¹ Robert T. DeBoy,¹ Ian T. Paulsen,¹ Derrick E. Fouts,¹
Jonathan A. Eisen,¹ Sean C. Daugherty,¹ Robert J. Dodson,¹ A. Scott Durkin,¹ Michelle Gwinn,¹
Daniel H. Haft,¹ James F. Kolonay,¹ William C. Nelson,¹ Tanya Mason,¹ Luke Tallon,¹
Jessica Gray,¹ David Granger,¹ Hervé Tettelin,¹ Hong Dong,² Jamie L. Galvin,²
Margaret J. Duncan,² Floyd E. Dewhirst,² and Claire M. Fraser¹

*The Institute for Genomic Research, Rockville, Maryland 20850,¹ and Department of
Molecular Genetics, The Forsyth Institute, Boston, Massachusetts 02115²*

Received 5 November 2002/Accepted 24 June 2003

The complete 2,343,479-bp genome sequence of the gram-negative, pathogenic oral bacterium *Porphyromonas gingivalis* strain W83, a major contributor to periodontal disease, was determined. Whole-genome comparative analysis with other available complete genome sequences confirms the close relationship between the *Cytophaga-Flavobacteria-Bacteroides* (CFB) phylum and the green-sulfur bacteria. Within the CFB phyla, the genomes most similar to that of *P. gingivalis* are those of *Bacteroides thetaiotaomicron* and *B. fragilis*. Outside of the CFB phyla the most similar genome to *P. gingivalis* is that of *Chlorobium tepidum*, supporting the previous phylogenetic studies that indicated that the *Chlorobia* and CFB phyla are related, albeit distantly. Genome analysis of strain W83 reveals a range of pathways and virulence determinants that relate to the novel biology of this oral pathogen. Among these determinants are at least six putative hemagglutinin-like genes and 36 previously unidentified peptidases. Genome analysis also reveals that *P. gingivalis* can metabolize a range of amino acids and generate a number of metabolic end products that are toxic to the human host or human gingival tissue and contribute to the development of periodontal disease.

Periodontal diseases are a group of infections that affect the structures surrounding teeth. If allowed to progress, periodontal disease can cause the destruction of supporting connective tissue and bone, ultimately resulting in tooth loss. Initiation and progression of periodontal diseases is the result of a complex interaction between the bacteria colonizing the gingival crevice and the hosts' immune and inflammatory responses. Since the species most strongly implicated in periodontal disease pathogenesis are also usually present in low numbers in healthy people, the distinction between pathogenic and commensal bacteria in the human host is not clearly defined.

The gram-negative anaerobe *Porphyromonas gingivalis* belongs to the family *Porphyromonadaceae*, order *Bacteroidales* in the phylum *Bacteroidetes*, previously known as the *Cytophaga-Flavobacteria-Bacteroides* (CFB) group (6). The bacterium is a major causative agent in the initiation and progression of severe forms of periodontal disease. *P. gingivalis* is a late or secondary colonizer of the oral cavity, a process that is facilitated by other microbial species that provide attachment sites, as well as supply growth substrates, and reduce oxygen tension to levels optimal for growth of *P. gingivalis*. Among the early plaque organisms that *P. gingivalis* adheres to are the oral streptococci (35, 36) and *Actinomyces naeslundii* (19). Adherence is facilitated by a variety of bacterial surface proteins, including fimbriae, hemagglutinins, and proteinases. *P. gingivalis* also binds to late colonizers such as *Fusobacterium nucleatum*, *Treponema denticola*, and *Bacteroides forsythus* (now

renamed *Tanerella forsythensis*) (20, 30, 72). The use of a variety of metabolic strategies appears to enable the success of this microbial community. Once established, *P. gingivalis* cells participate in intercellular communication networks with other oral prokaryotic cells, as well as with eukaryotic cells (37).

P. gingivalis is the third oral pathogen (1, 28) and the second member of the CFB group to be sequenced. The genome of *P. gingivalis* strain W83 (also known as strain HG66) is presented here. Strain W83 was isolated in the 1950s by H. Werner (Bonn, Germany) from an undocumented human oral infection (41, 46) and was brought to The Pasteur Institute by Madeleine Sebald during the 1960s. The strain was subsequently obtained by Christian Mouton (Quebec, Canada) during the late 1970s. It is anticipated that the availability of the complete genome sequence from this oral pathogen will give tremendous insight into the mechanisms that result in disease progression in an ecological niche deep within the oral cavity.

MATERIALS AND METHODS

Library preparation and random sequencing of *P. gingivalis*. *P. gingivalis* strain W83 was obtained from Christian Mouton, Laval University, Quebec City, Quebec, Canada. Genomic DNA was extracted twice with buffered phenol and once with 25:24:1 phenol-chloroform-isoamyl alcohol and precipitated with alcohol. Cloning, sequencing, and assembly were as described previously for genomes sequenced by The Institute for Genomic Research (TIGR) (13). One small-insert plasmid library (1.5 to 2.5 kb) was generated by random mechanical shearing of genomic DNA. One large-insert library was generated by partial *Tsp5091* digestion and ligation to the λ -DASHIII/*EcoRI* vector (Stratagene). In the initial random sequencing phase, ~8-fold sequence coverage was achieved with 39,623 sequences (average read length, 534 bases). The plasmid and λ sequences were jointly assembled by using TIGR Assembler. Sequences from both ends of 506 λ clones served as a genome scaffold, verifying the orientation,

* Corresponding author. Mailing address: The Institute for Genomic Research, 9712 Medical Center Dr., Rockville, MD 20850. Phone: (301) 838-3565. Fax: (301) 838-0200. E-mail: kenelson@tigr.org.




FIG. 1. Circular representation of the *P. gingivalis* genome. The outer circle shows the predicted coding regions on the plus strand color-coded by role categories as follows: violet, amino acid biosynthesis; light blue, biosynthesis of cofactors, prosthetic groups, and carriers; light green, cell envelope; red, cellular processes; brown, central intermediary metabolism; yellow, DNA metabolism; light gray, energy metabolism; magenta, fatty acid and phospholipid metabolism; pink, protein synthesis and fate; orange, purines, pyrimidines, nucleosides, and nucleotides; olive, regulatory functions and signal transduction; dark green, transcription; teal, transport and binding proteins; gray, unknown function; salmon, other categories; blue, hypothetical proteins. The second circle shows the predicted coding regions on the minus strand. The third circle presents the χ^2 analysis of atypical nucleotide composition; χ^2 values of >600 are indicated in red. The fourth circle shows the %G+C. The fifth circle shows atypical nucleotide composition (GC skew). The sixth circle shows the IS elements, indicated by color as follows: orange, ISPg1; light green, ISPg2; magenta, ISPg3; cyan, ISPg4; brown, ISPg5; gold, ISPg6; blue-green, ISPg7; pink, ISPg8; and violet, ISPg9; salmon, ISPg10; olive, ISPg11. The seventh circle shows MITE239 (magenta), MITE700 (cyan), and MITE464 (black). The eighth circle shows Tn4555 (blue), CTn (red), and other transposable elements (green). The ninth circle shows tRNA (green), rRNA (black), and sRNA (red).

order, and integrity of the contigs. Sequence gaps were closed by editing the ends of sequence traces and/or primer walking on plasmid clones. Physical gaps were closed by direct sequencing of genomic DNA or combinatorial PCR, followed by sequencing of the PCR product. The final molecule has $8.35\times$ sequence redundancy.

ORF prediction and gene family identification. An initial set of open reading frames (ORFs) that likely encode proteins was identified with GLIMMER (57), and those shorter than 90 bp, as well as some of those with overlaps, were eliminated. (For more details on the annotation process that was used to identify all of the ORFs in the *P. gingivalis* genome, see reference 70.) A region contain-

ing the likely origin of replication was identified, and bp 1 was designated adjacent to the *dnaA* gene that is in this region. ORFs were searched against a nonredundant protein database as previously described. Frameshifts and point mutations were detected and corrected where appropriate as described previously (51). Remaining frameshifts and point mutations are considered authentic, and corresponding regions were annotated as an "authentic frameshift" or an "authentic point mutation," respectively. ORF prediction and gene family identification was completed by using the methodology described previously (49). Two sets of hidden Markov models (HMMs) were used to determine ORF membership in families and superfamilies. These included 721 HMMs from

TABLE 1. Major families of contiguous repeats and their copy numbers in the genome

Designation ^a	No. of copies	No. of direct repeats (mean no.)	Sequence of direct repeat ^b
CR-41-A	12	2–16 (8)	TGCAGGACGCGATTGTCAA·CTGATTCTTGCTTCCTGCACGA
CR-41-B	1	4 (4)	GCTTGCAGGAA·GCGATTTCG·GCAAAAAAAAAATTCCTGTTTGCA
CR-41-C	1	4 (4)	TGCAGGAAAGCCGTTGTTCAC·CTGACAATCACTCCCTGCACGA
CR-23-A	14	3–14 (9)	TATCCTCTCAT·ATGAGACCATAG
CR-23-B	5	5 (5)	ACTAAACTAT·ATACTTTGGTNNN
CR-22-A	32	2–8 (5)	CGATTTATA·TATAAATCGAAAA
CR-22-B	4	4–6 (5)	TATTTCCGTAT·ATACGGAAATA
CR-22-C	2	4–5 (5)	AAACGTCT·AGACGTTTCGAGCA

^a The number indicates the length of the repetitive element; the letter indicates the family designation. Not included in table are CR-23-C and -D and CR-22-D, -E, -F, -G, and -H.

^b A dot indicates the location of dyad symmetry; underlined bases indicate palindromic match(es).

Pfam v2.0 and 631 HMMs from TIGR ortholog resource. TMHMM (33) was used to identify membrane-spanning domains in proteins.

Comparative genomics. All genes and predicted proteins from this genome, as well as from all other completed genomes, were compared by using the basic local alignment search tool (BLAST). For the identification of recent gene duplications, all genes from the *P. gingivalis* genome were compared to each other. A gene was considered recently duplicated if the most similar gene (as measured by *P* value) was another gene within the same genome (relative to genes from the two other genomes).

Database submission. The nucleotide sequence of the whole genome of *P. gingivalis* was submitted to GenBank under accession number AE015924.

RESULTS

Description of the genome. The genome of *P. gingivalis* is 2,343,479 bp, with an average G+C content of 48.3% (Fig. 1). There are four ribosomal operons (5S-23S-tRNA^{Ala}-tRNA^{Ile}-16S) and 2 structural RNA genes, as well as 53 tRNA genes with specificity for all 20 amino acids. A total of 1,990 ORFs could be identified in the genome (13). Of these, 1,075 (54%) could be assigned to biological role categories (54), 184 (9.2%) were conserved hypothetical proteins or conserved domain proteins, 208 (10.5%) were of unknown function, and 523 (26.3%) encoded hypothetical proteins. More than 85% of the genome encodes ORFs.

Repetitive elements occupy ca. 6% of the *P. gingivalis* genome and fall into two major classes: DNA repeats and transposable elements. The DNA repeats include uninterrupted direct repeats (Table 1), and a subclass of dispersed repeats known as clustered regularly interspaced short palindromic repeats (CRISPRs) (Table 2). Strain W83 does not appear to contain other classes of dispersed repetitive DNA sequence elements such as ERIC and REP elements. The transposable elements include insertion sequence (IS) elements and miniature inverted-repeat transposable elements (MITEs), which are summarized in Table 3, and large stretches of genes that resemble remnants of conjugable and mobilizable transposons

based on sequence similarity to elements previously described in *Bacteroides* species (11). The locations of transposon-associated genes are shown in Fig. 1. Although there are 96 complete or partial copies of IS elements and MITEs present in strain W83 that occupy more than 94 kb of the genome, the transposable elements are rarely found in a functional gene. Instead, these elements have inserted almost exclusively into intergenic regions and other copies of transposable elements, except for one insertion into a putative outer membrane protein (PG0176/PG0178) that is intact in at least four other strains of *P. gingivalis* (accession numbers AB069977 to AB069980). Analysis of the IS elements reveals two possible chromosomal inversions that most likely arose by homologous recombination between identical copies of elements at widely separated insertion sites. These potential DNA rearrangements are revealed by inspection of both the duplicated target site sequences (direct repeats) that flank some of the IS elements and a transposon gene disrupted by one IS insertion. The sites of these putative inversions are shown in Fig. 1. In one case, 821 kb (35%) of the chromosome appears to be inverted between one copy of *ISPg2* (PG1746) 512 kb before the origin and another copy of *ISPg2* (PG0277) 309 kb after the origin (red dots). In the second case, 103 kb (4.4%) of the chromosome appears to be inverted between two copies of *ISPg4*, which are located on opposite sides of the origin (PG2194 and PG0050) (green dots). Since inversions about the origin do not invert the direction of transcription relative to replication of genes on the segment, such inversions may be selectively neutral. It will be interesting to determine whether strain W83 and other strains of *P. gingivalis* share a common genetic structure, or whether the proposed chromosomal inversions are relatively recent events.

There are 21 areas of the genome that display an atypical nucleotide composition identified by χ^2 analysis (67) and that

TABLE 2. CRISPRs

Designation	Associated gene(s)	No. of direct repeats	Sequence of direct repeat ^b
CRISPR-30-36 ^a	PG2013 <i>cas2</i> , PG2014 <i>cas1</i> , PG2015 <i>cas4</i> , PG2016 <i>cas3</i>	23	GTTTTAATTCCTGTA·TGGTGCAATTGAAAT
CRISPR-36-30-A	PG0338	8	GTTGGATCTACCCCTCTAT·TCGAAGGGTACACACAAC
CRISPR-36-30-B	PG1164	8	GTTGGGAATACCCCTTAGT·TAGAAGGGTGGAGACAAC
CRISPR-37-35	PG1981, PG1982	8	GTCTTAATAGCCTTACGGACTGTGTATGTATAGTGAG

^a CRISPR-30-36 and *cas1* to *cas4* were described previously (26).

^b See Table 1, footnote b.

TABLE 3. IS elements and MITEs

Element ^a	Family	No. of copies (no. of fragments) ^b	Inverted repeat (length [bp]) ^c	Direct repeat length ^d	Length (bp)
ISPg1	IS5	25(7)	GAGACCTTTGCA (12)	5	1,334
ISPg2	ISAs1	5(11)	CAGGGCTGACGCATTAAA (18)	9	1,207
ISPg3	IS5	6(1)	ACGTCAGTTCTGA (12)	7	1,070
ISPg4	IS5	10(0)	CATTAGTGTCCTAAAAG (17)	9	1,394
ISPg5	IS3	11(1)	TTGACGTGTACTGAAAAAAGTTGACAGTTGG (31)	3	1,514
ISPg6	ISAs1	1(2)	CAGGGCAACCGCATCAA (18)	ND ^e	1,139
ISPg7	ISRM7	1(3)	GTCGTCCCCTTAAgAAcCTT (22)	6	1,428
ISPg8	IS5	0(1)	Unknown	ND	ND
ISPg9	IS5	2(2)	GAGACCATTGCA (12)	5	1,326
ISPg10	IS256	1	TGaCatTtTAAAAAaGAcATGG (24)	ND	~1,283
ISPg11	IS256	(1)	Unknown	ND	ND
MITE239	ISPg3	5	ACGysAGTTCGATmTAArsGAA	6–8	239
MITE464	ISPg1	13	GAGACTGTTGCA	5	464–1,071
MITE700	ISPg3	8	ACGTCAGTTCTGA	7	455–717

^a ISPg1, ISPg2, and ISPg3 were originally described as IS1126 (44), PGIS2 (69), and IS195 (40).

^b That is, the number of intact copies with the number of fragments in parentheses.

^c Lowercase lettering indicates sequence at a position which is not part of the perfect inverted repeat.

^d Direct repeat or duplicated target site length.

^e ND, not determined.

also correspond to regions of higher or lower G+C content than the rest of the genome. The areas range in size from 11 to 68 kb and range in G+C content from 29.4 to 61.6%. A variety of genes that could possibly have been acquired by this bacterium through lateral gene transfer are encoded in these regions. The genes include three restriction system proteins (PG0971, most similar to *Anabaena* sp. strain PCC 7120; PG0968, most similar to *Anabaena* sp. strain PCC 7120; and PG1469 most similar to *Agrobacterium tumefaciens*); hemagglutinin proteins B and C (HagB, PG1972, *P. gingivalis* specific; and HagC, PG1975, *P. gingivalis* specific); many capsular biosynthesis proteins, 20 transposase genes, two large mobile elements (PG1473 to PG1480, resembles only a conjugative element of *Bacteroides thetaiotaomicron*; and PG0868 to PG0875, whose sequence and gene organization most closely resembles the antibiotic-resistant mobilizable transposon Tn4555 from *Bacteroides fragilis* (68); and a thiamine biosynthesis operon (PG2107 to PG2111, which is most similar to the thiamine biosynthesis operon of *Escherichia coli*). These atypical regions in the *P. gingivalis* genome also encode many hypothetical and conserved hypothetical proteins, which undoubtedly contribute to the unique biology of this organism.

Comparative genomics. Comparison of the predicted proteome of *P. gingivalis* with that of other completely sequenced genomes confirms the close relationship of *P. gingivalis* to other members of the CFB, including *B. fragilis* and *B. thetaiotaomicron*. Outside of the CFB phyla, the genome most similar to that of *P. gingivalis* is the *Chlorobium tepidum* genome, supporting previous phylogenetic studies that indicated the chlorobia and CFB phyla are related, albeit distantly. The proteomes most similar to that of *P. gingivalis* (in terms of the number of proteins with the best scoring matches) were those of *B. thetaiotaomicron* and *B. fragilis* with 572 and 437 best-scoring matches ($P < 10^{-5}$), respectively.

A total of 332 genes were identified as being putatively duplicated in the *P. gingivalis* lineage. These duplicated genes are likely an indication that there is some selective evolutionary advantage to retaining these genes in the genome. Among these genes are 10 that encode DNA-binding histone-like pro-

teins that have a distinctive domain architecture compared to HU and related histone-like proteins. These DNA-binding proteins have been designated a superfamily (i.e., a set of proteins that share a given domain architecture; TIGRFAMs family TIGR01201). Outside of *P. gingivalis*, the single known example of a DNA-binding histone-like protein is found in the gut bacterium *B. fragilis*. All members of this superfamily are distantly related to the bacterial DNA-binding protein HU family (Pfam family PF00216, five of which are also found in the *P. gingivalis* genome) but differ in architecture, sharing both an N-terminal extension and a glycine-rich C terminus. HU has been shown, among other DNA-binding functions, to assist the unwinding of *oriC* DNA by the DNA replication initiation protein DnaA (4). Interestingly, all 10 members of the TIGR01201 family in *P. gingivalis* have direct repeats upstream of their genes that may act as binding sites for the DNA-binding proteins that are encoded by the nearby gene and perhaps regulate their own expression. Alternatively, the repeats may also coordinate expression of the other chromosomal genes that they flank.

Metabolism and transport. The microbial species that exist in supragingival plaque of the oral cavity are exposed to the host's dietary intake, and many of these bacteria, including the oral streptococci, ferment carbohydrates to acidic end products such as lactic acid for the purpose of energy production. On the other hand, anaerobic species in the subgingival plaque are exposed to crevicular fluid and to the host tissue proteins (61). The availability of the complete genome sequence of *P. gingivalis* W83 allowed for an analysis of the physiological potential of this species. Based on this analysis, the range of transport capabilities and metabolic pathways that could be identified is presented in Fig. 2.

Genome analysis suggests that *P. gingivalis* possesses a limited capacity for the uptake and metabolism of organic nutrients. Glucose utilization by *P. gingivalis* is known to be very poor, and carbohydrates in general do not appear to readily support growth (61). Strain W83 does, however, contain putative ORFs for all enzymes of the glycolytic pathway, as well as ORFs for a putative glucose/galactose transporter and glucose




FIG. 2. Overview of metabolism and transport in *P. gingivalis*. Primary substrates for energy metabolism are capitalized and underlined. End products of fermentation are highlighted by yellow boxes. Transporters are grouped by substrate specificity and indicated by color as follows: inorganic cations (green), organic nutrients (magenta), and drug efflux and other (black). Arrows indicate direction of transport for substrates (and coupling ions, where appropriate).

kinase. Sequence analysis shows that the glucose kinase is encoded in a split ORF generated by a missense mutation, and this is a likely explanation for the poor utilization of glucose to support growth. Four putative ORFs for the pentose phosphate pathway were identified, and it is likely that this pathway plays a role in the generation of precursor metabolites during anaerobic growth (Fig. 2).

Whole-genome analysis suggests that *P. gingivalis* can metabolize several sugars, including melibiose, galactose, starch, and maltodextrin. The bacterium also possesses enzymes for the degradation of complex amino sugars in the form of hexose aminidases. It is still unclear whether these complex sugars are metabolized, but one possibility is that the removal of amino sugars from host glycoproteins likely renders these proteins more susceptible to degradation by bacterial proteinases. In addition, at least 11 amino acids may serve as substrates for energy production (Fig. 2). These amino acids are most likely derived from the degradation of host tissues (see virulence section below) or from the breakdown of other bacterial cells in the oral cavity. Pathways for glutamate and aspartate utilization have been characterized by enzyme assays (65), and ORFs coding for all of these activities were found in the W83 genome. Intracellular glutamate is deaminated to 2-oxoglutarate by glutamate dehydrogenase and then decarboxylated to succinyl coenzyme A (succinyl-CoA) by a CoA-dependent 2-oxoglutarate oxidoreductase. The possession of this activity is somewhat unusual in bacterial species (23, 25). It has been established that two-thirds of the succinyl-CoA produced in this reaction is converted to butyryl-CoA and then to butyrate. The remaining third may be converted to propionate by a pathway that involves the enzymes methylmalonyl-CoA mutase and acyl-CoA:acetate-CoA transferase, as reported for other propionate-producing bacteria (18). This pathway appears to be unique to *P. gingivalis* since other anaerobes catabolize glutamate through the hydroxyglutarate, methylaspartate, and/or the aminobutyrate pathways (5, 17, 18). *P. gingivalis* did not possess activities for three key enzymes of these pathways: hydroxyglutarate dehydrogenase, 3-methylaspartate ammonia lyase, and 4-aminobutyrate aminotransferase (65). Peptide-derived aspartate is deaminated to fumarate by aspartate ammonia lyase and then either oxidized to acetate or reduced to propionate and butyrate (65).

Results from Takahashi et al. (65) suggest that *P. gingivalis* prefers to utilize arginine and lysine as free amino acids rather than in peptide form; thus, carboxy-terminal arginine and lysine residues could be released from proteins by carboxypeptidase activities. Masuda et al. (45) found such an activity in culture supernatants, and an ORF coding for an unspecified carboxypeptidase (PG0232) was identified in the genome. A report that *P. gingivalis* produces citrulline and ornithine from denatured protein (14) implies that the bacterium degrades arginine through the arginine deiminase pathway. Indeed, a gene with homology to arginine deiminase from *Bacillus licheniformis* (43) was identified. In addition, two genes—*pyrB* and *pyrI* (PG0357 and PG0358)—were contiguous in the genome and shared homology with aspartate/ornithine transcarbamylase catalytic and regulatory chains from *Vibrio* sp. strain 2693 and *Pyrococcus abyssi*, respectively.

The lysine catabolic pathways appear to be very similar to those found in *Clostridium* sp. ORFs were identified for the

first steps of both L- and D-lysine catabolism; thus, the isomers are apparently degraded by two different pathways that yield butyric acid, acetic acid, and ammonia. Lysine 2,3-aminomutase (KamA) catalyzes the interconversion of L-lysine and L-β-lysine, the first step in the lysine degradation pathway in *Clostridium subterminale* SB4 (56). In *P. gingivalis* W83, *kamA* was found clustered with the genes *kamD* and *kamE* (PG1070, PG1073, and PG1074) that encode subunits of D-lysine 5,6-aminomutase, the first enzyme of the D-lysine degradative pathway. Genes encoding enzymes for the subsequent conversion of lysine to butyrate and acetate were located 3' to *kamE*. It is not yet known whether these genes are transcribed as an operon.

Little is known about serine and threonine catabolism in *P. gingivalis*; however, an ORF was detected with homology to serine dehydratase (PG0084) that hydrolyzes serine to pyruvate, ammonia, and water. Threonine may be split to glycine and acetaldehyde by the activity of threonine aldolase, for which an ORF was detected (PG0474). In summary, *P. gingivalis* appears to catabolize amino acids through pathways that generate ammonia. The organism has a growth pH optimum of >7.5, and ammonia generation may have evolved as a strategy to shift the local pH to the favored alkaline range.

Several studies have shown that *P. gingivalis* preferentially uses peptides as sources of carbon and nitrogen (60, 65, 71) and, in addition to the previously described proteinases that are known to degrade host proteins, a number of peptidases that may be involved in the further digestion of protein fragments to smaller peptides and amino acids could be identified from the genome.

There are two carboxylate transporters possibly for lactate and formate, and no sugar transporters other than the aforementioned glucose/galactose importer. Although *P. gingivalis* possesses a broad assortment of secreted peptidases and pathways for the metabolism of amino acids, the bacterium appears to rely on two predicted peptide uptake systems and has only one amino acid transporter, the characterized sodium ion-driven serine/threonine uptake protein SstT (12). A LysE-type amino acid efflux protein is present that may protect the organism from toxic concentrations of amino acids.

The major fermentation products that can be produced based on whole-genome analysis and in vitro end product analyses are propionate, butyrate, isobutyrate, isovalerate, acetate, ethanol, and butanol (27). Many of these end products are probably toxic to human host tissues (see virulence section below).

Nucleosides and nucleobases may represent a hitherto-unsuspected important nutrient source for *P. gingivalis* and might be used either as building blocks for nucleic acid biosynthesis or may be catabolized as carbon and energy sources. There are three predicted purine uptake systems, a NupG nucleoside uptake system, and a homolog of the *Salmonella enterica* serovar Typhimurium nicotinamide mononucleotide transporter PnuC. In addition, there are four homologs of *E. coli* DinF, a DNA damage-induced protein related to sodium ion-driven drug efflux transporters, that are hypothesized to play a role in nucleoside and/or nucleotide efflux (8).

Common to most human pathogens, iron acquisition appears to be an important priority in *P. gingivalis*, and there are two iron chelate ABC uptake systems, two TonB-dependent

iron receptors, and two FeoB ferrous iron uptake systems. There is an array of metal ion homeostasis transporters, including three sodium ion/proton exchangers, which may be important since a significant number of *P. gingivalis* transporters are predicted to be sodium ion driven.

Virulence and *P. gingivalis*. The availability of the complete genome sequence of *P. gingivalis* facilitates the identification of putative virulence factors associated with the establishment and survival of the bacterium in the gingival crevice and subsequent penetration into host cells (Table 4). Initially, the bacterium must navigate the oral cavity where, as an obligate anaerobe, it is exposed to limited amounts of oxygen before it establishes itself in an anaerobic environment. A cluster of genes (PG1582 to PG1586) was identified with high levels of similarity to the recently described aerotolerance operon of *B. fragilis* (66). These functions promote the survival of *B. fragilis* upon exposure to oxygen, and their presence in *P. gingivalis* suggests that this system may also ensure tolerance to oxygen in the oral cavity. The genome also encodes a superoxide dismutase (PG1545), genes for an alkyl hydroperoxide reductase (PG0618 and PG0619) (55), a thiol peroxidase (PG1729), and a Dps homolog (PG0090) that is involved in the repair of oxidatively damaged nucleic acids (33).

The bacterium uses fimbriae to adhere to other bacterial species and host tissues. Hemagglutinins and various proteases (gingipains) are also involved in tissue colonization through adhesion to extracellular matrix proteins (38, 53, 59). Hemagglutinins in particular may mediate the binding of bacteria to receptors on human cells (21), and the gene sequences for six newly identified putative hemagglutinin-like proteins (PG0411, PG1326, PG1674, PG1427, PG1548, and PG2198) could be identified. Four of these are recent duplications in the genome of HagA and HagD adhesin domain-related sequences. A total of 42 proteinases were identified in the genome sequence that may enable adherence of the bacterium to host tissues, as well as to other bacterial cells, and that may also degrade host proteins (as discussed above). In vitro experiments have demonstrated that proteases attack a range of host proteins, including extracellular matrix proteins (32, 38, 53, 59) and cell adhesion molecules (29), the destruction of which leads to a loss of cell surface receptors (59) and tissue integrity (29). Protease destruction of cytokines (15, 42, 74) and gamma interferon (73) can result in disruption of polymorphonuclear leukocyte function (48) and ultimately affect the host immune response.

A single hemolysin for the release of iron and protophorin IX (PG1875) was identified. This sequence has full-length homology only to the characterized hemolysin gene of another periodontal pathogen, *Prevotella melaninogenica* (3). These two hemolysins show absolutely no homology to any other biochemically characterized hemolysin and have weak homology to a conserved hypothetical protein/putative hemolysin fusion protein sequence from *Vibrio cholerae*.

In *P. gingivalis*, metabolic end products from the catabolism of various substrates include short-chained carboxylic acids that can affect the host defense system in a variety of ways. When applied directly to healthy human gingiva tissue, short-chain carboxylic acids have been shown to stimulate a gingival inflammatory response and inflammatory cytokine release (50). Short-chain carboxylic acids have also been shown to alter

cell function and gene expression and may also contribute to the initiation and prolongation of gingival inflammation (50).

The capsule of *P. gingivalis* is most likely involved in the evasion of the host response and has been shown to be one of the important virulence determinants in this bacterium (34). Whole-genome analysis reveals at least four capsular biosynthesis gene clusters (PG0106 to PGPG0120, PG0435 to PG0437, PG1140 to PG1149, and PG1560 to PG1565) that are located across the genome. Closer investigation of these gene clusters suggests that mannose, glucose, and rhamnose may be some of the sugars that are present in the capsule of *P. gingivalis* strain W83. In several pathogens the secretion of virulence factors targeted to the host cells is mediated by type III protein secretion systems. The complete genome of *P. gingivalis* was searched for the presence of a cluster of nine genes Sct (Hrc/Ysc) that are known to be components of type III protein secretion systems (24). No BLAST matches with these motifs were found. Although several *sec* gene homologs are present in the genome, including SecA, SecY, SecD, and SecF, the main terminal branch of the general secretory pathway (type II) could not be identified, suggesting that this pathway is not functional in this bacterium.

DISCUSSION

It is estimated that 35% of the U.S. population has some form of periodontitis (2). Traditional methods for the prevention or treatment of periodontal disease include the mechanical removal of plaque and the use of antimicrobial agents. The availability of the genome sequences of at least three oral pathogens and recent investigations into the microbial composition of the human oral microbiome will afford new opportunities to investigate ways to alter the composition of the subgingival biofilm. Potential strategies include those that would decrease opportunities for biofilm formation, reduce attachment among species in the biofilm, or limit the availability of required nutrients. Inhibition of the primary colonizers could prevent the successful establishment of late colonizers such as *P. gingivalis*.

Recently, specific bacteria have been found to be associated with systemic diseases, e.g., *Helicobacter pylori*, as etiological agent in gastric ulcers, and the tentative association between the presence of *Chlamydia pneumoniae* in atherosclerotic plaques and cardiovascular disease (9). Stimulated by these studies and the detection of *P. gingivalis* and other oral pathogens in atherosclerotic plaques (10, 62), new research is assessing associations between periodontal infection and cardiovascular disease (22, 47, 52, 58). That the pathogenicity of oral bacteria may extend beyond their known ecological niche to other organ systems introduces an important and exciting new dimension to defining the genetic complement of these organisms. Although the genome sequence has not revealed many of the classical virulence factors that are associated with pathogens, it is anticipated that the newly described putative virulence factors of this bacterium, as well as from the others that have been recently sequenced or whose sequencing is nearing completion, will enable the development of antimicrobial agents that can be used against one of the major causative agents of periodontal disease. Ultimately, the genome sequence of *P. gingivalis* will facilitate an increased understanding of the virulence

TABLE 4. Previously characterized and newly identified putative virulence agents from the *P. gingivalis* genome sequence^a

Function and virulence agent	Description (reference)	Identification
Acquisition of iron		
PG1875	Hemolysin	New identification
Adhesion		
PG0411	Hemagglutinin, putative	New identification
PG0443	Hemagglutinin-related protein	New identification
PG1326	Hemagglutinin, putative	New identification
PG1427	Thiol protease/hemagglutinin PrtT precursor, putative	New identification
PG1548	Thiol protease/hemagglutinin PrtT precursor	New identification
PG1674	Hemagglutinin protein HagB, degenerate	New identification
PG1837	Hemagglutinin protein HagA (21)	Previously known
PG1844	Hemagglutinin protein HagD (64)	Previously known
PG1972	Hemagglutinin protein HagB	Previously sequenced
PG1975	Hemagglutinin protein HagC (39)	Previously known
PG2024	Hemagglutinin protein HagE (64)	Previously known
PG2132	Fimbrilin (16)	Previously known
PG2198	Hemagglutinin protein, truncation	New identification
Evasion		
PG0011	Glycosyl hydrolase, family 3	New identification
PG0032	Beta-mannosidase, putative	New identification
PG0043	Beta-hexosaminidase	New identification
PG0070	Acyl-(acyl-carrier-protein)-UDP- <i>N</i> -acetylglucosamine acyltransferase	New identification
PG0071	UDP-3- <i>O</i> -acyl-GlcNAc deacetylase	New identification
PG0072	UDP-3- <i>O</i> -3-hydroxy-myristoyl glucosamine <i>N</i> -acyltransferase	New identification
PG0076	<i>N</i> -acetylmuramoyl-L-alanine amidase, family 4	New identification
PG0106	Glycosyl transferase, group 4 family protein	New identification
PG0108	UDP- <i>N</i> -acetyl-D-mannosaminuronic acid dehydrogenase	New identification
PG0110	Glycosyl transferase, group 1 family protein	New identification
PG0111	Capsular polysaccharide biosynthesis gene, putative	New identification
PG0112	Conserved hypothetical protein, authentic frameshift	New identification
PG0113	Conserved domain protein	New identification
PG0115	Hexapeptide transferase family protein	New identification
PG0117	Polysaccharide transport protein, putative	New identification
PG0118	Glycosyl transferase, group 2 family protein	New identification
PG0119	Glycosyl transferase, WecB/TagA/CpsF family	New identification
PG0120	UDP- <i>N</i> -acetylglucosamine 2-epimerase	New identification
PG0129	Mannosyltransferase	New identification
PG0139	Membrane-bound lytic murein transglycosylase D, putative	New identification
PG0264	Glycosyl transferase, group 2 family protein	New identification
PG0294	Glycosyl transferase, group 2 family protein	New identification
PG0311	Glycosyl transferase, group 2 family protein	New identification
PG0334	Glycosyl transferase, group 2 family protein	New identification
PG0352	Sialidase, putative	New identification
PG0369	Phosphopantetheine adenyltransferase	New identification
PG0435	Capsular polysaccharide biosynthesis protein, putative	New identification
PG0436	Capsular polysaccharide transport protein, putative	New identification
PG0437	Polysaccharide export protein, BexD/CtrA/VexA family	New identification
PG0638	Tetraacyldisaccharide 4'-kinase	New identification
PG0750	Glycosyl transferase, group 2 family protein	New identification
PG0902	Alpha-1,2-mannosidase family protein	New identification
PG0920	Glycosyl transferase, group 2 family protein	New identification
PG0973	Alpha-1,2-mannosidase family protein	New identification
PG1048	<i>N</i> -acetylmuramoyl-L-alanine amidase, family 3	New identification
PG1058	Outer membrane protein	New identification
PG1135	Bacterial sugar transferase	New identification
PG1140	Glycosyl transferase, group 2 family protein	New identification
PG1141	Glycosyl transferase, group 1 family protein	New identification
PG1143	Sugar dehydrogenase, UDP-glucose/GDP-mannose dehydrogenase family	New identification
PG1149	Glycosyl transferase, group 1 family protein	New identification
PG1155	ADP-heptose-LPS heptosyltransferase, putative	New identification
PG1184	Alginate <i>O</i> -acetyltransferase, putative	New identification
PG1277	UDP-glucose 6-dehydrogenase, putative	New identification
PG1288	GDP-mannose 4,6-dehydratase	New identification
PG1289	GDP-fucose synthetase	New identification
PG1345	Glycosyl transferase, group 1 family protein	New identification
PG1346	Glycosyl transferase, group 1 family protein	New identification
PG1560	dTDP-glucose 4,6-dehydratase	New identification
PG1561	dTDP-4-dehydrorhamnose reductase	New identification
PG1562	dTDP-4-dehydrorhamnose 3,5-epimerase	New identification
PG1563	Glucose-1-phosphate thymidyltransferase	New identification
PG1565	3-Deoxy-D-manno-octulosonic-acid transferase, putative	New identification
PG1682	Glycosyl transferase, group 1 family protein	New identification
PG1711	Alpha-1,2-mannosidase family protein	New identification
PG1712	Alpha-1,2-mannosidase family protein	New identification
PG1743	2-Dehydro-3-deoxyphosphooctonate aldolase	New identification
PG1750	Alpha-1,3/4-fucosidase, putative	New identification
PG1783	Glycosyl transferase, group 2 family protein	New identification
PG1815	3-Deoxy-D-manno-octulosonate cytidyltransferase	New identification
PG1880	Glycosyl transferase, group 2 family protein	New identification
PG1884	Alpha-L-fucosidase precursor, putative	New identification
PG1964	Bacterial sugar transferase	New identification
PG2038	<i>N</i> -acetylmuramoyl-L-alanine amidase, putative	New identification

Continued on following page

TABLE 4—Continued

Function and virulence agent	Description (reference)	Identification
PG2162	Lipid A disaccharide synthase	New identification
PG2215	Mannose-1-phosphate guanylyltransferase	New identification
PG2223	Glycosyl transferase, group 2 family protein	New identification
Invasion		
PG0151	Signal recognition particle-docking protein FtsY	New identification
PG0293	Secretion activator protein, putative	New identification
PG0485	Preprotein translocase, YajC subunit	New identification
PG0514	Preprotein translocase, SecA subunit	New identification
PG1115	Signal recognition particle protein	New identification
PG1598	Lipoprotein signal peptidase, putative	New identification
PG1762	Protein-export membrane protein SecD/protein-export membrane protein SecF	New identification
PG1918	Preprotein translocase, SecY subunit	New identification
PG2000	Signal peptidase-related protein	New identification
PG2001	Signal peptidase I	New identification
Stress response		
PG0090	Dps family protein	New identification
PG0245	Universal stress protein family	New identification
PG0618	Alkyl hydroperoxide reductase, C subunit	New identification
PG0619	Alkyl hydroperoxide reductase, F subunit	New identification
PG0876	Thiophene and furan oxidation protein ThdF	New identification
PG1330	Large conductance mechanosensitive channel protein	New identification
PG1545	Superoxide dismutase, Fe-Mn	New identification
PG1582	BatA protein	New identification
PG1583	BatB protein	New identification
PG1584	BatC protein	New identification
PG1585	BatD protein	New identification
PG1586	BatE protein	New identification
PG1648	RelA/SpoT family protein	New identification
PG1729	Thiol peroxidase	New identification
PG1808	Guanosine-3',5'-bis(diphosphate) 3'-pyrophosphohydrolase	New identification
PG2163	Stationary-phase survival protein SurE	New identification
Antibiotic resistance		
PG0135	Dimethyladenosine transferase	New identification
PG0540	AcrB/AcrD/AcrF family protein	New identification
PG0604	Penicillin tolerance protein LytB	New identification
PG1407	Nitroimidazole resistance protein, putative	New identification
PG1538	Undecaprenol kinase, putative	New identification
Virulence and acquisition of peptides		
PG1754	Conserved domain protein	New identification
PG2115	Protease PrtT, degenerate	New identification
PG0010	ATP-dependent Clp protease, ATP-binding subunit ClpC	New identification
PG0088	Peptidase, M16 family	New identification
PG0137	Aminoacyl-histidine dipeptidase	New identification
PG0159	Endopeptidase PepO	New identification
PG0196	Peptidase, M16 family	Previously sequenced
PG0232	Zinc carboxypeptidase, putative	New identification
PG0235	Carboxyl-terminal protease	New identification
PG0317	Peptidase, M49 family	New identification
PG0383	Membrane-associated zinc metalloprotease, putative	New identification
PG0417	ATP-dependent Clp protease, ATP-binding subunit ClpX	New identification
PG0418	ATP-dependent Clp protease, proteolytic subunit	New identification
PG0445	Peptidase T	New identification
PG0503	Dipeptidyl aminopeptidase IV(31)	Previously known
PG0506	Arginine-specific cysteine proteinase(63)	Previously known
PG0537	Aminoacyl-histidine dipeptidase	New identification
PG0553	Extracellular protease, putative	New identification
PG0561	Peptidase, M20/M25/M40 family	New identification
PG0593	HtrA protein	New identification
PG0620	ATP-dependent protease La	New identification
PG0639	Signal peptide peptidase SppA, 67K type	New identification
PG0724	Prolyl oligopeptidase family protein	New identification
PG0753	Protease	Previously sequenced
PG0758	Peptidyl-dipeptidase Dcp	New identification
PG0889	Peptidase, M24 family protein	New identification
PG0956	Peptidase, M23/M37 family, putative	New identification
PG1004	Prolyl oligopeptidase family protein	New identification
PG1055	Thiol protease (7)	Previously known
PG1060	Carboxyl-terminal protease	New identification
PG1118	ClpB protein	New identification
PG1210	Peptidase, M24 family protein	New identification
PG1361	Dipeptidyl aminopeptidase IV, putative	New identification
PG1605	Aminopeptidase C	New identification
PG1654	D-alanyl-D-alanine dipeptidase	New identification
PG1701	trp-G type glutamine amidotransferase/dipeptidase	New identification
PG1724	O-sialoglycoprotein endopeptidase	New identification
PG1788	Cysteine peptidase, putative	New identification
PG1789	Peptidyl-dipeptidase Dcp	New identification
PG1855	Carboxyl-terminal protease	New identification
PG2192	Peptidase, M23/M37 family	New identification

^a The newly identified virulence agents were selected based on the presence of signal peptide sequences or transmembrane domains or by homology to previously described virulence agents in *P. gingivalis*.

of this periodontal pathogen and will enable the development of improved diagnostics and therapeutics.

ACKNOWLEDGMENTS

We thank Michael Heaney, Michael Holmes, Vadim Sapiro, and Emmanuel Mongodin for informatics support at TIGR. We thank J. F. Tomb for assistance in the early stages of the project.

This work was supported by the National Institute of Dental and Craniofacial Research at the National Institutes of Health (R01DE-13914).

REFERENCES

- Ajdic, D., W. M. McShan, R. E. McLaughlin, G. Savic, J. Chang, M. B. Carson, C. Primeaux, R. Tian, S. Kenton, H. Jia, S. Lin, Y. Qian, S. Li, H. Zhu, F. Najjar, H. Lai, J. White, B. A. Roe, and J. J. Ferretti. 2002. Genome sequence of *Streptococcus mutans* UA159, a cariogenic dental pathogen. *Proc. Natl. Acad. Sci. USA* **99**:14434–14439.
- Albandar, J. M., J. A. Brunelle, and A. Kingman. 1999. Destructive periodontal disease in adults 30 years of age and older in the United States, 1988–1994. *J. Periodontol.* **70**:13–29.
- Allison, H. E., and J. D. Hillman. 1997. Cloning and characterization of a *Prevotella melaninogenica* hemolysin. *Infect. Immun.* **65**:2765–2771.
- Bahloul, A., F. Boubrik, and J. Rouviere-Yaniv. 2001. Roles of *Escherichia coli* histone-like protein HU in DNA replication: HU-beta suppresses the thermosensitivity of dnaA46ts. *Biochimie* **83**:219–229.
- Barker, H. A. 1981. Amino acid degradation by anaerobic bacteria. *Annu. Rev. Biochem.* **50**:23–40.
- Boone, D. R., and R. W. Castenholz (ed.). 2001. *Bergey's manual of systematic bacteriology*, 2nd ed., vol. 1. Springer-Verlag, New York, N.Y.
- Bourgeau, G., H. Lapointe, P. Peloquin, and D. Mayrand. 1992. Cloning, expression, and sequencing of a protease gene (*mpr*) from *Porphyromonas gingivalis* W83 in *Escherichia coli*. *Infect. Immun.* **60**:3186–3192.
- Brown, M. H., I. T. Paulsen, and R. A. Skurray. 1999. The multidrug efflux protein NorM is a prototype of a new family of transporters. *Mol. Microbiol.* **31**:394–395.
- Byrne, G. L., S. I. Skarlotos, C. Grunfeld, M. V. Kalayoglu, P. Libby, P. Saikku, J. T. Summersgill, and P. Wyrick. 2000. Collaborative multidisciplinary workshop report: interface of lipid metabolism, atherosclerosis, and *Chlamydia* infection. *J. Infect. Dis.* **181**(Suppl. 3):S490–S491.
- Chiu, B. 1999. Multiple infections in carotid atherosclerotic plaques. *Am. Heart J.* **138**:S534–S536.
- Churchward, G. 2002. Conjugative transposons and related elements, p. 177–191. *In* N. Craig, R. Craigie, M. Gellert, and A. Lambowitz (ed.), *Mobile DNA II*. ASM Press, Washington, D.C.
- Dashper, S. G., L. Brownfield, N. Slakeski, P. S. Zilm, A. H. Rogers, and E. C. Reynolds. 2001. Sodium ion-driven serine/threonine transport in *Porphyromonas gingivalis*. *J. Bacteriol.* **183**:4142–4148.
- Delcher, A. L., D. Harmon, S. Kasif, O. White, and S. L. Salzberg. 1999. Improved microbial gene identification with GLIMMER. *Nucleic Acids Res.* **27**:4636–4641.
- Endo, J., M. Otsuka, E. Ohara, M. Sato, and R. Nakamura. 1989. Cleavage action of a trypsin-like protease from *Bacteroides gingivalis* 381 on reduced egg-white lysozyme. *Arch. Oral Biol.* **34**:911–916.
- Fletcher, J., K. Reddi, S. Poole, S. Nair, B. Henderson, P. Tabona, and M. Wilson. 1997. Interactions between periodontopathogenic bacteria and cytokines. *J. Periodontol. Res.* **32**:200–205.
- Fujiwara, T., S. Morishima, I. Takahashi, and S. Hamada. 1993. Molecular cloning and sequencing of the fimbriin gene of *Porphyromonas gingivalis* strains and characterization of recombinant proteins. *Biochem. Biophys. Res. Commun.* **197**:241–247.
- Gharbia, S. E., and H. N. Shah. 1991. Pathways of glutamate catabolism among *Fusobacterium* species. *J. Gen. Microbiol.* **137**(Pt. 5):1201–1206.
- Gottschalk, G. 1986. Bacterial fermentations, p. 208–282. *In* G. Gottschalk (ed.), *Bacterial metabolism*, 2nd ed. Springer-Verlag, New York, N.Y.
- Goulbourne, P. A., and R. P. Ellen. 1991. Evidence that *Porphyromonas (Bacteroides) gingivalis* fimbriae function in adhesion to *Actinomyces viscosus*. *J. Bacteriol.* **173**:5266–5274.
- Grenier, D. 1992. Demonstration of a bimodal coaggregation reaction between *Porphyromonas gingivalis* and *Treponema denticola*. *Oral Microbiol. Immunol.* **7**:280–284.
- Han, N., J. Whitlock, and A. Progulsk-Fox. 1996. The hemagglutinin gene A (*hagA*) of *Porphyromonas gingivalis* 381 contains four large, contiguous, direct repeats. *Infect. Immun.* **64**:4000–4007.
- Herzberg, M. C., and M. W. Meyer. 1998. Dental plaque, platelets, and cardiovascular diseases. *Ann. Periodontol.* **3**:151–160.
- Hoffman, P. S., A. Goodwin, J. Johnsen, K. Magee, and S. J. Veldhuyzen van Zanten. 1996. Metabolic activities of metronidazole-sensitive and -resistant strains of *Helicobacter pylori*: repression of pyruvate oxidoreductase and expression of isocitrate lyase activity correlate with resistance. *J. Bacteriol.* **178**:4822–4829.
- Hueck, C. J. 1998. Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiol. Mol. Biol. Rev.* **62**:379–433.
- Hughes, N. J., C. L. Clayton, P. A. Chalk, and D. J. Kelly. 1998. *Helicobacter pylori* porCDAB and oorDABC genes encode distinct pyruvate:flavodoxin and 2-oxoglutarate:acceptor oxidoreductases which mediate electron transport to NADP. *J. Bacteriol.* **180**:1119–1128.
- Jansen, R., J. D. Embden, W. Gaastra, and L. M. Schouls. 2002. Identification of genes that are associated with DNA repeats in prokaryotes. *Mol. Microbiol.* **43**:1565–1575.
- Kaczmarek, F. S., and A. L. Coykendall. 1980. Production of phenylacetic acid by strains of *Bacteroides asaccharolyticus* and *Bacteroides gingivalis* (sp. nov.). *J. Clin. Microbiol.* **12**:288–290.
- Kapatral, V., I. Anderson, N. Ivanova, G. Reznik, T. Los, A. Lykidis, A. Bhattacharyya, A. Bartman, W. Gardner, G. Grechkin, L. Zhu, O. Vasieva, L. Chu, Y. Kogan, O. Chaga, E. Goltsman, A. Bernal, N. Larsen, M. D'Souza, T. Walunas, G. Pusch, R. Haselkorn, M. Fonstein, N. Kyrpides, and R. Overbeek. 2002. Genome sequence and analysis of the oral bacterium *Fusobacterium nucleatum* strain ATCC 25586. *J. Bacteriol.* **184**:2005–2018.
- Katz, J., V. Sambandam, J. H. Wu, S. M. Michalek, and D. F. Balkovetz. 2000. Characterization of *Porphyromonas gingivalis*-induced degradation of epithelial cell junctional complexes. *Infect. Immun.* **68**:1441–1449.
- Kinder, S. A., and S. C. Holt. 1989. Characterization of coaggregation between *Bacteroides gingivalis* T22 and *Fusobacterium nucleatum* T18. *Infect. Immun.* **57**:3425–3433.
- Kiyama, M., M. Hayakawa, T. Shiroza, S. Nakamura, A. Takeuchi, Y. Masamoto, and Y. Abiko. 1998. Sequence analysis of the *Porphyromonas gingivalis* dipeptidyl peptidase IV gene. *Biochim. Biophys. Acta* **1396**:39–46.
- Kontani, M., H. Ono, H. Shibata, Y. Okamura, T. Tanaka, T. Fujiwara, S. Kimura, and S. Hamada. 1996. Cysteine protease of *Porphyromonas gingivalis* 381 enhances binding of fimbriae to cultured human fibroblasts and matrix proteins. *Infect. Immun.* **64**:756–762.
- Krogh, A., B. Larsson, G. von Heijne, and E. L. Sonnhammer. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J. Mol. Biol.* **305**:567–580.
- Laine, M. L., and A. J. van Winkelhoff. 1998. Virulence of six capsular serotypes of *Porphyromonas gingivalis* in a mouse model. *Oral Microbiol. Immunol.* **13**:322–325.
- Lamont, R. J., S. Gil, D. R. Demuth, D. Malamud, and B. Rosan. 1994. Molecules of *Streptococcus gordonii* that bind to *Porphyromonas gingivalis*. *Microbiology* **140**(Pt. 4):867–872.
- Lamont, R. J., G. W. Hsiao, and S. Gil. 1994. Identification of a molecule of *Porphyromonas gingivalis* that binds to *Streptococcus gordonii*. *Microb. Pathog.* **17**:355–360.
- Lamont, R. J., and H. F. Jenkinson. 2000. Subgingival colonization by *Porphyromonas gingivalis*. *Oral Microbiol. Immunol.* **15**:341–349.
- Lantz, M. S., R. D. Allen, L. W. Duck, J. L. Blume, L. M. Switalski, and M. Hook. 1991. Identification of *Porphyromonas gingivalis* components that mediate its interactions with fibronectin. *J. Bacteriol.* **173**:4263–4270.
- Lepine, G., and A. Progulsk-Fox. 1996. Duplication and differential expression of hemagglutinin genes in *Porphyromonas gingivalis*. *Oral Microbiol. Immunol.* **11**:65–78.
- Lewis, J. P., and F. L. Macrina. 1998. *IS195*, an insertion sequence-like element associated with protease genes in *Porphyromonas gingivalis*. *Infect. Immun.* **66**:3035–3042.
- Loos, B. G., D. W. Dyer, T. S. Whittam, and R. K. Selander. 1993. Genetic structure of populations of *Porphyromonas gingivalis* associated with periodontitis and other oral infections. *Infect. Immun.* **61**:204–212.
- Madianos, P. N., P. N. Papapanou, and J. Sandros. 1997. *Porphyromonas gingivalis* infection of oral epithelium inhibits neutrophil transepithelial migration. *Infect. Immun.* **65**:3983–3990.
- Maghnoji, A., T. F. de Sousa Cabral, V. Stalon, and C. Vander Wauven. 1998. The arcABDC gene cluster, encoding the arginine deiminase pathway of *Bacillus licheniformis*, and its activation by the arginine repressor *argR*. *J. Bacteriol.* **180**:6468–6475.
- Maley, J., and I. S. Roberts. 1994. Characterisation of *IS1126* from *Porphyromonas gingivalis* W83: a new member of the IS4 family of insertion sequence elements. *FEMS Microbiol. Lett.* **123**:219–224.
- Masuda, K., H. Hayashi, M. Yoshioka, and R. Nakamura. 1995. Presence of an arginine carboxypeptidase in culture supernatants of *Porphyromonas gingivalis*. *J. Dent. Res.* **74**:974.
- Menard, C., and C. Mouton. 1995. Clonal diversity of the taxon *Porphyromonas gingivalis* assessed by random amplified polymorphic DNA fingerprinting. *Infect. Immun.* **63**:2522–2531.
- Meyer, D. H., and P. M. Fives-Taylor. 1998. Oral pathogens: from dental plaque to cardiac disease. *Curr. Opin. Microbiol.* **1**:88–95.
- Nakayama, K., T. Kadowaki, K. Okamoto, and K. Yamamoto. 1995. Construction and characterization of arginine-specific cysteine proteinase (Arg-gingipain)-deficient mutants of *Porphyromonas gingivalis*: evidence for significant contribution of Arg-gingipain to virulence. *J. Biol. Chem.* **270**:23619–23626.
- Nelson, K. E., R. A. Clayton, S. R. Gill, M. L. Gwinn, R. J. Dodson, D. H. Haft, E. K. Hickey, J. D. Peterson, W. C. Nelson, K. A. Ketchum, L. Mc-

- Donald, T. R. Utterback, J. A. Malek, K. D. Linher, M. M. Garrett, A. M. Stewart, M. D. Cotton, M. S. Pratt, C. A. Phillips, D. Richardson, J. Heidelberg, G. G. Sutton, R. D. Fleischmann, J. A. Eisen, C. M. Fraser, et al. 1999. Evidence for lateral gene transfer between *Archaea* and bacteria from genome sequence of *Thermotoga maritima*. *Nature* **399**:323–329.
50. Niederman, R., J. Zhang, and S. Kashket. 1997. Short-chain carboxylic-acid-stimulated, PMN-mediated gingival inflammation. *Crit. Rev. Oral Biol. Med.* **8**:269–290.
51. Nierman, W. C., T. V. Feldblyum, M. T. Laub, I. T. Paulsen, K. E. Nelson, J. A. Eisen, J. F. Heidelberg, M. R. Alley, N. Ohta, J. R. Maddock, I. Potocka, W. C. Nelson, A. Newton, C. Stephens, N. D. Phadke, B. Ely, R. T. DeBoy, R. J. Dodson, A. S. Durkin, M. L. Gwinn, D. H. Haft, J. F. Kolonay, J. Smit, M. B. Craven, H. Khouri, J. Shetty, K. Berry, T. Utterback, K. Tran, A. Wolf, J. Vamathevan, M. Ermolaeva, O. White, S. L. Salzberg, J. C. Venter, L. Shapiro, and C. M. Fraser. 2001. Complete genome sequence of *Caulobacter crescentus*. *Proc. Natl. Acad. Sci. USA* **98**:4136–4141.
52. Offenbacher, S., P. N. Madianos, C. M. Champagne, J. H. Southerland, D. W. Paquette, R. C. Williams, G. Slade, and J. D. Beck. 1999. Periodontitis-atherosclerosis syndrome: an expanded model of pathogenesis. *J. Periodontol. Res.* **34**:346–352.
53. Pike, R. N., J. Potempa, W. McGraw, T. H. Coetzer, and J. Travis. 1996. Characterization of the binding activities of proteinase-adhesin complexes from *Porphyromonas gingivalis*. *J. Bacteriol.* **178**:2876–2882.
54. Riley, M. 1993. Functions of the gene products of *Escherichia coli*. *Microbiol. Rev.* **57**:862–952.
55. Rocha, E. R., and C. J. Smith. 1999. Role of the alkyl hydroperoxide reductase (*ahpCF*) gene in oxidative stress defense of the obligate anaerobe *Bacteroides fragilis*. *J. Bacteriol.* **181**:5701–5710.
56. Ruzicka, F. J., K. W. Lieder, and P. A. Frey. 2000. Lysine 2,3-aminomutase from *Clostridium subterminale* SB4: mass spectral characterization of cyanogen bromide-treated peptides and cloning, sequencing, and expression of the gene *kamA* in *Escherichia coli*. *J. Bacteriol.* **182**:469–476.
57. Salzberg, S. L., A. L. Delcher, S. Kasif, and O. White. 1998. Microbial gene identification using interpolated Markov models. *Nucleic Acids Res.* **26**:544–548.
58. Scannapieco, F. A., and R. J. Genco. 1999. Association of periodontal infections with atherosclerotic and pulmonary diseases. *J. Periodontol. Res.* **34**:340–345.
59. Scragg, M. A., S. J. Cannon, M. Rangarajan, D. M. Williams, and M. A. Curtis. 1999. Targeted disruption of fibronectin-integrin interactions in human gingival fibroblasts by the RI protease of *Porphyromonas gingivalis* W50. *Infect. Immun.* **67**:1837–1843.
60. Seddon, S. V., H. N. Shah, J. M. Hardie, and J. P. Robinson. 1988. Chemically defined and minimal media for *Bacteroides gingivalis*. *Curr. Microbiol.* **17**:147–149.
61. Shah, H. N., and R. A. D. Williams. 1987. Utilization of glucose and amino acids by *Bacteroides intermedius* and *Bacteroides gingivalis*. *Curr. Microbiol.* **15**:241–246.
62. Shelburne, C. E., N. P. VanPoperin, R. M. Gleason, A. Prabhu, and D. E. Lopatin. 2000. Assay of bacterial DNA in atherosclerotic lesions by quantitative PCR. *J. Dent. Res.* **79**:226.
63. Slakeski, N., P. S. Bhogal, N. M. O'Brien-Simpson, and E. C. Reynolds. 1998. Characterization of a second cell-associated Arg-specific cysteine proteinase of *Porphyromonas gingivalis* and identification of an adhesin-binding motif involved in association of the prtR and prtK proteinases and adhesins into large complexes. *Microbiology* **144**(Pt. 6):1583–1592.
64. Slakeski, N., S. M. Cleal, P. S. Bhogal, and E. C. Reynolds. 1999. Characterization of a *Porphyromonas gingivalis* gene *prtK* that encodes a lysine-specific cysteine proteinase and three sequence-related adhesins. *Oral Microbiol. Immunol.* **14**:92–97.
65. Takahashi, N., T. Sato, and T. Yamada. 2000. Metabolic pathways for cytotoxic end product formation from glutamate- and aspartate-containing peptides by *Porphyromonas gingivalis*. *J. Bacteriol.* **182**:4704–4710.
66. Tang, Y. P., M. M. Dallas, and M. H. Malamy. 1999. Characterization of the *BatI* (*Bacteroides* aerotolerance) operon in *Bacteroides fragilis*: isolation of a *B. fragilis* mutant with reduced aerotolerance and impaired growth in in vivo model systems. *Mol. Microbiol.* **32**:139–149.
67. Tettelin, H., K. E. Nelson, I. T. Paulsen, J. A. Eisen, T. D. Read, S. Peterson, J. Heidelberg, R. T. DeBoy, D. H. Haft, R. J. Dodson, A. S. Durkin, M. Gwinn, J. F. Kolonay, W. C. Nelson, J. D. Peterson, L. A. Umayam, O. White, S. L. Salzberg, M. R. Lewis, D. Radune, E. Holtzapple, H. Khouri, A. M. Wolf, T. R. Utterback, C. L. Hansen, L. A. McDonald, T. V. Feldblyum, S. Angiuoli, T. Dickinson, E. K. Hickey, I. E. Holt, B. J. Loftus, F. Yang, H. O. Smith, J. C. Venter, B. A. Dougherty, D. A. Morrison, S. K. Hollingshead, and C. M. Fraser. 2001. Complete genome sequence of a virulent isolate of *Streptococcus pneumoniae*. *Science* **293**:498–506.
68. Tribble, G. D., A. C. Parker, and C. J. Smith. 1999. Genetic structure and transcriptional analysis of a mobilizable, antibiotic resistance transposon from *Bacteroides*. *Plasmid* **42**:1–12.
69. Wang, C. Y., V. C. Bond, and C. A. Genco. 1997. Identification of a second endogenous *Porphyromonas gingivalis* insertion element. *J. Bacteriol.* **179**:3808–3812.
70. White, O. Bacterial genome annotation at The Institute for Genomic Research. In C. M. Fraser, T. D. Read, and K. E. Nelson (ed.), *Microbial genomics*, in press. Humana Press, Inc., New York, N.Y.
71. Wyss, C. 1992. Growth of *Porphyromonas gingivalis*, *Treponema denticola*, *T. pectinovorum*, *T. socranskii*, and *T. vincentii* in a chemically defined medium. *J. Clin. Microbiol.* **30**:2225–2229.
72. Yao, E. S., R. J. Lamont, S. P. Leu, and A. Weinberg. 1996. Interbacterial binding among strains of pathogenic and commensal oral bacterial species. *Oral Microbiol. Immunol.* **11**:35–41.
73. Yun, P. L., A. A. DeCarlo, and N. Hunter. 1999. Modulation of major histocompatibility complex protein expression by human gamma interferon mediated by cysteine proteinase-adhesin polyproteins of *Porphyromonas gingivalis*. *Infect. Immun.* **67**:2986–2995.
74. Zhang, J., H. Dong, S. Kashket, and M. J. Duncan. 1999. IL-8 degradation by *Porphyromonas gingivalis* proteases. *Microb. Pathog.* **26**:275–280.