

RESEARCH PAPER

Endothelium-dependent metabolism by endocannabinoid hydrolases and cyclooxygenases limits vasorelaxation to anandamide and 2-arachidonoylglycerol

W-SV Ho and MD Randall

School of Biomedical Sciences, University of Nottingham Medical School, Queen's Medical Centre, Nottingham, UK

Background and purpose: The endocannabinoids, *N*-arachidonylethanolamide (anandamide) and 2-arachidonoylglycerol (2-AG) are rapidly degraded by fatty acid amide hydrolase (FAAH) and monoacylglycerol lipase (MGL). Whilst these lipid mediators are known to modulate vascular tone, the extent to which they are inactivated via local metabolism in the vasculature remains unclear.

Experimental approach: In rat isolated small mesenteric arteries, the regulatory role of FAAH, MGL and cyclooxygenase (COX) in relaxant responses to anandamide and 2-AG was evaluated by using inhibitors of these enzymes. Relaxations to non-hydrolysable analogues of endocannabinoids and arachidonic acid were also examined.

Key results: Relaxation to anandamide but not 2-AG was potentiated by the selective FAAH inhibitor, URB597 (1 μ M). In contrast, MAFP (10 μ M; an inhibitor of FAAH and MGL) enhanced responses to both anandamide and 2-AG. Inhibition of COX-1 by indomethacin (10 μ M) potentiated relaxations to 2-AG, whereas inhibition of COX-2 by nimesulide (10 μ M) potentiated anandamide-induced relaxation. With the exception of MAFP, effects of FAAH and COX inhibitors were dependent on the endothelium. Relaxation to methanandamide and noladin ether, the non-hydrolysable analogues of anandamide and 2-AG respectively, were insensitive to the enzyme inhibitors.

Conclusion and implications: This study shows that local activity of FAAH, MGL and COX, which is present largely in the endothelium, limits the vasodilator action of endocannabinoids in rat small mesenteric arteries. Despite the differential roles played by these enzymes on relaxation to anandamide versus 2-AG, our results suggest that inhibitors of these enzymes enhance the vascular impact of endocannabinoids.

British Journal of Pharmacology (2007) **150**, 641–651. doi:10.1038/sj.bjp.0707141; published online 22 January 2007

Keywords: anandamide; 2-arachidonoylglycerol; fatty acid amide hydrolase; monoacylglycerol lipase; cyclooxygenase; rat mesenteric artery; endothelium

Abbreviations: MAFP, methyl arachidonyl fluorophosphonate; URB597, 3'-carbamoyl-biphenyl-3-yl-cyclohexylcarbamate

Introduction

Endocannabinoids, in particular the prototypical endocannabinoid, *N*-arachidonylethanolamide (anandamide), have been shown to induce vasodilatation, modulate regional blood flow and arterial blood pressure and reduce heart rate (Randall *et al.*, 1996; Lake *et al.*, 1997; Gardiner *et al.*, 2002; Batkai *et al.*, 2004). Although the precise mechanisms of action may depend on the vascular regions, species and state of anaesthesia, anandamide and its metabolically stable

analogue, methanandamide, often act via activation of the cannabinoid (CB)₁ receptor or the transient receptor potential vanilloid 1 (TRPV1) receptor, or both (Lake *et al.*, 1997; Gebremedhin *et al.*, 1999; Zygmunt *et al.*, 1999; Ralevic *et al.*, 2000; Gardiner *et al.*, 2002; Ho and Hiley, 2003). In the rat mesenteric artery, relaxations to anandamide and methanandamide are largely mediated by TRPV1 receptors on perivascular sensory nerves (Zygmunt *et al.*, 1999; Ralevic *et al.*, 2000; Ho and Hiley, 2003). In addition, activation of CB₁ receptors, a novel endothelial cannabinoid receptor coupled to Ca²⁺-activated K⁺ channels and/or direct inhibition of voltage-dependent calcium entry might also play a role (Ho and Hiley, 2003; Offertaler *et al.*, 2003; O'Sullivan *et al.*, 2004). However, in comparison with anandamide, little is known about the vasorelaxant effects

Correspondence: Dr W-S Vanessa Ho, School of Biomedical Sciences, University of Nottingham Medical School, Queen's Medical Centre, Nottingham NG7 2UH, UK.

E-mail: vanessa.ho@nottingham.ac.uk

Received 2 November 2006; revised 29 November 2006; accepted 30 November 2006; published online 22 January 2007

of the other major endocannabinoid, 2-arachidonoylglycerol (2-AG) despite observations that tissue content of 2-AG is often more than 200-fold higher than that of anandamide (e.g., hippocampus: Makara *et al.*, 2005; heart: Pacher *et al.*, 2005; cerebral artery: Rademacher *et al.*, 2005). Several studies have shown that 2-AG induces hypotension (Mechoulam *et al.*, 1998; Jarai *et al.*, 2000) and vasorelaxation (Kagota *et al.*, 2001; Ho and Hiley, 2004; Gauthier *et al.*, 2005). By activating the CB₁ receptor, 2-AG and its non-hydrolysable analogue noladin ether could reduce blood pressure (Jarai *et al.*, 2000). In mesenteric arteries, relaxant responses to 2-AG, which are mimicked by noladin ether, might involve CB₁ receptors and K⁺ channels (Kagota *et al.*, 2001; Ho and Hiley, 2004). However, there have been conflicting reports regarding its action as a vasodilator (Wagner *et al.*, 1999; Zygmunt *et al.*, 1999; Kagota *et al.*, 2001; Ho and Hiley, 2004), perhaps owing to its greater susceptibility to degradation than anandamide (Jarai *et al.*, 2000; Gauthier *et al.*, 2005).

It is now recognized that endocannabinoids are taken up by various cell types, followed by rapid metabolism by intracellular enzymes. Catabolism of both anandamide and 2-AG occurs via hydrolysis to arachidonic acid, and ethanolamine and glycerol, respectively. Anandamide hydrolysis is mediated primarily by fatty acid amide hydrolase (FAAH; Cravatt *et al.*, 1996, 2001). Activity of this serine hydrolase has been found, for example, in blood vessels, heart, liver and brain (Deutsch and Chin, 1993; Desarnaud *et al.*, 1995; Pratt *et al.*, 1998). Development of effective inhibitors has enabled the characterization of FAAH and its role in endocannabinoid signalling. For instance, the selective and potent FAAH inhibitor 3'-carbamoyl-biphenyl-3-yl-cyclohexylcarbamate (URB597) has been shown to increase tissue content of anandamide, but not 2-AG, in the heart and brain (Kathuria *et al.*, 2003; Batkai *et al.*, 2004). Application of URB597 alone has also been shown to lower blood pressure in anaesthetized, hypertensive rats (Batkai *et al.*, 2004). 2-AG might also serve as a substrate for FAAH (Goparaju *et al.*, 1998); indeed systemic and local treatment with FAAH inhibitors has been shown to elevate 2-AG content (Bifulco *et al.*, 2004; de Lago *et al.*, 2005; Maione *et al.*, 2006). Moreover, a distinct serine hydrolase called the monoacylglycerol lipase (MGL) also plays an important role in the hydrolysis of 2-AG, especially in the brain (Karlsson *et al.*, 2001; Cravatt and Lichtman, 2002; Dinh *et al.*, 2002; Kathuria *et al.*, 2003). As for FAAH, MGL appears to have a relatively ubiquitous tissue distribution; for example, its activity has been found in the heart, macrophages, adipose tissue and brain (Tornqvist and Belfrage, 1976; Di Marzo *et al.*, 1999; Dinh *et al.*, 2002). Since the identification of MGL, its sensitivity to some of the known serine hydrolase inhibitors has been investigated (see Ho and Hillard, 2005 for review). Notably, the compound methyl arachidonyl fluorophosphonate (MAFP), which is commonly used to inhibit FAAH, has also been found to be a potent inhibitor of MGL activity in cytosolic and membrane fractions of the brain (Goparaju *et al.*, 1999; Dinh *et al.*, 2002; Saario *et al.*, 2004). The crucial role of FAAH and MGL in the inactivation of anandamide and 2-AG suggests that inhibitors of these enzymes could be utilized experimentally and, perhaps also

clinically, to enhance endocannabinoid activity. However, it should be pointed out that other metabolic pathways might also play a role in the metabolism of endocannabinoids. In particular, cyclooxygenases (both COX-1 and COX-2 isoforms) could act on arachidonic acid subsequent to the hydrolysis of endocannabinoids (Grainger and Boachie-Ansah, 2001; Wahn *et al.*, 2005); both anandamide and 2-AG are potential substrates for COX-2 (Yu *et al.*, 1997; Kozak *et al.*, 2000). This raises the possibility that a combined application of endocannabinoid hydrolases and COX inhibitors could further increase the life-span of endocannabinoids.

Depending on the vascular regions and species, effects of endocannabinoids on local vascular control could be regulated or mediated or both, by metabolism. This is possibly in part due to differences in the expression of metabolizing enzymes. Where degradation is likely to regulate, rather than mediate, anandamide relaxations (Gebremedhin *et al.*, 1999; Zygmunt *et al.*, 1999; Ho and Hiley, 2003), the extent to which endocannabinoids are inactivated via local metabolism in the vasculature, remains unclear. Importantly, the potential role of FAAH- and MGL-mediated hydrolysis in the vascular wall has not been systematically studied and compared. In this study, we have investigated the effects of known inhibitors of FAAH and MGL on the vasorelaxant responses to anandamide and 2-AG in rat isolated small mesenteric arteries. In addition, the involvement of the two isoforms of COX was also examined.

Methods

Myograph studies

Male Wistar rats (200–300 g; Charles River UK Ltd, Kent, UK) were stunned by a blow to the back of their neck and killed by cervical dislocation. All animal care and use was in accordance with the UK Animal (Scientific Procedures) Act 1986. The third-order branches of the superior mesenteric artery were removed and cleaned of adherent tissue. Segments (2 mm in length) were mounted in a Mulvany-Halpern type wire myograph (Model 610M, Danish Myo Technology, Aarhus, Denmark) and maintained at 37°C in gassed (95% O₂/5% CO₂) Krebs-Henseleit solution of the following composition (mM): NaCl 118, KCl 4.7, MgSO₄ 1.2, KH₂PO₄ 1.2, NaHCO₃ 25, CaCl₂ 2, D-glucose 10, as described previously (Ho and Hiley, 2003). Vessels were equilibrated and set to a basal tension of 2–2.5 mN. The integrity of the endothelium was assessed by precontracting the vessel with 10 μM methoxamine (an α₁ adrenoceptor agonist), followed by relaxation with 10 μM carbachol (a muscarinic receptor agonist); vessels showing relaxations of >90% were designated as endothelium-intact. When endothelium was not required, it was removed by rubbing the intima with a human hair; carbachol-induced relaxation of less than 10% indicated successful removal.

Experimental protocols

After the test for endothelial integrity, vessels were left for 30 min and then precontracted with 10 μM methoxamine.

This was followed by construction of a cumulative concentration–relaxation curve to a cannabinoid or arachidonic acid. Each vessel was exposed only to one relaxant agent. In this study, most experiments were performed in matched vessels; effects of putative inhibitors or endothelial removal were compared with the control responses obtained in separate vessels of the same rat.

To investigate the role of metabolism in the relaxant responses to endocannabinoids, inhibitors of FAAH (URB597, arachidonyl trifluoromethyl ketone (ATFMK), MAFP), MGL (ATFMK, MAFP, 6-methyl-2-[(4-methylphenyl)amino]-4H-3,1-benzoxazin-one (URB754) and COX (indomethacin, flurbiprofen and nimesulide) were used. One or more of these enzyme inhibitors were added to the myograph bath 30 or 45 min before, and were kept present during construction of the concentration–response curve of an endocannabinoid or its metabolically stable analogue (methanandamide and noladin ether). In a separate set of experiments, effects of MAFP and COX inhibitors on the relaxant responses to arachidonic acid, a likely breakdown product of the endocannabinoids, were also examined.

Incubation of the mesenteric arteries with MAFP or ATFMK often reduced the contractions to methoxamine, possibly owing to inhibition of cytosolic phospholipase A₂ (LaBelle and Polyak, 1998). Therefore, an increased concentration (up to 30 μ M) of methoxamine was used to precontract vessels to obtain a similar level of tone to that evoked in the absence of MAFP and ATFMK. The tension generated in the test for endothelial integrity was 11.3 ± 0.5 mN, as compared with 10.1 ± 0.5 mN (40 vessels) in the presence of MAFP. Tension was 10.6 ± 0.8 mN in the endothelial test, as compared with 9.3 ± 0.8 mN in the presence of ATFMK (22 vessels).

Data and statistical analysis

All relaxation responses are expressed as percentage relaxation of the tone induced by methoxamine. Values are given as mean \pm s.e.m. and *n* represents the number of animals used. E_{\max} represents the maximum effect and pEC_{50} the negative logarithm of the concentration of relaxant giving half the maximal relaxation; these values were determined directly from individual log concentration–response curves. Statistical comparisons of concentration–response curves were made by two-way analysis of variance (Prism 4, GraphPad Software Inc.) of the whole data set. *P*-values of less than 0.05 were taken as statistically significant.

Drugs

Methoxamine and carbachol (Sigma Chemical Co., Poole, UK) were dissolved in deionized water. Anandamide (*N*-arachidonoyl ethanolamide) and *R*-(+)-methanandamide (Tocris Bioscience, Bristol, UK) were supplied in Tocrisolve 100 (1:4 soya:water emulsion) and diluted with deionized water. 2-AG, noladin ether (2-AG ether; Tocris) and arachidonic acid (Cayman Chemical, Ann Arbor, MI, USA) were supplied in 100% ethanol and diluted with deionized water. URB597 (Cayman Chemical), MAFP (Tocris), nimesulide and indomethacin and (\pm)-flurbiprofen (Sigma) were dissolved in 100% ethanol. ATFMK (Alexis Corporation, Nottingham,

UK) and URB754 (Cayman Chemical) were dissolved in 100% dimethyl sulphoxide.

Results

Relaxation to anandamide and 2-AG

Anandamide induced concentration-dependent relaxation of rat isolated mesenteric arteries ($pEC_{50} = 6.7 \pm 0.1$, $E_{\max} = 101 \pm 1\%$, $n = 5$). Removal of the endothelium caused a significant rightward shift in the concentration–response curve to anandamide ($pEC_{50} = 6.5 \pm 0.1$, $E_{\max} = 101 \pm 1\%$, $n = 5$; $P < 0.01$).

Another endocannabinoid, 2-AG also caused vasorelaxation in mesenteric arteries, albeit with lower potency than anandamide ($pEC_{50} = 5.5 \pm 0.1$, $E_{\max} = 75 \pm 10\%$, $n = 5$). Endothelial removal significantly reduced the potency and the maximal effect (at 10 μ M) of 2-AG-induced relaxations ($pEC_{50} = 5.4 \pm 0.1$, $E_{\max} = 41 \pm 11\%$, $n = 5$; $P < 0.001$).

Effects of FAAH and MGL inhibitors on relaxation to anandamide

The compound URB597 is an irreversible, selective inhibitor of FAAH and displays no activity at MGL (Kathuria *et al.*, 2003). It inhibits FAAH with IC_{50} values as low as 0.5 nM (Kathuria *et al.*, 2003) and 1 μ M URB597 has been shown to enhance the endogenous anandamide content in rat brain slices (Makara *et al.*, 2005). Here, in endothelium-intact vessels, the presence of URB597 (1 μ M) potentiated the relaxant responses to anandamide, resulting in a leftward displacement in the concentration–response curve (control, $pEC_{50} = 6.7 \pm 0.1$, $E_{\max} = 101 \pm 1\%$, $n = 4$; with URB597, $pEC_{50} = 7.2 \pm 0.1$, $E_{\max} = 98 \pm 4\%$, $n = 4$; $P < 0.01$; Figure 1a). However, URB597 had no significant effect on anandamide relaxations in endothelium-denuded vessels (with URB597, $pEC_{50} = 6.7 \pm 0.2$, $E_{\max} = 101 \pm 1\%$, $n = 5$; Figure 1b).

In another set of experiments, the effects of other FAAH inhibitors, namely MAFP and ATFMK which could also inhibit MGL (see Ho and Hillard, 2005 for review) were tested. In membrane preparations, the enzymes MAFP and ATFMK have been shown to inhibit FAAH at nanomolar concentrations (De Petrocellis *et al.*, 1997; Deutsch *et al.*, 1997). However, in view of the large variation in their potency at MGL (inhibition occurs at concentrations from low nM to μ M; Ho and Hillard, 2005), they were used at 10 μ M to ensure inhibition of both FAAH and MGL in isolated vessels. In endothelium-intact mesenteric arteries, the presence of MAFP (10 μ M) potentiated the relaxations to anandamide (control, $pEC_{50} = 6.7 \pm 0.2$, $E_{\max} = 100 \pm 2$, $n = 5$; with MAFP, $pEC_{50} = 7.2 \pm 0.1$, $E_{\max} = 100 \pm 1\%$, $n = 5$; $P < 0.01$). Similar results were also obtained with 10 μ M ATFMK (with endothelium, control, $pEC_{50} = 6.9 \pm 0.2$, $E_{\max} = 100 \pm 1\%$, $n = 5$; with ATFMK, $pEC_{50} = 7.6 \pm 0.1$, $E_{\max} = 99 \pm 1\%$, $n = 5$; $P < 0.001$). It was noted that MAFP, ATFMK or URB597 induced a similar leftward displacement of anandamide concentration–response curves, with about threefold increase in the potency of anandamide.

The presence of 1 μ M URB597 had no significant relaxant effect *per se* (data not shown). However, incubation of vessels with ATFMK and especially MAFP reduced methoxamine-




Figure 1 Effects of FAAH inhibitors on anandamide-induced relaxation in mesenteric arteries. Relaxation was elicited by anandamide (a, b) or methanandamide (c) alone, or in the presence of $1 \mu\text{M}$ URB597 in endothelium-intact or -denuded vessels. Values are shown as means and vertical lines represent s.e.m.; $n=4-7$. The results with URB597 for endothelium-intact mesenteric arteries were also seen with two other less selective FAAH inhibitors, MAFP ($10 \mu\text{M}$) and ATFMK ($10 \mu\text{M}$) (data not shown).

induced contractions (data not shown; see Methods). This is possibly owing to their additional inhibitory effect on cytosolic phospholipase A_2 and thereby reduces methoxamine-induced production of contractile mediators (LaBelle and Polyak, 1998), and/or an increase in the vascular content of vasodilator endocannabinoids.

The putative MGL inhibitor URB754 ($3 \mu\text{M}$; Makara *et al.*, 2005) had no effect on relaxations induced by anandamide (data not shown).

Effects of URB597 on relaxation to methanandamide

Methanandamide, a metabolically stable analogue of anandamide (Abadji *et al.*, 1994), induced endothelium-dependent relaxation in mesenteric arteries (with endothelium, $p\text{EC}_{50} = 6.5 \pm 0.2$, $E_{\text{max}} = 93 \pm 3\%$, $n = 6$; without endothelium, $p\text{EC}_{50} = 5.9 \pm 0.2$, $E_{\text{max}} = 91 \pm 6\%$, $n = 5$; $P < 0.001$). The presence of URB597 ($1 \mu\text{M}$) had no effect on relaxations to methanandamide (control, $p\text{EC}_{50} = 6.9 \pm 0.2$, $E_{\text{max}} = 96 \pm 3\%$, $n = 7$; with URB597, $p\text{EC}_{50} = 6.9 \pm 0.2$, $E_{\text{max}} = 94 \pm 4\%$, $n = 7$; Figure 1c).

Effects of FAAH and MGL inhibitors on relaxation to 2-AG

The selective FAAH inhibitor URB597 ($1 \mu\text{M}$) had no effect on 2-AG induced relaxation in endothelium-intact vessels (control, $p\text{EC}_{50} = 5.4 \pm 0.1$, $E_{\text{max}} = 89 \pm 4\%$, $n = 4$; with URB597, $p\text{EC}_{50} = 5.5 \pm 0.1$, $E_{\text{max}} = 98 \pm 1\%$, $n = 4$; Figure 2a). In contrast, the presence of MAFP ($10 \mu\text{M}$) enhanced 2-AG relaxations (with endothelium, control, $p\text{EC}_{50} = 5.6 \pm 0.1$, $E_{\text{max}} = 85 \pm 8\%$, $n = 5$; with MAFP, $p\text{EC}_{50} = 5.9 \pm 0.2$, $E_{\text{max}} = 100 \pm 1\%$, $n = 5$; $P < 0.001$; Figure 2b).

On the other hand, ATFMK appeared to modify 2-AG relaxations in a more complex fashion. In the presence of $10 \mu\text{M}$ ATFMK, relaxations tended to be potentiated at lower concentrations but inhibited at higher concentrations of 2-AG (with endothelium, control, $p\text{EC}_{50} = 5.6 \pm 0.1$, $E_{\text{max}} = 89 \pm 6\%$, $n = 7$; with ATFMK, $p\text{EC}_{50} = 6.1 \pm 0.2$, $E_{\text{max}} = 81 \pm 8\%$, $n = 7$; Figure 2c). Based on two-way analysis of variance of the concentration-response curves, there was a significant interaction ($P < 0.05$) between the ATFMK treatment and 2-AG concentrations. A similar tendency was observed in endothelium-denuded vessels (control, $p\text{EC}_{50} = 5.4 \pm 0.2$, $E_{\text{max}} = 62 \pm 11\%$, $n = 5$; with ATFMK, $p\text{EC}_{50} = 5.5 \pm 0.2$, $E_{\text{max}} = 48 \pm 11\%$, $n = 5$; Figure 2d).




Figure 2 Effects of FAAH inhibitors on 2-AG-induced relaxation in mesenteric arteries. Relaxation was elicited by 2-AG alone, or in the presence of 1 μM URB597 (a), 10 μM MAFP (b) or 10 μM ATFMK (c, d) in endothelium-intact or -denuded vessels. Values are shown as means and vertical lines represent s.e.m. $n=4-7$.

In contrast, URB754 (3 μM) applied either alone or in combination with URB597 (1 μM) had no effect on the 2-AG-induced relaxation (data not shown).

Effects of COX inhibitors on relaxation to anandamide

The COX inhibitor indomethacin (10 μM) had no significant effect on anandamide-induced relaxation, either in the presence (with indomethacin, $\text{pEC}_{50}=6.9\pm 0.1$, $E_{\text{max}}=98\pm 3\%$, $n=4$; Figure 3a) or absence of the endothelium (control, $\text{pEC}_{50}=6.5\pm 0.1$, $E_{\text{max}}=101\pm 1\%$, $n=5$; with indomethacin, $\text{pEC}_{50}=6.5\pm 0.2$, $E_{\text{max}}=100\pm 1\%$, $n=4$).

In endothelium-intact vessels, the selective COX-2 inhibitor nimesulide (10 μM ; Warner *et al.*, 1999) caused a small, but significant, potentiation of anandamide-induced relaxation (control, $\text{pEC}_{50}=6.6\pm 0.1$, $E_{\text{max}}=97\pm 2\%$, $n=5$; with nimesulide, $\text{pEC}_{50}=6.9\pm 0.2$, $E_{\text{max}}=102\pm 1\%$, $n=5$; $P<0.01$; Figure 3b). In the absence of endothelium, nimesulide had no effect on anandamide relaxations (control, $\text{pEC}_{50}=6.6\pm 0.2$, $E_{\text{max}}=98\pm 2\%$, $n=5$; with nimesulide, $\text{pEC}_{50}=6.5\pm 0.2$, $E_{\text{max}}=100\pm 1\%$, $n=5$; Figure 3c). Interestingly, the combined treatment of the FAAH inhibitor, URB597 and nimesulide had similar potentiation effects compared to those of URB597 alone (control, $\text{pEC}_{50}=7.0\pm 0.1$, $E_{\text{max}}=100\pm 2\%$, $n=6$; with nimesulide

and URB597, $\text{pEC}_{50}=7.4\pm 0.2$, $E_{\text{max}}=102\pm 1\%$, $n=7$; $P<0.001$; Figure 3d, c.f. Figure 1a). In contrast, the relaxant effects of methanandamide were unaffected by nimesulide (control, $\text{pEC}_{50}=6.5\pm 0.3$, $E_{\text{max}}=92\pm 4$, $n=6$; with nimesulide, $\text{pEC}_{50}=6.4\pm 0.3$, $E_{\text{max}}=93\pm 2\%$, $n=6$).

Effects of COX inhibitors on relaxation to 2-AG

Relaxations to 2-AG were significantly potentiated by 10 μM indomethacin (with endothelium, control, $\text{pEC}_{50}=5.5\pm 0.1$, $E_{\text{max}}=78\pm 5\%$, $n=7$; with indomethacin, $\text{pEC}_{50}=5.9\pm 0.1\%$, $E_{\text{max}}=86\pm 5\%$, $n=6$; $P<0.05$; Figure 4a). However, in endothelium-denuded vessels, indomethacin had no significant effect (control, $\text{pEC}_{50}=5.4\pm 0.1\%$, $E_{\text{max}}=51\pm 14\%$, $n=5$; with indomethacin, $\text{pEC}_{50}=5.3\pm 0.1$, $E_{\text{max}}=62\pm 13\%$, $n=5$; Figure 4b).

The COX inhibitor flurbiprofen, which displays a greater preference for COX-1 over COX-2 isoform than indomethacin (Warner *et al.*, 1999), was also used. The presence of flurbiprofen (10 μM) also caused a leftward displacement of the concentration-response curve to 2-AG (with endothelium, control, $\text{pEC}_{50}=5.5\pm 0.1$, $E_{\text{max}}=82\pm 8\%$, $n=4$; with flurbiprofen, $\text{pEC}_{50}=5.9\pm 0.2$, $E_{\text{max}}=95\pm 2\%$, $n=5$; $P<0.01$; Figure 4c). However, nimesulide (10 μM) had no effect on 2-AG-induced relaxation (with endothelium,




Figure 3 Effects of COX inhibitors on anandamide-induced relaxation in mesenteric arteries. Relaxation was elicited by anandamide alone, or in the presence of $10\ \mu\text{M}$ indomethacin (a), $10\ \mu\text{M}$ nimesulide (b, c) or $10\ \mu\text{M}$ nimesulide plus $1\ \mu\text{M}$ URB597 (d) in endothelium-intact or -denuded vessels. Values are shown as means and vertical lines represent s.e.m. $n = 4-6$.

control, $\text{pEC}_{50} = 5.4 \pm 0.1$, $E_{\text{max}} = 96 \pm 2\%$, $n = 5$; with nimesulide, $\text{pEC}_{50} = 5.6 \pm 0.3$, $E_{\text{max}} = 81 \pm 10\%$, $n = 5$; Figure 4d).

Interestingly, the combined treatment of indomethacin and the putative MGL inhibitor, MAFP ($10\ \mu\text{M}$ each) greatly enhanced relaxations to 2-AG; an apparent additive effect of the two inhibitors was observed (control, $\text{pEC}_{50} = 5.4 \pm 0.1$, $E_{\text{max}} = 77 \pm 15\%$, $n = 5$; with indomethacin and MAFP, $\text{pEC}_{50} = 6.3 \pm 0.3$, $E_{\text{max}} = 100 \pm 1\%$, $n = 5$; $P < 0.001$; Figure 5a; cf. Figures 2b and 4a). In endothelium-denuded vessels, MAFP also caused a significant potentiation effect on 2-AG relaxations (control, $\text{pEC}_{50} = 5.3 \pm 0.1$, $E_{\text{max}} = 63 \pm 15\%$, $n = 4$; with indomethacin and MAFP, $\text{pEC}_{50} = 5.4 \pm 0.1$, $E_{\text{max}} = 92 \pm 4\%$, $n = 4$; $P < 0.001$; Figure 5b).

Effects of indomethacin and MAFP on relaxation to noladin ether
The non-hydrolysable analogue of 2-AG, noladin ether (Sugiura *et al.*, 1999) induced endothelium-dependent relaxation in the mesenteric arteries (with endothelium, $\text{pEC}_{50} = 6.0 \pm 0.2$, $E_{\text{max}} = 90 \pm 5\%$, $n = 5$; without endothelium, $\text{pEC}_{50} = 5.3 \pm 0.1$, $E_{\text{max}} = 55 \pm 14\%$, $n = 5$; $P < 0.001$). Unlike the case for 2-AG, relaxations induced by noladin ether were not affected by either indomethacin (control, $\text{pEC}_{50} = 6.2 \pm 0.2$, $E_{\text{max}} = 94 \pm 4\%$, $n = 4$; with $10\ \mu\text{M}$ indo-

methacin, $\text{pEC}_{50} = 6.0 \pm 0.2$, $E_{\text{max}} = 94 \pm 5$, $n = 4$), or MAFP (control, $\text{pEC}_{50} = 6.3 \pm 0.2$, $E_{\text{max}} = 99 \pm 1\%$, $n = 5$; with $10\ \mu\text{M}$ MAFP, $\text{pEC}_{50} = 6.6 \pm 0.2$, $E_{\text{max}} = 90 \pm 2\%$, $n = 5$).

Effects of COX inhibitors and MAFP on relaxation to arachidonic acid

For comparison, vasoactive effects of arachidonic acid, the common product from hydrolysis of both anandamide and 2-AG were also examined. In endothelium-intact mesenteric arteries, arachidonic acid caused concentration-dependent relaxation ($\text{pEC}_{50} = 5.7 \pm 0.2$, $E_{\text{max}} = 99 \pm 2\%$; $n = 4$; Figure 6a). Removal of the endothelium greatly reduced arachidonic acid relaxations ($\text{pEC}_{50} = 4.8 \pm 0.2$, $E_{\text{max}} = 85 \pm 6\%$; $n = 4$; $P < 0.001$; Figure 6a).

Indomethacin ($10\ \mu\text{M}$; $P < 0.01$), but not nimesulide ($10\ \mu\text{M}$), significantly enhanced arachidonic acid-induced relaxation (control, $\text{pEC}_{50} = 5.7 \pm 0.1$, $E_{\text{max}} = 98 \pm 1\%$; $n = 7$; with indomethacin, $\text{pEC}_{50} = 6.5 \pm 0.2$, $E_{\text{max}} = 99 \pm 1\%$; $n = 6$; with nimesulide, $\text{pEC}_{50} = 6.0 \pm 0.3$, $E_{\text{max}} = 99 \pm 1\%$; $n = 5$; Figure 6b). On the other hand, relaxations to arachidonic acid were not affected by treatment with $10\ \mu\text{M}$ MAFP (control, $\text{pEC}_{50} = 5.7 \pm 0.2$, $E_{\text{max}} = 97 \pm 2\%$; $n = 4$; with MAFP, $\text{pEC}_{50} = 5.9 \pm 0.2$, $E_{\text{max}} = 89 \pm 4\%$; $n = 4$).




Figure 4 Effects of COX inhibitors on 2-AG-induced relaxation in mesenteric arteries. Relaxation was elicited by 2-AG alone, or in the presence of 10 μ M indomethacin (a, b), 10 μ M flurbiprofen (c) or 10 μ M nimesulide (d) in endothelium-intact or -denuded vessels. Values are shown as means and vertical lines represent s.e.m. $n = 4-7$.




Figure 5 Effects of the combined treatment of indomethacin (10 μ M) and MAFP (10 μ M) on relaxation to 2-AG in endothelium-intact (a) and -denuded (b) mesenteric arteries. Values are shown as means and vertical lines represent s.e.m. $n = 4-5$.

Discussion and conclusions

The present study suggests that in the rat small mesenteric artery, relaxant responses to anandamide and 2-AG are limited by hydrolysis to arachidonic acid and subsequent production of COX metabolites. For anandamide, catabolism by FAAH and COX-2 play a significant role in its inactivation. In contrast, MGL-like activity and COX-1 play

a more important role in the metabolism of 2-AG in this vascular preparation.

The primary route for anandamide catabolism is FAAH-mediated hydrolysis into arachidonic acid and ethanolamine (Cravatt *et al.*, 1996, 2001). Indeed, we found that the selective FAAH inhibitor, URB597 significantly potentiated anandamide-induced relaxation in rat mesenteric arteries. Two other FAAH inhibitors, ATFMK and MAFP, which are




Figure 6 Effects of endothelium removal (a), indomethacin ($10 \mu\text{M}$; b) and nimesulide ($10 \mu\text{M}$; b) on relaxation to arachidonic acid in mesenteric arteries. Values are shown as means and vertical lines represent s.e.m. $n=4-6$.

structurally distinct from URB597 (Deutsch *et al.*, 1997; Kathuria *et al.*, 2003), similarly potentiated anandamide relaxations. These results agree with the initial findings of others, that phenylmethylsulphonyl fluoride, a nonspecific serine hydrolase inhibitor, enhances mesenteric relaxations to anandamide (White and Hiley, 1997; Mendizabal *et al.*, 2001). Moreover, methanandamide, the metabolically stable analogue of anandamide (Abadji *et al.*, 1994), induced similar mesenteric relaxations that were insensitive to URB597. Together, our results point to a regulatory role for FAAH-mediated hydrolysis of anandamide in the vascular wall. It has been shown previously that FAAH are expressed and functional in endothelial cells of bovine coronary artery (Pratt *et al.*, 1998) and mouse cerebral microcirculation (Chen *et al.*, 2004). Here, URB597 had no significant effect on anandamide relaxations in endothelium-denuded vessels, suggesting that FAAH activity largely resides in the endothelium of rat small mesenteric arteries.

The other major endocannabinoid 2-AG is also liable to degradation. In fact, 2-AG might be more susceptible to degradation than anandamide both *in vitro* and *in vivo* (Pratt *et al.*, 1998; Jarai *et al.*, 2000; Savinainen *et al.*, 2001; Gauthier *et al.*, 2005). Recent evidence suggests that 2-AG is primarily hydrolysed by MGL to arachidonic acid and glycerol (Cravatt and Lichtman, 2002; Dinh *et al.*, 2002; Kathuria *et al.*, 2003). Nevertheless, as 2-AG could also serve as a substrate for FAAH (Goparaju *et al.*, 1998), FAAH-mediated hydrolysis might also play a role in the inactivation of 2-AG (Bifulco *et al.*, 2004; de Lago *et al.*, 2005; Maione *et al.*, 2006). In this study, the lack of effect of URB597 on 2-AG relaxations indicates that FAAH has little impact on 2-AG metabolism in the isolated mesenteric preparation. Interestingly, however, MAFP significantly potentiated responses to 2-AG in endothelium-intact and -denuded vessels. We propose that this potentiation occurs as a result of the inhibition of MGL by MAFP. A number of studies have also shown that MAFP is a combined FAAH and MGL inhibitor (Di Marzo *et al.*, 1999; Goparaju *et al.*, 1999; Dinh *et al.*, 2002; Saario *et al.*, 2004); it probably acts by targeting the arachidonyl substrate site of the two enzymes. In

membrane and cytosolic fractions of the brain, MAFP inhibits MGL with an IC_{50} as low as 2 nM (Goparaju *et al.*, 1999; Saario *et al.*, 2004), which is similar to IC_{50} values found for FAAH inhibition in enzyme assays (De Petrocellis *et al.*, 1997; Deutsch *et al.*, 1997). Thus, the observed differential effects of MAFP and URB597 on 2-AG relaxations could suggest the involvement of MGL. It was noted that MAFP is also known to inhibit cytosolic phospholipase A_2 (Lio *et al.*, 1996), which by unknown mechanisms, could also contribute to the relaxant responses to 2-AG. However, this seems unlikely based on the pharmacological profile of relaxations induced by 2-AG, noladin ether and arachidonic acid. First, ATFMK is also an inhibitor of cytosolic phospholipase A_2 (Street *et al.*, 1993) but it only tended to potentiate relaxations to lower concentrations ($\leq 1 \mu\text{M}$) of 2-AG. One possible explanation is that ATFMK is less effective than MAFP at reducing MGL activity, as has been shown in the brain (Goparaju *et al.*, 1999; Dinh *et al.*, 2002; Saario *et al.*, 2004). Second, noladin ether, a metabolically stable analogue of 2-AG, mimicked the endothelium-dependent mesenteric relaxation to 2-AG, but its effects were not affected by MAFP. Third, MAFP had no effect on arachidonic acid-induced relaxation. This argues against the possibility that inhibition of cytosolic phospholipase A_2 by MAFP somehow potentiated responses to the hydrolysis product of 2-AG, arachidonic acid. Taken together, the present results are consistent with 2-AG catabolism via MGL-like activity in the vascular wall, although involvement of other esterases cannot be ruled out. Given that the potentiation effect of MAFP was observed in vessels with and without endothelium, MGL activity is probably present in both endothelial and smooth muscle cells. In an attempt to characterize further the MGL-like activity in the mesenteric artery pharmacologically, we also tested the effects of URB754, which has recently been suggested to act as a selective inhibitor of MGL with no activity towards FAAH (Makara *et al.*, 2005). We found that URB754 had no detectable effect on relaxation to 2-AG. This may seem contradictory to our proposal that MGL activity (MAFP-sensitive) limits the relaxant effects of 2-AG. However, during the course of this




Figure 7 Schematic diagrams showing the proposed metabolic pathways for anandamide (a) and 2-AG (b) in rat small mesenteric arteries. The postulated locations (in parentheses) and inhibitors of the metabolizing enzymes are also shown. FAAH, fatty acid amide hydrolase; MGL, monacylglycerol lipase; COX, cyclooxygenase; ec, endothelium; vsm, vascular smooth muscle.

study, other researchers have independently found that the commercially available URB754 is ineffective in inhibiting 2-AG hydrolysis and thus its ability to target MGL has been questioned (e.g. Saario *et al.*, 2006).

An increasing number of reports indicate that metabolism of endocannabinoid by COX might be implicated in the cardiovascular actions of endocannabinoids (Jarai *et al.*, 2000; Gauthier *et al.*, 2005; Wahn *et al.*, 2005). Therefore, in this study, we further examined the role of COX-1 and COX-2 in the relaxation to endocannabinoids. The COX inhibitor, indomethacin had no significant effect on relaxations to anandamide, consistent with results from previous studies (Ho and Hiley, 2003; O'Sullivan *et al.*, 2004). Interestingly, selective inhibition of COX-2 with nimesulide resulted in a small but significant enhancement in anandamide-induced relaxation in endothelium-intact vessels. Nimesulide did not cause additional potentiation when co-applied with the FAAH inhibitor URB597, so it is likely that the metabolism mediated by COX-2 occurs downstream of anandamide hydrolysis (Figure 7a). Nevertheless, it remains possible that COX-2 catalyses a direct oxidation reaction with anandamide producing prostamides (Yu *et al.*, 1997; Figure 7a). This might contribute to the small inhibitory effect of nimesulide on anandamide relaxations, as the putative prostamides are inactive at prostanoid EP and FP receptors and hence likely to display no vasorelaxant activity (Matias *et al.*, 2004). Recently, Chen *et al.* (2005) showed that prolonged treatment with anandamide and methanandamide (after ≥ 1 h incubation) increases COX-2 expression in mouse cerebral endothelial cells. However, our observations that these cannabinoids had much faster relaxant responses (minutes), and that methanandamide induced a nimesulide-insensitive relaxation, argue against a significant contribution by this pathway to the mesenteric relaxations. It is noteworthy that, although the role for COX-2 (inducible) as compared to COX-1 (constitutively active) in vascular control under physiological conditions is still undefined, the detection of basal COX-2 activity in rat mesenteric arteries is consistent with recent findings that COX-2 is expressed in mesenteric arteries of healthy mice (Guo *et al.*, 2005), rabbits (Zhang *et al.*, 2005) and humans (Tabernero *et al.*, 2003).

In contrast to anandamide, indomethacin and the more COX-1-selective inhibitor, flurbiprofen (Warner *et al.*, 1999) but not nimesulide potentiated 2-AG relaxations. These

results were interpreted as evidence for a more important role for COX-1 compared to COX-2 in regulating relaxations elicited by 2-AG. Alternatively, simultaneous inhibition of both COX isoforms (by indomethacin and flurbiprofen at $10 \mu\text{M}$) was required to potentiate 2-AG relaxations. However, the lack of effect of nimesulide alone indicates that COX-2 metabolism plays a very small part, if any, in 2-AG degradation. The effect of indomethacin was absent in denuded vessels and thus points to COX-1 metabolism of 2-AG in the endothelium. Given that COX-2 but not COX-1 could directly oxidize 2-AG (Kozak *et al.*, 2000), COX-1 metabolism is presumed to occur downstream of 2-AG hydrolysis (Figure 7b). Interestingly, relaxant responses to arachidonic acid were also potentiated by indomethacin, suggesting that COX-1-derived contractile metabolites might be produced from 2-AG. This might, at least partly, explain the enhanced 2-AG relaxations in the presence of COX-1 inhibitors (Figure 7b). Indeed, relaxations to noladin ether were unaffected by indomethacin. At present, the mechanisms underlying the differential effects of indomethacin (presumed COX-1 inhibition) and nimesulide (COX-2 inhibition) on the relaxations to anandamide and 2-AG remain unclear. It is possible that 2-AG hydrolysis, mediated by MGL and perhaps other esterases, is strongly coupled with COX-1 activity, which might also explain the finding that 2-AG relaxations were greatly enhanced by the combined treatment of MAFP and indomethacin. Hypothetically, such functional coupling can be achieved by some form of physical association between the enzymes, so that arachidonic acid produced from MGL-mediated hydrolysis is in close proximity to COX.

To conclude, the present study provides pharmacological evidence suggesting that, in the rat small mesenteric artery, hydrolysis via FAAH and MGL-like activity limits the vasodilator actions of anandamide and 2-AG, respectively. For 2-AG, a combined inhibition of MGL and COX could further enhance its vascular impact. Our results also indicate that the endothelium represents a major metabolic barrier in the regulation of endocannabinoid actions, in as much as it determines the presence of significant FAAH, MGL, COX-1 and COX-2 activity, although MGL-mediated hydrolysis of 2-AG might also occur in the mesenteric smooth muscle (Figure 7). In support of this proposal, it has recently been found that the vascular content of both anandamide and

2-AG increases after endothelial removal in rat middle cerebral arteries (Rademacher *et al.*, 2005). It should be pointed out that, in some other vascular preparations, vasoactive effects of endocannabinoids might be largely mediated, rather than regulated, by degradation to arachidonic acid derivatives (bovine coronary: Gauthier *et al.*, 2005; rabbit pulmonary: Wahn *et al.*, 2005). Further experimentation is therefore warranted to fully understand the haemodynamic effects of inhibitors of endocannabinoid hydrolases. Nonetheless, in conditions where an increase in local concentration and life-span of endocannabinoids in the circulation is beneficial, for instance in hypertension (Batkai *et al.*, 2004), inhibitors of endocannabinoid inactivation including those used in this study might prove useful.

Acknowledgements

We are grateful to the British Heart Foundation (PG/06/064/20985) and the University of Nottingham (Anne McLaren Fellowship, WSVH) for financial support.

Conflict of interest

The authors state no conflict of interest.

References

- Abadji V, Lin S, Taha G, Griffin G, Stevenson LA, Pertwee RG *et al.* (1994). (R)-methanandamide: a chiral novel anandamide possessing higher potency and metabolic stability. *J Med Chem* **37**: 1889–1893.
- Batkai S, Pacher P, Osei-Hyiaman D, Radaeva S, Liu J, Harvey-White J *et al.* (2004). Endocannabinoids acting at cannabinoid-1 receptors regulate cardiovascular function in hypertension. *Circulation* **110**: 1996–2002.
- Bifulco M, Laezza C, Valenti M, Ligresti A, Portella G, Di Marzo V (2004). A new strategy to block tumor growth by inhibiting endocannabinoid inactivation. *FASEB J* **18**: 1606–1608.
- Chen P, Hu S, Harmon SD, Moore SA, Spector AA, Fang X (2004). Metabolism of anandamide in cerebral microvascular endothelial cells. *Prostaglandins Other Lipid Mediat* **73**: 59–72.
- Chen P, Hu S, Yao J, Moore SA, Spector AA, Fong X (2005). Induction of cyclooxygenases-2 by anandamide in cerebral microvascular endothelium. *Microvasc Res* **69**: 28–35.
- Cravatt BF, Demarest K, Patricelli MP, Bracey MH, Giang DK, Martin BR *et al.* (2001). Supersensitivity to anandamide and enhanced endogenous cannabinoid signaling in mice lacking fatty acid amide hydrolase. *Proc Natl Acad Sci USA* **98**: 9371–9376.
- Cravatt BF, Giang DK, Mayfield SP, Boger DL, Lerner RA, Gilula NB (1996). Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* **384**: 83–87.
- Cravatt BF, Lichtman AH (2002). The enzymatic inactivation of the fatty acid amide class of signaling lipids. *Chem Phys Lipids* **121**: 135–148.
- De Lago E, Petrosino S, Valenti M, Morera E, Ortega-Gutierrez S, Fernandez-Ruiz J *et al.* (2005). Effect of repeated systemic administration of selective inhibitors of endocannabinoid inactivation on rat brain endocannabinoid levels. *Biochem Pharmacol* **70**: 446–452.
- De Petrocellis L, Melck D, Ueda N, Maurelli S, Kurahashi Y, Yamamoto S *et al.* (1997). Novel inhibitors of brain, neuronal, and basophilic anandamide amidohydrolase. *Biochem Biophys Res Commun* **231**: 82–88.
- Desarnaud F, Cadas H, Piomelli D (1995). Anandamide amidohydrolase activity in rat brain microsomes. Identification and partial characterization. *J Biol Chem* **270**: 6030–6035.
- Deutsch D, Chin S (1993). Enzymatic synthesis and degradation of anandamide, a cannabinoid receptor agonist. *Biochem Pharmacol* **46**: 791–796.
- Deutsch DG, Omeir R, Arreaza G, Salehani D, Prestwich GD, Huang Z *et al.* (1997). Methyl arachidonoyl fluorophosphonate: a potent irreversible inhibitor of anandamide amidase. *Biochem Pharmacol* **53**: 255–260.
- Di Marzo V, Bisogno T, De Petrocellis L, Melck D, Orlando P, Wagner JA *et al.* (1999). Biosynthesis and inactivation of the endocannabinoid 2-arachidonoylglycerol in circulating and tumoral macrophages. *Eur J Biochem* **264**: 258–267.
- Dinh TP, Freund TF, Piomelli D (2002). A role for monoglyceride lipase in 2-arachidonoylglycerol inactivation. *Chem Phys Lipids* **121**: 149–158.
- Gardiner SM, March JE, Kemp PA, Bennett T (2002). Complex regional haemodynamic effects of anandamide in conscious rats. *Br J Pharmacol* **135**: 1889–1896.
- Gauthier KM, Baewer DV, Hittner S, Hillard CJ, Nithipatikom K, Reddy DS *et al.* (2005). Endothelium-derived 2-arachidonoylglycerol: an intermediate in vasodilatory eicosanoid release in bovine coronary arteries. *Am J Physiol Heart Circ Physiol* **288**: H1344–H1351.
- Gebremedhin D, Lange AR, Campbell WB, Hillard CJ, Harder DR (1999). Cannabinoid CB1 receptor of cat cerebral arterial muscle functions to inhibit L-type Ca²⁺ channel current. *Am J Physiol* **276**: H2085–H2093.
- Goparaju SK, Ueda N, Taniguchi K, Yamamoto S (1999). Enzymes of porcine brain hydrolyzing 2-arachidonoylglycerol, an endogenous ligand of cannabinoid receptors. *Biochem Pharmacol* **57**: 417–423.
- Goparaju SK, Ueda N, Yamaguchi H, Yamamoto S (1998). Anandamide amidohydrolase reacting with 2-arachidonoylglycerol, another cannabinoid receptor ligand. *FEBS Lett* **422**: 69–73.
- Grainger J, Boachie-Ansah G (2001). Anandamide-induced relaxation of sheep coronary arteries: the role of the vascular endothelium, arachidonic acid metabolites and potassium channels. *Br J Pharmacol* **134**: 1003–1012.
- Guo Z, Su W, Allen S, Pang H, Daugherty A, Smart E *et al.* (2005). COX-2 up-regulation and vascular smooth muscle contractile hyperreactivity in spontaneous diabetic db/db mice. *Cardiovasc Res* **67**: 723–735.
- Ho WS, Hiley CR (2003). Endothelium-independent relaxation to cannabinoids in rat-isolated mesenteric artery and role of Ca²⁺ influx. *Br J Pharmacol* **139**: 585–597.
- Ho WS, Hiley CR (2004). Vasorelaxant effects of 2-arachidonoylglycerol and noladin ether. In: *International Cannabinoid Research Society Symposium*, Abstract 195p.
- Ho WS, Hillard CJ (2005). Modulators of endocannabinoid enzymic hydrolysis and membrane transport. In: Roger Pertwee (eds). *Handb Exp Pharmacol Cannabinoids*. Springer-Verlag: Germany, **168** pp 187–207.
- Jarai Z, Wagner JA, Goparaju SK, Wang L, Razdan RK, Sugiura T *et al.* (2000). Cardiovascular effects of 2-arachidonoyl glycerol in anesthetized mice. *Hypertension* **35**: 679–684.
- Kagota S, Yamaguchi Y, Nakamura K, Sugiura T, Waku K, Kunitomo M (2001). 2-Arachidonoylglycerol, a candidate of endothelium-derived hyperpolarizing factor. *Eur J Pharmacol* **415**: 233–238.
- Karlsson M, Reue K, Xia YR, Lusis AJ, Langin D, Tornqvist H *et al.* (2001). Exon-intron organization and chromosomal localization of the mouse monoglyceride lipase gene. *Gene* **272**: 11–18.
- Kathuria S, Gaetani S, Fegley D, Valino F, Duranti A, Tontini A *et al.* (2003). Modulation of anxiety through blockade of anandamide hydrolysis. *Nat Med* **9**: 76–81.
- Kozak KR, Rowlinson SW, Marnett LJ (2000). Oxygenation of the endocannabinoid, 2-arachidonoylglycerol, to glyceryl prostaglandins by cyclooxygenase-2. *J Biol Chem* **275**: 33744–33749.
- LaBelle EF, Polyak E (1998). Norepinephrine stimulates arachidonic acid release from vascular smooth muscle via activation of cPLA2. *Am J Physiol* **274**: C1129–C1137.
- Lake KD, Martin BR, Kunos G, Varga K (1997). Cardiovascular effects of anandamide in anesthetized and conscious normotensive and hypertensive rats. *Hypertension* **29**: 1204–1210.

- Lio YC, Reynolds LJ, Balsinde J, Dennis EA (1996). Irreversible inhibition of Ca(2+)-independent phospholipase A2 by methyl arachidonoyl fluorophosphonate. *Biochim Biophys Acta* **1302**: 55–60.
- Maione S, Bisogno T, de Novellis V, Palazzo E, Cristino L, Valenti M *et al.* (2006). Elevation of endocannabinoid levels in the ventrolateral periaqueductal grey through inhibition of fatty acid amide hydrolase affects descending nociceptive pathways via both cannabinoid receptor type 1 and transient receptor potential vanilloid type-1 receptors. *J Pharmacol Exp Ther* **316**: 969–982.
- Makara JK, Mor M, Fegley D, Szabo SI, Kathuria S, Astarita G *et al.* (2005). Selective inhibition of 2-AG hydrolysis enhances endocannabinoid signaling in hippocampus. *Nat Neurosci* **8**: 1139–1141.
- Matias J, Chen J, de Petrocellis L, Bisogno T, Ligresti A, Fezza F *et al.* (2004). Prostaglandin ethanolamides (prostamides): *in vitro* pharmacology and metabolism. *J Pharmacol Exp Ther* **309**: 745–757.
- Mechoulam R, Fride E, Ben-Shabat S, Meiri U, Horowitz M (1998). Carbachol, an acetylcholine receptor agonist, enhances production in rat aorta of 2-arachidonoyl glycerol, a hypotensive endocannabinoid. *Eur J Pharmacol* **362**: R1–R3.
- Mendizabal VE, Orliac ML, Adler-Graschinsky E (2001). Long-term inhibition of nitric oxide synthase potentiates effects of anandamide in the rat mesenteric bed. *Eur J Pharmacol* **427**: 251–262.
- O'Sullivan SE, Kendall DA, Randall MD (2004). Heterogeneity in the mechanisms of vasorelaxation to anandamide in resistance and conduit rat mesenteric arteries. *Br J Pharmacol* **142**: 435–442.
- Offertaler L, Mo FM, Batkai S, Liu J, Begg M, Razdan RK *et al.* (2003). Selective ligands and cellular effectors of a G protein-coupled endothelial cannabinoid receptor. *Mol Pharmacol* **63**: 699–705.
- Pacher P, Batkai S, Osei-Hyiaman D, Offertaler L, Liu J, Harvey-White J *et al.* (2005). Hemodynamic profile, responsiveness to anandamide, and baroreflex sensitivity of mice lacking fatty acid amide hydrolase. *Am J Physiol Heart Circ Physiol* **289**: H533–H541.
- Pratt PF, Hillard CJ, Edgemond WS, Campbell WB (1998). N-arachidonyl ethanolamide relaxation of bovine coronary artery is not mediated by CB1 cannabinoid receptor. *Am J Physiol* **274**: H375–H381.
- Rademacher DJ, Patel S, Ho WS, Savoie AM, Rusch NJ, Gauthier KM *et al.* (2005). U-46619 but not serotonin increases endocannabinoid content in middle cerebral artery: evidence for functional relevance. *Am J Physiol Heart Circ Physiol* **288**: H2694–H2701.
- Ralevic V, Kendall DA, Randall MD, Zygmunt PM, Movahed P, Hogestatt ED (2000). Vanilloid receptors on capsaicin-sensitive sensory nerves mediate relaxation to methanandamide in the rat isolated mesenteric arterial bed and small mesenteric arteries. *Br J Pharmacol* **130**: 1483–1488.
- Randall MD, Alexander SP, Bennett T, Boyd EA, Fry JR, Gardiner SM *et al.* (1996). An endogenous cannabinoid as an endothelium-derived vasorelaxant. *Biochem Biophys Res Commun* **229**: 114–120.
- Saario SM, Palomaki V, Lehtonen M, Nevalainen T, Jarvinen T, Laitinen JT (2006). URB754 has no effect on the hydrolysis or signaling capacity of 2-AG in the rat brain. *Chem Biol* **13**: 811–814.
- Saario SM, Savinainen JR, Laitinen JT, Jarvinen T, Niemi R (2004). Monoglyceride lipase-like enzymatic activity is responsible for hydrolysis of 2-arachidonoylglycerol in rat cerebellar membranes. *Biochem Pharmacol* **67**: 1381–1387.
- Savinainen JR, Jarvinen T, Laine K, Laitinen JT (2001). Despite substantial degradation, 2-arachidonoylglycerol is a potent full efficacy agonist mediating CB(1) receptor-dependent G-protein activation in rat cerebellar membranes. *Br J Pharmacol* **134**: 664–672.
- Street IP, Lin HK, Laliberte F, Ghomashchi F, Wang Z, Perrier H *et al.* (1993). Slow- and tight-binding inhibitors of the 85-kDa human phospholipase A2. *Biochemistry* **32**: 5935–5940.
- Sugiura T, Kodaka T, Nakane S, Miyashita T, Kondo S, Suhara Y *et al.* (1999). Evidence that the cannabinoid CB1 receptor is a 2-arachidonoylglycerol receptor. Structure-activity relationship of 2-arachidonoylglycerol, ether-linked analogues, and related compounds. *J Biol Chem* **274**: 2794–2801.
- Tabernero A, Reimund JM, Chasserot S, Muller CD, Andriantsitohaina R (2003). Cyclooxygenase-2 expression and role of vasoconstrictor prostanoids in small mesenteric arteries from patients with Crohn's disease. *Circulation* **107**: 1407–1410.
- Tornqvist H, Belfrage P (1976). Purification and some properties of a monoacylglycerol-hydrolyzing enzyme of rat adipose tissue. *J Biol Chem* **251**: 813–819.
- Wagner JA, Varga K, Jarai Z, Kunos G (1999). Mesenteric vasodilation mediated by endothelial anandamide receptors. *Hypertension* **33**: 429–434.
- Wahn H, Wolf J, Kram F, Frantz S, Wagner JA (2005). The endocannabinoid arachidonoyl ethanolamide (anandamide) increases pulmonary arterial pressure via cyclooxygenase-2 products in isolated rabbit lungs. *Am J Physiol Heart Circ Physiol* **289**: H2491–H2496.
- Warner TD, Giuliano F, Vojnovic I, Bukasa A, Mitchell JA, Vane JR (1999). Nonsteroid drug selectivities for cyclo-oxygenase-1 rather than cyclo-oxygenase-2 are associated with human gastrointestinal toxicity: a full *in vitro* analysis. *Proc Natl Acad Sci USA* **96**: 7563–7568.
- White R, Hiley CR (1997). A comparison of EDHF-mediated and anandamide-induced relaxations in the rat isolated mesenteric artery. *Br J Pharmacol* **122**: 1573–1584.
- Yu M, Ives D, Ramesha CS (1997). Synthesis of prostaglandin E2 ethanolamide from anandamide by cyclooxygenase-2. *J Biol Chem* **272**: 21181–21186.
- Zhang DX, Gauthier KM, Chawengsub Y, Holmes BB, Campbell WB (2005). Cyclooxygenase- and lipoxygenase-dependent relaxation to arachidonic acid in rabbit small mesenteric arteries. *Am J Physiol Heart Circ Physiol* **288**: H302–H309.
- Zygmunt PM, Petersson J, Andersson DA, Chuang H, Sorgard M, Di Marzo V *et al.* (1999). Vanilloid receptors on sensory nerves mediate the vasodilator action of anandamide. *Nature* **400**: 452–457.