

Research article

Open Access

## Ca<sup>2+</sup> binding to complement-type repeat domains 5 and 6 from the low-density lipoprotein receptor-related protein

Olav M Andersen\*<sup>1</sup>, Henrik Vorum<sup>2</sup>, Bent Honoré<sup>2</sup> and Hans C Thøgersen<sup>1</sup>

Address: <sup>1</sup>Laboratory of Gene Expression, Department of Molecular and Structural Biology, University of Aarhus, DK-8000 Aarhus C, Denmark and <sup>2</sup>Department of Medical Biochemistry, University of Aarhus, DK-8000 Aarhus C, Denmark

Email: Olav M Andersen\* - o.andersen@mdc-berlin.de; Henrik Vorum - hv@biokemi.au.dk; Bent Honoré - bh@biokemi.au.dk; Hans C Thøgersen - hct@biobase.dk

\* Corresponding author

Published: 18 August 2003

Received: 27 April 2003

BMC Biochemistry 2003, 4:7

Accepted: 18 August 2003

This article is available from: <http://www.biomedcentral.com/1471-2091/4/7>

© 2003 Andersen et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** The binding of ligands to clusters of complement-type repeat (CR)-domains in proteins of the low-density lipoprotein receptor (LDLR) family is dependent on Ca<sup>2+</sup> ions. One reason for this cation requirement was identified from the crystal structure data for a CR-domain from the prototypic LDLR, which showed the burial of a Ca<sup>2+</sup> ion as a necessity for correct folding and stabilization of this protein module. Additional Ca<sup>2+</sup> binding data to other CR-domains from both LDLR and the LDLR-related protein (LRP) have suggested the presence of a conserved Ca<sup>2+</sup> cage within CR-domains from this family of receptors that function in endocytosis and signalling.

**Results:** We have previously described the binding of several ligands to a fragment comprising the fifth and the sixth CR-domain (CR56) from LRP, as well as qualitatively described the binding of Ca<sup>2+</sup> ions to this CR-domain pair. In the present study we have applied the rate dialysis method to measure the affinity for Ca<sup>2+</sup>, and show that CR56 binds 2 Ca<sup>2+</sup> ions with an average affinity of  $K_D = 10.6$  microM, and there is no indication of additional Ca<sup>2+</sup> binding sites within this receptor fragment.


**Conclusions:** Both CR-domains of CR56 bind a single Ca<sup>2+</sup> ion with an affinity of 10.6 microM within the range of affinities demonstrated for several other CR-domains.

### Background

The ligand-binding domains of the entire LDLR family of cell surface receptors are comprised of imperfect repeats of about 40 amino acids, the CR-domains [1–3]. Each repeat contains six cysteine residues which are disulphide linked in the pattern one to three, two to five, and four to six, and each repeat harbors a Ca<sup>2+</sup> binding site [4,5]. Furthermore, the modular domain architecture of LRP comprises several epidermal growth factor (EGF)-precursor homology domains, a segment spanning the plasma membrane and a cytoplasmic domain. The entire LRP molecule might contain as many as 39 Ca<sup>2+</sup> binding sites, one located in

each of the 31 CR-domains [5–7], and two binding sites present in each of 4 EGF-domain pairs [7–11] (see legend to Figure 1A).

The understanding of how LRP binds Ca<sup>2+</sup> ions is important. The binding of all ligands is dependent on the presence of Ca<sup>2+</sup> [12–14] and ligand dissociation within the endocytic pathway has been suggested to occur as a possible consequence of the decrease in pH and the accompanying loss of affinity for Ca<sup>2+</sup> in the acidic endosomes [8].



**Figure 1**

(A) Schematic representation of the domain architecture of the LRP, with domains suggested to contain a Ca<sup>2+</sup> binding site as shown in gray. For estimation of the stoichiometric binding of Ca<sup>2+</sup> to the entire LRP molecule, each CR-domain is a potential cation carrier based on the fact that all CR-domain structures described so far include a bound Ca<sup>2+</sup> ion [5–7]. The crystal structure of the LDLR YWTD-repeated β-propeller does not show any cations and we assume similar lack of Ca<sup>2+</sup> binding to these fragments within LRP. In contrast, some EGF-domains do bind Ca<sup>2+</sup> and some do not. For LDLR the EGF-domain pair amino-terminal to the β-propeller binds 2 Ca<sup>2+</sup> ions, and accordingly, it is assumed that the four EGF-domain pairs within LRP are also Ca<sup>2+</sup> binding. The carboxy-terminal EGF-domain is not chelating a Ca<sup>2+</sup> ion [7–11]. By these assumptions, one LRP molecule could bind a total of at least 39 Ca<sup>2+</sup> ions. Rectangles (□) represent CR-domains, pentagons (◑) represent EGF-domains, and hexagonals (◒) for each blade of the β-propellers. (B) Ribbon representation of the canonical CR-domain folding and Ca<sup>2+</sup> binding site. Left, the backbone folding of CR5<sup>LDLR</sup> (Protein Data Bank accession code 1ajj [5]) showing the location of the Ca<sup>2+</sup> ion indicated as the sphere. Right, zoom of the Ca<sup>2+</sup> cage showing the octahedral cation coordination. (C) Alignment of the primary structures of Ca<sup>2+</sup>-binding CR-domains with a known 3D-structure, the first, the second, the fifth, and the sixth CR-domain from LDLR, and the third, the seventh, and the eighth CR-domain from LRP. The symbols below the alignment indicate residues involved in Ca<sup>2+</sup> chelation (‡, coordination by backbone carbonyl; †, coordination by side chain carboxyl), and the six conserved cysteines are indicated above the sequences with roman numbers.

Ligand recognition requires key residues in the CR-domains of the LDLR-like proteins as well as the presence of Ca<sup>2+</sup> ions. One important residue for this interaction is located at the center position between the fourth and the fifth cysteine residue, where an acidic side chain is impor-

tant for the efficient recognition of multiple protein ligands [15,16]. The backbone carbonyl group of residues located at identical positions in several domains homologous to CR5 and CR6 is involved in the coordination of a Ca<sup>2+</sup> ion [5,6]. The coordination sphere of the bound Ca<sup>2+</sup>

ion is well defined in octahedral geometry, with four carboxylate oxygen atoms from the acidic motif in one plane and two carbonyl oxygen atoms completing the coordination sphere at the apices (Figure 1B).

We have previously demonstrated the specific binding of LRP ligands to a minimum receptor fragment comprising only 2 CR-domains [15], and focused on the ligand interaction with the tandem domain CR56 fragment [17,18]. The affinity for a bound  $\text{Ca}^{2+}$  ion has been reported for several CR-domains, and in order to better understand the  $\text{Ca}^{2+}$ -dependent ligand binding properties of CR56 we have determined the stoichiometry and affinity for  $\text{Ca}^{2+}$  binding to this CR-domain pair. Furthermore, we undertook a stringent/direct method of affinity determination using the microchamber rate dialysis method [19] independent of the local molecular environment at the  $\text{Ca}^{2+}$  localization site, and could compare the determined affinity with data obtained by less direct methods, e.g. fluorescence analysis. We conclude that in general the affinities do not vary significantly among the CR-domains investigated.

## Results and Discussion


The demonstration of  $\text{Ca}^{2+}$  binding to the ubiquitin-fused CR56 protein [15] strongly suggested that CR56 contains at least one efficient  $\text{Ca}^{2+}$  binding site similar to other CR-domains [1,5]. However, from the previously adopted approach we could not determine the stoichiometry and affinity between  $\text{Ca}^{2+}$  and CR56. This was the main objective of the current study. Furthermore, we wanted to test whether the coupling of a CR-domain to a neighboring domain would influence the affinity of the single CR-domain, compared to the increasing pool of data describing the binding of  $\text{Ca}^{2+}$  to isolated CR-domains [6,20–23]. After affinity purification of ubiquitin-fused CR56 containing the authentic amino acid sequence, we liberated the CR56 domain pair from its fusion partner by factor  $X_a$  cleavage (as described in ref. [15]). 2D gel analysis showed a high degree of purity, since the recombinant CR56 protein migrated as a single major spot (not shown).  $\text{Ca}^{2+}$  binding was measured using the microchamber rate dialysis technique [19,24].

The  $\text{Ca}^{2+}$  binding data is shown in Figure 2A, and from Scatchard analysis of the data (Figure 2B), we conclude that the folded CR56 domain pair binds two  $\text{Ca}^{2+}$  ions with an average affinity of  $K_D = 10.6 \mu\text{M}$  assuming two identical binding sites. Even though there is a small derivation from the exact 1:2 stoichiometry ( $n = 2.0$  sites), the observed  $n = 2.2$  is within experimental error not significantly different from 2.0. Secondly, the number is in accord with other data determining  $\text{Ca}^{2+}$  affinity to tandem CR-domain pair [25]. Bieri et al. found a similar stoichiometry ( $n = 2.2$ ) when they measured the  $\text{Ca}^{2+}$

binding to a concatemer of  $\text{CR1}^{\text{LDLR}}\text{-CR2}^{\text{LDLR}}$ , and neither solution nor crystal structure indicate that there is more than two  $\text{Ca}^{2+}$  binding sites within this fragment [7,26]. Furthermore, the recently solved crystal structure of the LDLR luminal domain shows a single bound cation for each CR-domain without any additional interdomain binding sites [7]. The determined affinity is very close to the reported  $\text{Ca}^{2+}$  affinities for homologous LRP CR-domains,  $\text{CR3}^{\text{LRP}}$ ,  $\text{CR7}^{\text{LRP}}$ ,  $\text{CR8}^{\text{LRP}}$  and for the two LDLR CR-domains  $\text{CR1}^{\text{LDLR}}$  and  $\text{CR2}^{\text{LDLR}}$  (listed in Table 1), which indicates that the binding sites might be similar. Apparently, most CR-domains bind  $\text{Ca}^{2+}$  with an approximate affinity  $K_D \sim 10\text{--}20 \mu\text{M}$ , except  $\text{CR5}^{\text{LDLR}}$  and  $\text{CR6}^{\text{LDLR}}$  which show a higher affinity. Furthermore, our data suggests that the various methods used to measure the affinities are reliable.

The  $\text{Ca}^{2+}$  ions in the hitherto solved domain structures are located in identical  $\text{Ca}^{2+}$  cages as for  $\text{CR5}^{\text{LDLR}}$  described previously by Fass and colleagues [5] (Figure 1B). From a high level of sequence conservation for the sequences of  $\text{CR5}^{\text{LRP}}$  and  $\text{CR6}^{\text{LRP}}$  compared to the sequences of the CR-domains with a solved domain structure (Figure 1C), we suggest that the binding site for  $\text{Ca}^{2+}$  within both CR5 and CR6 are very similar to these. This is very important information for the assignment of nuclear magnetic resonances for the solution structure determination of CR56 (ongoing project).

The demonstration of independent folding of each CR-domain in tandem CR-domain pairs is substantiated by the reports of a total lack of interdomain interactions, and that  $\text{Ca}^{2+}$  coordination does not involve chelates from adjacent CR-domains [27,28]. In line with this our data suggest that two and only two  $\text{Ca}^{2+}$  are bound per tandem fragment as also reported for domain pairs from LDLR comprising  $\text{CR1}^{\text{LDLR}}\text{-CR2}^{\text{LDLR}}$  and  $\text{CR5}^{\text{LDLR}}\text{-CR6}^{\text{LDLR}}$  [25,27]. Since the  $\text{Ca}^{2+}$  affinity for CR5 and CR6 is similar to other known binding sites, it is tempting to believe that the  $\text{Ca}^{2+}$  ion is located in a similar  $\text{Ca}^{2+}$  cage as for these domains [1,5,6], and therefore the chelating residues located at identical positions within the primary structure is also cation coordinating in these domains. If this indeed is the case, we have recently demonstrated that the two residues speculated to provide electrons for  $\text{Ca}^{2+}$  coordination via their backbone functional group (Trp-953/Asp-958 in CR5 and Trp-994/Asp-999 in CR6) both contribute significantly to ligand binding [17,18]. The possibility that  $\text{Ca}^{2+}$  most likely are intimately linked to these residues suggest that  $\text{Ca}^{2+}$  binding exerts influence on ligand binding to CR-domains, because of a lack of dynamic and flexibility of residues at this particular position. In addition, especially the acidic residues is also involved in the intramolecular binding of CR-domains to the EGF-precursor homology domain at low pH, speculated to



**Figure 2**

(A) The  $Ca^{2+}$ -affinity of CR56 was determined by rate dialysis  $[Ca^{2+}]_{bound}/[CR56]$  versus the free  $Ca^{2+}$  concentration. The concentration of CR56 was  $15 \mu M$  throughout the analysis. (B) Scatchard analysis of the data shown in panel A. The solid line represents the best fit by least-squares analysis as described in the materials and methods section. The fact that the experimental points are located on a straight line indicates absence of cooperativity between the two  $Ca^{2+}$  binding sites. Strong positive cooperativity would result in upward curvature, while strong negative cooperativity would result in downward curvature of the Scatchard plot. The average affinity is determined to,  $K_D = 10.6 \mu M$  assuming two identical sites ( $n = 2.2$ ).

**Table 1: Ca<sup>2+</sup> binding properties of various CR-domains**

CR-domain	Ca <sup>2+</sup> affinity (K <sub>D</sub> ) μM	References	Method <sup>a)</sup>	Conditions
CR1 <sup>LDLR</sup>	7 10	[22] [30]	Fluorescence	25°C
CR2 <sup>LDLR</sup>	14	[25]		
CR5 <sup>LDLR</sup>	0.040 0.070 0.500 13	[27] [20,27] [6] [6]	Fluorescence Fluorescence ITC ITC	pH 7.0 pH 7.0 pH 7.4, 30°C pH 5.0, 30°C
CR6 <sup>LDLR</sup>	0.200	[27]	Fluorescence	pH 7.0
CR3 <sup>LRP</sup>	8 24	[6] [22]	ITC Fluorescence	pH 7.4, 30°C 25°C
CR5 <sup>LRP</sup>	11	The present data	Rate dialysis	pH 7.4, 37°C
CR6 <sup>LRP</sup>	11	The present data	Rate dialysis	pH 7.4, 37°C
CR7 <sup>LRP</sup>	13	[6]	ITC	pH 7.4, 30°C
CR8 <sup>LRP</sup>	6 13	[6] [22]	ITC Fluorescence	pH 7.4, 30°C 25°C
CR <sup>Tva</sup>	40	[23]	ITC	pH 7.4, 30°C

<sup>a)</sup> ITC; Isothermal titration calorimetry CRX<sup>LDLR</sup>, CRX<sup>LRP</sup>, and CR<sup>Tva</sup> denote the Xth CR-domain counting from the amino-terminal end of LDLR, LRP and the Tva receptor of avian leucosis and sarcoma virus type A, respectively.

result in structural rearrangement and ligand release within endosomes, underscoring the importance of understanding the local environment around the Ca<sup>2+</sup> binding site [7].

## Conclusions

Both CR-domains of the CR56-domain pair bind a single Ca<sup>2+</sup> ion with an average affinity, K<sub>D</sub> ~ 10.6 μM.

## Methods

### Proteins

Production and RAP-affinity purification of the ubiquitin-fused CR-domain pair comprising the fifth and the sixth CR-domain from LRP (see Figure 1) was described previously [15]. Purity was verified by two-dimensional gel electrophoresis showing the sample to be highly homogeneous (not shown).

### Calcium binding analysis

Qualitative <sup>45</sup>Ca blotting analysis to ubiquitin-fused CR56 has been described [15]. The quantitative rate dialysis method [19] was applied to determine the Ca<sup>2+</sup> binding constants for CR56. The binding experiments were performed in a medium containing 10 mM HEPES pH. 7.0, 150 mM NaCl and a final CR56 concentration at 15 μM. Buffers and protein solutions were passed through a Chelex 100 column (BioRAD) in order to obtain cation free solutions before use. The resin was pre-equilibrated with HEPES binding buffer before use. We have previously shown that this procedure is able to bring the Ca<sup>2+</sup> content of the solutions to negligible levels as determined by atomic absorption spectrometry [29]. The dialysis

membrane was of cellulose, Type 14.10, molecular cut-off 5000, from Diachema (Munich, Germany). The following equation was used to calculate the free Ca<sup>2+</sup> concentration from the total Ca<sup>2+</sup> concentration:

$$[Ca^{2+}]_{free} = - [Ca^{2+}]_{total} [k^*(t+t_0)]^{-1} \ln [(A_{left} - A_{right}) / (A_{left} + A_{right})]$$

where k is a pre-determined rate constant, t is time of dialysis and t<sub>0</sub> is an experimentally determined value which is dependent upon the procedure of filling, withdrawal, and rinsing of chambers and varies with the substance dialysed [19]. A<sub>left</sub> and A<sub>right</sub> denote the radioactivity in the left and right solution, respectively, after dialysis measured by liquid scintillation counting in an LKB Wallac 1218 Rackbeta scintillation counter. The values of k and t<sub>0</sub> for the Ca<sup>2+</sup> ligand used were 0.04650 min<sup>-1</sup> and 0.18 min. They were determined in separate experiments with no protein present and using varying dialysis times as described in detail [19]. In short, values of ln [(A<sub>left</sub> - A<sub>right</sub>) / (A<sub>left</sub> + A<sub>right</sub>)] are plotted versus the dialysis time. The rate constant k is then determined from the slope of the curve while t<sub>0</sub> is determined as the numerical value of the intercept with the time-axis (x-axis). The average number of Ca<sup>2+</sup> ions bound per protein molecule, r, was calculated from

$$r = ([Ca^{2+}]_{total} - [Ca^{2+}]_{free}) / [CR56]$$

Under the presumption that CR56 contains a number, n, of identical and independent Ca<sup>2+</sup>-binding sites the bind-

ing isotherm was fitted to the Scatchard equation by linear regression:

$$r/[Ca^{2+}]_{free} = -r/K_D + n/K_D,$$

where  $K_D$  is the dissociation constant.

### Abbreviations

CR-domain, complement-type repeat domain; EGF, epidermal growth factor; LDLR, low-density lipoprotein receptor; LRP, LDLR-related protein

### Authors' contributions

OMA produced and purified recombinant protein, prepared figures and drafted the manuscript. HV and BH performed the quantitative rate dialysis measurements and participated in drafting the manuscript. HCT conceived the study as well as participated in its design and coordination. All authors read and approved the final manuscript.

### Acknowledgements

We thank Kirsten Peterslund for excellent technical assistance. This work was supported by Grant 9901730 from the Danish National Science Research Council (HCT), the Danish Medical Research Council and the Novo Nordisk Foundation.

### References

- Brown MS, Herz J and Goldstein JL: **LDL-receptor structure. Calcium cages, acid baths and recycling receptors.** *Nature* 1997, **388**:629-630.
- Neels JG, Horn IR, van den Berg BMM, Pannekoek H and van Zonneveld A-J: **Ligand-receptor interactions of the low density lipoprotein receptor-related protein, a multi-ligand endocytic receptor.** *Fibrinolysis & Proteolysis* 1998, **12**:219-240.
- Herz J: **Deconstructing the LDL receptor – a rhapsody in pieces.** *Nat Struct Biol* 2001, **8**:476-478.
- Bieri S, Djordjevic JT, Daly NL, Smith R and Kroon PA: **Disulfide bridges of a cysteine-rich repeat of the LDL receptor ligand-binding domain.** *Biochemistry* 1995, **34**:13059-13065.
- Fass D, Blacklow S, Kim PS and Berger JM: **Molecular basis of familial hypercholesterolaemia from structure of LDL receptor module.** *Nature* 1997, **388**:691-693.
- Simonovic M, Dolmer K, Huang W, Strickland DK, Volz K and Gettins PGW: **Calcium coordination and pH dependence of the calcium affinity of ligand-binding repeat CR7 from the LRP. Comparison with related domains from the LRP and the LDL receptor.** *Biochemistry* 2001, **40**:15127-15134.
- Rudenko G, Henry L, Henderson K, Ichtchenko K, Brown MS, Goldstein JL and Deisenhofer J: **Structure of the LDL receptor extracellular domain at endosomal pH.** *Science* 2002, **298**:2353-8.
- Malby S, Pickering R, Saha S, Smallridge R, Linse S and Downing AK: **The first epidermal growth factor-like domain of the low-density lipoprotein receptor contains a noncanonical calcium binding site.** *Biochemistry* 2001, **40**:2555-2563.
- Kurniawan ND, Aliabadzadeh K, Brereton IM, Kroon PA and Smith R: **NMR structure and backbone dynamics of a concatemer of epidermal growth factor homology modules of the human low-density lipoprotein receptor.** *J Mol Biol* 2001, **311**:341-356.
- Saha S, Boyd J, Werner JM, Knott V, Handford PA, Campbell ID and Downing AK: **Solution structure of the LDL Receptor EGF-AB pair: A paradigm for the assembly of tandem calcium binding EGF domains.** *Structure* 2001, **9**:451-456.
- Jeon H, Meng W, Takagi J, Eck MJ, Springer TA and Blacklow SC: **Implications for familial hypercholesterolemia from the structure of the LDL receptor YWTD-EGF domain pair.** *Nat Struct Biol* 2001, **8**:499-504.
- Krieger M and Herz J: **Structures and functions of multiligand lipoprotein receptors: macrophage scavenger receptors and LDL receptor-related protein (LRP).** *Annu Rev Biochem* 1994, **63**:601-637.
- Goldstein JL, Brown MS, Anderson RG, Russell DW and Schneider WJ: **Receptor-mediated endocytosis: concepts emerging from the LDL receptor system.** *Annu Rev Cell Biol* 1985, **1**:1-39.
- Nykjær A, Olivecrona-Bengtsson G, Lookene A, Moestrup SK, Petersen CM, Weber W, Beisiegel U and Gliemann J: **The  $\alpha_2$ -macroglobulin receptor/low density lipoprotein receptor-related protein binds lipoprotein lipase and beta-migrating very low density lipoprotein associated with the lipase.** *J Biol Chem* 1993, **268**:15048-15055.
- Andersen OM, Christensen LL, Christensen PA, Sørensen ES, Jacobsen C, Moestrup SK, Etzerodt M and Thøgersen HC: **Identification of the Minimal Functional Unit in the Low Density Lipoprotein Receptor-related Protein for Binding the Receptor-associated Protein (RAP).** *J Biol Chem* 2000, **275**:21017-21024.
- Andersen OM, Christensen PA, Christensen LL, Jacobsen C, Moestrup SK, Etzerodt M and Thøgersen HC: **Specific Binding of  $\alpha$ -Macroglobulin to Complement-type Repeat CR4 of the Low Density Lipoprotein Receptor-related Protein.** *Biochemistry* 2000, **39**:10627-10633.
- Andersen OM, Petersen HH, Jacobsen C, Moestrup SK, Etzerodt M, Andreasen PA and Thøgersen HC: **Analysis of a two-domain binding site for the urokinase-type plasminogen activator – plasminogen activator inhibitor-1 complex in low density lipoprotein receptor-related protein.** *Biochem J* 2001:289-296.
- Andersen OM, Schwarz FP, Eisenstein E, Jacobsen C, Moestrup SK, Etzerodt M and Thøgersen HC: **Dominant thermodynamic role of the third independent receptor binding site in the receptor-associated protein RAP.** *Biochemistry* 2001, **40**:15408-15417.
- Honoré B: **Equilibrium dialysis and rate dialysis.** In: *Protein-Ligand Interactions: Hydrodynamics and Calorimetry. A Practical Approach* Edited by: Harding SE, Chowdhry BZ. Oxford: Oxford University Press; 2001:19-46.
- Blacklow SC and Kim PS: **Protein folding and calcium binding defects arising from familial hypercholesterolemia mutations of the LDL receptor.** *Nat Struct Biol* 1996, **3**:758-762.
- Atkins AR, Brereton IM, Kroon PA, Lee HT and Smith R: **Calcium is essential for the structural integrity of the cysteine-rich, ligand-binding repeat of the low-density lipoprotein receptor.** *Biochemistry* 1998, **37**:1662-1670.
- Dolmer K, Huang W and Gettins PGW: **Characterization of the calcium site in two complement-like domains from the low-density lipoprotein receptor-related protein (LRP) and comparison with a repeat from the low-density lipoprotein receptor.** *Biochemistry* 1998, **37**:17016-17023.
- Wang QY, Dolmer K, Huang W, Gettins PG and Rong L: **Role of calcium in protein folding and function of Tva, the receptor of subgroup A avian sarcoma and leukosis virus.** *J Virol* 2001, **75**:2051-2058.
- Vorum H, Liu X, Madsen P, Rasmussen HH and Honoré B: **Molecular cloning of a cDNA encoding human calumenin, expression in *Escherichia coli* and analysis of its  $Ca^{2+}$ -binding activity.** *Biochim Biophys Acta* 1998, **1386**:121-131.
- Bieri S, Atkins AR, Lee HT, Winzor DJ, Smith R and Kroon PA: **Folding, calcium binding, and structural characterization of a concatemer of the first and second ligand-binding modules of the low-density lipoprotein receptor.** *Biochemistry* 1998, **37**:10994-11002.
- Kurniawan ND, Atkins AR, Bieri S, Brown CJ, Brereton IM, Kroon PA and Smith R: **NMR structure of a concatemer of the first and second ligand-binding modules of the human low-density lipoprotein receptor.** *Protein Sci* 2000, **9**:1282-1293.
- North CL and Blacklow SC: **Structural independence of ligand-binding modules five and six of the LDL receptor.** *Biochemistry* 1999, **38**:3926-3935.
- North CL and Blacklow SC: **Evidence that familial hypercholesterolemia mutations of the LDL receptor cause limited local misfolding in an LDL-A module pair.** *Biochemistry* 2000, **39**:13127-13135.
- Vorum H, Madsen P, Rasmussen HH, Etzerodt M, Svendsen I, Celis JE and Honoré B: **Expression and divalent cation binding properties of the novel chemotactic inflammatory protein psoriasin.** *Electrophoresis* 1996, **17**:1787-1796.

30. van Driel IR, Goldstein JL, Südhof TC and Brown MS: **First cysteine-rich repeat in ligand-binding domain of low density lipoprotein receptor binds  $\text{Ca}^{2+}$  and monoclonal antibodies, but not lipoproteins.** *J Biol Chem* 1987, **262**:17443-17449.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

