

Tengchuan Jin,^a Andrew Howard,^a Erica A. Golemis,^b Yingtong Wang^a and Yu-Zhu Zhang^{a*}

^aDepartment of Biology, Illinois Institute of Technology, Chicago, IL 60616, USA, and

^bFox Chase Cancer Center, Philadelphia, PA 19111, USA

Correspondence e-mail: yuzhu.zhang@iit.edu

Received 6 March 2005

Accepted 19 April 2005

Online 28 April 2005

Overproduction, purification, crystallization and preliminary X-ray diffraction studies of the human transcription repressor ERH

The human gene coding for the enhancer of rudimentary homologue (ERH) protein was overexpressed in *Escherichia coli*. The ERH protein was purified by anion-exchange, hydrophobic interaction and gel-filtration chromatography. Well diffracting single crystals were obtained by the vapour-diffusion method in hanging drops. The crystals belong to the trigonal space group $P3_121$ or its enantiomorph $P3_221$, with unit-cell parameters $a = b = 53.74$, $c = 67.45$ Å, $\alpha = \beta = 90$, $\gamma = 120^\circ$. They diffract to at least 1.75 Å. A selenomethionine derivative of the protein was prepared and crystallized for multiwavelength anomalous diffraction (MAD) phasing.

1. Introduction

Regulation of gene expression at the transcriptional level involves a well controlled balance between transcription factors and their positive and negative cofactors (reviewed in Kadonaga, 2004; Martinez, 2002). The enhancer of rudimentary homologue (ERH) gene encodes the ERH protein, which is evolutionarily highly conserved (Gelsthorpe *et al.*, 1997; Wojcik *et al.*, 1994). ERH was first found as a genetic interactor of the *Drosophila* rudimentary gene, which encodes a protein with enzymatic activities in the pyrimidine-biosynthesis pathway (Jarry & Falk, 1974; Norby, 1973; Rawls & Fristrom, 1975). Subsequently, RNAi experiments in *Caenorhabditis elegans* showed that disruption of ERH resulted in an embryonic lethal phenotype (Gonczy *et al.*, 2000; <http://worm-srv1.mpi-cbg.de/dbScreen/>). Thus, ERH was believed to have evolutionarily conserved functions associated with a basic cellular process. Recently, in a two-hybrid study aiming at identifying protein interactors with the dimerization cofactor of HNF1/pterin-4-carbinolamine dehydratase (DCoH/PCD), Pogge von Strandmann *et al.* (2001) found that the *Xenopus* ERH may interfere with HNF-1-dependent gene regulation through its interaction with DCoH/PCD and proposed ERH to be a cell-type-specific transcriptional repressor. Here, we describe the details of the overexpression, purification and crystallization of human ERH, as well as the preliminary crystallographic data.

2. Methods, results and discussion

2.1. Cloning, expression and purification

The coding region of an ERH cDNA was PCR-amplified from a HeLa cDNA library. The PCR was performed with CCCCAG-ATCCATATGTCTCACACCATTTTGCTGGTA and CCCCTC-GAGGATCCTTATTTCCCAGCCTGTTGGGCC as 5' and 3' primers, respectively. The PCR product was cloned into the pBlue-script II SK vector *via* an *EcoRI* and an *XhoI* site introduced by the 5' and 3' primer, respectively, to create pBlueERH. The ERH coding region was then released from pBlueERH with *NdeI* and *XhoI* and cloned into the vector pET29b to create pET29ERH. The insert of pET29ERH was confirmed by sequencing.

Escherichia coli BL21(DE3) cells (Novagen, Madison, WI, USA) transformed with pET29ERH were grown in 1 l of LB medium



© 2005 International Union of Crystallography
All rights reserved

containing 50 mg l⁻¹ kanamycin at 310 K. At an OD₆₀₀ of 1.2, expression of ERH was induced by adding IPTG to the medium to a final concentration of 1 mM. The cells were collected by centrifugation after 4 h induction. An SDS-PAGE analysis of the total cell lysate showed an intense band with the expected molecular weight for ERH.

For purification, the cell pellet was resuspended in 50 ml buffer A (10 mM phosphate pH 7.9, 5 mM DTT) plus a protease-inhibitor cocktail. The cocktail contained aprotinin, antipain, leupeptin and pepstatin. Their final concentrations were 100 nM, 50 μM, 50 μM and 0.5 μg ml⁻¹, respectively. Cells were sonicated for 10 min in a beaker surrounded by a mixture of ice and water. The cell extract was subjected to centrifugation at 25 000g for 30 min at 277 K. The supernatant was filtered with 0.20 μm syringe filters and the sample was loaded onto a Source 15Q (Pharmacia, now GE Healthcare, Piscataway, NJ, USA) ion-exchange column with a bed volume of 10 ml, pre-equilibrated with buffer A. The column was washed with three bed volumes of buffer A and then run with buffer A and a linear 0–0.5 M NaCl gradient (five bed volumes) with a flow rate of 1 ml min⁻¹. The ERH protein eluted at 0.12 M NaCl. ERH-containing fractions were collected and loaded onto two coupled 5 ml Hitrap phenyl HP columns (Pharmacia) after adding an equal volume of 2 M (NH₄)₂SO₄. After washing with five bed volumes of buffer A plus 1 M (NH₄)₂SO₄, ERH was eluted with a linear 1–0 M (NH₄)₂SO₄ gradient (20 bed volumes) with a flow rate of 1 ml min⁻¹. ERH eluted at 0.26 M (NH₄)₂SO₄ and the ERH-containing fractions were concentrated to ~5 ml using an Ultrafree-5K centrifugal filter device (Millipore, MA, USA). The concentrated sample was loaded onto a Superdex75 gel-filtration column (diameter 26 mm, height 65 cm). The column was pre-equilibrated and eluted with buffer A at a flow rate of 1 ml min⁻¹. ERH eluted from the gel column with an apparent molecular weight close to that of an ERH tetramer, as assessed by comparison with gel-filtration runs of standard proteins. Gel filtration of ERH was also performed at lower pH using the above Superdex75 column. At pH 5.6, at which ERH was crystallized (see below), ERH eluted from the gel column with an apparent molecular weight close to that of an ERH monomer. This procedure yielded approximately 20 mg of homogeneous ERH from a 1 l culture.

All chromatographic steps were carried out at 277 K using an FPLC system (Pharmacia). The molecular weight of the pure protein was determined to be 12 127 Da by electrospray mass spectrometry, consistent with the ERH protein lacking the N-terminal methionine.




Figure 1
Crystal of ERH obtained by vapour diffusion in hanging drops.

Table 1
X-ray data-collection statistics.

Values in parentheses are for the outer shell.

Resolution (Å)	50–1.75 (1.81–1.75)
Wavelength (Å)	1.0
Space group	<i>P</i> ₃ <i>2</i> ₁ or <i>P</i> ₃ <i>2</i> ₁
Unit-cell parameters	
<i>a</i> (Å)	53.74
<i>b</i> (Å)	53.74
<i>c</i> (Å)	67.45
α (°)	90
β (°)	90
γ (°)	120
Data-collection temperature (K)	110
No. of observed reflections	124880
No. of unique reflections	11671
Completeness (%)	99.9 (100)
$\langle I/\sigma(I) \rangle$	46.7 (9.7)
R_{sym}^{\dagger} (%)	4.5 (25.1)

$$\dagger R_{\text{sym}} = \sum |I - \langle I \rangle| / \sum I.$$

2.2. Crystallization

The pure protein was desalted and concentrated to 90 mg ml⁻¹ in Milli-Q H₂O using an Ultrafree-5K centrifugal filter device. The sample was filtered with a 0.2 μm filter and crystallization experiments were performed at 277 K in Linbro plates using the hanging-drop vapour-diffusion technique. An initial crystallization screen was performed using Hampton Research Crystal Screen and Crystal Screen Lite kits. 1 μl protein sample was mixed with 1 μl reservoir solution and sealed against 0.5 ml reservoir solution. All reservoir solutions contained 5 mM DTT and 0.02% (w/v) sodium azide. Several crystals appeared in the well containing solution No. 40 of the Crystal Screen Lite kit [0.1 M trisodium citrate dihydrate pH 5.6, 10% (v/v) isopropanol, 10% (w/v) polyethylene glycol 4000] in 3 d. Single crystals of dimensions ~0.2 × 0.05 × 0.05 mm were obtained within a month using this condition (Fig. 1). No crystals were obtained at 295 K and crystals grown at 277 K redissolved in 10 min after being moved to 295 K. Currently, we are still screening for conditions to grow ERH crystals at higher pH values where ERH would assume a tetramer in solution.

2.3. X-ray diffraction experiments and crystal characterization

To collect data at low temperature, single crystals were transferred with nylon loops to a cryoprotectant, which was a mixture of an equal volume of the reservoir solution and 50% (v/v) ethylene glycol solution. The crystals were then picked up with nylon loops and flash-frozen in liquid nitrogen. Data were collected at IMCA-CAT beamline 17-ID (Advanced Photon Source, US Argonne National Laboratory) equipped with an ADSC Quantum 210 detector. A complete data set of 1° frames with 4 s exposures was collected at 12 398 eV. The ERH crystals diffracted to 1.75 Å and the data were processed with the *HKL2000* suite of programs (Otwinowski & Minor, 1997) and *X-GEN* (Howard, 2000), revealing a trigonal crystal system with unit-cell parameters $a = b = 53.74$, $c = 67.45$ Å, $\alpha = \beta = 90$, $\gamma = 120$ °. From systematic absences in specific reflections in the diffraction, the space group was determined to be *P*₃*2*₁ or its enantiomorph *P*₃*2*₁ (Table 1). Assuming one molecule in the asymmetric unit, the Matthews coefficient V_M was 2.32 Å³ Da⁻¹, corresponding to a solvent content of 47%.

2.4. Purification and crystallization of selenomethionine ERH

To facilitate multiwavelength anomalous diffraction (MAD) phasing, the met⁻ *E. coli* strain B834(DE3) (Novagen, Madison, WI,

USA) transformed with pET29ERH was grown in 1 l M9 medium supplemented with 0.15 mM thiamine and 50 mg L-methionine at 310 K in the presence of 50 mg l⁻¹ kanamycin. When the OD₆₀₀ of the culture reached 1.0, cells were spun down by centrifugation and resuspended in 1 l of the above medium devoid of methionine. After 30 min, 15 mg L-selenomethionine was added to the culture. After an additional 30 min, induction of selenomethionine ERH expression was initiated by adding IPTG to a final concentration of 1 mM. Induction was allowed to continue for 4 h. Purification of the selenomethionine ERH was identical to that of the native protein. The incorporation of selenomethionine was confirmed by electrospray mass spectrometry. Assuming equal efficiency of selenomethionine incorporation at all three sites, the success rate of selenomethionine labelling was determined to be 70%. Selenomethionine ERH crystals were grown under conditions identical to those for growing native ERH crystals. Selenomethionine ERH single crystals were obtained within a month that were morphologically identical and grew to the same size as the native ERH crystals. Selenomethionine ERH crystals diffracted equally well and had the same unit-cell parameters as crystals of natural ERH. For selenomethionine ERH, a complete data set has been collected at the IMCA-CAT beamline 17-ID and attempts to solve the structure of ERH by multiwavelength anomalous diffraction (MAD) methods are currently under way.

Use of the IMCA-CAT beamline is supported by the companies of the Industrial Macromolecular Crystallography Association through a contract with Illinois Institute of Technology. Use of the Advanced Photon Source was supported by US DOE, Office of Science, Office of Basic Energy Sciences under Contract No. W-31-109-Eng-38.

References

- Gelsthorpe, M., Pulumati, M., McCallum, C., Dang-Vu, K. & Tsubota, S. I. (1997). *Gene*, **186**, 189–195.
- Gonczy, P. *et al.* (2000). *Nature (London)*, **408**, 331–336.
- Howard, A. J. (2000). In *Crystallographic Computing 7*, edited by P. E. Bourne & K. D. Watenpaugh. Oxford University Press.
- Jarry, B. & Falk, D. (1974). *Mol. Gen. Genet.* **135**, 113–122.
- Kadonaga, J. T. (2004). *Cell*, **116**, 247–257.
- Martinez, E. (2002). *Plant Mol. Biol.* **50**, 925–947.
- Norby, S. (1973). *Hereditas*, **73**, 11–16.
- Otwinowski, Z. & Minor, W. (1997). *Methods Enzymol.* **276**, 307–326.
- Pogge von Strandmann, E., Senkel, S. & Ryffel, G. U. (2001). *Biol. Chem.* **382**, 1379–1385.
- Rawls, J. M. & Fristrom, J. W. (1975). *Nature (London)*, **255**, 738–740.
- Wojcik, E., Murphy, A. M., Fares, H., Dang-Vu, K. & Tsubota, S. I. (1994). *Genetics*, **138**, 1163–1170.