

Common Functions for Diverse Small RNAs of Land Plants

Michael J. Axtell,^{a,1} Jo Ann Snyder,^a and David P. Bartel^{b,c}

^aDepartment of Biology and Huck Institutes of the Life Sciences, Pennsylvania State University, University Park, Pennsylvania 16802

^bWhitehead Institute, Howard Hughes Medical Institute, Cambridge, Massachusetts 02142

^cDepartment of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02142

Endogenous small RNAs, including microRNAs (miRNAs) and short interfering RNAs (siRNAs), are critical components of plant gene regulation. Some abundant miRNAs involved in developmental control are conserved between anciently diverged plants, while many other less-abundant miRNAs appear to have recently emerged in the *Arabidopsis thaliana* lineage. Using large-scale sequencing of small RNAs, we extended the known diversity of miRNAs in basal plants to include 88 confidently annotated miRNA families in the moss *Physcomitrella patens* and 44 in the lycopod *Selaginella moellendorffii*. Cleavage of 29 targets directed by 14 distinct *P. patens* miRNA families and a *trans*-acting siRNA (ta-siRNA) was experimentally confirmed. Despite a core set of 12 miRNA families also expressed in angiosperms, weakly expressed and apparently lineage-specific miRNAs accounted for the majority of miRNA diversity in both species. Nevertheless, the molecular functions of several of these lineage-specific small RNAs matched those of angiosperms, despite dissimilarities in the small RNA sequences themselves, including small RNAs that mediated negative feedback regulation of the miRNA pathway and miR390-dependent ta-siRNAs that guided the cleavage of *AUXIN RESPONSE FACTOR* mRNAs. Diverse, lineage-specific, small RNAs can therefore perform common biological functions in plants.

INTRODUCTION

Most eukaryotic organisms are capable of producing small RNAs, typically between 21 and 24 nucleotides in length, that serve as sequence-specific posttranscriptional regulators. Many short interfering RNAs (siRNAs) are initially produced as short duplexes containing two-nucleotide, 3' overhangs via the enzymatic activity of endonucleases of the Dicer family acting upon long double-stranded RNA (dsRNA) (Baulcombe, 2005). One of the two siRNA strands is stably incorporated into an ARGONAUTE (AGO) protein that then functions to repress the expression of target RNAs, which are selected via base-pairing interactions with the siRNA (Hammond, 2005). This repression can take several forms, including AGO-catalyzed target cleavage (Liu et al., 2004; Meister et al., 2004) and translational repression (Doench et al., 2003). Some endogenous siRNAs also direct DNA and histone modifications to homologous genomic loci (Zilberman et al., 2003). Such siRNAs are especially numerous and diverse in flowering plants (Lu et al., 2005; Rajagopalan et al., 2006; Kasschau et al., 2007).

MicroRNAs (miRNAs) are a distinct type of small RNA that also function as negative regulators of gene expression. Primary


miRNA transcripts (pri-miRNAs) adopt stem-loop RNA secondary structures from which a specific ~21-nucleotide duplex is excised by a Dicer endonuclease (Bartel, 2004). Similar to siRNA duplexes, the resulting miRNA/miRNA* duplex possesses characteristic two-nucleotide 3' overhangs and is unwound coincident with the association of the mature, single-stranded miRNA with an AGO protein. The mature miRNA then serves to guide the bound AGO protein to target mRNAs based upon complementarity. In both plants and animals, the regulation of target gene expression via miRNAs is critical for numerous biological processes (Wienholds and Plasterk, 2005; Jones-Rhoades et al., 2006; Mallory and Vaucheret, 2006). In plants, miRNA-directed target cleavage can sometimes stimulate the production of dsRNA using the cleaved product as a template; the resulting dsRNA is subsequently processed by a Dicer protein to produce siRNAs referred to as *trans*-acting siRNAs (ta-siRNAs) because of their miRNA-like ability to repress mRNAs from loci distinct from the originating locus (Peragine et al., 2004; Vazquez et al., 2004; Allen et al., 2005; Yoshikawa et al., 2005; Axtell et al., 2006).

Although the miRNAs characterized in angiosperms have numerous distinct biological functions, there is a striking tendency for many of them to regulate developmental processes (Jones-Rhoades et al., 2006). Previous studies have shown that some miRNA–target interactions first identified in *Arabidopsis thaliana* are deeply conserved among land plants (Floyd and Bowman, 2004; Arazi et al., 2005; Axtell and Bartel, 2005). Interestingly, the ancient miRNA regulatory interactions are disproportionately those thought to be involved in development, suggesting that these units of posttranscriptional gene control have been indispensable during the diversification of land plants.

¹To whom correspondence should be addressed. E-mail mja18@psu.edu.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Michael J. Axtell (mja18@psu.edu).




www.plantcell.org/cgi/doi/10.1105/tpc.107.051706

The sequencing of a moss (*Physcomitrella patens*) genome and a lycopod (*Selaginella moellendorffii*) genome provides an opportunity to explore the diversity of small RNA functions in nonseed plants and to more rigorously address the trajectories of small RNA evolution in three widely separated plant lineages. To date, >40 distinct miRNA families have been annotated from *P. patens* based upon analysis of sequenced small RNAs or bioinformatic predictions (Arazi et al., 2005; Talmor-Neiman et al., 2006a; Fattash et al., 2007). In lycopods, the expression of miR166 has been confirmed in *S. moellendorffii* (Floyd and Bowman, 2004), and the expression of two additional families proposed based upon microarray experiments using RNA from *Selaginella uncinata* (Axtell and Bartel, 2005).

In this study, we applied high-throughput small RNA sequencing to the discovery and analysis of miRNAs from the moss *P. patens* and the lycopod *S. moellendorffii*. Our results directly demonstrated the expression of a much larger number of distinct miRNA families in *P. patens* and *S. moellendorffii* than has previously been reported. Only a few of these miRNAs were expressed in common between mosses, lycopods, and angiosperms; these core conserved plant miRNAs were abundantly expressed in both angiosperms and basal plants, and all, save one, appeared to be involved in the regulation of developmental processes. The majority of miRNA diversity in both lycopods and mosses appeared lineage specific and generally expressed at lower levels than were the core conserved miRNAs. The experimental isolation of 29 cleavage products directed by 14 distinct *P. patens* miRNAs and a ta-siRNA demonstrated that the core conserved plant miRNAs have generally retained homologous target interactions during the diversification of land plants. Surprisingly, experimentally validated targets also included several instances in which lineage-specific *P. patens* miRNAs and ta-siRNAs shared common functions with sequence-distinct *Arabidopsis* small RNAs.

RESULTS

Identification of miRNAs from *P. patens* and *S. moellendorffii*

Direct sequencing of small RNAs and computational predictions have resulted in the annotation of 44 distinct *P. patens* miRNA families (Arazi et al., 2005; Talmor-Neiman et al., 2006a; Fattash et al., 2007). We have recently reported a data set of 127,135 unique *P. patens* small RNAs from three libraries, represented by a total of 561,102 reads, which provided insights into siRNA biogenesis in plants (Axtell et al., 2006). Because of the >100-fold increase in sequencing depth over previous *P. patens* small RNA sequencing efforts, we reasoned that additional less-abundant miRNAs might be found by analysis of these data. In addition, a single small RNA library prepared from the above-ground tissues of *S. moellendorffii* was sequenced to yield 36,582 unique small RNAs, represented by 149,586 reads that matched the genome (whole-genome shotgun traces).

To produce a list of possible miRNA loci, consensus genomic sequences surrounding each small RNA were derived by assembly of the relevant whole-genome shotgun (WGS) traces,

followed by RNA secondary structure prediction. As a preliminary screen, the minimum free energy structure for each locus was analyzed using MIRcheck (Jones-Rhoades and Bartel, 2004), which evaluates paired regions of a predicted RNA secondary structure for characteristic features of miRNA stem loops. The candidate miRNA loci that passed MIRcheck were then analyzed for evidence of miRNA* accumulation. A miRNA/miRNA* duplex, with two-nucleotide 3' overhangs indicative of DICER-LIKE1 (DCL1) cleavage, would be expected if a moss DCL1 ortholog excised the miRNA from the hairpin precursor as indicated by a previous study (Talmor-Neiman et al., 2006a). Indeed, the sequencing of both a miRNA and a miRNA* constitutes evidence of DCL1-like processing from a hairpin precursor, which is evidence that can supplement, or even substitute for, other criteria for confidently distinguishing miRNAs from endogenous siRNAs and other RNAs that should not be annotated as miRNAs (Rajagopalan et al., 2006; Ruby et al., 2006; Fahlgren et al., 2007).

In total, 88 distinct *P. patens* miRNA families, arising from 205 miRNA genes, were recovered by this analysis (Tables 1 and 3; see Supplemental Table 1 online). This set included all previously recognized *P. patens* miRNA families except for 13 of the 22 families uniquely reported by Fattash et al. (2007). Analysis of the deep sequencing data from *P. patens* also suggested refinements to a small number of previous miRNA annotations; these suggestions included reversals of miRNA/miRNA* designations, changes in the annotated positions of the dominant mature miRNA within stem-loop precursors, and consolidations of redundantly reported families (see Supplemental Table 1 online). Our resulting roster of confidently identified *P. patens* miRNAs (see Supplemental Table 1 online) included 12 families conserved with angiosperms and/or lycopods (Table 3). An additional six families of *P. patens* miRNAs annotated on the basis of computational comparisons to angiosperm miRNAs (miR167, miR172, miR395, miR414, miR418, and miR419; Fattash et al., 2007) were not observed in our sequence libraries. The expression of the remaining 76 sequence-distinct families was, to the best of our current knowledge, bryophyte specific. These included 57 novel *P. patens* miRNA families that have not previously been reported (Table 1).

In *S. moellendorffii*, 44 distinct miRNA families were found to originate from 58 miRNA genes (see Supplemental Table 2 online), including miR156, miR159/319, and miR166, as expected (Floyd and Bowman, 2004; Axtell and Bartel, 2005), as well as five additional deeply conserved miRNA families (miR160, miR171, miR396, miR408, and miR536; Table 3). However, the majority of *S. moellendorffii* miRNA diversity, comprising 36 novel families, was to the best of our present knowledge restricted to the lycopod lineage (Table 2). The deeply conserved *S. moellendorffii* miRNAs were often encoded by fewer paralogs than in other species (Table 3), and most of the novel *S. moellendorffii* miRNAs appeared to arise from only a single locus (Table 2). This reduction in paralogs was consistent with the reduced genome size of *S. moellendorffii* compared with most land plants with fully sequenced genomes (Wang et al., 2005). Collectively, the results from *P. patens* and *S. moellendorffii* were consistent with recent results from angiosperms, which demonstrated a large diversity of recently evolved miRNAs expressed at low levels that only became apparent after deep

Table 1. Novel *P. patens* miRNAs

miRNA	Sequence (5'–3') ^b	Loci	Length	Normalized Abundance (TPQ) ^a			Arm ^c
				P	PG	GS	
ppt-miR1023(5')	aCACUCUCUCCaUUUCuCUaC	4	21	513 (1696)	4341 (12373)	392 (929)	Amb
ppt-miR1023(3')	AGaGAAUuGgAgAGAGUGcAu		21	983 (1696)	6666 (12373)	328 (929)	Amb
ppt-miR1024	UCUGGUUGGAUUGUAGGCCUC	2	21	1102 (1258)	1236 (1744)	841 (1157)	3'
ppt-miR1025	UGCCACAACAAAGCUAAUAAC	1	21	8486 (9486)	8690 (10624)	9370 (10954)	3'
ppt-miR1026	UGAGAAAGACUUGAGAGGACA	2	21	480 (534)	174 (187)	42 (44)	5'
ppt-miR1027	UUUCUAUCUUCUCUCCAAUC	2	21	102 (383)	129 (503)	156 (406)	5'
ppt-miR1028(5')	UCUUAGAUCUACAAGuCaCa	3	21	934 (1744)	235 (582)	933 (3607)	Amb
ppt-miR1028(3')	uGgCAUUGuGguuUaAGagC		21	491 (1744)	216 (582)	2437 (3607)	Amb
ppt-miR1029	UCUCUCUCAACCAACCAUAC	1	20	16 (113)	31 (158)	104 (234)	3'
ppt-miR1030	uCUGCAUCUGCACCCUGCACCa	10	21	64 (367)	35 (220)	140 (638)	5'
ppt-miR1031	UCAAUGCUCUCUGGAGCUUCU	2	21	421 (491)	233 (264)	458 (490)	5'
ppt-miR1032	AGGUGACUGCCUGGAAUUGGG	1	21	86 (118)	147 (176)	174 (222)	5'
ppt-miR1033	UGACGGGUCGUGAUGGGCACU	5	21	5 (10)	24 (239)	276 (767)	3'
ppt-miR1034	UUACUUUGGCAGCGCUGUGCU	1	21	0 (5)	22 (37)	98 (188)	3'
ppt-miR1035	CGUCUUAGCCACAAAACGAA	1	21	54 (91)	74 (143)	94 (98)	5'
ppt-miR1036(5')	UGUGGAGUCCGUAAUJAGCUG	1	21	16 (54)	18 (49)	14 (42)	Amb
ppt-miR1036(3')	AGCUAAUUAAGGAUUCUACAC		21	21 (54)	12 (49)	4 (42)	Amb
ppt-miR1037	AGCCUUUUJAGGAUUUGAUGG	1	21	0 (0)	4 (12)	14 (24)	3'
ppt-miR1038(5')	UAGGUGCGUUUCACCAAAG	1	20	37 (86)	22 (70)	40 (94)	Amb
ppt-miR1038(3')	CAUGGUGGAAUCGCAUCCAGG		21	27 (86)	8 (70)	30 (94)	Amb
ppt-miR1039(5')	UCUUUGGGUCUUUCUCUCCUG	1	21	0 (5)	10 (20)	2 (4)	Amb
ppt-miR1039(3')	GGGGAGACGGCUCAAGGAUC		21	0 (5)	6 (20)	0 (4)	Amb
ppt-miR1040	UGAACGCAAAUGAACAUUUC	1	21	21 (48)	37 (52)	40 (50)	3'
ppt-miR1041	UUUUCGGGUGAGAGUGGUCCU	1	21	16 (54)	2 (4)	6 (22)	5'
ppt-miR1042(5')	GUCUGGCAGGAACUAGAUAGG	1	21	0 (0)	4 (8)	0 (4)	Amb
ppt-miR1042(3')	UGUCUAGUCUCUCCACGGCCCG		22	0 (0)	2 (8)	2 (4)	Amb
ppt-miR1043(5')	CCUUCACUUAUCUGCGUGCA	1	20	0 (5)	8 (14)	0 (18)	Amb
ppt-miR1043(3')	UUGCGCGUGAAUUUGAAGGCU		21	0 (5)	4 (14)	10 (18)	Amb
ppt-miR1044(5')	GCACAAAUUGCAGCACUACAAAC	1	21	0 (0)	16 (47)	0 (8)	Amb
ppt-miR1044(3')	UUGUAGUGCAUUAUUUGUUUU		20	0 (0)	16 (47)	0 (8)	Amb
ppt-miR1045	AAGUGCUGGCUUUUUGACGUU	1	21	0 (0)	20 (37)	64 (196)	5'
ppt-miR1046(5')	UGGAUUUCAUUUUUUCACG	1	20	16 (54)	8 (14)	6 (46)	Amb
ppt-miR1046(3')	UGGUGAAAAAUUGAAAAUUC		21	5 (54)	2 (14)	12 (46)	Amb
ppt-miR1047(5')	CUUGAUUACAAAGGCCUUAAU	1	21	5 (27)	12 (62)	14 (72)	Amb
ppt-miR1047(3')	UGAUCAGCGAUGACCUAGUUG		21	10 (27)	22 (62)	42 (72)	Amb
ppt-miR1048(5')	UAGAACAUGAGUGUAGACGAC	1	21	27 (59)	14 (29)	6 (14)	Amb
ppt-miR1048(3')	UGUCUACACUCAUGUUCUAGA		21	16 (59)	12 (29)	2 (14)	Amb
ppt-miR1049	UCUCUCUUAGCCAAACAGUCU	1	21	108 (124)	10 (18)	2 (2)	3'
ppt-miR1050	UGACCACUUGAUUCCGGCCU	1	21	10 (59)	60 (114)	474 (661)	5'
ppt-miR1051(5')	GGUUCAGUGAACAGGAAGA	1	20	0 (5)	27 (47)	2 (24)	Amb
ppt-miR1051(3')	UUCCUGUCUACUUGAAGCCAC		21	0 (5)	14 (47)	20 (24)	Amb
ppt-miR1052	UUCCUUUUUJUGAUUGUGGUA	1	21	189 (216)	647 (907)	372 (470)	3'
ppt-miR1053(5')	GAUGGGUUUUCUCAAAGUGG	1	20	0 (0)	2 (4)	0 (2)	Amb
ppt-miR1053(3')	CUCUACUUUGACAUAGCCACU		22	0 (0)	0 (4)	2 (2)	Amb
ppt-miR1054	UAAACCCUCUCUCUUAUCCUG	1	21	43 (59)	79 (118)	288 (326)	3'
ppt-miR1055	UUAGGGGUGUUUCCAGUGACU	1	22	5 (5)	27 (56)	0 (6)	3'
ppt-miR1056	UGGAUCUUUGCAUCAUAACCC	1	21	37 (102)	27 (52)	10 (50)	3'
ppt-miR1057	UUUUUGUGUUCUAGCUGGGGU	1	22	0 (0)	0 (2)	4 (8)	3'
ppt-miR1058	AGAGAUUCCAUCACGAAGCAC	1	21	10 (32)	27 (93)	174 (252)	3'
ppt-miR1059	UGAAAGUCCUUCACAACAAC	1	21	81 (86)	108 (120)	62 (64)	5'
ppt-miR1060	UUUGUCAUAGGAUUACACACG	1	21	16 (21)	4 (8)	8 (10)	5'
ppt-miR1061(5')	UUUUUUCAUAGACUACUAAA	1	21	5 (5)	2 (4)	2 (6)	Amb
ppt-miR1061(3')	UGAGUAGUCAUUGGAUUAUG		21	0 (5)	2 (4)	2 (6)	Amb
ppt-miR1062	UCCUCACGUGGUGUUUGCAGC	1	22	5 (5)	10 (18)	16 (18)	5'
ppt-miR1063	CAUCUUGGAGUACUGCAUCUU	8	21	378 (615)	235 (345)	34 (62)	5'
ppt-miR1064(5')	CAUUCUGAGAAUAUCAUCCG	1	21	0 (10)	0 (4)	4 (4)	Amb
ppt-miR1064(3')	GGGUGAUUUUUCUCAGAGUUC		21	5 (10)	2 (4)	0 (4)	Amb

(Continued)

Table 1. (continued).

miRNA	Sequence (5'–3') ^b	Loci	Length	Normalized Abundance (TPQ) ^a			Arm ^c
				P	PG	GS	
ppt-miR1065	ACAGUCUCUGACUUCUCGCAG	1	21	0 (10)	8 (18)	218 (296)	5'
ppt-miR1066	ACAUGUUGCAGAGCGGGGUAC	1	21	16 (64)	22 (58)	46 (112)	3'
ppt-miR1067	ACAUACUGAAGUUUGAUGCCA	1	21	27 (32)	16 (29)	2 (6)	5'
ppt-miR1068	UAGCCAUUUGCUUGAAGGUCA	1	21	21 (27)	10 (14)	76 (92)	5'
ppt-miR1069(5')	UCUCAAGUCUUUCAUUGGAU	1	21	5 (10)	4 (14)	0 (4)	Amb
ppt-miR1069(3')	UGAUAAAUCAAAGUGCUCACU		21	5 (10)	2 (14)	4 (4)	Amb
ppt-miR1070	UCGGUUUCUAGUAAAACUUGC	1	21	21 (21)	10 (16)	0 (0)	5'
ppt-miR1071(5')	GCGCAAUUCCCGAACGGGUAG	1	21	5 (10)	0 (0)	0 (0)	Amb
ppt-miR1071(3')	ACCCGUUCUGGAAUUGCGCGG		21	5 (10)	0 (0)	0 (0)	Amb
ppt-miR1072	UGCAUUGUGUUUUUGAAGCUUGA	1	24	43 (54)	37 (77)	56 (108)	3'
ppt-miR1073(5')	UGAAUGAUUUACGUCCACG	1	21	189 (659)	239 (1041)	138 (418)	Amb
ppt-miR1073(3')	UGGGCGUUAAUACAUUUCUAU		21	383 (659)	587 (1041)	232 (418)	Amb
ppt-miR1074	AGGGUUGUUAGUUGUGUUGAU	1	21	70 (232)	66 (154)	20 (106)	5'
ppt-miR1075	UGUUUCAGUCAUGGUUUCUAC	1	21	48 (81)	20 (49)	18 (24)	3'
ppt-miR1076(5')	UAUCACAAGGUGCGAUAAUUG	1	21	27 (172)	56 (331)	20 (224)	Amb
ppt-miR1076(3')	CCAAGCACUUUUCGCACCCUG		21	21 (172)	45 (331)	60 (224)	Amb
ppt-miR1077(5')	UUCAAAUGUUCGGAUGCGGUGC	1	21	0 (0)	0 (6)	8 (24)	Amb
ppt-miR1077(3')	AGGGAUGGACGUUACGAGCC		21	0 (0)	4 (6)	10 (24)	Amb
ppt-miR1078	UUUGGAUGAUUCAUUUGUGAU	1	21	91 (124)	160 (195)	28 (36)	5'
ppt-miR1079(5')	CACCGAGCUCUUCUUUUUGGAGG	1	23	0 (5)	2 (6)	0 (0)	Amb
ppt-miR1079(3')	ACCAAAGAAUAGCUAUGUGGU		22	0 (5)	4 (6)	0 (0)	Amb

^a To enable comparison between samples, the number of reads corresponding to the mature miRNA and all reads corresponding to the miRNA hairpin (in parentheses) were converted into transcripts per quarter million (TPQ) by dividing the raw number of reads by the total number of reads that matched the Phypa1_1 genome assembly, multiplying by 250,000, and rounding to the nearest integer. P, protonemata library (GSM115095); PG, protonemata and young gametophores library (GSM115096); GS, mature gametophores and sporophytes library (GSM115097).

^b Sequence of the most abundant mature miRNA is shown, with lowercase letters indicating positions where the sequence of less-abundant paralogs differ.

^c Arm of stem loop where the mature miRNA resides. Amb, ambiguous.

sequencing of small RNA libraries (Rajagopalan et al., 2006; Fahlgren et al., 2007).

A few plant pri-miRNA transcripts have been observed to produce more than one miRNA/miRNA* duplex, including *Arabidopsis* miR161 (Allen et al., 2004) and miR163 (Kurihara and Watanabe, 2004) as well as *P. patens* miR319d and miR1219a (Talmor-Neiman et al., 2006a; Fattash et al., 2007). Certain miRNAs from the unicellular green alga *Chlamydomonas reinhardtii* also produce multiple miRNA/miRNA* duplexes from a single pri-miRNA transcript (Molnár et al., 2007; Zhao et al., 2007). We found that two novel *S. moellendorffii* pri-miRNAs produced two distinct miRNA/miRNA* duplexes, in both cases separated from each other by distances that were rough multiples of 21 (see Supplemental Figure 1A and Supplemental Table 2 online). This spatial separation was consistent with sequential DCL1-mediated processing of the pri-miRNA, as reported for *Arabidopsis* miR163 (Kurihara and Watanabe, 2004), although independent recognition and processing of the two miRNA/miRNA* duplexes could not be excluded. Production of three sequential miRNA/miRNA* duplexes is a conserved feature of plant miR159 and miR319 loci (see Supplemental Figure 1B and Supplemental Tables 1 and 2 online; Rajagopalan et al., 2006; Talmor-Neiman et al., 2006a; Fattash et al., 2007). The previously annotated mature miR159 and miR319 miRNAs are the best

conserved of these sequences, likely reflecting their conserved functions in targeting messages in *trans*. However, as noted previously by comparing *Arabidopsis* and *Oryza sativa* miR159 and miR319 pri-miRNAs (Li et al., 2005), the loop-proximal alternative miRNA/miRNA* duplex was also relatively well conserved among land plants (see Supplemental Figure 1B online). Although some potential targets could be predicted for the *Arabidopsis* and *P. patens* alternative miR319 species, we were not able to verify them using RNA ligase-mediated 5' rapid amplification of cDNA ends (RLM 5'-RACE; Llave et al., 2002a). Thus, these data are inconclusive as to whether any of these conserved alternative miR319 miRNAs are in fact incorporated into a functional silencing complex.

***P. patens* miRNA Loci Are Often Clustered and Frequently Overlap with Protein-Coding Genes**

The availability of a draft assembly and transcript annotations of the *P. patens* genome allowed the detailed examination of miRNA loci. Most miRNA loci characterized in angiosperms appear to arise from independent transcription units, each encoding a single miRNA-producing stem loop that does not overlap with annotated protein-coding loci (Jones-Rhoades et al., 2006). Four *P. patens* miR1219 loci are contained in two clusters, strongly suggesting that they are

Table 2. Novel *S. moellendorffii* miRNAs

miRNA	Sequence (5'–3') ^b	No. of Loci	Length	Abundance ^a		Arm ^c
				miRNA	Total	
smo-miR1080	UUCACUAUCUGCAAACACCCUCU	1	22	79	110	3'
smo-miR1081	UGAGGCUUGCCUUGAUUCUC	1	21	5001	5469	5'
smo-miR1082	GcGUUUGGCCUGCUGGCCGGuG	2	21	6188	8320	5'
smo-miR1083	UAGCCUGGAACGAAGCACGGU	1	21	5	8	5'
smo-miR1084	UAACUCAGGUGGUUAGUUCCCA	1	22	1346	1430	5'
smo-miR1085(5')	CCCCUACAAGAGAUUCAAUUC	1	21	1	3	Amb
smo-miR1085(3')	UUGAUUUGUCUUGUAGGGAC	1	20	1	3	Amb
smo-miR1086	UAGUGCCGUGGUCCUUUGGC	1	21	19	22	5'
smo-miR1087	JACAGAUUGGUGUJAGUGCUU	1	21	5	11	3'
smo-miR1088(5')	CAGAAAGAAAGAGAGCACGCAU	1	21	22	56	Amb
smo-miR1088(3')	GCGUGCUCUUUUUCUUCUGUC	1	21	29	56	Amb
smo-miR1089	AGAUCAUUCJAGGAUUGUUUGC	1	21	12	18	3'
smo-miR1090	UGAAAGCCAUUCUCAACAAA	1	21	7	11	5'
smo-miR1091	CGGCAGUGAGGGAGGAUUUGC	1	21	191	294	5'
smo-miR1092	UGACAGGAAUGCAUUGGUGUU	1	21	10	23	5'
smo-miR1093	UGGAGGUGUCGUUGCCAAGGA	1	21	17	30	5'
smo-miR1094	UACUUCuCcGUgCCACAGUA	3	20	785	1406	3'
smo-miR1095	UAGGUUCCCUUGUUCCAAAU	2	21	1694	2381	3'
smo-miR1096	CUGCCUCUUUGCUUCAGGAU	1	21	11	37	3'
smo-miR1097	UAGCCAUUGUUGUUGUUGGAA	1	21	11	19	5'
smo-miR1098	UGAUGACGUUUGUGCUGAAAU	1	21	2415	2835	5'
smo-miR1099	UAUJAGCAAUGGUGUUUUGUC	1	21	23	41	5'
smo-miR1100	UGUCACGGACAGAACCCACUC	1	22	41	68	5'
smo-miR1101(5')	GGCAUUCUCUCGAAACAAGUG	1	21	1	2	Amb
smo-miR1101(3')	UUGUUUGCAGAGAAUGCCCAA	1	21	1	2	Amb
smo-miR1102	AGCUAUUGCCACGGUCAAGCA	1	21	196	210	5'
smo-miR1103(5')	AAGAGUGCACCCCUUCCAAU	1	22	1	4	Amb
smo-miR1103(3')	UGGAAAAAGGAGGUGCAUUCUUGU	1	24	1	4	Amb
smo-miR1104	CGCAGCUGUUCUUUUUCCUUC	1	21	692	862	3'
smo-miR1105	UUCGGAUGUGAAGAUCGCUGC	1	21	48	87	3'
smo-miR1106	UUUAAAGGUGUUAAUGUGUGA	1	21	30	38	3'
smo-miR1107	GGUGCUUGGUUCCAAUUCAGgU	1	21	2405	2556	5'
smo-miR1108	UGAGCUGAGAGACAUAACCCC	1	21	10	32	3'
smo-miR1109	JAGUGGGAGAUUUUGUGUAAC	1	21	46	129	5'
smo-miR1110	GCUAGGGGCAGUGGUCAAGGA	1	21	12	22	5'
smo-miR1111	GCCAUGACUAAGCAGGACCAG	1	21	4	10	3'
smo-miR1112(5')	ACUUUGUUAUGGCAUUCUCG	1	21	33	61	Amb
smo-miR1112(3')	AGGAAUGCUAUACAAGUCA	1	21	26	61	Amb
smo-miR1113	UGAGCAGUCAUAAGGUAGCCU	1	21	5443	5919	5'
smo-miR1114	AAGGGACAAAACAUUGAGCAG	1	21	6	14	3'
smo-miR1115(5')	UGAGCUCAGGCACUUUGGUGG	1	21	1	3	Amb
smo-miR1115(3')	UCACCAAAGUGCCUGAGCUCA	1	21	1	3	Amb

^a Number of times the mature miRNA was sequenced (miRNA) or any small RNA matching the hairpin was sequenced. A total of 149,586 *S. moellendorffii* small RNAs that matched at least one WGS trace were sequenced from a single library.

^b Sequence of the most abundant mature miRNA is shown, with lowercase letters indicating positions where the sequence of less-abundant paralogs differ.

^c Arm of stem loop where the mature miRNA resides. Amb, ambiguous.

expressed from two polycistronic precursors (Talmor-Neiman et al., 2006a). We found that clustered miRNA expression was common for *P. patens* miRNAs (Figure 1A; see Supplemental Table 1 online). A total of 48 of the 205 *P. patens* miRNA loci were arranged in closely linked clusters of two or three miRNA stem loops. Clustered miRNA stem loops were observed for both ancient miRNAs, such as miR160 and miR166, and for novel bryophyte-specific miRNAs (Figure 1A). With only a few excep-

tions, these clusters contained paralogs encoding mature miRNA families in the same family, rather than encoding distinct mature miRNAs unrelated in sequence as is frequently the case for animal miRNA clusters (see Supplemental Table 1 online). Similarly, recent results in maize (*Zea mays*) have demonstrated the expression of a polycistronic miRNA precursor for miR156 (Chuck et al., 2007), and a miRNA from the green alga *C. reinhardtii* is also arranged in a tandem cluster (Zhao et al., 2007).

Table 3. Deeply Conserved Plant miRNA Families

Family	At Loci ^a	Pt Loci ^a	Os Loci ^a	Sm Loci ^b	Pp Loci ^c
miR156/157	12	11	12	4	3
miR160	3	8	6	2	9
miR165/166	9	17	14	3	13
miR170/171	4	11	9	4	2
miR159/319	6	15	8	2	5
miR390	2	4	1		3
miR396	2	7	5	1	
miR408	1	1	1	1	2
miR473/477		2			8
miR529(5') ^d			1		7
miR535			1		4
miR536				1	6

^aData from miRBase version 9.2.

^bData from this work.


^cData from miRBase version 9.2 and this work.

^dBased upon the sequencing frequencies obtained in *P. patens*, we suggest the previously annotated miR529 from *O. sativa* actually represents the miRNA*, and the sequence from the 5' arm represents the miRNA. miR529(5') may also be related to miR156.

Strikingly, the majority of the *P. patens* miRNA hairpins overlapped with the positions of annotated protein-coding loci in the *P. patens* version 1_1 (Phypa1_1) draft genome assembly (Figure 1B). Of the miRNAs that overlapped with protein-coding annotations, approximately two-thirds were in the sense orientation with respect to the annotated gene, and approximately equal numbers overlapped exonic compared with intronic sequences

(Figure 1B). The preponderance of *P. patens* miRNA loci that overlapped with annotated protein-coding loci was more reminiscent of miRNA expression in the unicellular green alga *C. reinhardtii* than that of angiosperms (Zhao et al., 2007). The Phypa1_1 genome release contains 35,938 annotated protein coding genes, the majority of which were annotated using a combination of 5' and 3' EST sequences and de novo gene-finding algorithms. We therefore considered the possibility that many of the miRNA-overlapping protein-coding loci represented erroneous annotations triggered by the fact that, except for protein-coding potential, miRNA loci have typical features of RNA polymerase II-transcribed regions that might be recognized by automated gene annotation software (Lee et al., 2004; Xie et al., 2005). If this were the case, then the lengths of the miRNA-overlapping annotated proteins would be predicted to be much shorter than the typical length of all of the annotated proteins. However, this was not the case: The median length for all annotated proteins in the draft *P. patens* Phypa1_1 assembly was 277 amino acids compared with a median of 267 for the subset of annotated protein-coding genes that overlapped mature miRNAs. The median lengths of proteins that overlapped with miRNA loci in introns and exons was similar (medians of 269.5 and 256, respectively), suggesting that there is no consistent bias for the overlapping genes of either intronic or exonic miRNAs to be misannotated. We conclude that many *P. patens* miRNAs are indeed expressed from loci that also have protein-coding potential.

Some lineage-specific miRNA hairpins expressed in *Arabidopsis* possess extended complementarity to protein-coding transcripts that are frequently, though not always, the targets of

**Figure 1.** Properties of miRNA Expression in *P. patens*.

(A) Schematic depictions of representative clustered *P. patens* miRNAs that are potentially processed from polycistronic pri-miRNAs, indicated by black lines. In total, 48 out of the 205 *P. patens* miRNA loci were contained in 21 clusters containing two or three miRNA stem loops each (see Supplemental Table 1 online). Distances refer to the number of nucleotides separating the regions annotated as miRNA foldbacks.

(B) Mapping of mature *P. patens* miRNAs with respect to annotated protein-coding genes. The numbers of mature miRNAs that overlapped various annotations of the *P. patens* Phypa1_1 genome assembly are indicated.




Figure 2. Validated *P. patens* miRNA and ta-siRNA Targets.

such miRNAs (Allen et al., 2004; Rajagopalan et al., 2006; Fahlgren et al., 2007). This suggests that the initiation of novel miRNA–target interactions at least sometimes involves the inverted duplication of transcribed regions. Similar analyses for *P. patens* miRNA loci were complicated by the fact that more than half of the miRNA hairpins overlapped annotated protein-coding regions; thus, any extended homology found between such hairpins and other protein-coding transcripts would be more likely due to conserved protein domains rather than indicative of the origin of the miRNA locus. Nonetheless, of the 87 loci that did not overlap with annotated protein-coding loci, five were found to have extended complementarity with annotated transcripts (FASTA search $E < 0.05$): miR1026a, miR1072, miR1073, miR1220a, and miR1220b. Interestingly, none of the presumed originating transcripts of these five loci were also predicted as targets of their mature miRNAs, suggesting that these loci might have been selected to have targets distinct from their originating gene families as appears to have happened for a few *Arabidopsis* miRNAs (Fahlgren et al., 2007). However, the majority of *P. patens* miRNA loci that don't themselves overlap with annotated protein-coding transcripts did not appear to have extended complementarity to annotated transcripts.

P. patens miRNA Targets

Several targets of *P. patens* miRNAs and ta-siRNAs have been predicted and/or validated using EST data, WGS trace data, or directed cDNA cloning (Floyd and Bowman, 2004; Arazi et al., 2005; Axtell et al., 2006; Floyd et al., 2006; Talmor-Neiman et al., 2006a, 2006b; Fattash et al., 2007). The availability of a large set of annotated mRNAs derived from the draft genome assembly coupled with the expanded number of *P. patens* miRNAs discovered in this study prompted us to revisit miRNA and ta-siRNA target prediction. Targets were predicted using the methodology described by Allen et al. (2005), except that conservation of targeting in other plant genomes was not considered (see Supplemental Table 3 online). This methodology performed reasonably well, as assessed by a signal-to-noise ratio of 2.78 to 1 when compared with a set of shuffled miRNA sequences controlled for di- and trinucleotide composition.

Computational target predictions were based upon the full contingent of annotated transcripts that accompanied the *Phypha1_1* genome release and thus are expected to be more comprehensive than those based upon EST or WGS sequence data. However, we recognized that this database of *P. patens* transcripts was probably not as accurate as those derived from

the current annotations of the more thoroughly studied *Arabidopsis* and *O. sativa* genomes; thus, failure to annotate *P. patens* transcripts could have led to false negatives (missed predictions), while the presence of inaccurately annotated transcripts and inability to consider conservation in a related genome could have led to false positive predictions. We therefore used RLM 5'-RACE (Llave et al., 2002a) to test many of these predictions by assaying for the existence of cDNAs corresponding to miRNA-mediated cleavage products. A total of 29 targets of 14 miRNA families and a ta-siRNA were validated (Figure 2). As expected based upon previous observations (Floyd and Bowman, 2004; Arazi et al., 2005; Axtell and Bartel, 2005; Floyd et al., 2006; Fattash et al., 2007), most of the miRNAs conserved between *Arabidopsis* and *P. patens* had similar targets in both organisms (Figure 2A; see Supplemental Table 3 online), supporting the hypothesis that these *trans*-regulatory interactions have remained constant during the diversification of land plants. For instance, miR156 directed the cleavage of two *SBP* box genes (including *Pp SBP3*, which had previously been demonstrated to be a miR156 target; Arazi et al., 2005), miR166 directed the cleavage of the five class III *HD-ZIP* genes that had been previously observed to possess miR166 complementary sites (Floyd et al., 2006), miR171 directed the cleavage of two *GRAS* domain transcription factor genes, and miR408 directed the cleavage of a plastocyanin domain-containing gene. In flowering plants, the miR159/319 family targets subsets of the *MYB* and *TCP* transcription factor gene families (Rhoades et al., 2002; Palatnik et al., 2003); none of the initially predicted *P. patens* miR319 targets had recognizable homology to *MYB* or *TCP* genes (see Supplemental Table 3 online). However, a gene with a cyclin domain was among the initially predicted *P. patens* miR319 targets and was indeed cleaved *in vivo* at the predicted miR319 complementary site (Figure 2A). Reducing the stringency of the search for *P. patens* miR319 targets allowed the identification of two homologs of *MYB* transcription factors that were similar to the angiosperm targets of the sequence-related miR159 (see Supplemental Table 3 online); RLM 5'-RACE demonstrated that these two genes were also cleaved by a miR319-guided activity in *P. patens* (Figure 2A). Thus, the initial search parameters were too stringent to identify all bona fide cleavage targets. This sensitivity trade-off when predicting plant miRNA targets with acceptable specificity without recourse to evolutionary conservation was expected. For example, the target prediction score cutoffs would have been too stringent to identify the 3' miR390 complementary site of *Pp TAS3a* and also would have failed to identify the 5' miR390 complementary sites of

Figure 2. (continued).

(A) Cleaved targets of conserved *P. patens* miRNAs. Schematics of target mRNAs along with the relative positions of the 5' residues of all sequenced uncapped cDNAs are shown. The positions of the miRNA complementary sites are indicated with red vertical lines. Gray boxes indicate the positions of regions coding for conserved protein domains, whose abbreviated names are listed below. Boxes correspond to open reading frames and horizontal black lines to untranslated regions. Shaded pixels indicate the positions of 5' residues of individual cDNA clones, which are shaded as indicated in the key below to indicate their proximity to the site predicted by miRNA-mediated cleavage between the tenth and eleventh nucleotide of complementarity. An asterisk indicates that *Pp SBP3* was included as a positive control (Arazi et al., 2005). Alignments of target sites with small RNAs are provided in Supplemental Table 3 online. nt, nucleotides.

(B) Cleaved targets of nonconserved *P. patens* miRNAs as in **(A)**. Hatch marks in *Phypha1_1 109598* indicate a region where the splicing of the observed cDNA fragments differed from the annotation.

angiosperm *TAS3* loci, even though these have been confirmed by direct experimentation to be functional (Axtell et al., 2006; Talmor-Neiman et al., 2006b).

The predicted and experimentally confirmed targets of the *P. patens*-specific miRNAs included many potential developmental regulatory genes and a set of genes whose sequences indicated more diverse functions (Figure 2B; see Supplemental Table 3 online). For instance, miR534 directed the cleavage of two mRNAs encoding ankyrin repeat proteins, while miR536 directed the cleavage of an F-box encoding mRNA. The *P. patens*-specific miR538 was demonstrated to target three *MADS* box transcription factors: *PPM1*, *Pp MADS1*, and a close relative of *PPM2*, *Phypa1_1 109598* (Krogan and Ashton, 2000; Henschel et al., 2002). Among these miR538 targets, *PPM1* has been demonstrated to play a role in moss development; antisense knockdown of *PPM1* results in delayed gametangia development and aberrant leaf development (Singer et al., 2007). Interestingly, the *Arabidopsis*-specific miR824 also directs the cleavage of a *MADS* box mRNA (Fahlgren et al., 2007) as does the *O. sativa*-specific miR444 (Sunkar et al., 2005). There is no detectable sequence similarity among miR538, miR824, and miR444 despite the fact that all three regulate *MADS* box-containing genes in their respective lineages. The two small RNAs comprising the miRNA/miRNA* duplex of miR902 were roughly equivalent in sequencing abundance, and both had several predicted targets with favorable scores (see Supplemental Table 3 online). miR902(5') was predicted to target several potential *helix-loop-helix* (*HLH*) transcription factor genes, while the targets of miR902(3') were predicted to include PHD/SET domain genes potentially involved in the regulation of chromatin structure. Although the predictions suggested that miR902(5') and miR902(3') might both have posttranscriptional regulatory functions, we were only able to experimentally validate the *HLH* targets of miR902(5') (Figure 2B).

Recurrence of miRNA-Mediated Negative Feedback Loops in Plants

The gene encoding the DCL protein chiefly responsible for miRNA biogenesis in *Arabidopsis*, *DCL1*, is itself regulated by miRNAs: miR162 is expressed from two distinct loci and can target *DCL1* mRNA via a complementary site formed by the splicing of exon 12 to exon 13 (Xie et al., 2003), while miR838 is excised from a stem loop present in intron 14 of the *DCL1* pre-mRNA (Rajagopalan et al., 2006). Target cleavage mediated by miR162 and excision of the miR838/miR838* duplex, both of which would lead to repression of *DCL1* mRNA expression, have been hypothesized to require the activity of the DCL1 protein and thereby constitute negative feedback circuits. We found that the *P. patens* *MIR1047* locus resided in the sense orientation relative to intron seven of a gene whose predicted protein product had strong overall similarities to DCL proteins (Figure 3A). RLM 5'-RACE experiments detected RNA fragments terminating at positions consistent with the excision of the miR1047/miR1047* duplex from this pre-mRNA (Figure 3B). Evidence for miR1047 excision from pre-mRNA in which intron seven was not spliced was found from both poly(A)⁺ and poly(A)⁻ RNA samples. We also noted that approximately half of these truncated pre-mRNAs

terminated at the loop-proximal cleavage position, indicating that the first DCL-mediated cleavage of the stem loop must have occurred at the loop-proximal position (Figure 3B). This was in contrast with the processing of animal pri-miRNAs (Lee et al., 2003) and certain other plant pri-miRNAs (Kurihara and Watanabe, 2004; Vaucheret et al., 2006), for which the first cleavage events can occur proximal to the base of the hairpins. Nevertheless, similar results for *Arabidopsis* miR838 processing (Rajagopalan et al., 2006) support the conclusion that the loop-proximal cleavage event can sometimes occur first during the biogenesis of some plant miRNAs.

Phylogenetic reconstruction of the relationships between *Arabidopsis*, *O. sativa*, *C. reinhardtii*, and *P. patens* DCL proteins clearly demonstrated that Pp DCL1a (*Phypa1_1 205895*), within whose pre-mRNA *MIR1047* is embedded, was part of the land plant DCL1 clade (Figure 3C). Neither miR1047(5'), miR1047(3'), nor the entire Pp *DCL1a* intron seven sequence possessed detectable similarity with miR838 or with the entire At *DCL1* intron 14 sequence, which encodes miR838. *MIR838* and *MIR1047* also differed with respect to their positions within their host genes, in that the intron encoding *MIR838* is flanked by exons coding for the PAZ region of DCL1 (Rajagopalan et al., 2006), whereas the *MIR1047*-encoding intron is flanked by exons coding for the helicase region (Figure 3A). Because the number and positions of introns were identical between At *DCL1* and Pp *DCL1a*, this difference suggested that miR838 and miR1047 arose independently rather than having diverged from a common ancestral miRNA. Nonetheless, despite their sequence and positional differences, miR838 and miR1047 appeared to have analogous functions in the negative feedback regulation of *DCL1* pre-mRNAs. We did not find any *O. sativa* small RNAs that corresponded to the *Os DCL1* locus among two large data sets of sequenced rice small RNAs (Johnson et al., 2007; Nobuta et al., 2007).

The founding member of the AGO gene family, *Arabidopsis* *AGO1*, encodes a miRNA-guided endonuclease required for the *trans*-regulatory effects of many miRNAs (Vaucheret et al., 2004; Baumberger and Baulcombe, 2005; Qi et al., 2005). *AGO1* is required for the stability and function of miR168, which targets the *AGO1* mRNA; this homeostatic feedback loop is critical for proper *Arabidopsis* development and the normal functioning of many other *Arabidopsis* miRNAs (Vaucheret et al., 2004, 2006). We found that the three validated targets of the *P. patens*-specific miR904 were homologous to AGO genes (see Supplemental Table 3 online; Figures 2B and Figure 3D). The three *P. patens* AGO proteins whose mRNAs were validated targets of miR904 were clearly embedded within the AGO1 clade of land plant AGO proteins (Figure 3D), suggesting that one or more of these miR904 targets might be the major miRNA-guided endonucleases in *P. patens*; to reflect this similarity, we named these genes Pp *AGO1a* (*Phypa1_1 205541*), Pp *AGO1b* (*Phypa1_1 158832*), and Pp *AGO1c* (*Phypa1_1 141045*). Despite having homologous targets, no similarity between the sequences of miR168 and miR904 was detected. This suggested that the miRNA-AGO feedback loops either appeared independently in angiosperms and bryophytes by convergent evolution or that the sequences of the miRNA and AGO target complementary sites have diverged from those of a common ancestor. Both miR168




Figure 3. Analogous miRNA-Mediated Feedback Loops Despite Divergent Small RNA Sequences.

(A) Schematic of the *PpDCL1a* pre-mRNA, which contains an expressed miRNA within intron seven. Thick rectangles represent predicted coding exons, thin rectangles represent untranslated regions, and thin lines represent predicted introns. Colored exonic regions correspond to conserved protein domains as indicated. Gray indicates the location of the *MIR1047* hairpin within intron seven.

(B) Cleavage at miR1047 in intron seven of the *PpDCL1a* pre-mRNA. The predicted secondary structure of *MIR1047* is shown, with the miRNA/miRNA* duplex highlighted in red. Fractions indicate the number of 5'-RACE products observed to terminate at the indicated positions over the total number of 5'-RACE clones sequenced from three different experiments from the indicated combinations of RNA samples and RT primers.



(C) An unrooted phylogenetic reconstruction of relationships among *Arabidopsis*, *O. sativa*, and *P. patens* DCL proteins using *C. reinhardtii* DCL1 as an outgroup. The known miRNA-mediated feedback loops between *AtDCL1* pre-mRNA, mRNA, protein, and DCL1-dependent miRNAs are indicated in blue. The hypothesized feedback loop between *PpDCL1a* pre-mRNA, protein, and ppt-MIR1047 is indicated in red. Nodes with <100% bootstrap support are noted with the percentage of supporting replicates. Accession numbers are given in Methods.

(D) An unrooted phylogenetic reconstruction of relationships among *Arabidopsis*, *O. sativa*, and *P. patens* AGO proteins using *C. reinhardtii* AGO1 and AGO2 as outgroups as displayed in **(B)**. miRNA-mediated control of *Arabidopsis* AGO genes is shown in blue, and miRNA-mediated control of *P. patens* AGO genes by ppt-miR904 is shown in red. Accession numbers are given in Methods.

and miR904 are released from the 5' arms of their respective precursor RNAs, and their cleavage sites in both *Arabidopsis* and *P. patens* AGO mRNAs lie just upstream of the region encoding the conserved PAZ domain. These observations leave open the possibility that the miR168-AGO and miR904-AGO regulatory interactions could have descended from a common ancestor. The observation that negative feedback interactions between miRNAs, DCL genes, and AGO genes exist in both bryophytes

and angiosperms suggests that these interactions are critical for control of miRNA functions in both lineages.

The phylogenetic comparison of the DCL and AGO families in *Arabidopsis*, *O. sativa*, *P. patens*, and *C. reinhardtii* provided the opportunity to consider the evolution of these families during the divergence of land plants. The DCL family appeared to have diverged and specialized early, with at least three members present in the last common ancestor of these land plants and



B Block 1 (+) predicted and validated targets




Figure 4. Similar ARF Targets for Diverse ta-siRNAs and miRNAs.

rather modest duplication since then. We failed to observe a DCL2 homolog in *P. patens*, though the node supporting the placement of Pp DCL4 in the DCL4 clade was weak (Figure 3C). Whether this reflected the first appearance of DCL2 in the seed plant lineage after the divergence of the bryophytes, selective loss of DCL2 in the bryophyte lineage, or simply a failure to annotate the Pp *DCL2* transcript in the Phypa1_1 draft genome was unclear. Because the genome of *S. moellendorffii* has not yet been assembled and annotated, we were unable to include lycopod DCL proteins in this analysis. Evolution of the AGO family appears to have been more dynamic, with much more expansion, and perhaps loss, of paralogs since the last common ancestor. None of the six identified *P. patens* AGO proteins fell in the AGO2/3/7-like clade (Figure 3D); whether this reflected selective loss in the bryophyte lineage, diversification of the AGO2/3/7 clade of AGO proteins in the seed plant lineage after divergence from the bryophytes, or a failure to annotate some *P. patens* AGO loci in the Phypa1_1 assembly was unclear. It is also possible that some of the apparent duplications and extinctions in the AGO genes were artifacts derived from errors in phylogenetic reconstruction; as for the DCL proteins, the *S. moellendorffii* AGO proteins were not available to assist in resolving this issue.

Pp *TAS3a-d* Targets Include Genes Similar to *Arabidopsis* *ETT/ARF3* and *ARF4*

We previously described four ta-siRNA loci targeted by miR390 in *P. patens*, which we referred to as Pp *TAS1-4* (Axtell et al., 2006). The existence of what we called Pp *TAS1* was first suggested by Arazi et al. (2005) and was functionally characterized by Talmor-Neiman et al. (2006b), who named it *TAS4*. We suggest that these four *P. patens* ta-siRNA loci would be better referred to as Pp *TAS3a-d* for three reasons: All four loci possess dual miR390 complementary sites, similar to angiosperm *TAS3* loci (Axtell et al., 2006; Howell et al., 2007), an unrelated *Arabidopsis* ta-siRNA locus has also recently been named *TAS4* (Rajagopalan et al., 2006), and targets identified and validated in this study (see below) demonstrate that ta-siRNAs derived from these moss loci function to regulate *AUXIN RESPONSE FACTOR (ARF)* genes similar to those regulated by the angiosperm *TAS3* loci.

Pp *TAS3a* produces abundant phased siRNAs that are dependent upon the RNA-dependent RNA polymerase Pp *RDR6* (Talmor-Neiman et al., 2006b). Interestingly, Pp *rdi6* mutants also display accelerated differentiation of leafy gametophores

from protonemata, suggesting that the Pp *TAS3* loci are important for the timing of gametophyte development (Talmor-Neiman et al., 2006b); in *Arabidopsis*, *rdi6* mutants display an At *TAS3*-dependent acceleration of sporophytic vegetative phase change (Peragine et al., 2004; Adenot et al., 2006; Fahlgren et al., 2006; Garcia et al., 2006). Alignment of angiosperm *TAS3* genes from multiple species allowed the identification of two conserved, nearly identical ta-siRNAs that function in *Arabidopsis* to regulate *ETT/ARF3* and *ARF4* (Allen et al., 2005; Fahlgren et al., 2006; Hunter et al., 2006), but these ta-siRNAs did not have any recognizable homology to the Pp *TAS3a-d* loci (Axtell et al., 2006).

Given their similar complementarity to miR390, we reasoned that Pp *TAS3a-d* might be paralogs derived from duplication and divergence of an ancestral locus, and perhaps the areas producing biologically relevant ta-siRNAs may have been preferentially conserved. This appears to have been the case, as alignment of the ta-siRNA-producing regions of Pp *TAS3a-d* revealed two ~21-nucleotide blocks of strong conservation (Figure 4A). Sense ta-siRNAs whose 5' residues were coincident with the beginning of block 1 were two to three nucleotides offset from the phasing register set by cleavage at the 3' miR390 complementary site, while antisense ta-siRNAs whose 5' ends were coincident with the beginning of block 2 were exactly coincident with the phasing register set by the 3' miR390 complementary site (Figure 4A). We found no evidence for miR390 expression in *S. moellendorffii*, suggesting that the miR390-*TAS3* cascade of small RNA production might have been lost in that lineage.

Target predictions using the block 1(+) ta-siRNAs recovered three predicted targets of Pp *TAS3a-d*. These included *Phypa1_1 129196*, an AP2 domain-containing gene that has been directly demonstrated to be a target of a block 1(+) ta-siRNA by RLM 5'-RACE (Talmor-Neiman et al., 2006b), and two other AP2 domain-containing genes (Figure 4B). Interestingly, the Pp *TAS3a-d*, block 1(+) ta-siRNAs that could optimally pair with these predicted targets were all offset by two or three nucleotides from the phasing register predicted by the 3' miR390 cleavage site and were closer to the register predicted by the 5' miR390 cleavage site. The experimentally determined cleavage site of *Phypa1_1 129196* is consistent with the offset ta-siRNAs that are optimal for interaction with these targets (Talmor-Neiman et al., 2006b; Figure 4B). One of these predicted block 1(+) targets was also a validated target of miR529(5') (*Phypa1_1 65352*; Figure 2A). Our failure to detect cleavage products corresponding to the upstream Pp *TAS3a-d* block 1(+) complementary site in *Phypa1_1 65352* could have

Figure 4. (continued).

(A) T-COFFEE alignment of the regions of Pp *TAS3a-d* between the two miR390 complementary sites. Two ~21-nucleotide blocks conserved between the four paralogs are indicated.

(B) Alignments of the block 1 (+) siRNAs and the block 2 (-) RNAs with their predicted and validated targets. Lowercase letters indicate positions that cannot form either Watson-Crick or G-U pairs with the target. Arrows indicate cleavage-validated target sites. The asterisk indicates the cleavage site of the AP2 domain-containing gene *Phypa1_1 129196* by Pp *TAS3a-d* block 1 (+)-derived siRNAs as demonstrated by Talmor-Neiman et al. (2006b).

(C) An unrooted phylogenetic reconstruction of the relationships between *Arabidopsis* and *P. patens* ARF proteins. Nodes with <100% bootstrap support are noted with the percentage of supporting replicates. ARF genes known or predicted to be under small RNA-mediated regulation are indicated. Accession numbers are given in Methods.

been due to highly efficient cleavage at the 3' miR529(5') site. No secondary siRNAs that corresponded to the region of *Phypa1_1* 65352 between the two sites were sequenced in our study.

The block 2(-) ta-siRNAs were predicted to target four distinct transcripts (see Supplemental Table 3 online), two of which encoded proteins homologous to angiosperm ARFs (Figures 4B and 4C). Cleavage products were detected for both *ARF* messages, thereby demonstrating that these *P. patens* *ARF* mRNAs were indeed regulated by Pp *TAS3*-derived ta-siRNAs (Figure 2B). Unexpectedly, these RLM 5'-RACE experiments revealed a second, longer species of uncapped mRNA for both of the *P. patens* *ARF* genes, leading us to investigate whether the longer products also corresponded to small RNA-mediated cleavage products. The putative upstream small RNA complementary sites did not appear to correspond to any ta-siRNA produced from Pp *TAS3a-d* nor were these two *ARF* genes predicted as targets of any *P. patens* miRNAs using established protocols (Allen et al., 2005; Rajagopalan et al., 2006). However, relaxing the stringency of the target prediction parameters revealed that the upstream cleavage sites corresponded with the bryophyte-specific miR1219 such that the observed cleavage product termini aligned exactly with the tenth nucleotide of complementarity (Figure 2B). We concluded that the two *P. patens* *ARF* genes were targeted at distinct sites by miR390-initiated, Pp *TAS3a-d*-derived ta-siRNAs and by the bryophyte-specific miR1219. We did not sequence any small RNAs that corresponded to the regions of the two *P. patens* *ARF* genes between the Pp *TAS3a-d* block 2(-) and miR1219 cleavage sites.

Phylogenetic comparison of *P. patens* *ARF* proteins to *Arabidopsis* *ARF* proteins revealed that the Pp *TAS3a-d* block 2(-) and miR1219-targeted *ARFs* were closely related to At *ETT/ARF3* and At *ARF4* (Figure 4C). At *ETT/ARF3* and At *ARF4* are targeted by miR390-phased, At *TAS3*-derived ta-siRNAs at two distinct complementary sites located downstream of the region coding for the *ARF* domain (Allen et al., 2005). Our data thus demonstrated that the arrangement of dual small RNA target sites within *ARF* mRNAs was shared between bryophytes and angiosperms despite divergence of both the sequence composition and of the biogenesis of the small RNAs themselves, providing a striking example of common regulatory functions being fulfilled by lineage-specific small RNAs. Moreover, the Pp *TAS3a-d* genes appear to transmute the expression of one miRNA (miR390) into two biologically relevant sets of related small RNAs that are capable of the coordinated regulation of a diverse set of regulatory transcripts. One set of Pp *TAS3a-d* ta-siRNAs regulates *AP2* domain genes in concert with miR529(5'), while another set acts in concert with a lineage-specific miRNA, miR1219, to regulate the expression of *ETT/ARF3* and *ARF4*-like genes.

Other Small RNAs in Basal Plants

The entire set of 561,102 *P. patens* small RNA reads that matched at least one WGS trace (Axtell et al., 2006) were categorized based upon genomic and length characteristics (Figure 5A). miRNAs and ta-siRNAs accounted for 28 and 3% of the reads, respectively. Both subpopulations were dominated by 21-nucleotide sequences, with a considerable minority of

22-nucleotide sequences deriving from the Pp *TAS3a-d* ta-siRNAs (Figure 5A); these properties were consistent with those of angiosperm miRNAs and ta-siRNAs (Reinhart et al., 2002; Rajagopalan et al., 2006). Strikingly, 47% of the reads that initially matched at least one raw WGS trace did not match anywhere along the *Phypa1_1* draft genome assembly. This subpopulation of reads included many short sequences of sizes inconsistent with the activities of any known Dicer endonuclease and some small RNAs between 20 and 24 nucleotides (Figure 5A). A substantial number of these reads were only sequenced once and disproportionately tended to match only a single WGS trace (data not shown), a statistically unlikely occurrence considering that shotgun coverage of the *P. patens* genome is $\sim 8\times$. This suggested that this subpopulation of reads contains some convergent sequencing errors (i.e., a sequencing error in a single small RNA and a single WGS trace that causes a spurious match). It might also contain breakdown products of RNAs and Dicer-derived small RNAs originating from genomic regions that are difficult to sample and assemble using shotgun sequencing and thus were left out of the draft genome assembly.

The remaining 22% of the *P. patens* reads that matched the version 1_1 genome assembly, but did not appear to originate from recognizable miRNA or ta-siRNA loci, were classified as unknown reads (Figure 5A). Repetitive unknown reads were those that corresponded to >10 loci in the draft genome assembly. Both repetitive and nonrepetitive subpopulations of unknown small RNA reads had clear size peaks at 21, 23, and 24 nucleotides, consistent with the hypothesis that many corresponded to endogenous siRNAs produced by DCL activities. In *Arabidopsis*, many endogenous 24-nucleotide siRNAs are dependent on the DCL3, AGO4, and AGO6 proteins (Qi et al., 2006; Kasschau et al., 2007; Zheng et al., 2007). Phylogenetic reconstructions of DCL proteins indicated that the four recognizable *P. patens* DCL proteins fell into three clades: a DCL1-like clade, a DCL3-like clade, and a DCL4-like clade (Figure 3C). Similarly, *P. patens* AGO proteins also fell into two distinct groups: an AGO1-like clade and an AGO4/6-like clade (Figure 3D). Taken together, these results suggest that, similar to angiosperms, *P. patens* produces at least some miRNA-independent (i.e., non-TAS) siRNAs that might participate in DNA silencing.

A similar analysis of the *S. moellendorffii* small RNA reads found that 35% of them corresponded to miRNA hairpins, most of which were 21 nucleotides in length (Figure 5B). The remaining 65% of the reads that matched at least one *S. moellendorffii* WGS trace were binned into repetitive and nonrepetitive subpopulations, which comprised 11 and 54% of the total, respectively (Figure 5B). The length distributions of the *S. moellendorffii* unknown reads were dispersed and overall were not consistent with the small RNA sizes produced by the biochemically or genetically characterized DCL proteins from other species. However, based upon the experience with *P. patens* small RNAs, we expect that approximately one-half of these unknown small RNAs will not be matched to the forthcoming *S. moellendorffii* draft genome sequence. As with *P. patens*, subtracting these reads may reveal the signature of a small population of potential miRNA-independent siRNAs in *S. moellendorffii*.

The proportion of *P. patens* and *S. moellendorffii* potential endogenous siRNAs was lower than that observed in similar




Figure 5. Small RNA Populations in *P. patens* and *S. moellendorffii*.

(A) Categorization of 561,102 *P. patens* small RNA reads that matched at least one WGS trace. Reads were categorized as miRNAs (matching one or more of the 205 annotated miRNA hairpins), ta-siRNAs (from *PpTAS3a-d*), unknown (corresponding to 10 or fewer loci in the Phypha1_1 genome assembly), unknown-repetitive (corresponding to 11 or more loci in the Phypha1_1 genome assembly), or those not matching any loci in the Phypha1_1 genome assembly. Histograms display the length distributions of each of these small RNA populations. Reads that corresponded to the nuclear rRNAs and the chloroplast genome were not included.

(B) Categorization of 149,586 *S. moellendorffii* small RNA reads that matched at least one WGS trace. Reads are categorized as miRNAs (matching one or more of the 58 annotated miRNA hairpins), unknown (matching 100 or fewer WGS traces), or unknown-repetitive (corresponding to 101 or more WGS traces). Histograms display the length distributions of each of these small RNA populations. Reads that corresponded to nuclear rRNA were not included.

deep sequencing projects that examined *Arabidopsis* small RNAs (Lu et al., 2005; Rajagopalan et al., 2006; Kasschau et al., 2007), raising the possibility that endogenous siRNAs play a less prominent role in genome regulation in basal plants. However, it is also possible that basal plant endogenous siRNAs possess biochemical characteristics that render them recalcitrant to the sequencing strategy employed in this study, which is highly selective for molecules with both a 5' monophosphate and a 3' hydroxyl. This would not be without precedent. In *Caenorhabditis elegans*, secondary siRNAs from an exogenous trigger and a subset of endogenous siRNAs are recalcitrant to this procedure because they possess 5'-triphosphates, reflecting their likely origins as short unprimed RdRp transcripts (Ruby et al., 2006; Pak and Fire, 2007; Sijen et al., 2007). Angiosperm siRNA and miRNA/miRNA* duplexes are methylated by the HEN1 protein on the 2' hydroxyl of the 3'-most residue (Yu et al., 2005; Yang et al., 2006). Periodate treatment of *P. patens* total RNA demonstrated that the 3'-most residue of miR160 does not possess both free 2' and 3' hydroxyls, which was strongly suggestive of a conservation of HEN1-mediated small RNA methylation in bryophytes (Figure 6). Although methylation of the terminal 2' hydroxyl would not have had a major effect on sequencing frequency, other modifications of this terminal ribose might have, especially if different classes of basal plant small RNAs possess different modifications.

DISCUSSION

Recent Diversification of Many Weakly Expressed miRNAs in Three Major Land Plant Lineages

The first small RNA sequencing efforts in both animals (Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001) and plants (Llave et al., 2002b; Park et al., 2002; Reinhart et al., 2002) were performed at relatively low sequencing depths comprising hundreds of reads and therefore captured only the most abundant of the miRNAs expressed in the sampled tissues. Within both lineages, these initially discovered and most abundant miRNAs were very well conserved, leading to the hypothesis that miRNAs are generally ancient posttranscriptional regulators. The use of high-throughput DNA sequencing for small




Figure 6. The 3'-Most Residue of *P. patens* miR160 Is Modified.

The indicated RNA samples were oxidized with periodate, blotted, and probed for miR160. Markers (M) from top to bottom are 24, 21, and 18 nucleotides, respectively.

RNA discovery has allowed a 100- to 1000-fold increase in sequencing depth for various *Arabidopsis* small RNA samples (Lu et al., 2005, 2006; Henderson et al., 2006; Rajagopalan et al., 2006; Kasschau et al., 2007; Zhang et al., 2007). Analyses of these large data sets have revealed that the majority of *Arabidopsis* miRNA diversity (but not numerical abundance) is accounted for by weakly expressed miRNAs that lack identifiable homologs in other species. These nonconserved miRNAs tend to be encoded by only a single locus and also tend to have few, if any, confidently predicted targets. These properties have led to the suggestion that many of the weakly expressed, less well-conserved miRNAs represent a pool of evolutionarily transient regulators with minimal fitness consequences that may be occasionally recruited into biologically relevant circuits of post-transcriptional control but are more likely to decay and die due to mutational drift of the DCL1 recognized stem loops of their pri-miRNAs (Rajagopalan et al., 2006; Fahlgren et al., 2007). Many of the miRNAs discovered from the lycopod *S. moellendorffii* and *P. patens* in this study are reminiscent of the recently evolved miRNAs of *Arabidopsis*. Weakly expressed miRNAs that tend to originate from a single genomic locus and have few predicted mRNA targets comprise the majority of miRNA diversity in all three lineages. Thus, our data strongly support the hypothesis that miRNA-producing loci are frequently born and can rapidly die over relatively short evolutionary time scales, with only a select few being stabilized during their otherwise brief existence by recruitment into a beneficial regulatory interaction (Rajagopalan et al., 2006; Fahlgren et al., 2007). This process clearly occurs in all three deeply branching lineages of land plants (angiosperms, lycopods, and bryophytes), whose small RNAs have been sampled by deep sequencing, and might also occur in animals (Bentwich et al., 2005; Berezikov et al., 2006).

Common Functions for Plant Small RNAs, with and without Conservation of Sequence Identity

The targets of the deeply conserved *P. patens* miRNAs were clearly homologous with the known targets of the same miRNAs in *Arabidopsis*, as observed previously for deeply conserved miRNAs (Floyd and Bowman, 2004; Arazi et al., 2005; Axtell and Bartel, 2005; Floyd et al., 2006). The sequences of these targets strongly suggest that most of these deeply conserved regulatory circuits function to control transcriptional regulators that influence multicellular development and morphology. It will be fascinating to experimentally dissect how these conserved molecular modules have been recruited to control the very different developmental programs of angiosperms and bryophytes.

Analysis of the *P. patens* miRNA loci and their targets strongly suggests that miRNA-mediated feedback control on the miRNA pathway is also shared between bryophytes and angiosperms through the action of diverse miRNA loci generated through convergent and perhaps divergent evolutionary pathways. *P. patens* *MIR1047* is expressed from the intron of a *DCL1* homolog, and just as for *Arabidopsis* *MIR838*, processing of the stem loop by DCL1 cleaves the *DCL1* pre-mRNA, implying a negative feedback loop. These two miRNAs appear to have arisen independently yet serve a common function and thus appear to constitute an example of convergent evolution for miRNA functions.

Considering that large reservoirs of newly emergent and continuously evolving miRNAs have now been described in an angiosperm, a lycopod, and a moss, we expect that many additional examples of convergent miRNA functions will be found in land plants. Whether miR904 targeting of *AGO1*-like genes in *P. patens* and miR168 targeting of *AGO1* in angiosperms was achieved through convergent or divergent emergence of these diverse miRNAs was less clear. Biogenesis from the same arm of the miRNA hairpins and cleavage at similar locations within the homologous *AGO1* messages were both consistent with divergence from a common ancestor, but in either evolutionary scenario, our data demonstrated that analogous regulatory specificity of plant miRNAs does not always entail conservation of the mature miRNA sequences. Similar observations have been made for animal miRNAs, most notably the vertebrate miR-196 and insect miR-iab-4, which both target *Ubx* orthologs in their respective lineages (Yekta et al., 2004; Ronshaugen et al., 2005).

Just as for the miRNAs that appeared to mediate feedback regulation upon *DCL* and *AGO* genes in plants, the biologically relevant ta-siRNAs emanating from plant *TAS3* loci differed in sequence but were found to perform analogous molecular functions in mosses and seed plants. The two *P. patens* *ARF* homologs possessed dual cleavage sites just downstream of the conserved *ARF* domain (Figure 2B), a location and arrangement closely mirroring that of the cleavage sites within *Arabidopsis* *ETT/ARF3* and *ARF4* (Allen et al., 2005). These data involving miR390-triggered siRNA regulation of similar angiosperm and bryophyte *ARF* transcripts could thus be regarded as evidence that the *At TAS3* and *Pp TAS3a-d* genes might have descended from a common molecular ancestor. However, *Arabidopsis* *ETT/ARF3* and *ARF4* both possessed dual complementary sites to identical *At TAS3*-derived ta-siRNAs, whereas the *P. patens* *ARFs* possessed only one site complementary to *Pp TAS3* ta-siRNAs. Cleavage at the other site in the *P. patens* *ARF* mRNAs was guided by the lineage-specific miR1219, which did not have any detectable sequence similarity to the *ARF*-targeting *At TAS3* ta-siRNAs nor to any *Pp TAS3a-d*-derived ta-siRNAs. These observations add to the examples of convergent evolutionary origins for similarly functioning small RNAs, while illustrating that cleavage of a homologous target at an analogous location does not always indicate descent from a common ancestor. Moreover, the *ARF*-targeting ta-siRNAs are derived from the (+) strand of the angiosperm *TAS3* loci, whereas they derived from the (-) strands of the moss *TAS3* loci, suggesting a more complex evolutionary path for plant *TAS3* loci than simple sequence divergence from a common ancestor. Data from additional plant lineages might provide important clues for distinguishing between these interesting possibilities.

The *AGO1*-targeting miRNAs and the miR390-phased ta-siRNAs might be the descendants of anciently derived small RNA-target interactions that have coordinately diverged from common molecular ancestors over long periods of time. This process would thus have hidden the ancestral relationships of the functionally homologous small RNA-target interactions in divergent contemporary organisms, making them appear to be nonconserved when considering primary sequence alone. In other cases, such as miRNAs excised from *DCL1* pre-mRNAs, independently derived, nonconserved miRNAs appear to have

converged upon an identical regulatory function. We conclude that the sequence diversity of miRNAs and ta-siRNAs expressed in various plants can sometimes obfuscate analogous regulatory functions.

METHODS

Small RNA sequencing

Sequencing of *Physcomitrella patens* small RNAs has been described (Axtell et al., 2006). *Selaginella moellendorffii* was purchased from Plant Delights Nursery. A total RNA sample derived from the above-ground parts of *S. moellendorffii* was isolated using the method of Chang et al. (1993). Small RNAs from this sample were sequenced as described (Axtell et al., 2006).

Annotation of miRNAs and Target Predictions

Sequenced small RNAs were filtered to remove contaminant sequences derived from rRNA or the synthetic small RNAs used as guides during the small RNA excision procedure. The remaining sequences were then compared against the relevant WGS traces available from the National Center for Biotechnology Information (NCBI) trace archive and the corresponding matches recorded. After discarding small RNAs that were repetitive (matched >1000 different WGS traces), genomic sequences surrounding small RNA hits were examined for their potential to form an RNA secondary structure typical of pri-miRNA stem loops. A 600-nucleotide window centered upon a given small RNA hit was analyzed using MIRcheck (Jones-Rhoades and Bartel, 2004). Consensus genomic sequences were derived for each potential miRNA locus that passed MIRcheck by use of BLAT (Kent, 2002) to capture redundant WGS traces followed by consensus assembly using CAP3 (Huang and Madan, 1999). All sequenced small RNAs that matched either polarity of the assembled consensus sequences of potential miRNA loci were captured and aligned. Candidate loci where two dominant sequences accumulated from only a single polarity and that, when analyzed in the context of a predicted RNA secondary structure, could form a typical miRNA/miRNA* duplex were annotated as miRNAs. In cases where no sequence corresponding to the presumed miRNA* was sequenced, a locus was still annotated as a miRNA if there was a paralogous locus from which a miRNA/miRNA* was isolated. After the *P. patens* draft genome Phypa1_1 assembly became available, the WGS-derived consensus miRNA loci were all successfully mapped.

The transcripts annotated from the draft *P. patens* genome assembly were used to predict the potential targets of *P. patens* miRNAs. The list of mature *P. patens* miRNAs (see Supplemental Table 1 online) was first collapsed to generate a nonredundant set of queries, and targets were predicted as described (Allen et al., 2005; Rajagopalan et al., 2006). To assess the signal-to-noise ratio of these predictions, a set of randomized sequences with mononucleotide compositions identical to the query and di- and trinucleotide compositions similar to the transcript database (Farh et al., 2005) was produced for each miRNA query and used to predict targets (see Supplemental Table 3 online).

Molecular Phylogenetic Analyses

P. patens proteins homologous to *Arabidopsis thaliana* *DCL1* (At1g01040.1), *AGO1* (At1g48410.1), and *ETT/ARF3* (At2g33860.1) were found using BLASTP and manually examined to identify hits with the domain architectures expected of the respective protein families. Alignments were constructed using the T-COFFEE server (Poirot et al., 2003) under default settings (alignment computations using Mlalign_id_pair and Mslow_pair).

Alignment scores were 50, 63, and 57 for the DCL, AGO, and ARF alignments, respectively. Phylogenetic analyses were performed within MEGA 3.1 (Kumar et al., 2004) using neighbor joining. Options common to all three analyses were as follows: phylogeny test and options = bootstrap, 1000 replicates; model = amino: Poisson correction; substitutions to include = all; pattern among lineages = same (homogenous); rates among sites = uniform rates. The DCL1 and AGO1 analyses used gaps/missing data = pairwise deletion and the ARF analyses used gaps/missing data = complete deletion. The sequence alignments used to construct these phylogenies are available as Supplemental Figures 2 to 4 online.

Isolation of Small RNA-Mediated Cleavage Products

Isolation of uncapped mRNAs using RLM 5'-RACE was performed essentially as described (Kasschau et al., 2003) using *P. patens* RNA from 17-d-old protonemata cultures with young leafy gametophores. Experiments used cDNA produced by reverse transcription of poly(A)⁺ RNA with the GeneRacer oligo(dT) oligo (5'-GCTGTCAACGATACGCTACGTAACGGCATGACAGTGTGTTTTTTTTTTTTTTTTTTTTTTT-3'), except those indicated otherwise in Figure 3. All cleavage products were amplified using two rounds of nested PCR with the GeneRacer 5' oligo (5'-CGACTGGAGCAGGAGACTGA-3') and the GeneRacer 5' nested oligo (5'-GGACACTGACATGGACTGAAGGAGTA-3') in conjunction with outer and nested gene-specific oligos. Products were then cloned and sequenced to determine the 5' ends. Pp *DCL1a* oligos were as follows: exon 8 RT oligo, 5'-GGCTCCGAGCAGCACCACGAGACTGAA-3'; Pp *DCL1a* outer (spans intron seven/exon eight junction), 5'-GCAACGTCACCTAGCCCAACCAACTT-3'; and *DCL1a* nested (intron seven), 5'-GCTGAGCAAACCAGCTCGACAAGATG-3'. All other gene-specific oligos are listed in Supplemental Table 4 online.

Periodate treatments

Total RNA from wild-type 7-d-old *Arabidopsis* seedlings and 7-d-old *P. patens* protonemata was isolated using TRI-Reagent (Ambion). Synthetic miR160 RNAs (5'-UGCCUGGCUCUCCUGUAUGCCA-3') with a 5' monophosphate and either 2', 3' hydroxyls or a 2'-O-methyl, 3' hydroxyl on the 3'-most residue were purchased from Dharmacon. In separate reactions, 90 μg of *Arabidopsis* and *P. patens* total RNA and 10 fmols of the synthetic miR160 RNAs were incubated for 45 min in 10 mM NaIO₄ at 20° in the dark, followed by addition of L-Lys, pH 8.5, to a concentration of 1 M and incubation at 45° for 90 min. Reactions were then phenol-chloroform extracted, desalted through Sephadex G-25 spin columns (GE Healthcare), and ethanol precipitated. Plant RNA reactions were resuspended in 15 μL water, while reactions with synthetic miR160 were resuspended in 4 μL water. The samples were then fractionated, blotted, and hybridized with an oligonucleotide probe against miR160 as described (Axtell and Bartel, 2005). Lanes containing plant RNA had 5 μL of the relevant sample, while lanes containing synthetic RNAs had 1 μL of the relevant sample.

Databases and Gene Family Annotations

All referenced *Arabidopsis* sequences were obtained from The Arabidopsis Information Resource version 6 genome annotation; in cases where multiple versions of a transcript and/or protein were annotated, version 1 was used (e.g., At1g01040.1). The nomenclature of the *Arabidopsis* ARF, DCL, and AGO gene families was derived from Okushima et al. (2005), Margis et al. (2006), and Fagard et al. (2000), respectively. All referenced *Oryza sativa* sequences were obtained from The Institute for Genomic Research (TIGR) version 4 genome annotation; in cases where multiple versions of a transcript and/or protein were annotated, version 1 was used (e.g., Os06g39640.1). The nomenclature of the *O. sativa* DCL

and AGO gene families was derived from Margis et al. (2006) and Shi et al. (2007), respectively. Annotations of the *S. moellendorffii* small RNAs made use of the raw WGS traces available from the NCBI trace server. Analyses of *P. patens* small RNAs made use of the Phypa1_1 draft assembly of the genome from the Joint Genome Institute and the associated transcript and protein models, except where noted. *C. reinhardtii* sequences were derived from the Joint Genome Institute's genome version 2.0; the nomenclature of the Cr *DCL1*, Cr *AGO1*, and Cr *AGO2* genes follows Schroda (2006).

Accession Numbers

P. patens and *S. moellendorffii* small RNA data sets have been deposited with the Gene Expression Omnibus under accession numbers GSE5103 and GSE7320, respectively. Partial cDNAs corresponding to *P. patens* miRNA-mediated cleavage products have been deposited with GenBank under accession numbers ES492157 to ES492185. All novel miRNAs have been registered with miRBase. The Arabidopsis Genome Initiative identifiers for the *Arabidopsis* genes/proteins mentioned by other names in this article are as follows: At *RDR6*, At3g49500; At *TAS3a*, At3g17185; At *DCL1*, At1g01040; At *DCL2*, At3g03300; At *DCL3*, At3g43920; At *DCL4*, At5g20320; At *AGO1*, At1g48410; At *AGO2*, At1g31280; At *AGO3*, At1g31290; At *AGO4*, At2g27040; At *AGO5*, At2g27880; At *AGO6*, At2g32940; At *AGO7*, At1g69440; At *AGO8*, At5g21030; At *AGO9*, At5g21150; At *ZLL/PNH*, At5g43810; At *ARF1*, At1g59750; At *ARF2*, At5g6200; At *ETT/ARF3*, At2g33860; At *ARF4*, At5g60450; At *MP/IAA24/ARF5*, At1g19850; At *ARF6*, At1g30330; At *ARF7*, At5g20730; At *ARF8*, At5g37020; At *ARF9*, At4g23980; At *ARF10*, At2g28350; At *ARF11*, At2g46520; At *ARF12*, At1g34310; At *ARF13*, At1g34170; At *ARF14*, At1g35540; At *ARF15*, At1g35520; At *ARF16*, At4g30080; At *ARF17*, At1g77850; At *ARF18*, At3g61830; At *ARF19*, At1g19220; At *ARF20*, At1g35240; At *ARF21*, At1g34410; At *ARF22*, At1g34390; At *ARF23*, At1g43950. The TIGR locus identifiers for the *O. sativa* genes/proteins mentioned by other names in this article are as follows: Os *DCL1*, Os03g02970; Os *DCL2a*, Os03g38740; Os *DCL2b*, Os09g14610; Os *DCL3a*, Os01g68120; Os *DCL3b*, Os10g34430; Os *DCL4*, Os04g43050; Os *AGO7*, Os03g33550. Based on the results of phylogenetic analysis (Figure 3), we have designated *Phypa1_1 205895*, *Phypa1_1 163666*, *Phypa1_1 130425*, and *Phypa1_1 234670* as Pp *DCL1a*, Pp *DCL1b*, Pp *DCL3*, and Pp *DCL4*, respectively. Similarly, *Phypa1_1 205541*, *Phypa1_1 158832*, and *Phypa1_1 141045* have been designated as Pp *AGO1a*, Pp *AGO1b*, and Pp *AGO1c*, respectively. Most *P. patens* genes mentioned or used for analyses were as annotated from the Phypa1_1 data set, except the following: Pp *RDR6*, DQ531710; Pp *TAS3a*, BK005825; Pp *TAS3b*, BK005826; Pp *TAS3c*, BK005827; Pp *TAS3d*, BK005828; *PPM1*, AF150932; Pp *MADS1*, AB067688; Pp *SBP3*, AJ968318; Pp *HB10*, DQ385518; Pp *C3HDZ1*, DQ385516; Pp *C3HDZ2*, DQ385517. The Joint Genome Institute gene model names for the *C. reinhardtii* proteins referenced in the text are as follows: Cr *DCL1*, C_130110; Cr *AGO1*, C_130206; Cr *AGO2*, C_1700017.

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Figure 1. Alternative miRNA Production.

Supplemental Figure 2. T-COFFEE Alignment of *Arabidopsis*, *O. sativa*, *C. reinhardtii*, and *P. patens* DCL Proteins.

Supplemental Figure 3. T-COFFEE Alignment of *Arabidopsis*, *O. sativa*, *C. reinhardtii*, and *P. patens* AGO Proteins.

Supplemental Figure 4. T-COFFEE Alignment of *Arabidopsis* and *P. patens* ARF Proteins.

Supplemental Table 1. *P. patens* miRNA Loci.

Supplemental Table 2. *S. moellendorffii* miRNA Loci.

Supplemental Table 3. Predicted and Validated Targets of *P. patens* miRNAs and ta-siRNAs.

Supplemental Table 4. RLM 5'-RACE Gene-Specific Oligo Sequences.

ACKNOWLEDGMENTS

We thank the Joint Genome Institute and the *Physcomitrella patens* sequencing consortium members for the availability of the draft genome assembly. We thank Graham Ruby for many stimulating discussions, Ramya Rajagopalan for discussions and critical comments, and Nina Federoff and John Carlson for constructive comments on this manuscript. This work was supported by awards from the National Institutes of Health, the Prix Louis D. from the Institut de France (D.P.B.), the Helen Hay Whitney Foundation, and Pennsylvania State University (M.J.A.).

Received March 19, 2007; revised June 6, 2007; accepted June 8, 2007; published June 29, 2007.

REFERENCES

- Adenot, X., Elmayan, T., Laussergues, D., Boutet, S., Bouche, N., Gascioli, V., and Vaucheret, H. (2006). DRB4-dependent *TAS3* trans-acting siRNAs control leaf morphology through AGO7. *Curr. Biol.* **16**: 927–932.
- Allen, E., Xie, Z., Gustafson, A.M., and Carrington, J.C. (2005). MicroRNA-directed phasing during trans-acting siRNA biogenesis in plants. *Cell* **121**: 207–221.
- Allen, E., Xie, Z., Gustafson, A.M., Sung, G.H., Spatofora, J.W., and Carrington, J.C. (2004). Evolution of microRNA genes by inverted duplication of target gene sequences in *Arabidopsis thaliana*. *Nat. Genet.* **36**: 1282–1290.
- Arazi, T., Talmor-Neiman, M., Stav, R., Riese, M., Huijser, P., and Baulcombe, D.C. (2005). Cloning and characterization of micro-RNAs from moss. *Plant J.* **43**: 837–848.
- Axtell, M.J., and Bartel, D.P. (2005). Antiquity of microRNAs and their targets in land plants. *Plant Cell* **17**: 1658–1673.
- Axtell, M.J., Jan, C., Rajagopalan, R., and Bartel, D.P. (2006). A two-hit trigger for siRNA biogenesis in plants. *Cell* **127**: 565–577.
- Bartel, D.P. (2004). MicroRNAs: Genomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Baulcombe, D. (2005). RNA silencing. *Trends Biochem. Sci.* **30**: 290–293.
- Baumberger, N., and Baulcombe, D.C. (2005). *Arabidopsis* ARGONAUTE1 is an RNA Slicer that selectively recruits microRNAs and short interfering RNAs. *Proc. Natl. Acad. Sci. USA* **102**: 11928–11933.
- Bentwich, I., et al. (2005). Identification of hundreds of conserved and nonconserved human microRNAs. *Nat. Genet.* **37**: 766–770.
- Berezikov, E., Thummler, F., van Laake, L.W., Kondova, I., Bontrop, R., Cuppen, E., and Plasterk, R.H. (2006). Diversity of microRNAs in human and chimpanzee brain. *Nat. Genet.* **38**: 1375–1377.
- Chang, S., Puryear, J., and Cairney, J. (1993). A simple and efficient method for isolating RNA from pine trees. *Plant Mol. Biol. Rep.* **11**: 113–116.
- Chuck, G., Cigan, A.M., Saeteurn, K., and Hake, S. (2007). The heterochronic maize mutant *Corngrass1* results from overexpression of a tandem microRNA. *Nat. Genet.* **39**: 544–549.
- Doench, J.G., Petersen, C.P., and Sharp, P.A. (2003). siRNAs can function as miRNAs. *Genes Dev.* **17**: 438–442.
- Fagard, M., Boutet, S., Morel, J.B., Bellini, C., and Vaucheret, H. (2000). AGO1, QDE-2, and RDE-1 are related proteins required for post-transcriptional gene silencing in plants, quelling in fungi, and RNA interference in animals. *Proc. Natl. Acad. Sci. USA* **97**: 11650–11654.
- Fahlgren, N., Howell, M.D., Kasschau, K.D., Chapman, E.J., Sullivan, C.M., Cumbie, J.S., Givan, S.A., Law, T.F., Grant, S.R., Dangl, J.L., and Carrington, J.C. (2007). High-throughput sequencing of *Arabidopsis* microRNAs: Evidence for frequent birth and death of *MIRNA* genes. *PLoS ONE* **2**: e219.
- Fahlgren, N., Montgomery, T.A., Howell, M.D., Allen, E., Dvorak, S.K., Alexander, A.L., and Carrington, J.C. (2006). Regulation of *AUXIN RESPONSE FACTOR3* by *TAS3* ta-siRNA affects developmental timing and patterning in *Arabidopsis*. *Curr. Biol.* **16**: 939–944.
- Farh, K.K., Grimson, A., Jan, C., Lewis, B.P., Johnston, W.K., Lim, L.P., Burge, C.B., and Bartel, D.P. (2005). The widespread impact of mammalian microRNAs on mRNA repression and evolution. *Science* **310**: 1817–1821.
- Fattash, I., Voss, B., Reski, R., Hess, W.R., and Frank, W. (2007). Evidence for the rapid expansion of microRNA-mediated regulation in early land plant evolution. *BMC Plant Biol.* **7**: 13.
- Floyd, S.K., and Bowman, J.L. (2004). Gene regulation: Ancient microRNA target sequences in plants. *Nature* **428**: 485–486.
- Floyd, S.K., Zalewski, C.S., and Bowman, J.L. (2006). Evolution of class III homeodomain-leucine zipper genes in streptophytes. *Genetics* **173**: 373–388.
- Garcia, D., Collier, S.A., Byrne, M.E., and Martienssen, R.A. (2006). Specification of leaf polarity in *Arabidopsis* via the trans-acting siRNA pathway. *Curr. Biol.* **16**: 933–938.
- Hammond, S.M. (2005). Dicing and slicing: The core machinery of the RNA interference pathway. *FEBS Lett.* **579**: 5822–5829.
- Henderson, I.R., Zhang, X., Lu, C., Johnson, L., Meyers, B.C., Green, P.J., and Jacobsen, S.E. (2006). Dissecting *Arabidopsis thaliana* DICER function in small RNA processing, gene silencing and DNA methylation patterning. *Nat. Genet.* **38**: 721–725.
- Henschel, K., Kofuji, R., Hasebe, M., Saedler, H., Munster, T., and Theissen, G. (2002). Two ancient classes of MIKC-type MADS-box genes are present in the moss *Physcomitrella patens*. *Mol. Biol. Evol.* **19**: 801–814.
- Howell, M.D., Fahlgren, N., Chapman, E.J., Cumbie, J.S., Sullivan, C.M., Givan, S.A., Kasschau, K.D., and Carrington, J.C. (2007). Genome-wide analysis of the RNA-DEPENDENT RNA POLYMERASE6/DICER-LIKE4 pathway in *Arabidopsis* reveals dependency on miRNA- and ta-siRNA-directed targeting. *Plant Cell* **19**: 926–942.
- Huang, X., and Madan, A. (1999). CAP3: A DNA sequence assembly program. *Genome Res.* **9**: 868–877.
- Hunter, C., Willmann, M.R., Wu, G., Yoshikawa, M., de la Luz Gutierrez-Nava, M., and Poethig, R.S. (2006). Trans-acting siRNA-mediated repression of *ETTIN* and *ARF4* regulates heteroblasty in *Arabidopsis*. *Development* **133**: 2973–2981.
- Johnson, C., Bowman, L., Adai, A.T., Vance, V., and Sundaresan, V. (2007). CSRDB: A small RNA integrated database and browser resource for cereals. *Nucleic Acids Res.* **35**: D829–D833.
- Jones-Rhoades, M.W., and Bartel, D.P. (2004). Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Mol. Cell* **14**: 787–799.
- Jones-Rhoades, M.W., Bartel, D.P., and Bartel, B. (2006). MicroRNAs and their regulatory roles in plants. *Annu. Rev. Plant Biol.* **57**: 19–53.
- Kasschau, K.D., Fahlgren, N., Chapman, E.J., Sullivan, C.M., Cumbie, J.S., Givan, S.A., and Carrington, J.C. (2007). Genome-wide profiling and analysis of *Arabidopsis* siRNAs. *PLoS Biol.* **5**: e57.
- Kasschau, K.D., Xie, Z., Allen, E., Llave, C., Chapman, E.J., Krizan, K.A., and Carrington, J.C. (2003). P1/HC-Pro, a viral suppressor of RNA silencing, interferes with *Arabidopsis* development and miRNA function. *Dev. Cell* **4**: 205–217.

- Kent, W.J.** (2002). BLAT—The BLAST-like alignment tool. *Genome Res.* **12**: 656–664.
- Krogan, N.T., and Ashton, N.W.** (2000). Ancestry of plant MADS-box genes revealed by bryophyte (*Physcomitrella patens*) homologues. *New Phytol.* **147**: 505–517.
- Kumar, S., Tamura, K., and Nei, M.** (2004). MEGA3: Integrated software for Molecular Evolutionary Genetics Analysis and sequence alignment. *Brief. Bioinform.* **5**: 150–163.
- Kurihara, Y., and Watanabe, Y.** (2004). *Arabidopsis* micro-RNA biogenesis through Dicer-like 1 protein functions. *Proc. Natl. Acad. Sci. USA* **101**: 12753–12758.
- Lagos-Quintana, M., Rauhut, R., Lendeckel, W., and Tuschl, T.** (2001). Identification of novel genes coding for small expressed RNAs. *Science* **294**: 853–858.
- Lau, N.C., Lim, L.P., Weinstein, E.G., and Bartel, D.P.** (2001). An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* **294**: 858–862.
- Lee, R.C., and Ambros, V.** (2001). An extensive class of small RNAs in *Caenorhabditis elegans*. *Science* **294**: 862–864.
- Lee, Y., Ahn, C., Han, J., Choi, H., Kim, J., Yim, J., Lee, J., Provost, P., Radmark, O., Kim, S., and Kim, V.N.** (2003). The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**: 415–419.
- Lee, Y., Kim, M., Han, J., Yeom, K.H., Lee, S., Baek, S.H., and Kim, V.N.** (2004). MicroRNA genes are transcribed by RNA polymerase II. *EMBO J.* **23**: 4051–4060.
- Li, Y., Li, W., and Jin, Y.X.** (2005). Computational identification of novel family members of microRNA genes in *Arabidopsis thaliana* and *Oryza sativa*. *Acta Biochim. Biophys. Sin. (Shanghai)* **37**: 75–87.
- Liu, J., Carmell, M.A., Rivas, F.V., Marsden, C.G., Thomson, J.M., Song, J.J., Hammond, S.M., Joshua-Tor, L., and Hannon, G.J.** (2004). Argonaute2 is the catalytic engine of mammalian RNAi. *Science* **305**: 1437–1441.
- Llave, C., Kasschau, K.D., Rector, M.A., and Carrington, J.C.** (2002b). Endogenous and silencing-associated small RNAs in plants. *Plant Cell* **14**: 1605–1619.
- Llave, C., Xie, Z., Kasschau, K.D., and Carrington, J.C.** (2002a). Cleavage of *Scarecrow-like* mRNA targets directed by a class of *Arabidopsis* miRNA. *Science* **297**: 2053–2056.
- Lu, C., Kulkarni, K., Souret, F.F., MuthuValliappan, R., Tej, S.S., Poethig, R.S., Henderson, I.R., Jacobsen, S.E., Wang, W., Green, P.J., and Meyers, B.C.** (2006). MicroRNAs and other small RNAs enriched in the *Arabidopsis* RNA-dependent RNA polymerase-2 mutant. *Genome Res.* **16**: 1276–1288.
- Lu, C., Tej, S.S., Luo, S., Haudenschild, C.D., Meyers, B.C., and Green, P.J.** (2005). Elucidation of the small RNA component of the transcriptome. *Science* **309**: 1567–1569.
- Mallory, A.C., and Vaucheret, H.** (2006). Functions of microRNAs and related small RNAs in plants. *Nat. Genet.* **38**(Suppl): S31–S36.
- Margis, R., Fusaro, A.F., Smith, N.A., Curtin, S.J., Watson, J.M., Finnegan, E.J., and Waterhouse, P.M.** (2006). The evolution and diversification of Dicers in plants. *FEBS Lett.* **580**: 2442–2450.
- Meister, G., Landthaler, M., Patkaniowska, A., Dorsett, Y., Teng, G., and Tuschl, T.** (2004). Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol. Cell* **15**: 185–197.
- Molnár, A., Schwach, F., Studholme, D.J., Thuenemann, E.C., and Baulcombe, D.C.** (2007). miRNAs control gene expression in the single-cell alga *Chlamydomonas reinhardtii*. *Nature* <http://dx.doi.org/10.1038/nature05903>.
- Nobuta, K., Venu, R.C., Lu, C., Belo, A., Vemaraju, K., Kulkarni, K., Wang, W., Pillay, M., Green, P.J., Wang, G.L., and Meyers, B.C.** (2007). An expression atlas of rice mRNAs and small RNAs. *Nat. Biotechnol.* **25**: 473–477.
- Okushima, Y., et al.** (2005). Functional genomic analysis of the *AUXIN RESPONSE FACTOR* gene family members in *Arabidopsis thaliana*: Unique and overlapping functions of *ARF7* and *ARF19*. *Plant Cell* **17**: 444–463.
- Pak, J., and Fire, A.** (2007). Distinct populations of primary and secondary effectors during RNAi in *C. elegans*. *Science* **315**: 241–244.
- Palatnik, J.F., Allen, E., Wu, X., Schommer, C., Schwab, R., Carrington, J.C., and Weigel, D.** (2003). Control of leaf morphogenesis by microRNAs. *Nature* **425**: 257–263.
- Park, W., Li, J., Song, R., Messing, J., and Chen, X.** (2002). CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in micro-RNA metabolism in *Arabidopsis thaliana*. *Curr. Biol.* **12**: 1484–1495.
- Peragine, A., Yoshikawa, M., Wu, G., Albrecht, H.L., and Poethig, R.S.** (2004). *SGS3* and *SGS2/SDE1/RDR6* are required for juvenile development and the production of *trans*-acting siRNAs in *Arabidopsis*. *Genes Dev.* **18**: 2368–2379.
- Poirot, O., O'Toole, E., and Notredame, C.** (2003). Tcoffee@igs: A web server for computing, evaluating and combining multiple sequence alignments. *Nucleic Acids Res.* **31**: 3503–3506.
- Qi, Y., Denli, A.M., and Hannon, G.J.** (2005). Biochemical specialization within *Arabidopsis* RNA silencing pathways. *Mol. Cell* **19**: 421–428.
- Qi, Y., He, X., Wang, X.J., Kohany, O., Jurka, J., and Hannon, G.J.** (2006). Distinct catalytic and non-catalytic roles of ARGONAUTE4 in RNA-directed DNA methylation. *Nature* **443**: 1008–1012.
- Rajagopalan, R., Vaucheret, H., Trejo, J., and Bartel, D.P.** (2006). A diverse and evolutionarily fluid set of microRNAs in *Arabidopsis thaliana*. *Genes Dev.* **20**: 3407–3425.
- Reinhart, B.J., Weinstein, E.G., Rhoades, M.W., Bartel, B., and Bartel, D.P.** (2002). MicroRNAs in plants. *Genes Dev.* **16**: 1616–1626.
- Rhoades, M.W., Reinhart, B.J., Lim, L.P., Burge, C.B., Bartel, B., and Bartel, D.P.** (2002). Prediction of plant microRNA targets. *Cell* **110**: 513–520.
- Ronshaugen, M., Biemar, F., Piel, J., Levine, M., and Lai, E.C.** (2005). The *Drosophila* microRNA *iab-4* causes a dominant homeotic transformation of halteres to wings. *Genes Dev.* **19**: 2947–2952.
- Ruby, J.G., Jan, C., Player, C., Axtell, M.J., Lee, W., Nusbaum, C., Ge, H., and Bartel, D.** (2006). Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in *Caenorhabditis elegans*. *Cell* **127**: 1193–1207.
- Schroda, M.** (2006). RNA silencing in *Chlamydomonas*: Mechanisms and tools. *Curr. Genet.* **49**: 69–84.
- Shi, Z., Wang, J., Wan, X., Shen, G., Wang, X., and Zhang, J.** (2007). Over-expression of rice *OsAGO7* gene induces upward curling of the leaf blade that enhanced erect-leaf habit. *Planta* **226**: 99–108.
- Sijen, T., Steiner, F.A., Thijssen, K.L., and Plasterk, R.H.** (2007). Secondary siRNAs result from unprimed RNA synthesis and form a distinct class. *Science* **315**: 244–247.
- Singer, S.D., Krogan, N.T., and Ashton, N.W.** (2007). Clues about the ancestral roles of plant MADS-box genes from a functional analysis of moss homologues. *Plant Cell Rep.* <http://dx.doi.org/10.1007/s00299-007-0312-0>.
- Sunkar, R., Girke, T., Jain, P.K., and Zhu, J.K.** (2005). Cloning and characterization of microRNAs from rice. *Plant Cell* **17**: 1397–1411.
- Talmor-Neiman, M., Stav, R., Frank, W., Voss, B., and Arazi, T.** (2006a). Novel micro-RNAs and intermediates of micro-RNA biogenesis from moss. *Plant J.* **47**: 25–37.
- Talmor-Neiman, M., Stav, R., Klipcan, L., Buxdorf, K., Baulcombe, D.C., and Arazi, T.** (2006b). Identification of *trans*-acting siRNAs in moss and an RNA-dependent RNA polymerase required for their biogenesis. *Plant J.* **48**: 511–521.

- Vaucheret, H., Mallory, A.C., and Bartel, D.P.** (2006). AGO1 homeostasis entails coexpression of *MIR168* and *AGO1* and preferential stabilization of miR168 by AGO1. *Mol. Cell* **22**: 129–136.
- Vaucheret, H., Vazquez, F., Crete, P., and Bartel, D.P.** (2004). The action of *ARGONAUTE1* in the miRNA pathway and its regulation by the miRNA pathway are crucial for plant development. *Genes Dev.* **18**: 1187–1197.
- Vazquez, F., Vaucheret, H., Rajagopalan, R., Lepers, C., Gascioli, V., Mallory, A.C., Hilbert, J.L., Bartel, D.P., and Crete, P.** (2004). Endogenous *trans*-acting siRNAs regulate the accumulation of *Arabidopsis* mRNAs. *Mol. Cell* **16**: 69–79.
- Wang, W., et al.** (2005). Construction of a bacterial artificial chromosome library from the spikemoss *Selaginella moellendorffii*: A new resource for plant comparative genomics. *BMC Plant Biol.* **5**: 10.
- Wienholds, E., and Plasterk, R.H.** (2005). MicroRNA function in animal development. *FEBS Lett.* **579**: 5911–5922.
- Xie, Z., Allen, E., Fahlgren, N., Calamar, A., Givan, S.A., and Carrington, J.C.** (2005). Expression of *Arabidopsis* *MIRNA* genes. *Plant Physiol.* **138**: 2145–2154.
- Xie, Z., Kasschau, K.D., and Carrington, J.C.** (2003). Negative feedback regulation of *Dicer-Like1* in *Arabidopsis* by microRNA-guided mRNA degradation. *Curr. Biol.* **13**: 784–789.
- Yang, Z., Ebright, Y.W., Yu, B., and Chen, X.** (2006). HEN1 recognizes 21–24 nt small RNA duplexes and deposits a methyl group onto the 2' OH of the 3' terminal nucleotide. *Nucleic Acids Res.* **34**: 667–675.
- Yekta, S., Shih, I.H., and Bartel, D.P.** (2004). MicroRNA-directed cleavage of *HOXB8* mRNA. *Science* **304**: 594–596.
- Yoshikawa, M., Peragine, A., Park, M.Y., and Poethig, R.S.** (2005). A pathway for the biogenesis of *trans*-acting siRNAs in *Arabidopsis*. *Genes Dev.* **19**: 2164–2175.
- Yu, B., Yang, Z., Li, J., Minakhina, S., Yang, M., Padgett, R.W., Steward, R., and Chen, X.** (2005). Methylation as a crucial step in plant microRNA biogenesis. *Science* **307**: 932–935.
- Zhang, X., Henderson, I.R., Lu, C., Green, P.J., and Jacobsen, S.E.** (2007). Role of RNA polymerase IV in plant small RNA metabolism. *Proc. Natl. Acad. Sci. USA* **104**: 4536–4541.
- Zhao, T., Li, G., Mi, S., Li, S., Hannon, G.J., Wang, X.J., and Qi, Y.** (2007). A complex system of small RNAs in the unicellular green alga *Chlamydomonas reinhardtii*. *Genes Dev.* **21**: 1190–1203.
- Zheng, X., Zhu, J., Kapoor, A., and Zhu, J.K.** (2007). Role of *Arabidopsis* AGO6 in siRNA accumulation, DNA methylation and transcriptional gene silencing. *EMBO J.* **26**: 1691–1701.
- Zilberman, D., Cao, X., and Jacobsen, S.E.** (2003). ARGONAUTE4 control of locus-specific siRNA accumulation and DNA and histone methylation. *Science* **299**: 716–719.