

Is the *IS1* Transposase, InsAB', the Only *IS1*-Encoded Protein Required for Efficient Transposition?

J. M. ESCOUBAS,† D. LANE, AND M. CHANDLER*

Molecular Genetics and Microbiology, UPR9007 du Centre National de la Recherche Scientifique, 31062 Toulouse Cedex, France

Received 2 May 1994/Accepted 2 July 1994

The transposase of the bacterial insertion sequence *IS1* is normally expressed by inefficient translational frameshifting between an upstream reading frame which itself specifies a transposition inhibitor, InsA, and a second consecutive reading frame located immediately downstream. A fused-frame mutant which carries an additional base pair inserted at the point of frameshifting was constructed. This mutant exhibits high transposition activity and should express the transposase, InsAB', constitutively without frameshifting. Unexpectedly, a second protein species was observed to be expressed from this mutant. We demonstrate here that this protein, InsA*, results from continued frameshifting on the modified frameshift motif. The protein retains the activities of the repressor InsA. Its elimination, by further modification of the frameshift motif, results in a further increase in various transposition activities of *IS1*. These results support the hypothesis that a single *IS1*-encoded protein, InsAB', is necessary for transposition.

IS1 is a small, genetically compact, insertion sequence present in the chromosomes of several enterobacterial species (10). It exhibits two partially overlapping open reading frames, *insA* and *insB'*, located respectively in relative reading phases 0 and -1, whose integrity is essential for transposition (5, 9). The product of the first, InsA, binds specifically to both left and right *IS1* terminal inverted repeats, IRL and IRR (16) (Fig. 1A), repressing its own transcription from a promoter located partially in IRL and simultaneously inhibiting transposition (8, 17). A second protein, InsAB', carries domains contributed by both reading frames and is produced by inefficient translational frameshifting between *insA* and *insB'* (3, 13). Frameshifting has been shown to occur on an A₆C sequence (3, 13) (Fig. 1B), as found also in the synthesis of the Gag-Pol polyprotein in some retroviruses (for a review, see reference 15). The InsAB' protein can be produced "constitutively" by introduction of a single additional A residue into the A₆C run to form A₇C (2, 13). This insertion mutation physically fuses *insA* and *insB'* and increases the frequency of *IS1* transposition by a factor of between 10² and 10⁴ (2, 13). The mutation should eliminate expression of the InsA repressor since it places the InsA termination codon out of phase (Fig. 1C). The large stimulation in transposition frequency suggested that the InsAB' fusion protein is the *IS1* transposase and led to the idea that it is the only *IS1*-encoded protein necessary for transposition.

It has been shown previously (3), by using a phage T7 expression system, that the A₇C frameshift mutant, carried by plasmid pMET11, specifies a protein of the size expected for InsAB' and that this protein is expressed at a high level compared with that achieved by natural frameshifting which occurs in the wild-type *IS1* coding sequences carried by plasmid pZBT29 (3) (Fig. 2B, lanes 1 and 2). The natural frequency of translational frameshifting, estimated from mea-

surements of the relative amounts of InsAB' and InsA for the wild-type *IS1* coding sequences, was about 1% under these conditions. Since the InsA termination codon is placed out of phase in the frameshift mutant (Fig. 1C), only one protein species, InsAB', should be produced. However, inspection of the results (Fig. 2B, lane 2) indicated the presence of a significant amount of a second protein species. This species had an apparent molecular mass of 10.7 kDa, slightly larger than that observed for InsA (as seen with plasmid pZBT29 [Fig. 2B, lane 1]). This raised the possibility that a third protein is normally made in small amounts from wild-type *IS1* and that its presence had been obscured in previous experiments by high levels of InsA. The unexplained presence of this additional protein brings into question the contention that InsAB' alone is necessary for *IS1* transposition. It was therefore important to determine its origin and its possible effects on transposition activity.

Origin of the additional *IS1*-specified protein species. One possibility is that the additional protein reflects the occurrence of an alternative translational frameshift. Experiments with retroviral frameshift signals have demonstrated that several heptanucleotide sequences are capable of inducing -1 translational frameshifting (15). Among these is an A₇C signature similar to that created in the fused-frame mutant, pMET11. One explanation for the appearance of the 10.7-kDa protein from pMET11 is that it results from frameshifting on the A₇C motif. This explanation is supported by inspection of the nucleotide sequence (Fig. 1C). Following a -1 frameshift on the A₇C sequence, the ribosome would encounter a termination codon (UGA) approximately 16 codons downstream and generate a protein of 100 amino acids with a predicted molecular mass of 10.7 kDa. The resulting protein would carry the NH₂-terminal 85 amino acids of InsA, including the α -helix-turn- α -helix motif thought to be involved in recognizing and binding to the *IS1* terminal repeats (16) (Fig. 1D). To test this possibility, an *IS1* derivative in which the sequence GA₂GA₃C replaces the A₆C motif was generated by site-directed mutagenesis. Like the A₇C mutation, this also fuses the *insA* and *insB'* reading frames, but the GA₂GA₃C mutation disrupts the presumed alternative interactions between tRNAs and mRNA in the -1 reading phase (6, 7) and should reduce

* Corresponding author. Mailing address: Molecular Genetics and Microbiology, UPR9007 du CNRS, 118 Rte de Narbonne, 31062 Toulouse Cedex, France. Phone: (33) 613355858. Fax: (33) 613355886. Electronic mail address: mike@biotoul.ibcg.fr.

† Present address: Oceanographic and Atmospheric Science Division, Department of Applied Science, Brookhaven National Laboratory, Upton, NY 11973.




FIG. 1. Organization of IS1. (A) Structure of IS1. Left (IRL) and right (IRR) inverted terminal repeats are shown as solid boxes. Relative positions of the *insA* and *insB'* reading frames, together with their overlap region, are also shown within the open box representing IS1. The InsA protein is represented by a hatched box. The InsA and InsB' components of the InsAB' frameshift product are shown as hatched and stippled boxes, respectively. Arrows indicate the probable region of action of InsA and InsAB' proteins. (B and C) RNA and protein sequence in the crossover region between the two open reading frames. Codons shown above the RNA sequence show the product of direct translational readout. Those below show the product of a -1 translational frameshift. Presumed amino acids at the point of frameshift are shown in parentheses. In panel B, the heptanucleotide A₆C frameshift sequence involved in production of InsAB' from the wild-type IS1 coding sequence (upper sequence) and the A₇C derivative (lower sequence) are indicated in boldface type as is the UAA termination codon for InsA. In panel C, the A₇C fused-frame sequence is shown in boldface type, as is the UGA termination codon for InsA*.

(D) Schematic representation of InsA and InsA*. The solid boxes indicate the position of the potential helix-turn-helix motif in the InsA and InsA* proteins. The difference in the amino acid sequence of the carboxy-terminal end of the protein is distinguished by a stippled box and a hatched box for InsA and InsA*, respectively. The numbers represent amino acid residues.




FIG. 2. IS1 proteins produced in the phage T7 expression system. (A) Structure of the plasmids used. Mutant IS1 coding sequences are indicated in parentheses below the plasmid name. The construction of pZBT29 and pMET11 has been described previously. Plasmid pMET24 was generated in the same way as pMET11, except that the oligonucleotide used in the site-directed mutagenesis carried the appropriate GA₂GA₃C mutation. Transcription from the phage T7 promoter (pT7) is from left to right. Open reading frames are indicated as hatched and open boxes within IS1 (larger box). The pointed end of this box represents IRR. T ϕ indicates the presence of a phage T7 transcription terminator. The β -lactamase gene (Ap) is also shown as an open box. (B) Expression of IS1 proteins in the T7 expression system. The methods used were described previously (3). Lanes: 1, pZBT29; 2, pMET11; 3, pMET24. (C) Gel retardation with crude-cell extracts. The methods used were described previously (18). DNA fragments were prepared by PCR with pMP20 (11) as a matrix. Lanes: 1, no extract; 2 and 3, pZBT29; 4 and 5, pMET24; 6 and 7, pMET11. Lanes 2, 4, and 6 show results obtained with a 1:10 dilution of the crude extracts. Lanes 3, 5, and 7 show results obtained with a dilution of 1:100. Band A, unbound DNA fragment; band B (and C), InsA (InsA*)-specific band.

or eliminate frameshifting over these nucleotides. The amino acid sequence of the InsAB' protein produced by this mutant should be identical to that produced from the A₇C frameshift mutant. The proteins produced from this plasmid, pMET24, in the phage T7 expression system are shown in Fig. 2B, lane 3. It



PLASMID	STRUCTURE	TRANSF. FREQ.	β -gal
pZPT2		$2.9 \pm 0.2 \times 10^{-7}$ (4)	88 ± 4 (6)
pMET13 (A ₆ C)		$7.3 \pm 1.4 \times 10^{-7}$ (8)	77 ± 7 (6)
pMET12 (A ₇ C)		$2.5 \pm 0.5 \times 10^{-3}$ (10)	523 ± 66 (8)
pMET23 (GA ₂ GA ₃ C)		$1.2 \pm 0.2 \times 10^{-2}$ (5)	766 ± 58 (4)

FIG. 3. Transposition frequencies and SOS-inducing capacities. The structure of the plasmids used is indicated on the left of the figure together with the wild-type and two mutant *IS1* coding sequences (in parentheses). The construction of pMET13 and pMET12 has been described previously (3), as has that of pZPT2 (17). Plasmid pMET23 was generated in the same way as pMET12 except that the oligonucleotide used in the site-directed mutagenesis carried the appropriate GA₂GA₃C mutation. Transcription from the phage λ promoter (p_L) is from right to left. Open reading frames are indicated as hatched and stippled boxes within *IS1* (larger box). The pointed end of this box represents IRR. T ϕ indicates the presence of a phage T7 transcription terminator. The β -lactamase gene (*Ap*) is also shown. In addition, the presence of a synthetic *IS1*-based transposon, Ω -on, composed of inverted copies of IRL flanking a gene specifying resistance to streptomycin (*Sm*) and spectinomycin (*Sp*) is also shown. Transposition frequencies were determined by using a standard mating-out assay as previously described and represent the sum of the frequencies of cointegrate formation and direct transposition (3). The method used in determining the capacity for SOS induction by measuring the level of β -galactosidase (β -gal.) production from a *lacZ* gene driven from an SOS inducible promoter has also been described (7a). The numbers in parentheses show the number of independent determinations for each value. Standard errors were less than 10% for both transposition and β -galactosidase activities.

is clear that the mutation eliminates the 10.7-kDa species while maintaining constitutive *InsAB'* synthesis, supporting the notion that the 10.7-kDa species is produced by frameshifting on the A₇C motif. The protein is called *InsA** in the following discussion.

Effect of *InsA on intermolecular transposition activity.** In previous experiments, transposition activity was assessed primarily for the wild-type A₆C coding sequences and the A₇C mutant. It was therefore important to determine the influence of *InsA** on these results and the effect of its absence. In a first set of experiments, we have compared transposition frequencies obtained by using wild-type A₆C and the A₇C and GA₂GA₃C mutant *IS1* coding sequences in a standard mating-out assay (3). Plasmids pMET13 (A₆C), pMET12 (A₇C), and pMET23 (GA₂GA₃C) carry the *IS1* coding sequences under control of the phage λ p_L promoter. A single IRR is located at its natural position at the 3' end of *insB'*. In addition, the plasmids carry an artificial *IS1*-based transposon, Ω -on, composed of an Sp^r Sm^r cassette flanked by inverted copies of IRL (11). As discussed elsewhere (3), use of the p_L promoter and elimination of the endogenous IRL promoter and *InsA*-binding site eliminate any *InsA* repression of transposase expression. The results of mating-out assays, by using the conjugative plasmid pOX38Km (1) as a recipient replicon, are presented in Fig. 3. As expected, use of the A₇C mutation results in a substantial increase in overall transposition fre-

quency compared with the wild-type A₆C sequence. Furthermore, substitution of GA₂GA₃C for the A₇ mutation results in a further fourfold increase to a transposition frequency of 1%. These results suggest that *InsA**, produced by the A₇C mutant, is capable of inhibiting intermolecular transposition.

Effect of *InsA on induction of an SOS signal.** Another property associated with the *IS1* transposase, like that of *IS10* (4, 12), is its capacity, in the presence of suitable transposon ends, to induce the host SOS response (7a). This presumably results from the endonucleolytic activity associated with the transposase. To determine whether *InsA** influences the ability of *InsAB'* to induce the SOS response, the activity of pMET13, pMET12, and pMET23 was determined in a standard SOS induction assay. Briefly, the assay uses an SOS indicator strain in which a *lacZ* gene is driven by the p_L promoter of a defective lambdaoid 434 prophage (2). Induction of the SOS response results in cleavage of the 434 repressor and consequent induction of β -galactosidase synthesis. The results obtained with the three plasmids by using a standard assay (7) are presented in Fig. 3. SOS induction by pMET13 is low. A significant increase is observed with pMET12, while pMET23 gives a further stimulation in SOS activity. Thus, as in the case of intermolecular transposition, *InsA** interferes with transposase activity.

Sequence-specific DNA binding. The results presented above are consistent with the idea that the effects of *InsA**, like those of *InsA*, are due to its capacity to bind to the ends of *IS1*. To determine the nucleic acid-binding properties of *InsA**, extracts of cultures containing high levels of *IS1*-derived proteins were used in gel retardation assays with labeled DNA fragments of 85 bp carrying 28 bp of IRL (18). The results are presented in Fig. 2C. It can be seen that, while extracts derived from strains carrying the A₇C mutant (pMET11) exhibit a distinct retarded band (lanes 6 and 7), those derived from cells carrying the GA₂GA₃C mutant (pMET24) showed little retardation activity (lanes 4 and 5). This suggests that *InsA** binding is responsible for the retarded species. It is interesting that although *InsAB'* should also carry the *InsA*-specific nucleic acid-binding domain, we could not observe specific complexes of higher apparent molecular mass expected to arise from *InsAB'* binding. This inability to detect efficient *InsAB'* binding to the terminal inverted repeats is under investigation. The electrophoretic mobility of the retarded band obtained in the presence of *InsA** is similar to that obtained with an extract enriched for *InsA* itself (plasmid pZB29; lanes 2 and 3). The *InsA** protein may therefore behave in a similar way to *InsA* itself in influencing *IS1* transposition.

The results presented here provide evidence that a significant level of translational frameshifting occurs on the mutant A₇ heptamer derived from the A₆(C) wild-type *IS1* sequence and designed to physically fuse the *insA* and *insB'* reading frames of *IS1*. This mutation results in high-level synthesis of the fusion protein *InsAB'* without requiring a -1 translational frameshift. Frameshifting on the mutant A₇ produces a small protein, *InsA**, related to the *IS1* transposition inhibitor/repressor, *InsA*. The level of frameshifting on the A₇ heptamer cannot be assessed accurately from our present data. Estimates from the intensities of the relevant protein species from autoradiographs such as that shown in Fig. 2B suggest that it might be as high as 55% (compared with less than 1% on the wild-type A₆C motif) (3). However, this value is certainly an overestimate, since it has been demonstrated that, at least in vitro, the T7 RNA polymerase specifically amplifies frameshifting on an A₇ sequence (14). We have not attempted to assess the "normal" in vivo rate by more reliable techniques.

Inclusion of the GA₂GA₃C sequence in place of A₇C

eliminates InSA*, confirming that translational frameshifting occurs on the A₇ heptamer. InSA* has a molecular mass slightly larger than that of InSA itself on sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Calculation of the predicted molecular mass of InSA* gives a value of 10.7 kDa compared with 9.8 kDa for InSA. The InSA* protein carries sufficient information to bind specifically to the left terminal inverted repeat of IS1. Since the amino acid sequences of InSA and InSA* presumably diverge at amino acid 86 (Fig. 1B to D), the specific DNA sequence recognition domain must be included within the first 85 amino acids of the proteins. We have also provided evidence which suggests that InSA*, like InSA, inhibits IS1 transposition. This inhibitory activity of InSA* is also illustrated by its ability to reduce the transposase-mediated induction of the SOS response. The most important aspect of these results is that elimination of InSA* production, by use of the GA₂GA₃C motif, leads to an increase in the various activities of InsAB'. The data thus further support the idea that InsAB' is the IS1 transposase and is the only IS1-encoded protein required for efficient transposition.

We thank all members of the mobile genetic elements group for discussions. We are also grateful to Pierre Prentki for providing the initial stimulus for this analysis.

This work was supported by CNRS grant UPR9007 and NATO travel grant 0.1669.

REFERENCES

1. Chandler, M., and D. J. Galas. 1983. Cointegrate formation by Tn9: activity of IS1 is modulated by external sequences. *J. Mol. Biol.* **170**:61-91.
2. Elespuru, R. K., and M. B. Yarmolinsky. 1979. A colorimetric assay of lysogenic induction designed for screening potential carcinogenic and carcinostatic agents. *Environ. Mutagen.* **1**:65-78.
3. Escoubas, J. M., M.-F. Prère, O. Fayet, I. Salvagnol, D. J. Galas, D. Zerbib, and M. Chandler. 1991. Translational control of transposition activity of the bacterial insertion sequence IS1. *EMBO J.* **10**:705-712.
4. Haniford, D. B., A. R. Chelouche, and N. Kleckner. 1989. A specific class of IS10 transposase mutants are blocked for target site interactions and promote formation of an excised transposon fragment. *Cell* **59**:385-394.
5. Jackowec, M., P. Prentki, M. Chandler, and D. J. Galas. 1988. Mutational analysis of the open reading frames in the transposable element IS1. *Genetics* **120**:47-55.
6. Jacks, T., M. D. Power, F. R. Masiarz, P. A. Luciw, P. J. Barr, and H. E. Varmus. 1988. Characterization of ribosomal frameshifting in HIV-1 gag-pol expression. *Nature (London)* **331**:280-283.
7. Jacks, T., K. Townsley, H. E. Varmus, and J. Majors. 1987. Two efficient ribosomal frameshifting events are required for synthesis of mouse mammary tumor virus gag-related polyproteins. *Proc. Natl. Acad. Sci. USA* **84**:4298-4302.
- 7a. Lane, D., J. Cavallé, and M. Chandler. *J. Mol. Biol.*, in press.
8. Machida, C., and Y. Machida. 1989. Regulation of IS1 transposition by the *insA* gene product. *J. Mol. Biol.* **208**:567-574.
9. Machida, Y., C. Machida, and E. Ohtsubo. 1984. Insertion element IS1 encodes two structural genes required for its transposition. *J. Mol. Biol.* **177**:229-245.
10. Nyman, K., H. Ohtsubo, D. Davison, and E. Ohtsubo. 1983. Distribution of insertion element IS1 in natural isolates of *Escherichia coli*. *Mol. Gen. Genet.* **189**:516-518.
11. Prentki, P., M.-H. Pham, P. Gamas, M. Chandler, and D. Galas. 1987. Artificial transposable elements in the study of the ends of IS1. *Gene* **61**:91-101.
12. Roberts, D., and N. Kleckner. 1988. Tn10 transposition promotes RecA-dependent induction of λ prophage. *Proc. Natl. Acad. Sci. USA* **85**:6037-6041.
13. Sekine, Y., and E. Ohtsubo. 1989. Frameshifting is required for production of the transposase encoded by insertion sequence 1. *Proc. Natl. Acad. Sci. USA* **86**:4609-4613.
14. Tsuchihashi, Z., and P. O. Brown. 1992. Sequence requirements for efficient translational frameshifting in the *Escherichia coli* *dnaX* gene and the role of an unstable interaction between tRNA^{Lys} and an AAG lysine codon. *Genes Dev.* **6**:511-519.
15. Weiss, R. B., D. M. Dunn, M. Shuh, J. F. Atkins, and R. F. Gesteland. 1989. *E. coli* ribosomes rephase on retroviral frameshift signals at rates ranging from 2 to 50%. *New Biol.* **1**:159-169.
16. Zerbib, D., M. Jakowec, P. Prentki, D. Galas, and M. Chandler. 1987. Expression of proteins essential for IS1 transposition: specific binding of InSA to the ends of IS1. *EMBO J.* **6**:3163-3169.
17. Zerbib, D., P. Polard, J.-M. Escoubas, D. Galas, and M. Chandler. 1990. The regulatory role of the IS1-encoded InSA protein in transposition. *Mol. Microbiol.* **4**:471-477.
18. Zerbib, D., P. Prentki, P. Gamas, E. Freund, D. J. Galas, and M. Chandler. 1990. Functional organization of the ends of IS1: specific binding site for an IS1-encoded protein. *Mol. Microbiol.* **4**:1477-1486.