

Research article

Open Access

## Molecular cloning and analysis of zebrafish voltage-gated sodium channel beta subunit genes: implications for the evolution of electrical signaling in vertebrates

Sameer S Chopra<sup>1</sup>, Hiroshi Watanabe<sup>1</sup>, Tao P Zhong<sup>†2,3</sup> and Dan M Roden<sup>\*†1,3</sup>

Address: <sup>1</sup>Department of Pharmacology, Vanderbilt University School of Medicine, Nashville, TN, 37232, USA, <sup>2</sup>Cell & Developmental Biology, Vanderbilt University School of Medicine, Nashville, TN, 37232, USA and <sup>3</sup>Medicine, Vanderbilt University School of Medicine, Nashville, TN, 37232, USA

Email: Sameer S Chopra - sameer.chopra@vanderbilt.edu; Hiroshi Watanabe - hiroshi.watanabe@vanderbilt.edu; Tao P Zhong - tao.zhong@vanderbilt.edu; Dan M Roden\* - dan.roden@vanderbilt.edu

\* Corresponding author †Equal contributors

Published: 10 July 2007

Received: 22 December 2006

BMC Evolutionary Biology 2007, 7:113 doi:10.1186/1471-2148-7-113

Accepted: 10 July 2007

This article is available from: <http://www.biomedcentral.com/1471-2148/7/113>

© 2007 Chopra et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Action potential generation in excitable cells such as myocytes and neurons critically depends on voltage-gated sodium channels. In mammals, sodium channels exist as macromolecular complexes that include a pore-forming alpha subunit and 1 or more modulatory beta subunits. Although alpha subunit genes have been cloned from diverse metazoans including flies, jellyfish, and humans, beta subunits have not previously been identified in any non-mammalian species. To gain further insight into the evolution of electrical signaling in vertebrates, we investigated beta subunit genes in the teleost *Danio rerio* (zebrafish).

**Results:** We identified and cloned single zebrafish gene homologs for beta1-beta3 (*zbeta1-zbeta3*) and duplicate genes for beta4 (*zbeta4.1, zbeta4.2*). Sodium channel beta subunit loci are similarly organized in fish and mammalian genomes. Unlike their mammalian counterparts, *zbeta1* and *zbeta2* subunit genes display extensive alternative splicing. Zebrafish beta subunit genes and their splice variants are differentially-expressed in excitable tissues, indicating tissue-specific regulation of *zbeta1-4* expression and splicing. Co-expression of the genes encoding *zbeta1* and the zebrafish sodium channel alpha subunit  $Na_v1.5$  in Chinese Hamster Ovary cells increased sodium current and altered channel gating, demonstrating functional interactions between zebrafish alpha and beta subunits. Analysis of the synteny and phylogeny of mammalian, teleost, amphibian, and avian beta subunit and related genes indicated that all extant vertebrate beta subunits are orthologous, that beta2/beta4 and beta1/beta3 share common ancestry, and that beta subunits are closely related to other proteins sharing the V-type immunoglobulin domain structure. Vertebrate sodium channel beta subunit genes were not identified in the genomes of invertebrate chordates and are unrelated to known subunits of the *para* sodium channel in *Drosophila*.

**Conclusion:** The identification of conserved orthologs to all 4 voltage-gated sodium channel beta subunit genes in zebrafish and the lack of evidence for beta subunit genes in invertebrate chordates together indicate that this gene family emerged early in vertebrate evolution, prior to the divergence of teleosts and tetrapods. The evolutionary history of sodium channel beta subunits suggests that these genes may have played a key role in the diversification and specialization of electrical signaling in early vertebrates.

## Background

Coordinated electrical signals in the metazoan nervous system, heart, and skeletal muscle depend on the generation of action potentials, rapid changes in membrane potential mediated by the passage of ions through voltage-gated ion channels [1]. The initial upstroke of the action potential in most excitable cells is determined by sodium channels, membrane proteins characterized by rapid activation and inactivation and high ionic conductance [1-3]. Unlike evolutionarily-ancient, tetrameric potassium channels, sodium channels are comprised of four homologous domains that are encoded by a single polypeptide [1-3]. Sodium channels are suspected to have evolved from structurally-similar calcium channels, which in turn likely arose following the duplication of potassium channel genes [1-3]. The use of sodium as a conducting ion rather than calcium is thought to have permitted rapid conduction and high-frequency electrical signaling in the rudimentary nervous systems of early multicellular animals, without the numerous additional intracellular effects mediated by calcium's role as a second messenger [1,4].

In mammals, voltage-gated sodium channels are multi-protein complexes that include a pore-forming  $\alpha$  subunit and 1 or more modulatory  $\beta$  subunits [5-9]. While sodium channel  $\alpha$  subunits are large proteins with 24 membrane-spanning domains,  $\beta$  subunits are smaller proteins with a single transmembrane domain and an extracellular V-type immunoglobulin (IG)-like motif [10-14]. Ten distinct  $\alpha$  subunit genes (*SCNxA*) and 4  $\beta$  subunit genes (*SCN1B-4B*) have been cloned from mammals, and all have been functionally expressed with the exception of *SCN7A* [2,3,15]. While heterologous expression of  $\alpha$  subunit genes alone can reconstitute key properties of voltage-gated sodium channels observed in native tissues, co-expression with  $\beta$  subunit genes modifies channel gating and increases channel cell surface expression [10-13]. Studies of mice with targeted ablation of either  $\beta 1$  or  $\beta 2$  genes corroborated these roles for  $\beta$  subunits *in vivo* and revealed an additional function for  $\beta$  subunits in  $\alpha$  subunit localization [16,17]. Moreover, the unique expression profiles of each  $\beta$  subunit gene and their variable effects on  $\alpha$  subunit function, expression, and localization suggest that different subunit combinations in heart, brain, and muscle may contribute to diversity and specialization in electrical signaling [18,19].

The physiological importance of sodium channel  $\alpha$  and  $\beta$  subunits is reinforced by their role in human disease. Mutations that cause even subtle changes in sodium channel  $\alpha$  subunit function may result in severe human disease phenotypes including epilepsy and arrhythmia [20]. To date, mutations in 7  $\alpha$  subunit genes (*SCN1A*, *SCN2A*, *SCN3A*, *SCN4A*, *SCN5A*, *SCN8A*, *SCN9A*) have been

linked to human disease [21-27]. Given their important modulatory effects on the expression and function of sodium channel  $\alpha$  subunits,  $\beta$  subunits are also important candidate genes for human diseases related to sodium channel dysfunction.  $\beta 1$  subunit gene mutations are a cause of generalized epilepsy with febrile seizures plus (GEFS+), and variants in sodium channel  $\alpha$  subunits that disrupt their interaction with  $\beta$  subunits may be pathological in the heart [28-31]. More recently, a mutation in the  $\beta 4$  subunit gene was implicated as a cause of the Long QT syndrome, a congenital arrhythmia [32].

The relatively large number of distinct mammalian sodium channel  $\alpha$  subunit isoforms has generated great interest in the evolution of this gene family, particularly since invertebrate genomes contain far fewer conserved  $\text{Na}_v 1$  sodium channel genes [33-36]. The physical linkage of voltage-gated sodium channel genes to the 4 homeobox (HOX) gene clusters suggests that polyploidization at least partly contributed to the emergence of multiple ancestral sodium channel genes in early vertebrates [37-39]. Moreover, the clustering of closely-related sodium channel genes in mammalian genomes (e.g. *SCN1A*, *SCN2A*, and *SCN3A* on human chromosome 2) is consistent with the hypothesis that several of the 10 mammalian isoforms arose following tandem duplication of ancestral vertebrate genes [36-39]. Since fish and mammals possess different numbers of sodium channel isoforms, the duplication of ancestral vertebrate sodium channel genes is suspected to have occurred independently in teleost and tetrapod lineages [38]. Expansion of the sodium channel gene family by tandem duplication may also have occurred uniquely in tetrapods, since the complement of sodium channel genes in teleosts appears to have resulted from whole-genome duplication [40,41].

The unique electrophysiological properties, expression patterns, and physiological roles of mammalian sodium channel  $\alpha$  subunits suggest that the duplication of these genes may have been an adaptation that permitted the diversification and functional specialization of electrical signaling in vertebrates [1,39,42]. The physical interaction of  $\alpha$  subunits with auxiliary proteins such as  $\beta$  subunits that modify their expression or function may have been similarly adaptive [36,37]. Although sodium channel  $\beta$  subunits have not previously been identified in non-mammalian species, we hypothesize that a sodium channel macromolecular complex comprised of  $\alpha$  subunits and auxiliary  $\beta$  subunits is an evolutionarily-conserved structural entity. Evidence for the emergence of sodium channel  $\beta$  subunit genes early in vertebrate evolution would be consistent with the idea that these genes may have played a role in diversifying and fine-tuning electrical signaling not only in mammals but in all vertebrates.

Here we report the identification and molecular cloning of conserved orthologs of all 4 mammalian  $\beta$  subunit genes in *Danio rerio* (zebrafish), a teleost vertebrate and a popular developmental and physiological model system. By direct sequence analysis and alignment, we assessed whether zebrafish  $\beta$  subunits possess the structural hallmarks of their mammalian counterparts. We provide evidence for extensive splicing of zebrafish beta subunit genes and demonstrate their expression in excitable tissues. To determine whether zebrafish sodium channel  $\alpha$  and  $\beta$  subunits functionally interact, we studied sodium currents generated by the co-expression of the genes encoding  $\alpha$  and  $\beta$  subunits in a heterologous cell system. Finally, we assessed the synteny and phylogeny of  $\beta$  subunits and a group of closely-related genes with IG domains in several vertebrate species. Our findings strongly suggest that all 4 voltage-gated sodium channel  $\beta$  subunit genes emerged early in vertebrate evolution, and support the concept that the voltage-gated sodium channel  $\alpha$ - $\beta$  subunit macromolecular complex is a conserved and functionally-important vertebrate innovation.

## Results

### Identification and cloning of zebrafish sodium channel $\beta$ subunit genes


We identified 5 distinct zebrafish  $\beta$  subunit gene loci by mining the draft zebrafish genome sequence (ensembl Zv2.4) for genes homologous to the human and mouse  $\beta$ 1- $\beta$ 4 genes. *In silico* prediction of coding sequence on the relevant genomic DNA contigs enabled us to amplify partial  $\beta$  subunit gene sequences for each locus by RT-PCR and subsequently complete the full-length zebrafish  $\beta$  subunit clones by rapid amplification of cDNA ends (RACE)-PCR. Analysis of putative open reading frames within each cDNA sequence predicted proteins ranging from 220–232 amino acids in length. Deduced translation and alignment of the five zebrafish amino acid sequences with each mammalian  $\beta$  subunit protein sequence suggested the identities of these genes (Figs. 1A, 2A, 3A, 4A). While single homologs were identified for the genes encoding  $\beta$ 1- $\beta$ 3 ( $z\beta$ 1- $z\beta$ 3), zebrafish express two distinct but closely-related  $\beta$ 4 genes ( $z\beta$ 4.1,  $z\beta$ 4.2). Alignment of the amino acid sequences of zebrafish  $\beta$ 1,  $\beta$ 2,  $\beta$ 3,  $\beta$ 4.1, and  $\beta$ 4.2 with their human homologs indicated that these proteins share 53.4%, 49.6%, 51.1%, 43.2%, and 40.8% identity, respectively.

### Conserved features of zebrafish sodium channel $\beta$ subunits

Mammalian sodium channel  $\beta$  subunits possess a single transmembrane domain, a short intracellular carboxyl C-terminal tail, a cleavable amine (N)-terminal signal peptide, and a conserved V-type extracellular IG motif that most closely resembles the type found in cell-adhesion molecules [14,43]. Hydropathy analysis of zebrafish  $\beta$  subunit amino acid sequences using TMpred software


revealed that all 5 proteins are likely to possess cleavable N-terminal signal peptides and single transmembrane domains (additional file 1). To assess the presence of an extracellular V-type IG domain, we first analyzed each zebrafish beta subunit amino acid sequence with the NCBI conserved domain database. By this method, all 5 zebrafish  $\beta$  subunit genes are predicted to possess V-type extracellular IG domains (additional file 2). To more rigorously analyze these predictions, we manually examined the amino acid sequence of each zebrafish  $\beta$  subunit for the V-type IG domain consensus sequence (annotated beneath alignments, Figs. 1A, 2A, 3A, 4A). With the exception of a conservative tyrosine to phenylalanine (Y > F) substitution in  $z\beta$ 1 and a non-polar methionine to basic lysine residue substitution (M > K) in  $z\beta$ 3 (illustrated in red text in Figs. 1A and 3A), all 5 zebrafish  $\beta$  subunits exhibit 100% conservation of the consensus sequence for V-type IG domains [14,43]. Typical of this domain is the linkage of 2 cysteine residues in a disulfide bridge (DS\*, Figs. 1A, 2A, 3A, 4A), the structural importance of which was revealed by a mutation that results in familial epilepsy [28]. A second putative disulfide bridge that may further stabilize the extracellular IG domain in  $\beta$ 1 and  $\beta$ 3 is suggested of the crystal structure of myelin protein zero (MPZ), the primary structural protein of peripheral nerve myelin, whose IG domain closely resembles that found in sodium channel  $\beta$  subunits [12,43,44]. The cysteine residues that contribute to proposed second bridge are conserved in  $z\beta$ 1 and  $z\beta$ 3 (DS, Figs. 1A, 3A). Similar to mammalian  $\beta$ 2 and  $\beta$ 4,  $z\beta$ 2 and  $z\beta$ 4 proteins do not possess this second disulfide bridge but have an unpaired cysteine residue that may underlie covalent interactions with partnering sodium channel  $\alpha$  subunits (S-, Figs. 2A, 4A) [5,11,13].

In addition to V-type IG domains, our sequence analysis indicated that other functional and/or structural domains are conserved in zebrafish  $\beta$  subunits (Figs. 1A, 2A, 3A, 4A). These include a tyrosine in  $\beta$ 1 that is phosphorylated and regulates interactions with ankyrin G (Y181 or Y200 with signal peptide intact) [45], the C-terminal tyrosine-leucine-alanine-isoleucine (Y-L-A-I) internalization motif in  $\beta$ 1 and  $\beta$ 3 which may be recognized by clathrin-coated pits [12], and a juxtatransmembrane gamma-secretase cleavage site in  $\beta$ 2 that may play a role in cell adhesion or the movement of cells expressing this subunit [46]. Zebrafish  $\beta$  subunits are also likely to exhibit post-translational modification by amide nitrogen (N)-linked glycosylation, a key feature of all 4 mammalian  $\beta$  subunits [10-13]. The  $z\beta$ 1 protein possesses 2 of 4 predicted N-linked glycosylation sites found in human or rat  $\beta$ 1 in addition to exhibiting two unique sites (Fig. 1A). Similarly,  $z\beta$ 2 conserves 3 of 4 sites,  $z\beta$ 3 conserves 1 of 3 sites and displays 2 novel sites,  $z\beta$ 4.1 conserves 4 sites and displays 1




**Figure 1**

**Analysis of the cloned zebrafish β1 subunit gene and novel splice variants.** A) Alignment of cloned human, rat, and zebrafish β1 amino acid sequences. Black = identical in all three species; grey = identical in 2/3 species or conserved substitution. Shown for zebrafish is the most conserved β1 splice form (variant D). Hs = *Homo sapiens*, Rn = *Rattus norvegicus*, z = zebrafish; DS\* = cysteine residue predicted to participate in a disulfide bridge, based on the myelin P0 protein crystal structure; DS = predicted second disulfide bridge; N = predicted N-linked glycosylation site (N1 = human/rat, N2 = zebrafish); M1 = site of epilepsy-causing deletion (I70\_E74del) in Hsβ1; M2 = site of second epilepsy-causing mutation (C121W) in Hsβ1; M3/4 = site of third and fourth epilepsy-causing mutations (R85C, R85H); S1 = nonsynonymous Hsβ1 single nucleotide polymorphism (SNP, G/A > R85H); S2 = nonsynonymous Hsβ1 SNP (C/T > T189M); P = phosphorylation site (tyrosine Y181) that regulates ankyrin recruitment (NOTE: Y200 = Y181 following cleavage of 19 amino acid signal peptide); IN = putative internalization sequence. Consensus sequence for V-type IG domain is depicted beneath the alignment: G = glycine, x = any residue, ^ = hydrophobic residues, C = cysteine, - = gap in alignment with consensus sequence, W = tryptophan, \* = basic residue, L = leucine, D = aspartic acid, & = glycine, alanine, or aspartate, and Y = tyrosine. Red indicates zebrafish residues that deviate from the consensus sequence. See Results for references supporting sequence annotation. B) 5' and 3' RLM-RACE PCR and RT-PCR identified four distinct splice variants expressed from zβ1 locus on zebrafish chromosome 16 (*Ensembl*). C) Splice donor and acceptor sites of zebrafish β1 splice variants, derived from comparing cloned cDNA against genomic DNA sequences (*Ensembl*). Consensus GT-AG splice sites are labeled in red. A splice-site deviating from the consensus appears in grey. D) Schematic diagram of β1 splice variants A, B, and D, whose predicted proteins differ only in the length of their intracytoplasmic C-terminal tail. S-S = disulfide bridge. NH<sub>2</sub> = 5' amino terminus, CO<sub>2</sub> = 3' carboxyl terminus, β = putative N-linked glycosylation site. Alignment of C-terminal tail of variants zβ1A, zβ1B, and zβ1D (below). zβ1C is not shown as it is predicted to lack both extracellular IG and transmembrane domains.




**Figure 2**

**Analysis of the cloned zebrafish β2 subunit gene and novel splice variants.** Presentation and labeling as in Figure 1. A) Alignment of cloned human, rat, and zebrafish β2 amino acid sequences. Shown for zebrafish is the most conserved β2 splice form (variant B). S- = conserved cysteine in β2 that is a putative site of covalent linkage with a partner α subunit; S1 = nonsynonymous Hsβ2 single nucleotide polymorphism (SNP, C/T > R28W) (NCBI dbSNP, PharmGKB); S2 = nonsynonymous Hsβ2 SNP (G/A > R47H) (NCBI dbSNP, PharmGKB); GSCS = γ-secretase cleavage site. B) 5' and 3' RLM-RACE PCR and RT-PCR identified four distinct splice variants expressed from the zβ2 locus on zebrafish chromosome 15 (*Ensembl*). C) Splice donor and acceptor sites of zebrafish β2 splice variants. Zβ2 variants C and D both differ from the consensus sequences at the exon 4-intron 4 and intron 4-exon 5 splice junctions. D) Schematic diagram of β2 splice variants A-D. With the exception of variant A, the predicted proteins of zβ2 variants B-D differ only in the length of their intracytoplasmic C-terminal tail. Alignment of C-terminal tail of variants zβ2B, zβ2C, and zβ2D (below).



**Figure 3**  
**Analysis of the cloned zebrafish β3 subunit gene.** Presentation and labeling as in Figure 1. A) Alignment of cloned human, mouse, rat, and zebrafish β3 amino acid sequences. INT = putative internalization sequence. B) 5' and 3' RLM-RACE PCR and RT-PCR identified a single transcript expressed from the zβ3 locus on zebrafish chromosome 15 (*Ensembl*). zβ3 has six exons. C) All zβ3 splice sites adhere to GT-AG consensus sequences. D) Schematic diagram of the β3 protein.



**Figure 4**

**Analysis of cloned zebrafish β4.1 and β4.2 subunit genes.** Presentation and labeling as in Figure 1. A) Alignment of cloned human, rat, and zebrafish β4 amino acid sequences. S- = conserved cysteine in β4 that is a putative site of covalent linkage with a partner α subunit; N = predicted N-linked glycosylation site (N1 = human, N2 = rat, N3 = zebrafish β4.1, N4 = zebrafish β4.2); M = site of putative Long QT syndrome-causing mutation L179F; S1 = nonsynonymous Hsβ4 SNP (A/C > N210H). B) Genomic organization of zβ4.1 and zβ4.2 derived from comparing cloned cDNA sequences with genomic sequences of zebrafish chromosomes 15 and 5, respectively. zβ4.1 has five exons and zβ4.2 has six exons. C) All zβ4.1 and zβ4.2 splice sites exhibit consensus GT-AG donor/acceptor sequences. D) Schematic diagram of zβ4.1 and zβ4.2 proteins.

novel site, and zβ4.2 conserves 3 of four sites and displays 3 novel sites (Figs. 2A, 3A, 4A).

**Comparative genomics of β subunit gene variants**

The conservation of amino acid residues between homologous genes in distantly-related species may uncover previously unappreciated regions of functional importance and facilitate the evaluation of novel human mutations and polymorphisms. To determine whether previously identified mutations and polymorphisms are conserved in zebrafish β subunit genes, we mapped these variants onto the alignments of mammalian and zebrafish β subunit amino acid sequences. Four mutations in the human β1 gene have been linked to the heritable epilepsy syndrome GEFS+, the first resulting in a 5 amino acid deletion from isoleucine at position 70 to glutamic acid at position 74 (I70\_E74del), the second in substitution of a cysteine residue that participates in a disulfide bridge (C121W), and the third and fourth resulting in substitution of an arginine at position 85 for either a cysteine or a histidine, respectively (R85C, R85H) [28,47,48]. All of these mutations are expected to destabilize the extracellular IG domain of β1 and result in loss of function. In zβ1, 2 of 5 amino acids in the deleted segment (isoleucine, tyrosine), C121, and R85 are all conserved (M1, M2, M3/4, Fig. 1A). R85H was also previously reported as a non-synonymous SNP in NCBI and PharmGKB SNP databases (S1, Fig. 1A). A second reported SNP in human β1 results in a threonine to methionine substitution at position 189 (T189M), but T189 is not conserved in the amino acid sequence of zβ1 (S2, Fig. 1A).

Although disease-causing mutations are not currently associated with either *SCN2B* or *SCN3B*, nonsynonymous

SNPs have been reported for *SCN2B* that result in an arginine to tryptophan substitution at position 28 (R28W) and an arginine to histidine substitution at position 47 (R47H). R47 but not R28 is conserved in the amino acid sequence of zebrafish β2 (S1, S2, Fig. 2A). A mutation in *SCN4B* resulting in a leucine to phenylalanine substitution at position 179 (L179F) was recently implicated as a putative cause of the congenital Long QT syndrome [32]. Although this leucine is conserved in zebrafish β4.1, rat β4 and zebrafish β4.2 display a cysteine (C) and threonine (T) at this position, respectively (M1, Fig. 4A). A nonsynonymous *SCN4B* polymorphism has also been reported, resulting in an asparagine to histidine substitution at position 210 (N210H). N210 is conserved in the sequences of both zβ4.1 and zβ4.2 (S1, Fig. 4A).

**Alternative splicing of zebrafish β subunit genes**

The mammalian β1 subunit gene is alternatively-spliced in both rats (β1A) and humans (β1B), with retention of intron 3 being the primary event in both species that dramatically alters the transmembrane domain and intracellular C-terminus of the β1 subunit protein [49,50]. An alternative splice-variant of β1 that retains intron 5 and adds 86 nucleotides to the 3' untranslated region (UTR) of β1 mRNA transcripts has also been reported [51,52]. In zebrafish, we detected 4 alternatively-spliced variants of the β1 gene, designated here as A-D. Comparisons between complementary DNA and genomic DNA revealed that zβ1A transcripts have 5 exons while zβ1B and zβ1D each have 6 exons and zβ1C has only 2 (Fig. 1B, Table 1). Transcripts of zβ1A, zβ1B, and zβ1D all share exons 1 through 4 but not exons 5 and 6; and all splice junctions except exon 5-intron 5 of zβ1B display canonical GT-AG splice donor-acceptor sites (Figs. 1B, 1C; Table

**Table 1: Comparative genomics of the sodium channel β1 gene in zebrafish and mammals.**

gene	Genbank accession number	gene location	cDNA	exon1	exon2	exon3	exon4	exon5	exon6	protein	%ID (%SIM)	topology
zβ1A	DQ489722	Chr 16	1411	219 (52)	167	238	142	645 (31)	-	209	47.5% (57.0%)	IG(V): 37-149 TM: 164-185
zβ1B	DQ489723	Chr 16	1054	219 (52)	167	238	142	28	260 (9)	211	49.8% (59.6%)	IG(V): 37-149 TM: 164-185
zβ1C	DQ489724	Chr 16	574	219 (52)	355 (266)	-	-	-	-	105	13.0% (17.5%)	IG(V): none TM: none
zβ1D	DQ489725	Chr 16	1320	219 (52)	167	238	142	72 (67)	482 (0)	221	53.4% (63.7%)	IG(V): 37-149 TM: 164-185
hβ1	NM_001037	Chr 19	1521	231 (40)	167	241	142	72 (67)	668 (0)	218	100% (100%)	IG(V): 33-146 TM: 161-182
rβ1A	AF182949	Chr 1	850	40	167	643 (615)	-	-	-	273	55.1% (59.1%)	IG(V): 33-146 TM: 216-234
hβ1B	NM_199037	Chr 19	1170	231 (40)	167	772 (600)	-	-	-	268	58.0% (60.2%)	IG(V): 33-146 TM: 244-263

Gene: z = zebrafish, h = human, r = rat, and letters A-D refer to different splice variants. Accession numbers are identifiers for nucleotide sequences in NCBI GenBank. Values for cDNA and exons represent number of nucleotides. (Parentheses) indicate nucleotides within the open reading frame (difference = untranslated sequence or UTR). Protein values represent amino acid number with signal peptide intact. %ID (%SIM) refers to percentage identity and similarity with the human β1 protein sequence as determined by alignment. Zebrafish β subunit topology was determined as described in Methods. IG(V) = V-type immunoglobulin domain and TM = transmembrane domain. Numbering refers to β1 protein sequence with signal peptide intact. IG(V) domain and transmembrane regions for human β1 and human/rat splice variants were annotated based on previous reports [10, 14, 43, 49, 50].



1). The  $z\beta 1$  splice variant that shares the most identity with human  $\beta 1$  at the amino acid level ( $z\beta 1D$ ) also shares nearly identical genomic organization (Table 1). With the exception of differences in exons 5 and 6,  $z\beta 1A$  and  $z\beta 1B$  variants also share this conserved genomic architecture (Table 1).

Similar to splice variants of the mammalian  $\beta 1$  gene, splice variants of  $z\beta 1$  are all predicted to encode proteins with variable C-termini. Unlike the mammalian variants, however,  $z\beta 1A$ ,  $z\beta 1B$ , and  $z\beta 1D$  are predicted to possess identical transmembrane domains and differ only in their distal C-termini (Fig. 1D). As a result of alternative-splicing,  $z\beta 1A$  lacks the conserved tyrosine residue (Y200, Fig. 1A) that was found to regulate recruitment of ankyrin G by mammalian  $\beta 1$ , as well as the putative internalization sequence (Y-L-A-I) discussed above (Figs. 1A, 1D) [12,45,53]. These alterations may affect interactions between  $z\beta 1A$  and zebrafish ankyrin, as well as influence the cycling of the mature  $z\beta 1A$  protein from its membrane compartment. The  $z\beta 1C$  splice variant, which results from retention of intron 2 and has an open reading frame of only 105 amino acids, is predicted to lack both an extracellular IG motif and a membrane-spanning segment.

Although alternative splicing of the  $\beta 2$  gene has not previously been identified in mammals, we identified 4 unique  $\beta 2$  transcripts in zebrafish. While  $z\beta 2A$  and  $z\beta 2B$  are spliced at canonical splice donor and acceptor sites, variants  $z\beta 2C$  and  $z\beta 2D$  are assembled by splicing at non-canonical exon 4-intron 4 and intron 4-exon 5 splice donor and acceptor sites.  $z\beta 2A$  is assembled from only 3 exons,  $z\beta 2B$  from 6 exons, and  $z\beta 2C$  and  $z\beta 2D$  from 5 exons each (Fig. 2B). Mammalian and zebrafish  $\beta 2$  genes all share a second exon that is 167 nucleotides in length and encodes the initial segment of the subunit's extracellular IG domain. Otherwise, the genomic organization of  $z\beta 2$  splice variants diverges from that of human, mouse, and rat  $\beta 2$  genes which all have 4 coding exons (Table 2). Nevertheless, exons 1 and 3 are similar in length in all species, as is the coding segment of exon 4 in the mammalian  $\beta 2$  genes and the  $z\beta 2B$  and  $z\beta 2D$  splice variants (Table 2).  $z\beta 2B$ ,  $z\beta 2C$ , and  $z\beta 2D$  are unique in possessing a fifth exon that contributes to the gene's open reading frame. Similar to mouse  $\beta 2$ , the most conserved zebrafish  $\beta 2$  splice variant at the amino acid level ( $z\beta 2B$ ) has a lengthy terminal exon that is entirely non-coding.

As observed for  $z\beta 1$ , the conceptual translation of  $z\beta 2$  splice variants predict  $z\beta 2$  proteins that are of variable length and amino acid sequence at their intracellular C-terminal tail (Fig. 2D). The variant  $z\beta 2A$  results from retention of intron 3 and is predicted to result in a 160 amino acid protein that has an intact extracellular IG domain but no transmembrane domain (Fig. 2D).  $z\beta 2A$  is

thus unlikely to display canonical beta subunit activities such as modulation of the trafficking and/or function of voltage-gated sodium channels  $\alpha$  subunits, or mediation of cell-adhesion between cells expressing this protein and other cells or the extracellular matrix, which all depend on integration in the cell membrane. Transcripts for variants  $z\beta 2B$ -D result from alternative splicing of exons 4-6 and are expected to produce proteins of 223, 201, and 231 amino acids, respectively (Fig. 2D, Table 2). Since the function of the intracellular C-terminal tail of  $z\beta 2$  is not well-characterized, the predicted impact of these splice variants on  $z\beta 2$  function cannot be readily predicted.

Unlike the  $z\beta 1$  and  $z\beta 2$  genes, alternatively-spliced transcripts of  $z\beta 3$ ,  $z\beta 4.1$ , and  $z\beta 4.2$  genes were not detected. For each of these three genes, the genomic organization is well-conserved compared to that of its respective mammalian homolog (Tables 3, 4). The open reading frame of *SCN3B* in humans, mice, rats, and zebrafish is derived from 5 exons (exons 2-6), with a non-coding initial exon found in all 4 species and a non-coding terminal exon found only in the mammalian  $\beta 3$  gene (Table 3). Splicing of zebrafish  $\beta 3$  is determined by canonical GT-AG splice donor and acceptor sites, producing a putative protein whose secondary structure is similar to the  $\beta 3$  subunit found in mammals (Figs. 3C, D).  $z\beta 4.1$  and  $z\beta 4.2$  transcripts are comprised of 5 and 6 exons, respectively, all of which are assembled from splicing at canonical GT-AG splice donor and acceptor sites (Fig. 4B, C).  $z\beta 4.2$  is unique among  $\beta 4$  genes in possessing a non-coding exon 1 (Table 4). The predicted secondary structures of  $z\beta 4.1$  and  $z\beta 4.2$  are both similar to the  $\beta 4$  subunit found in mammals (Fig. 4D).

#### **Tissue-specific regulation of zebrafish $\beta$ subunit gene expression and splicing**

Mammalian sodium channel  $\beta$  subunit genes are expressed primarily in excitable tissues such as the heart, brain, and skeletal muscle. By RT-PCR, we observed that zebrafish  $\beta$  subunit genes are also expressed in excitable tissues where they may act to regulate the expression or function of sodium channel  $\alpha$  subunits (Fig. 5, Table 5). Moreover, we detected distinct expression patterns for  $z\beta 1$ -4, suggesting that the functional importance of each  $\beta$  subunit likely varies by tissue type. Transcripts of  $z\beta 2$  or  $z\beta 3$  are absent from skeletal muscle while  $z\beta 1$  and  $z\beta 4.2$  are expressed only at low levels in the heart. As might be expected for non-excitabile tissues, few  $\beta$  subunit transcripts were detected in the zebrafish liver and only  $z\beta 1$  and  $z\beta 2$  are expressed in the gill.

By designing splice variant-specific primers, we also found evidence for tissue-specific regulation of splicing of zebrafish  $\beta$  subunit genes (Fig. 5). While  $z\beta 1$  variants B and D are expressed primarily in the brain and eye

**Table 2: Comparative genomics of the sodium channel  $\beta 2$  gene in zebrafish and mammals.**

gene	Genbank accession number	gene location	cDNA	exon 1	exon 2	exon 3	exon 4	exon 5	exon 6	protein	%ID (%SIM)	topology
<b>z<math>\beta</math>2A</b>	<a href="#">DQ489726</a>	Chr 15	821	243 (82)	167	411 (234)	-	-	-	160	32.6% (42.5%)	IG(V): 47–152 TM: none
<b>z<math>\beta</math>2B</b>	<a href="#">DQ489727</a>	Chr 15	2449	243 (82)	167	217	196	68 (10)	1558 (0)	223	49.6% (64.7%)	IG(V): 47–152 TM: 162–186
<b>z<math>\beta</math>2C</b>	<a href="#">DQ489728</a>	Chr 15	851	243 (82)	167	217	124	100 (13)	-	201	45.2% (58.4%)	IG(V): 47–152 TM: 162–186
<b>z<math>\beta</math>2D</b>	<a href="#">DQ489729</a>	Chr 15	918	243 (82)	167	217	193	98 (34)	-	231	48.1% (62.8%)	IG(V): 47–152 TM: 162–186
<b>Hs<math>\beta</math>2</b>	<a href="#">NM_004588</a>	Chr 11	4939	260 (70)	167	211	4301 (200)	-	-	215	100% (100%)	IG(V): 43–146 TM: 156–180
<b>Mm<math>\beta</math>2</b>	<a href="#">NM_001014761</a>	Chr 9	3980	213 (70)	167	211	618 (200)	2771 (0)	-	215	92.1% (94.4%)	IG(V): 43–146 TM: 156–180
<b>Rn<math>\beta</math>2</b>	<a href="#">NM_012877</a>	Chr 8	873	236 (70)	167	211	259 (200)	-	-	215	93.0% (94.9%)	IG(V): 43–146 TM: 156–180

Presentation and labeling as in Table 1. %ID (%SIM) refers to percentage identity and similarity with the human  $\beta 2$  protein sequence as determined by alignment. Zebrafish  $\beta 2$  subunit topology was determined as described in Methods. Numbering refers to  $\beta 2$  protein sequence with signal peptide intact. IG(V) domain and transmembrane regions for human  $\beta 2$  and human/mouse/rat splice variants were annotated based on previous reports [11, 14, 43, 101].

(including the optic nerve), z $\beta$ 1A is expressed in the atrium, ventricle, skeletal muscle and gill. Similarly, while z $\beta$ 2 variant B is primarily expressed in the brain and eye, z $\beta$ 2 variants A, C, and D are also expressed in the atrium, ventricle, brain, eye, and variant D is additionally expressed in the gill. Interestingly, z $\beta$ 3 may be alternatively-spliced in the brain and eye but not in the atrium or ventricle. We were unable to recover the shorter variant because of its apparent low level of expression. Although z $\beta$ 1 variant C was identified by RACE-PCR using total embryonic RNA as template, we were unable to detect expression of this short splice variant in the adult tissues analyzed for this study. This suggests that z $\beta$ 1 variant C may be developmentally regulated, expressed in other tissues, unstable, or is the result of an infrequent alternative splicing event.

#### **Zebrafish $\beta 1$ functionally modifies sodium channel expression, function in vitro**

As the first sodium channel auxiliary subunit to be identified,  $\beta 1$  is the most widely-studied of the four known mammalian  $\beta$  subunit proteins. To evaluate whether functional  $\alpha$  and  $\beta$  subunit interactions are likely to occur in zebrafish, we heterologously co-expressed the most conserved variant of z $\beta$ 1 (variant D) with the gene encoding the zebrafish pore-forming sodium channel  $\alpha$  subunit zNa $_v$ 1.5, whose expression we detected in the adult zebrafish brain and heart (unpublished observations). Prior studies suggest that the mammalian  $\beta 1$  subunit may influence the current amplitude and possibly the gating of mammalian Na $_v$ 1.5 *in vitro* [54–57]. Co-expression of z $\beta$ 1 with *zscn5a* in Chinese Hamster Ovary (CHO) cells increased peak sodium current by 68% at a -30 mV depolarizing pulse ( $n = 5$ ,  $p < 0.001$ ) compared to *zscn5a* alone ( $n = 8$ ) (Fig. 6A–C, table 6). Co-expression of z $\beta$ 1 with

*zscn5a* also resulted in small but significant changes in the gating of the zNa $_v$ 1.5 channel protein. z $\beta$ 1 induced hyperpolarizing shifts in both the voltage-dependence of activation and inactivation of zNa $_v$ 1.5, without altering recovery from inactivation (Fig. 6D–F, table 6). These data suggest that the canonical modulatory effects of mammalian sodium channel  $\beta$  subunits on  $\alpha$  subunit function and expression are likely to be conserved in teleosts and other non-mammalian vertebrates.

#### **Evolutionary relationships of vertebrate $\beta$ subunit genes**

Although  $\beta$  subunit genes appear to be related both structurally and functionally to each other and to cell adhesion molecules with V-type IG domains such as myelin protein zero and contactin [11,12], the evolutionary relationships among these genes have not previously been formally studied. To investigate these relationships, we first identified an extended group of human genes that display sequence homology to  $\beta$  subunits. BLAST searches of the human genome with human  $\beta$  subunit nucleotide and amino acid sequences most frequently-identified the genes *myelin protein zero*, *myelin protein zero-like 1*, *isoform A (MPZL1 iso A)*, *epithelial V-like antigen (EVA1)*, and hypothetical protein LOC196264 which we named "EVA1-like gene" (*EVA1L*) for its similarity to *EVA1* and the proximity of these two genes in the human genome. Analysis of the amino acid sequence of each gene using the NCBI conserved domain database and Tmpred software revealed that all 4 proteins likely possess single V-type IG domains and transmembrane segments, respectively, similar to sodium channel  $\beta$  subunits (Fig. 7A). Moreover, comparison of the complementary DNA sequences of these genes against their respective genomic loci indicated that all 4 genes also possess genomic organization similar to sodium channel  $\beta$  subunits (Fig. 7B).

**Table 3: Comparative genomics of the sodium channel β3 gene in zebrafish and mammals.**

gene	Genbank accession number	gene location	cDNA	exon 1	exon 2	exon 3	exon 4	exon 5	exon 6	exon 7	protein	%ID (%SIM)	topology
zβ3	<a href="#">DQ489730</a>	Chr 15	1006	294 (0)	103 (55)	170	235	139	65 (64)	-	220	51.1% (65.2%)	IG(V): 38–150 TM: 165–186
Hsβ3	<a href="#">NM_018400</a>	Chr 11	4052	778 (0)	80 (55)	164	226	139	86 (64)	2579 (0)	215	100% (100%)	IG(V): 38–145 TM: 160–181
Mmβ3	<a href="#">NM_178227</a>	Chr 9	3549	206 (0)	80 (55)	164	226	139	80 (64)	2654 (0)	215	97.7% (97.7%)	IG(V): 38–145 TM: 160–181
Rnβ3	<a href="#">NM_139097</a>	Chr 8	3910	350 (0)	82 (55)	164	226	139	80 (64)	2837 (0)	215	98.1% (98.1%)	IG(V): 38–145 TM: 160–181

Presentation and labeling as in Table 1. %ID (%SIM) refers to percentage identity and similarity with the human β3 protein sequence as determined by alignment. Zebrafish β3 subunit topology was determined as described in Methods. Numbering refers to β3 protein sequence with signal peptide intact. IG(V) domain and transmembrane regions for human β3 and human/mouse/rat splice variants were annotated based on previous reports [12, 14].

We therefore included these genes in our analysis of the synteny and phylogeny of the extended family of vertebrate β subunit genes.

To analyze synteny, we first identified all β subunit-like genes in human, rat, zebrafish, frog (*Xenopus tropicalis*) and bird (*Gallus gallus*) genomes (Table 7). Notably, while humans and rats each have 4 and zebrafish have 5 β subunit genes, only 3 β subunit genes were identified in frogs and birds (β1 was not found in either genome). Next, we used reciprocal BLAST searches and *in silico* chromosome walking to assess physical relationships among β-subunit like genes (Fig. 8). Strikingly, *scn2b*, *scn4b*, and *EVA1/EVA1L* genes map to common locations in every vertebrate genome analyzed (human chromosome 11, rat chromosome 8, zebrafish chromosome 5/15, frog scaffold\_39, and chicken chromosome 24), suggestive of a close evolutionary relationship that may have resulted from gene duplication. Although physically more distant, *scn3b* is also syntenic with this group of genes in 4/5 genomes analyzed. Of the 4 sodium channel β subunits, only *scn1b* falls outside this syntenic block. The common ancestry of human, rat, and zebrafish *scn1b* is supported,

however, by the proximity of the genes encoding the 26S protease regulatory subunit 6B (*PSMC4*) and fibrillarin (*FBL*) to the *scn1b* locus in all three species (human chromosome 19, rat chromosome 1, zebrafish chromosome 16). The synteny of β subunit genes in mammalian, fish, amphibian, and avian lineages thus supports the hypothesis that extant vertebrate β subunits are orthologous. Moreover, these findings strongly suggest that the evolution of at least several members of this extended gene family (*scn2b*, *scn4b*, *EVA1*, *EVA1L*) may have arisen from duplication events that predated the divergence of teleosts and tetrapods.

To further analyze the evolution of vertebrate β subunit-like genes, we reconstructed the phylogeny of this gene family (Fig. 9). Neighbor-joining phylogenetic analysis of amino acid sequences demonstrated that all-identified sodium channel β subunits fall on one of four branches corresponding to β1, β2, β3, or β4, supporting the preliminary identity assigned to each cloned zebrafish β subunit gene (including the duplicated zebrafish β4 genes) by alignment with each individual mammalian subunit (Figs. 1A, 2A, 3A, 4A). Despite synteny among *scn2b*,

**Table 4: Comparative genomics of the sodium channel β4 gene in zebrafish and mammals.**

gene	Genbank accession number	gene location	cDNA	exon -1	exon 1	exon 2	exon 3	exon 4	exon 5	protein	%ID (%SIM)	topology
zβ4.1	<a href="#">DQ489731</a>	Chr 22	2072	-	597 (91)	164	220	130	961 (94)	232	43.2% (59.3%)	IG(V): 53–155 TM: 165–187
zβ4.2	<a href="#">DQ489732</a>	Chr 5	1490	233 (0)	410 (97)	164	223	130	330 (85)	232	40.8% (56.7%)	IG(V): 55–158 TM: 168–190
Hsβ4	<a href="#">NM_174934</a>	Chr 11	4489	-	214 (61)	173	229	130	3743 (94)	228	100% (100%)	IG(V): 46–151 TM: 162–183
Mmβ4	<a href="#">NM_001013390</a>	Chr 9	4244	-	61	173	229	130	3651 (94)	228	79.4% (88.2%)	IG(V): 46–151 TM: 162–183
Rnβ4	<a href="#">NM_001008880</a>	Chr 8	4272	-	61	173	229	130	3679 (94)	228	80.3% (88.6%)	IG(V): 46–151 TM: 162–183

Presentation and labeling as in Table 1. %ID (%SIM) refers to percentage identity and similarity with the human β4 protein sequence as determined by alignment. Zebrafish β4 subunit topology was determined as described in Methods. Numbering refers to β4 protein sequence with signal peptide intact. IG(V) domain and transmembrane regions for human β4 and human/mouse/rat splice variants were annotated based on previous reports [13, 14].

**Table 5: Primers used to detect expression of zebrafish  $\beta$  subunit genes and splice variants in different tissues of the adult zebrafish.**

Forward Primer (5'-3')	Reverse Primer (5'-3')	Gene	Variant	Amplicon Size (bp)
CTACACTTATGCAGAAATGACAGCCAGC	GATGGACAGAGCTTCAAGCTTTTGGCT	$z\beta 1$	A	422
		$z\beta 1$	D	494
GACAGAATCCTCATCTTCCCAACTATG	CTGCATTCTTCATTTAAACTCAGAGGT	$z\beta 1$	B	267
		$z\beta 1$	D	534
GTCCCTGCGCTGTTGTGTTTAAACACAT	GGGTGAACAATCCCTTTAAGCTGCACT	$z\beta 1$	C	382
CAGCTGACAGACGAGGGCATCTACAACCT	CAACACCTGCAGTGAGAAAACCCCAT	$z\beta 2$	A	201
CATCCTTGCTCTGCTCATTCTGTCCAT	TCGCTACACGATAATACCAGGGAGTGT	$z\beta 2$	B	304
		$z\beta 2$	C	169
		$z\beta 2$	D	236
CTGGTGTGTGTGGATGTGCCATCA	CTTGTTGGCGGAAAGCTTGAATGA	$z\beta 3$	-	357
AGGTGAGCACAGGGAAGGTCCATT	GGAGGCCATTTTCTGTGTTGTCGT	$z\beta 4$	loc1	543
TGTGTTGTGTTTCATGCTTTG	GACCACCTTATGTTCTCTCTA	$z\beta 4$	loc2	395

Note: several primer pairs detected multiple splice variants, as indicated above and displayed in Figure 5. BP = base pairs or nucleotide number.

*scn3b*, and *scn4b*, phylogenetic analysis revealed that  $\beta 3$  is more closely related to  $\beta 1$  than to either  $\beta 2$  or  $\beta 4$ . Moreover, our phylogenetic model incorporating  $\beta$  subunit-like genes additionally indicated that  $\beta 2$  and  $\beta 4$  are more closely-related to each other and to  $\beta$  subunit-like genes than to either  $\beta 1$  or  $\beta 3$ . Our analysis thus supports an evolutionary model where  $\beta 2/\beta 4/\beta$  subunit-like genes and  $\beta 1/\beta 3$  arose following duplications of 2 distinct precursor genes. The timing of such duplications cannot be gleaned from this model. The presence of gene orthologs to all 4  $\beta$  subunits in divergent vertebrates (mammals and teleosts) strongly suggests, however, that the early vertebrate common ancestor to these lineages already had four distinct  $\beta$  subunit genes. Duplication events that occurred in this gene family – aside from that which produced two zebrafish  $\beta 4$  genes – thus occurred earlier in vertebrate evolution or prior to the emergence of the vertebrates altogether.


#### Absence of $\beta$ subunit genes in invertebrates

The sequenced genomes of the ascidians *Ciona intestinalis* and *Ciona savignyi* provide an opportunity to identify putative precursors to vertebrate  $\beta$  subunits genes in invertebrate chordates [58,59]. Although both ascidian genomes contain several loci encoding  $Na_v 1$  voltage-gated sodium channel  $\alpha$  subunits, BLAST searches using the nucleotide and protein sequences of zebrafish and mammalian sodium channel  $\beta$  subunits and related proteins (MPZ, EVA1) did not identify any homologous genes. The absence of  $\beta$  subunit-like genes in sequenced *Ciona* genomes does not preclude the presence of numerous other genes with V-type and other IG-like domains ([60,61] and unpublished observations). These findings strongly suggest that sodium channel  $\beta$  subunit genes are an innovation of vertebrates. As predicted, searches of the genome of the echinoderm *Strongylocentrotus purpuratus* (sea urchin) [62], a non-chordate deuterostome, also did not reveal any sodium channel  $\beta$  subunits genes.

Despite the lack of evidence for  $\beta$  subunits genes outside of vertebrates, studies conducted in *Drosophila melanogaster* demonstrate that invertebrate sodium channels require additional subunits for normal function. Mutations in the *Drosophila tip-E* (temperature-induced paralysis) gene, which encodes a sodium channel auxiliary subunit, disrupt nerve conduction by perturbing the expression and function of the *para* voltage-gated sodium channel [63-67]. Moreover, 4 recently-identified *tip-E* related genes (*TEH1-4* or *tip-E* homologs 1-4) were found to modulate the density and kinetics of sodium currents of heterologously-expressed *para* sodium channels [68]. Based on these findings, we sought to assess whether vertebrate and invertebrate sodium channel auxiliary subunits share common structural elements or functional domains. Use of TMPred software and the NCBI conserved domain database demonstrated, however, that *Drosophila tip-E* and *tip-E* homologous genes (*TEH1-4*) each possess two predicted membrane-spanning segments (not one) and lack the canonical V-type IG domain found in vertebrate  $\beta$  subunits. Additionally, alignment of the amino acid sequences of cloned zebrafish and human sodium channel  $\beta$  subunit genes with the sequences of *Drosophila tip-E* and *tip-E* homologous genes also revealed only minimal homology (< 15% amino acid identity). BLAST searches of the *Drosophila* genome did not uncover any additional genes sharing close homology with vertebrate  $\beta$  subunits, despite the presence of numerous genes encoding proteins with immunoglobulin domains ([69] and unpublished observations). These results indicate that vertebrate and invertebrate sodium channel auxiliary subunits are dissimilar in structure and unrelated.

#### Discussion

Numerous studies of mammalian voltage-gated sodium channel  $\beta$  subunits have demonstrated that these small, single membrane-spanning proteins are integral components of sodium channel complexes in excitable tissues,



**Figure 5**  
**Zebrafish sodium channel β1-4 subunit genes and novel splice variants are differentially expressed in excitable tissues.** Total RNA was isolated from wild-type adult zebrafish tissues. RT-PCR with gene and splice variant-specific primers was used to detect expression (see Table 5 for primer sequences and amplicon details). Atr = atrium, Ven = ventricle, Skm = skeletal muscle, Brn = Brain, Eye = eye/optic nerve, Liv = liver, Gil = gill. + = enzyme added to reverse transcription step, - = no reverse transcriptase enzyme (negative control). Zebrafish β-actin was amplified from each template as a positive control.

modulators of the expression and function of pore-forming sodium channel α subunits, and candidate genes for clinical disorders linked to perturbed membrane excitability such as arrhythmia and epilepsy [18,70]. Despite the early origins of the Na<sub>v</sub>1 family of sodium channel α subunits and their cloning from diverse metazoans including eels, jellyfish, flies, and humans, the evolutionary history of sodium channel β subunits has remained obscure. The primary objective of this study was thus to investigate β subunit genes in *Danio rerio* (zebrafish), a pivotal vertebrate species whose teleost ancestors diverged from mammals over 400 million years ago, in order to gain further insight into the origin and regulation of the voltage-gated sodium channel macromolecular complex.

Using a combination of bioinformatics, molecular cloning, and phylogenetic analysis, we identified conserved orthologs of all 4 mammalian β subunit genes and 8 novel β1 and β2 splice variants in zebrafish. Using our cloned zebrafish sequences, we subsequently identified β subunit genes in other non-mammalian vertebrates including *Xenopus tropicalis* (Western clawed frog) and *Gallus gallus* (Red jungle fowl). The existence of conserved mammalian, teleost, amphibian, and avian sodium channel β subunit loci indicates that this gene family is likely to be found in most if not all vertebrate genomes. Moreover, our detection of zebrafish β subunit gene expression in tissues including the heart, muscle, and the brain suggests that the voltage-gated sodium channel α-β subunit macromolecular complex is likely to be a common structural feature of excitable membranes.

**Sodium channel complexes in non-mammalian vertebrates: an emerging concept**


Our data are the first to support the existence of voltage-gated sodium channel α-β subunit macromolecular complexes in non-mammalian vertebrates. Data from prior studies, however, suggest that sodium channels form α-β subunit complexes in mammals but not in other vertebrate lineages [5-7,71-79]. While sodium channels biochemically purified from rat neuronal membranes and rat and rabbit skeletal muscle membranes were found to be comprised of 1 α subunit and 1 or more auxiliary β subunits, for example, α subunits purified from the electroplax (electric organ) of the South American eel *Electrophorus electricus* and from chick cardiac muscle were unaccompanied by β subunits [5-7,71-79]. Further investigation into the existence of sodium channel α-β macromolecular complexes in non-mammalian vertebrates is thus needed to reconcile our current findings with the results of previous studies.

There are several possible reasons for the discord between biochemical and molecular genetic approaches. First, it is possible that some but not all voltage-gated sodium channel isoforms within a particular species form complexes by associating with auxiliary β subunits, or that pore-forming α subunits form complexes only in specific tissues. Evidence suggests that even in mammals, for example, the subunit composition of voltage-gated sodium

**Table 6: Biophysical properties of zNa<sub>v</sub>1.5 and zNa<sub>v</sub>1.5 plus zβ1 (variant D) in CHO cells.**

	Peak amplitude (pA)*	Activation V <sub>1/2</sub> (mV)	Inactivation V <sub>1/2</sub> (mV)	Recovery from inactivation, Tau (ms)
<b>zNa<sub>v</sub>1.5 alone</b>	-125.7 ± 16.9 pA/pF (n = 8)	-47.5 ± 1.6 mV (n = 8)	-79.3 ± 1.0 mV (n = 5)	127.9 ± 8.1 ms (n = 4)
<b>zNa<sub>v</sub>1.5 + zβ1</b>	-211.1 ± 11.2 pA/pF† (n = 5)	-53.0 ± 2.5 mV† (n = 5)	-82.6 ± 1.1 mV† (n = 6)	125.7 ± 4.1 ms (n = 6)

All values are mean ± standard error of the mean (SEM). \* at a -30 mV depolarizing pulse. † p < 0.001 vs. zNa<sub>v</sub>1.5 alone, Student's T-Test. pA = picoamperes; mV = millivolts; ms = milliseconds; pF = picofahrrads; V<sub>1/2</sub> = half maximal voltage.

**Figure 6****The zebrafish  $\beta 1$  subunit modulates the biophysical properties of the zebrafish sodium channel  $\alpha$  subunit**


**$zNa_v1.5$  in CHO cells.** A) Typical whole-cell sodium current trace of  $zNa_v1.5$  following expression of the pBK-CMV-*zscn5a* expression vector in CHO cells (n = 8). B) Typical whole-cell sodium current trace of  $zNa_v1.5 + z\beta 1$  (variant D) following co-expression of pBK-CMV-*zscn5a* and pGFP-IRES-*zβ1D*.  $z\beta 1$  significantly increased the peak amplitude of sodium current by 68% ( $p = 0.005$ ) at a -30 millivolt (mV) depolarizing pulse (n = 5). C) Current-voltage relationship demonstrating an increase in sodium current at every test potential between -50 and +50 mV. Filled circles =  $zNa_v1.5$  alone; open circles =  $zNa_v1.5 + z\beta 1$ . D) Voltage dependence of activation ( $zNa_v1.5$  alone, n = 8;  $zNa_v1.5 + z\beta 1$ , n = 5). E) Voltage-dependence of inactivation. ( $zNa_v1.5$  alone, n = 5;  $zNa_v1.5 + z\beta 1$ , n = 6). F) Recovery from inactivation ( $zNa_v1.5$  alone, n = 4;  $zNa_v1.5 + z\beta 1$ , n = 6). Pulse protocol in inset. Summary data is reported in table 6.

channels significantly varies both by  $\alpha$  subunit isoform and by tissue [9,72]. Second, at least for the eel electroplax, the function of this organ may depend on sodium channel  $\alpha$  subunits that have evolved unique properties which may preclude association with auxiliary subunits [80]. Third, it is possible that sodium channel  $\alpha$  subunit isoforms in certain tissues form complexes not with canonical  $\beta$  subunits but with yet unidentified proteins. In mammals, the interaction of sodium channel  $\alpha$  subunits with a number of additional components and modulators in macromolecular complexes suggest that  $\beta$  subunits may not always serve as obligatory functional partners (for review, see refs [18,81]). Our detection of conserved  $\beta$  subunit mRNA expression in excitable organs such as heart, brain, and skeletal muscle, however, suggests the strong possibility of  $\alpha$ - $\beta$  subunit complexes in these zebrafish tissues. Moreover, co-expression of zebrafish  $\alpha$

and  $\beta$  subunit genes in CHO cells demonstrated meaningful functional interactions between the two subunits, with significant differences observed in sodium current amplitude and channel gating when both subunit genes are expressed together versus when the  $\alpha$  subunit gene is expressed alone. Despite these findings, however, we cannot completely rule out the possibility that sodium channel  $\beta$  subunit genes play different *in vivo* functional roles in mammals than in non-mammalian vertebrates.

**Evolution of sodium channel  $\beta$  subunits and related genes**

Our analysis of the synteny and phylogeny of vertebrate sodium channel  $\beta$  subunit genes suggests a different evolutionary history for  $\alpha$  and  $\beta$  subunits. For sodium channel  $\alpha$  subunits, prevailing evolutionary models posit that the 10 mammalian isoforms arose from the tandem duplication of at least 2 of 4 ancestral sodium channel



**Figure 7**  
**Four additional human genes share homology with  $\beta$  subunits in both sequence and genomic organization.** A) BLASTP searches of the human genome using sodium channel  $\beta$  subunit amino acid sequences identified significant homology to myelin P<sub>0</sub> protein (MPZ), myelin P<sub>0</sub> protein-like protein isoform A (MPZL1 isoform A), epithelial V-like antigen I (EVA1), and epithelial V-like antigen I-like gene (EVA1L), all of which are single transmembrane proteins with extracellular V-type immunoglobulin domains and intracellular C-terminal tails as predicted by NCBI conserved domain database v2.09 and TMPred (see Methods for further details). B) This group of genes additionally shares similar genomic organization. Numbers in grey boxes refer to exon size (nucleotides) with untranslated sequence excluded for clarity.

genes, which in turn had duplicated from 1 or 2 precursor chordate Na<sub>v</sub>1 sodium channel genes by polyploidization [36-39]. As would be predicted by this model, the total number of Na<sub>v</sub>1 sodium channel genes varies between teleosts and mammals despite the phylogenetic clustering of all of these genes into 4 groups derived from ancestral vertebrate sodium channel genes [38,40]. Sodium channel  $\beta$  subunits, which are fewer in number, do not appear to have undergone tandem duplication in either mammals or in non-mammalian vertebrates. This is supported by our finding that 4 distinct vertebrate lineages (mammals, ray-finned fishes, amphibians and birds) all appear to share distinct orthologs to 3 or 4  $\beta$  subunit genes. Thus, it

is likely that the common ancestor to teleosts and tetrapods over 400 million years ago also possessed 4 distinct ancestral  $\beta$  subunit genes ( $\beta$ 1-4). The presence of 2  $\beta$ 4 orthologs on different zebrafish chromosomes is likely to be the remnant of an additional polyploidization event known to have occurred in teleost vertebrates but not in mammals [41]. It is difficult to determine why duplicate genes for  $\beta$ 1-3 have not been retained in zebrafish; it is possible that > 5  $\beta$  subunit genes did not offer teleosts any evolutionary advantage [42].

The close phylogenetic relationship of *SCN1B* and *SCN3B* and the physical proximity of *SCN2B* and *SCN4B* to each

**Table 7: List of cloned and predicted genes utilized for analysis of synteny and phylogeny of the extended  $\beta$  subunit gene family in vertebrates.**

Gene	Species	Analysis of Synteny Accession No. (Gene)	Physical Location	Analysis of Phylogeny Accession No. (Protein)
<b>SCN1B</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000105711</a>	chr. 19 (FWD, 40.21–40.22 Mb)	<a href="#">ENSP00000262631</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000021102</a>	chr. 1 (REV, 86.16–86.17 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000060222</a>	chr. 16 (REV, 47.82–47.83 Mb)	<a href="#">ABF47239</a>
<b>SCN2B</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000149575</a>	chr. 11 (REV, 117.54–117.55 Mb)	<a href="#">ENSP00000278947</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000016221</a>	chr. 8 (FWD, 48.07–48.08 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000041176</a>	chr. 15 (REV, 27.32 Mb)	<a href="#">ABF47241</a>
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETESTG00000009930</a>	*scaffold_39 (FWD, 0.58–0.59 Mb)	<a href="#">ENSXETESTP00000016971</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALG00000021272</a>	chr. 24 (REV, 5.10–5.11 Mb)	<a href="#">ENSGALP00000033652</a>
<b>SCN3B</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000166257</a>	chr. 11 (REV, 123.01–123.03 Mb)	<a href="#">ENSP00000299333</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000006937</a>	chr. 8 (FWD, 43.23–43.25 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARESTG00000011553</a>	chr. 15 (REV, 28.62–28.64 Mb)	<a href="#">ABF47244</a>
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETESTG00000002589</a>	*scaffold_298 (FWD, 0.94–0.96 Mb)	<a href="#">ENSXETESTP00000004357</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALESTG00000012587</a>	chr. 24 (FWD, 2.82–2.83 Mb)	<a href="#">ENSGALESTP00000019882</a>
<b>SCN4B.1</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000177098</a>	chr. 11 (REV, 117.51–117.53 Mb)	<a href="#">ENSP00000322460</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000026679</a>	chr. 8 (FWD, 48.09–48.11 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARESTG00000010162</a>	chr. 15 (REV, 27.26–27.29 Mb)	<a href="#">ABF47245</a>
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETESTG00000009931</a>	*scaffold_39 (FWD, 0.61–0.63 Mb)	<a href="#">ENSXETESTP00000016972</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALG00000007409</a>	chr. 24 (REV, 5.10 Mb)	<a href="#">ENSGALP00000011971</a>
<b>SCN4B.2</b>	<i>Danio rerio</i>	<a href="#">ENSDARESTG000000000878</a>	chr. 5 (FWD, 38.08–38.10 Mb)	<a href="#">ABF47246</a>
<b>MPZ</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000158887</a>	chr. 1 (REV, 159.54–159.55 Mb)	<a href="#">ENSP00000289928</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOESTG00000011377</a>	chr. 13 (FWD, 87.04–87.05 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000038609</a>	chr. 2 (REV, 46.60–46.62 Mb)	<a href="#">ENSDARP00000056371</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALESTG00000015635</a>	*chr. Un (REV, 60.79–60.80 Mb)	<a href="#">ENSGALP00000008810</a>
<b>MPZLI isoA</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000197965</a>	chr. 1 (FWD, 165.96–166.03 Mb)	<a href="#">ENSP00000352513</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000003248</a>	chr. 13 (REV, 81.32–81.36 Mb)	n/a
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETG000000023014</a>	*scaffold_195 (REV, 1.78–1.79 Mb)	<a href="#">ENSXETP000000049765</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALG00000015438</a>	chr. 1 (REV, 85.41–85.44 Mb)	<a href="#">ENSGALP000000024855</a>
<b>EVAI</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000149573</a>	chr. 11 (REV, 117.63–117.64 Mb)	<a href="#">ENSP00000278937</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000016085</a>	chr. 8 (FWD, 47.99–48.00 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000027345</a>	chr. 5 (FWD, 38.05–38.07 Mb)	<a href="#">ENSDARP000000032601</a>
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETESTG00000008504</a>	*scaffold_39 (FWD, 0.51–0.54 Mb)	<a href="#">ENSXETESTP00000014524</a>
	<i>Gallus gallus</i>	<a href="#">ENSGALG000000007412</a>	chr. 24 (REV, 5.12–5.13 Mb)	<a href="#">ENSGALP00000011976</a>
<b>"EVAI-like"</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000160588</a>	chr. 11 (REV, 117.60–117.63 Mb)	<a href="#">ENSP00000278949</a>
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000026753</a>	chr. 8 (FWD, 48.00–48.01 Mb)	n/a
	<i>Gallus gallus</i>	<a href="#">ENSGALG00000021271</a>	chr. 24 (REV, 5.11–5.12 Mb)	<a href="#">ENSGALP00000033649</a>
<b>PSMC4</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000013275</a>	chr. 19 (FWD, 45.17–45.18 Mb)	n/a
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000018994</a>	chr. 1 (REV, 83.15–83.16 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000027099</a>	chr. 16 (FWD, 47.99–48.00 Mb)	n/a
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETG00000014266</a>	*scaffold_1060 (REV, 0.17 Mb)	n/a
<b>FBL</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000105202</a>	chr. 19 (REV, 45.02–45.03 Mb)	n/a
	<i>Rattus norvegicus</i>	<a href="#">ENSRNOG00000019229</a>	chr. 1 (FWD, 83.27–83.28 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000053912</a>	chr. 16 (FWD, 47.49–47.50 Mb)	n/a
	<i>Xenopus tropicalis</i>	<a href="#">ENSXETG00000017830</a>	*scaffold_1073 (REV, 0.14 Mb)	n/a
<b>APOAI</b>	<i>Homo sapiens</i>	<a href="#">ENSG00000118137</a>	chr. 11 (REV, 116.21 Mb)	n/a
	<i>Danio rerio</i>	<a href="#">ENSDARG00000012076</a>	chr. 5 (REV, 38.04 Mb)	n/a
	<i>Gallus gallus</i>	<a href="#">ENSGALG00000007114</a>	chr. 24 (REV, 4.79–4.80 Mb)	n/a

All genes and proteins are identified by Ensembl accession numbers, except for cloned zebrafish beta subunit protein sequences (which begin with ABF and represent accession numbers assigned by GenBank). \* = unmapped scaffolds in Ensembl database. FWD = forward and REV = reverse and refers to gene orientation. n/a = not applicable (e.g. proteins for this species were not included in phylogenetic analysis).



other and to genes (e.g. *EVA1*) that are  $\beta$  subunit-like in sequence and genomic organization is strong evidence that duplication events gave rise to the  $\beta$  subunit gene family during the postulated large-scale expansion of the early vertebrate genome. Although we cannot accurately predict the early evolutionary history of sodium channel  $\beta$  subunit genes prior to the divergence of teleosts from other vertebrate lineages, the results of our phylogenetic analysis support the existence of 2 ancestral  $\beta$  subunit genes ( $\beta 1/\beta 3$  and  $\beta 2/\beta 4$ ) in early vertebrates. Somewhat surprisingly (but in agreement with a previous report [33]), BLASTN and BLASTP searches of available urochordate (*Ciona intestinalis*, *Ciona savignyi*) genomes did not identify any sodium channel  $\beta$  subunit-like genes, despite the clear existence of genes encoding proteins with immunoglobulin domains similar to those found in  $\beta$  subunits [60,61]. Moreover, we and others also found limited homology between vertebrate  $\beta$  subunit genes and the *tip-E/TEH* subunits of the *para* sodium channel in *Drosophila* [34,65]. These findings strongly suggest that sodium channel  $\beta$  subunits are a unique vertebrate innovation.

The similarity of vertebrate sodium channel  $\beta$  subunits to members of a more ancient family of IG domain-containing proteins suggests that  $\beta$  subunit genes may have arisen *de novo* in early vertebrates from genes encoding cell-adhesion molecules, membrane receptors, or components of the innate immunity. The property of modulation of the expression and function of sodium channel  $\beta$  subunits may thus represent a more evolutionarily recent function for this gene family. These findings support the idea that molecular mechanisms enhancing functional diversity and specialization in electrical signaling – either through  $\alpha$  subunit gene duplication, editing or alternative splicing of  $\alpha$  subunit RNA transcripts, or the emergence of auxiliary proteins that associate with  $\alpha$  subunits to modulate their expression and/or function – may have been adaptive and underwent selection in the evolving nervous system of early vertebrates and subsequently, in the increasingly complex excitable tissues of diverse vertebrate lineages including mammals. Despite the lack of common ancestry among invertebrate *tip-E/TEH* subunits and vertebrate IG-like  $\beta$  subunits, the apparent independent evolution of  $\text{Na}_v 1$  sodium channel-interacting proteins in distantly-related species demonstrates that the formation of sodium channel macromolecular complexes is an evolutionary advantageous and conserved mechanism for fine-tuning the properties of excitable membranes.


#### **Functional implications of alternative splicing of zebrafish $\beta$ subunit genes**

The evidence we present for the extensive C-terminal alternative splicing of zebrafish  $\beta$  subunit genes is intriguing because it may represent another conserved molecular

mechanism for modulating electrical signaling in vertebrates. Alternative splicing has been reported for both mammalian sodium channel  $\alpha$  and  $\beta$  subunits and invertebrate  $\alpha$  subunits, often with important functional consequences [49,50,82-88]. The intronic retention events that produce both the rat  $\beta 1A$  and human  $\beta 1B$  variants each result in a protein with a novel transmembrane domain and intracellular C-terminal tail. When stably expressed in Chinese Hamster Lung (CHL) fibroblasts, rat  $\beta 1$  and  $\beta 1A$  differentially modulate the function of the rat sodium channel  $\alpha$  subunit  $\text{Na}_v 1.2$  (*scn2a*) [49]. Similarly, the human  $\beta 1$  and  $\beta 1B$  subunits differentially modulate the function of human  $\text{Na}_v 1.2$  in *Xenopus* oocytes [50]. These studies suggest that the transmembrane domain and C-terminus of the human and rat  $\beta 1$  subunits contribute to the functional modulation of sodium channel  $\alpha$  subunits. In a subsequent study, the  $\beta 1$  intracellular C-terminal tail in particular was found to be required for both efficient physical association with  $\text{Na}_v 1.2$  and for the modulation of sodium channel function in both mammalian cells and *Xenopus* oocytes [89]. This raises the possibility that the zebrafish  $\beta 1$  and  $\beta 2$  subunit splice variants identified in this study may also differentially influence the function of sodium channel  $\alpha$  subunits. Our identification of distinctive expression patterns for different splice variants of the same gene underscores the complexity of this putative sodium channel regulatory mechanism.

#### **Conclusion**

We have identified conserved orthologs to all 4 mammalian  $\beta$  subunit genes (*z $\beta 1$ -z $\beta 4$* ) in zebrafish. Despite highly-conserved genomic organization in fish and mammals, zebrafish express 8 distinct mRNA transcripts for the  $\beta 1$  and  $\beta 2$  genes that are generated by alternative splicing. Zebrafish  $\beta$  subunit genes and their splice variants are differentially-expressed in excitable tissues, and co-expression of *z $\beta 1$*  (*variant D*) with the zebrafish sodium channel  $\alpha$  subunit gene *zscn5a* in CHO cells demonstrated functional  $\alpha$ - $\beta$  interactions that we predict may also occur in native tissues. Our evolutionary analysis of mammalian, teleost, amphibian, and avian  $\beta$  subunit and related genes indicated that all extant vertebrate  $\beta$  subunits are orthologous, that  $\beta 2/\beta 4$  and  $\beta 1/\beta 3$  share common ancestry, and that  $\beta$  subunits are closely-related to other proteins with V-type IG domains including myelin protein zero (MPZ) and epithelial V-like antigen 1 (EVA1). Homologs to vertebrate  $\beta$  subunit genes were not identified in the genomes of invertebrate chordates, and  $\beta$  subunits were found to be unrelated to the *tip-E/TEH* subunits of the *para* sodium channel in *Drosophila*. Taken together, these findings suggest that the family of sodium channel  $\beta$  subunit genes emerged early in vertebrate evolution, prior to the divergence of teleosts and tetrapods. The evolutionary history of vertebrate  $\beta$  subunits is thus consistent with the



**Figure 8**  
**SCN2B, SCN3B and SCN4B are syntenic with each other and with EVA1, EVA-1L, and APOA-1 in multiple vertebrate genomes, while SCN1B is syntenic with FBL and PSMC4 in humans, rats, and zebrafish.** To analyze synteny,  $\beta$  subunits and subunit-like genes were identified in human, rat, zebrafish, frog and bird genomes (see Table 7 for gene IDs and physical locations). While humans and rats have four and zebrafish have five  $\beta$  subunit genes, only three  $\beta$  subunit genes were identified in frogs and birds ( $\beta 1$  was not found in either genome). Reciprocal blast searches and *in silico* chromosome walking were used to assess physical relationships among vertebrate  $\beta$  subunit-like genes. HUGO gene nomenclature symbol IDs: *PSMC4* = 26S protease regulatory subunit 6B; *FBL* = fibrillarin; *APOA1* = apolipoprotein A-1; *MPZ* = myelin protein zero; *MPZL1* = myelin protein zero-like gene, isoform A; *EVA1* = epithelial V-like antigen 1; *EVA1L* = unannotated gene similar to *EVA1*. Red =  $\beta$  subunit genes, Blue =  $\beta$  subunit-like genes, Black = unrelated genes that are syntenic with  $\beta$  subunits.

hypothesis that voltage-gated sodium channel complexes are evolutionarily-conserved structural entities, and that  $\beta$  subunit genes may have played a role in the functional diversification and specialization of electrical signaling in early vertebrates.

## Methods

### Identification and cloning of zSCN1B-4B genes and splice variants

Mammalian *SCN1B-4B* gene sequences were used in BLASTN queries of the draft zebrafish genome (Ensembl Zv2-5) to identify DNA contigs containing putative zebrafish orthologs. Primers directed against predicted exons were then used to amplify partial  $\beta$  subunit gene sequences using the Titan RT-PCR enzyme system (Roche) and total day 2 embryonic zebrafish RNA as template. RNA was isolated using Trizol (GIBCO), digested with RQ1 DNase (Promega), and purified with the RNeasy Mini Kit (Qiagen). 5' and 3' ends and alternatively-spliced forms of each gene were identified using 5' and 3' RNA ligase mediated rapid amplification of cDNA ends (RLM-RACE) PCR (Ambion). Additional splice variants were identified by RT-PCR while examining tissue-specific expression. All amplicons were subcloned directly into the

pGEM-TEasy vector (Promega) and sequenced in their entirety. Consensus sequences for each gene were established after comparing the forward and reverse sequences of a minimum of 5 clones.

### Sequence data and annotation


Eleven novel zebrafish gene sequences were deposited in the NCBI GenBank database: z $\beta$ 1A [Genbank:DO489722], z $\beta$ 1B [Genbank:DO489723], z $\beta$ 1C [Genbank:DO489724], z $\beta$ 1D [Genbank:DO489725], z $\beta$ 2A [Genbank:DO489726], z $\beta$ 2B [Genbank:DO489727], z $\beta$ 2C [Genbank:DO489728], z $\beta$ 2D [Genbank:DO489729], z $\beta$ 3 [Genbank:DO489730], z $\beta$ 4 locus 1 [Genbank:DO489731], z $\beta$ 4 locus 2 [Genbank:DO489732]. Intron-exon boundaries for each gene and splice variant were determined by comparing cDNA and genomic sequences. Sequences for zebrafish  $\beta 1$ - $\beta 4$  were annotated by alignment with their mammalian orthologs using the AlignX function of Vector NTI v9.1 (Invitrogen) and by use of the following informatics resources: signal peptide (SMART) [90], IG-like domain (SMART [90], NCBI conserved domain database, CDD, v2.06 [91], and REF [14]), transmembrane (TMpred [92] and REF [93]), disulfide bridge (Prosite) [94], and putative N-linked glycosylation sites (NetNGlyc 1.0 server) [95].

### Gene expression

Tissues were dissected from wild type adult zebrafish (strain TuAB, 12–16 months old) and flash frozen on dry ice in ethanol. Tissue-specific total RNA was isolated and purified as described above. First strand cDNA synthesis was performed using 1  $\mu$ g of RNA from each tissue, random hexamer primers, and Transcriptor reverse transcriptase enzyme (Roche). 2  $\mu$ l of first strand cDNA was used in 25  $\mu$ l PCR reactions with Expand High Fidelity DNA polymerase (Roche). Amplicons were analyzed on a 1–2% agarose gel made with 1 x Tris-Acetate-EDTA (TAE) buffer. Primer pairs used to amplify each individual gene and splice variant are listed in Table 5. All amplicons were subcloned directly into the pGEM-TEasy vector (Promega) and sequenced in their entirety in both sense and anti-sense directions.

### Identification of $\beta$ subunit-like genes

Human sodium channel  $\beta$  subunit nucleotide and amino acid sequences were used in BLAST searches of the human genome to identify  $\beta$  subunit-like genes. Four human genes sharing the greatest homology as well as similar genomic organization and/or synteny with known sodium channel  $\beta$  subunit genes were included for further analysis: myelin protein zero (MPZ) [Genbank:NP\_000521], myelin protein zero-like gene, isoform A (MPZL1 isoform A) [Genbank:NP\_003944], epithelial V-like antigen (EVA) [Genbank:NP\_005788],



**Figure 9**  
**Phylogenetic analysis demonstrates that vertebrate sodium channel  $\beta$ 1-4 subunit genes are orthologous, that  $\beta$ 1/ $\beta$ 3 and  $\beta$ 2/ $\beta$ 4 are closely related, and that z $\beta$ 4.1 and z $\beta$ 4.2 resulted from a recent gene duplication in fish.**  
 Actual (human, zebrafish) and predicted (chicken, frog) amino acid sequences of  $\beta$  subunit and related genes were aligned using CLUSTALX (v1.83). Phylogenetic trees were reconstructed using the neighbor-joining method of Saitou and Nei and viewed with NJPlot software. Alignment gaps were excluded and the Kimura correction was made for multiple substitutions. Bootstrapping (n = 1000) was applied to test the robustness of each node. Tree is unrooted due to the lack of evidence for  $\beta$  subunit-like genes in invertebrate species. HUGO gene nomenclature symbol IDs: MPZ = myelin protein zero; MPZL1 = myelin protein zero-like gene isoform A; EVA1 = epithelial V-like antigen I; EVA1L = unannotated gene similar to EVA1.

and epithelial V-like antigen-like gene (hypothetical protein LOC196264 or EVA1-like gene) [Genbank:NP\_938016].

**Analysis of synteny**

Synteny between vertebrate  $\beta$  subunits and related genes was assessed by chromosomal walking and reciprocal BLAST searches of genes adjacent to  $\beta$  subunit loci in human, zebrafish, *Rattus norvegicus*, *Xenopus tropicalis*, and *Gallus gallus* genome databases (Ensembl).

**Analysis of phylogeny**

To estimate phylogeny, additional  $\beta$  subunit and  $\beta$  subunit-like gene sequences were identified in *Rattus norvegicus*,

*Xenopus tropicalis*, and *Gallus gallus* genome databases (Ensembl). Predicted amino acid sequences were aligned with cloned zebrafish sequences using CLUSTALX (v1.83). Phylogenetic trees were reconstructed using the neighbor-joining method of Saitou and Nei (Ref [96]) and viewed with NJplot software. Alignment gaps were excluded in the analysis, and the Kimura correction was made for multiple substitutions [97]. The robustness of each node in the phylogenetic tree was analyzed with bootstrap analysis (n = 1000). Trees were unrooted due to limited evidence for  $\beta$  subunit-like genes in non-vertebrate species.

### Genome databases

All genome resources utilized for this study were accessed through the Ensembl gateway [98] with the exception of the sea urchin genome, which is available for BLAST searches by the National Human Genome Sequencing Center at Baylor College of Medicine [99].

### Expression vectors

The full-length zebrafish  $\beta$ 1D splice variant was amplified in one step from total D2 embryonic RNA template using the Titan one-tube RT-PCR system (Roche) and the following primers: Fwd: *Spe* I-GACTCT-GAAAACAAAGCCTG, Rev: *Sac* II-AGAGCTTCAAGCTTTGGCT. The 733 base pair product was subcloned directly into the pGEM-TEasy vector. Multiple clones were purified, sequenced, and screened against the consensus sequence determined during the cloning of the  $z\beta$ 1 gene. Insert was digested out of a single  $z\beta$ 1D consensus clone and subcloned into the pGFP-IRES vector (Clontech) using *Spe* I and *Sac* II restriction sites. We have used similar methods to identify, clone and assemble a full-length *zscn5a* expression construct (pBK-CMV-*zscn5a*) which encodes zebrafish  $Na_v1.5$ . These methods are described elsewhere (manuscript in preparation).

### Transient transfection and electrophysiology

Cultured Chinese Hamster Ovary (CHO) cells were transiently transfected with pBK-CMV-*zscn5a* and pGFP-IRES-*z\beta*1D constructs using FuGENE6 (Roche). Cells were grown for 48 hours after transfection before electrophysiologic study. Whole-cell voltage clamp was performed at room temperature with 2-M $\Omega$  patch microelectrodes and an Axopatch 200A amplifier. To minimize the capacitive transients, we compensated for approximately 70% to 80% of the cell capacitance and series resistance [100]. Cells exhibiting very large currents (> 6 nA) were also excluded from further analysis. Cells were also excluded if voltage control was compromised. The extracellular bath solution contained (in mmol/L) NaCl 145, KCl 4.0, MgCl<sub>2</sub> 1.0, CaCl<sub>2</sub> 1.8, glucose 10, and HEPES 10; the pH was 7.4, adjusted with NaOH. The pipette (intracellular) solution contained (in mmol/L) NaF 10, CsF 110, CsCl 20, EGTA 10, and HEPES 10; the pH was 7.4, adjusted with CsOH. Cells were held at -120 mV, and activating currents were elicited with depolarizing pulses from -100 to +50 mV in 10 mV increments. Specific clamp protocols are indicated with the data. Data were acquired by pClamp8.0 (Axon Instruments Inc), sampled at 50 kHz, and low-pass filtered at 5 kHz. All currents were normalized to the cell capacitance calculated by Membrane Test (OUT O) in pClamp8.0.

### Abbreviations

BLAST Basic local alignment search tool

BLASTN BLAST nucleotide

BLASTP BLAST protein

dbSNP Database of single nucleotide polymorphisms

HUGO Human genome organization

IRES Internal ribosomal entry site

mRNA Message ribonucleic acid

$Na_v1$  Voltage-gated sodium channel, family 1

NCBI National Center for Biotechnology Information

*Para Paralytic* gene, *Drosophila*

PCR Polymerase chain reaction

PharmGKB Pharmacogenomics knowledge base

RACE Rapid amplification of cDNA ends

RT-PCR Reverse transcription polymerase chain reaction

SNPs Single nucleotide polymorphisms

*TEH1-4Tip-E* homologous genes 1–4

*Tip-E* Temperature-induced paralysis gene, *Drosophila*

TMPred Transmembrane prediction software

UTR Untranslated region

*z\beta*1 zebrafish sodium channel beta 1 subunit gene (zebrafish *scn1b*).

*z\beta*2 zebrafish sodium channel beta 2 subunit gene (zebrafish *scn2b*).

*z\beta*3 zebrafish sodium channel beta 3 subunit gene (zebrafish *scn3b*).

*z\beta*4.1 zebrafish sodium channel beta 4.1 subunit gene (zebrafish *scn4.1b*).

*z\beta*4.2 zebrafish sodium channel beta 4.2 subunit gene (zebrafish *scn4.2b*).

*zscn5a* zebrafish homolog to the mammalian sodium channel pore-forming  $\alpha$  subunit gene *scn5a*.

## Authors' contributions

SSC designed this study with input from DMR and TPZ. SSC collected, analyzed, and summarized the data for all figures and tables except Figure 6/Table 6, which represent electrophysiological analyses performed by HW using expression constructs built by SSC. DMR and TPZ were the co-principal investigators of this project, overseeing the design, interpretation, and reporting of this study. SSC wrote the manuscript and all authors read and approved of the final version.

## Additional material

### Additional file 1

*Hydropathy analysis of zebrafish  $\beta$  subunit amino acid sequences reveals conserved protein secondary structure. TMpred hydropathy plot based on the method of Kyte and Doolittle. Dotted red line drawn at 0 (neutral). SP = signal peptide and TM = transmembrane domain. Dotted line plot (black) indicates outside > inside orientation relative to membrane, while solid line plot (black) indicates inside > outside orientation. A) Hydropathy plots for  $\alpha\beta 1$  variants A, B, and D are nearly identical ( $\alpha\beta 1D$  shown). B)  $\alpha\beta 2A$  and  $\alpha\beta 2B-D$ . C)  $\alpha\beta 3$ . D)  $\alpha\beta 4.1$  and  $\alpha\beta 4.2$ .*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2148-7-113-S1.ppt>]

### Additional file 2

*Prediction of conserved extracellular V-type IG domains in the amino acid sequences of the most highly-conserved splice variants of  $\alpha\beta 1-4$ , NCBI Conserved Domain Database (CDD v2.09). Predicted V-type IG domain is shown in red, and the scale denotes the length of the full-length protein. A)  $\alpha\beta 1D$ , B)  $\alpha\beta 2B$ , C)  $\alpha\beta 3$ , D)  $\alpha\beta 4.1$ , E)  $\alpha\beta 4.2$ .*

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2148-7-113-S2.ppt>]

## Acknowledgements

This work was supported by Public Health Service award T32 GM07347 from the National Institute of General Medical Studies for the Vanderbilt Medical-Scientist Training Program. Portions of this work were also supported by HL46681 (DMR) and HL073348 (TPZ).

## References

- Hille B: *Ion channels of excitable membranes* 3rd edition. Sunderland, MA, Sinauer Associates; 2001.
- Yu FH, Catterall WA: **Overview of the voltage-gated sodium channel family.** *Genome Biol* 2003, **4**:207.
- Goldin AL: **Resurgence of sodium channel research.** *Annu Rev Physiol* 2001, **63**:871-894.
- Anderson PA, Greenberg RM: **Phylogeny of ion channels: clues to structure and function.** *Comp Biochem Physiol B Biochem Mol Biol* 2001, **129**:17-28.
- Hartshorne RP, Messner DJ, Coppersmith JC, Catterall WA: **The saxitoxin receptor of the sodium channel from rat brain. Evidence for two nonidentical beta subunits.** *J Biol Chem* 1982, **257**:13888-13891.
- Barchi RL: **Protein components of the purified sodium channel from rat skeletal muscle sarcolemma.** *J Neurochem* 1983, **40**:1377-1385.
- Roberts RH, Barchi RL: **The voltage-sensitive sodium channel from rabbit skeletal muscle. Chemical characterization of subunits.** *J Biol Chem* 1987, **262**:2298-2303.
- Isom LL, De Jongh KS, Catterall WA: **Auxiliary subunits of voltage-gated ion channels.** *Neuron* 1994, **12**:1183-1194.
- Maier SK, Westenbroek RE, McCormick KA, Curtis R, Scheuer T, Catterall WA: **Distinct subcellular localization of different sodium channel alpha and beta subunits in single ventricular myocytes from mouse heart.** *Circulation* 2004, **109**:1421-1427.
- Isom LL, De Jongh KS, Patton DE, Reber BF, Offord J, Charbonneau H, Walsh K, Goldin AL, Catterall WA: **Primary structure and functional expression of the beta 1 subunit of the rat brain sodium channel.** *Science* 1992, **256**:839-842.
- Isom LL, Ragsdale DS, De Jongh KS, Westenbroek RE, Reber BF, Scheuer T, Catterall WA: **Structure and function of the beta 2 subunit of brain sodium channels, a transmembrane glycoprotein with a CAM motif.** *Cell* 1995, **83**:433-442.
- Morgan K, Stevens EB, Shah B, Cox PJ, Dixon AK, Lee K, Pinnock RD, Hughes J, Richardson PJ, Mizuguchi K, Jackson AP: **Beta 3: an additional auxiliary subunit of the voltage-sensitive sodium channel that modulates channel gating with distinct kinetics.** *Proc Natl Acad Sci U S A* 2000, **97**:2308-2313.
- Yu FH, Westenbroek RE, Silos-Santiago I, McCormick KA, Lawson D, Ge P, Ferriera H, Lilly J, DiStefano PS, Catterall WA, Scheuer T, Curtis R: **Sodium channel beta4, a new disulfide-linked auxiliary subunit with similarity to beta2.** *J Neurosci* 2003, **23**:7577-7585.
- Isom LL, Catterall WA: **Na<sup>+</sup> channel subunits and Ig domains.** *Nature* 1996, **383**:307-308.
- Catterall WA, Goldin AL, Waxman SG: **International Union of Pharmacology. XLVII. Nomenclature and structure-function relationships of voltage-gated sodium channels.** *Pharmacol Rev* 2005, **57**:397-409.
- Chen C, Westenbroek RE, Xu X, Edwards CA, Sorenson DR, Chen Y, McEwen DP, O'Malley HA, Bharucha V, Meadows LS, Knudsen GA, Vilaythong A, Noebels JL, Saunders TL, Scheuer T, Shrager P, Catterall WA, Isom LL: **Mice lacking sodium channel beta1 subunits display defects in neuronal excitability, sodium channel expression, and nodal architecture.** *J Neurosci* 2004, **24**:4030-4042.
- Chen C, Bharucha V, Chen Y, Westenbroek RE, Brown A, Malhotra JD, Jones D, Avery C, Gillespie PJ III, Kazen-Gillespie KA, Kazarinova-Noyes K, Shrager P, Saunders TL, Macdonald RL, Ransom BR, Scheuer T, Catterall WA, Isom LL: **Reduced sodium channel density, altered voltage dependence of inactivation, and increased susceptibility to seizures in mice lacking sodium channel beta 2-subunits.** *Proc Natl Acad Sci U S A* 2002, **99**:17072-17077.
- Meadows LS, Isom LL: **Sodium channels as macromolecular complexes: implications for inherited arrhythmia syndromes.** *Cardiovasc Res* 2005, **67**:448-458.
- Isom LL: **Sodium channel beta subunits: anything but auxiliary.** *Neuroscientist* 2001, **7**:42-54.
- George AL Jr.: **Inherited disorders of voltage-gated sodium channels.** *J Clin Invest* 2005, **115**:1990-1999.
- Escayg A, MacDonald BT, Meisler MH, Baulac S, Huberfeld G, An-Gourfinkel I, Brice A, LeGuern E, Moulard B, Chaigne D, Buresi C, Malafosse A: **Mutations of SCN1A, encoding a neuronal sodium channel, in two families with GEFS+2.** *Nat Genet* 2000, **24**:343-345.
- Sugawara T, Tsurubuchi Y, Agarwala KL, Ito M, Fukuma G, Mazaki-Miyazaki E, Nagafuji H, Noda M, Imoto K, Wada K, Mitsudome A, Kaneko S, Montal M, Nagata K, Hirose S, Yamakawa K: **A missense mutation of the Na<sup>+</sup> channel alpha II subunit gene Na(v)1.2 in a patient with febrile and afebrile seizures causes channel dysfunction.** *Proc Natl Acad Sci U S A* 2001, **98**:6384-6389.
- Weiss LA, Escayg A, Kearney JA, Trudeau M, MacDonald BT, Mori M, Reichert J, Buxbaum JD, Meisler MH: **Sodium channels SCN1A, SCN2A and SCN3A in familial autism.** *Mol Psychiatry* 2003, **8**:186-194.
- Ptacek LJ, George AL Jr., Griggs RC, Tawil R, Kallen RG, Barchi RL, Robertson M, Leppert MF: **Identification of a mutation in the gene causing hyperkalemic periodic paralysis.** *Cell* 1991, **67**:1021-1027.
- Wang Q, Shen J, Splawski I, Atkinson D, Li Z, Robinson JL, Moss AJ, Towbin JA, Keating MT: **SCN5A mutations associated with an**

- inherited cardiac arrhythmia, long QT syndrome. *Cell* 1995, **80**:805-811.
26. Trudeau MM, Dalton JC, Day JW, Ranum LP, Meisler MH: **Heterozygosity for a protein truncation mutation of sodium channel SCN8A in a patient with cerebellar atrophy, ataxia, and mental retardation.** *J Med Genet* 2006, **43**:527-530.
  27. Yang Y, Wang Y, Li S, Xu Z, Li H, Ma L, Fan J, Bu D, Liu B, Fan Z, Wu G, Jin J, Ding B, Zhu X, Shen Y: **Mutations in SCN9A, encoding a sodium channel alpha subunit, in patients with primary erythralgia.** *J Med Genet* 2004, **41**:171-174.
  28. Wallace RH, Wang DW, Singh R, Scheffer IE, George AL Jr., Phillips HA, Saar K, Reis A, Johnson EW, Sutherland GR, Berkovic SF, Mulley JC: **Febrile seizures and generalized epilepsy associated with a mutation in the Na<sup>+</sup>-channel beta1 subunit gene SCN1B.** *Nat Genet* 1998, **19**:366-370.
  29. Spampinato J, Kearney JA, de HG, McEwen DP, Escayg A, Aradi I, MacDonald BT, Levin SI, Soltész I, Benna P, Montalenti E, Isom LL, Goldin AL, Meisler MH: **A novel epilepsy mutation in the sodium channel SCN1A identifies a cytoplasmic domain for beta subunit interaction.** *J Neurosci* 2004, **24**:10022-10034.
  30. An RH, Wang XL, Kerem B, Benhorin J, Medina A, Goldmit M, Kass RS: **Novel LQT-3 mutation affects Na<sup>+</sup> channel activity through interactions between alpha- and beta1-subunits.** *Circ Res* 1998, **83**:141-146.
  31. Wan X, Wang Q, Kirsch GE: **Functional suppression of sodium channels by beta(1)-subunits as a molecular mechanism of idiopathic ventricular fibrillation.** *J Mol Cell Cardiol* 2000, **32**:1873-1884.
  32. Medeiros-Domingo A, Kaku T, Tester DJ, Iturralde-Torres P, Itty A, Ye B, Valdivia C, Ueda K, Canales-Quinteros S, Tusie-Luna MT, Makielski JC, Ackerman MJ: **SCN4B-Encoded Sodium Channel (beta)4 Subunit in Congenital Long-QT Syndrome.** *Circulation* 2007, **116**:134-142.
  33. Okamura Y, Nishino A, Murata Y, Nakajo K, Iwasaki H, Ohtsuka Y, Tanaka-Kunishima M, Takahashi N, Hara Y, Yoshida T, Nishida M, Okado H, Watari H, Meinertzhagen IA, Satoh N, Takahashi K, Satou Y, Okada Y, Mori Y: **Comprehensive analysis of the ascidian genome reveals novel insights into the molecular evolution of ion channel genes.** *Physiol Genomics* 2005, **22**:269-282.
  34. Littleton JT, Ganetzky B: **Ion channels and synaptic organization: analysis of the Drosophila genome.** *Neuron* 2000, **26**:35-43.
  35. Blackshaw SE, Henderson LP, Malek J, Porter DM, Gross RH, Angstadt JD, Levasseur SM, Maue RA: **Single-cell analysis reveals cell-specific patterns of expression of a family of putative voltage-gated sodium channel genes in the leech.** *J Neurobiol* 2003, **55**:355-371.
  36. Goldin AL: **Evolution of voltage-gated Na(+) channels.** *J Exp Biol* 2002, **205**:575-584.
  37. Plummer NW, Meisler MH: **Evolution and diversity of mammalian sodium channel genes.** *Genomics* 1999, **57**:323-331.
  38. Lopreato GF, Lu Y, Southwell A, Atkinson NS, Hillis DM, Wilcox TP, Zakon HH: **Evolution and divergence of sodium channel genes in vertebrates.** *Proc Natl Acad Sci U S A* 2001, **98**:7588-7592.
  39. Piontkivska H, Hughes AL: **Evolution of vertebrate voltage-gated ion channel alpha chains by sequential gene duplication.** *J Mol Evol* 2003, **56**:277-285.
  40. Novak AE, Jost MC, Lu Y, Taylor AD, Zakon HH, Ribera AB: **Gene duplications and evolution of vertebrate voltage-gated sodium channels.** *J Mol Evol* 2006, **63**:208-221.
  41. Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, Ho RK, Langeland J, Prince V, Wang YL, Westerfield M, Ekker M, Postlethwait JH: **Zebrafish hox clusters and vertebrate genome evolution.** *Science* 1998, **282**:1711-1714.
  42. Hughes AL: **The evolution of functionally novel proteins after gene duplication.** *Proc Biol Sci* 1994, **256**:119-124.
  43. McCormick KA, Isom LL, Ragsdale D, Smith D, Scheuer T, Catterall WA: **Molecular determinants of Na<sup>+</sup> channel function in the extracellular domain of the beta1 subunit.** *J Biol Chem* 1998, **273**:3954-3962.
  44. Shapiro L, Doyle JP, Hensley P, Colman DR, Hendrickson WA: **Crystal structure of the extracellular domain from PO, the major structural protein of peripheral nerve myelin.** *Neuron* 1996, **17**:435-449.
  45. Malhotra JD, Koopmann MC, Kazen-Gillespie KA, Fettman N, Hortsch M, Isom LL: **Structural requirements for interaction of sodium channel beta 1 subunits with ankyrin.** *J Biol Chem* 2002, **277**:26681-26688.
  46. Kim DY, Ingano LA, Carey BW, Pettingell WH, Kovacs DM: **Presenilin/gamma-secretase-mediated cleavage of the voltage-gated sodium channel beta2-subunit regulates cell adhesion and migration.** *J Biol Chem* 2005, **280**:23251-23261.
  47. Scheffer IE, Harkin LA, Grinton BE, Dibbens LM, Turner SJ, Zielinski MA, Xu R, Jackson G, Adams J, Connellan M, Petrou S, Wellard RM, Briellmann RS, Wallace RH, Mulley JC, Berkovic SF: **Temporal lobe epilepsy and GEFs+ phenotypes associated with SCN1B mutations.** *Brain* 2007, **130**(Pt.1):100-109.
  48. Audenaert D, Claes L, Ceulemans B, Lofgren A, Van BC, De JP: **A deletion in SCN1B is associated with febrile seizures and early-onset absence epilepsy.** *Neurology* 2003, **61**:854-856.
  49. Kazen-Gillespie KA, Ragsdale DS, D'Andrea MR, Mattei LN, Rogers KE, Isom LL: **Cloning, localization, and functional expression of sodium channel beta1A subunits.** *J Biol Chem* 2000, **275**:1079-1088.
  50. Qin N, D'Andrea MR, Lubin ML, Shafaei N, Codd EE, Correa AM: **Molecular cloning and functional expression of the human sodium channel beta1B subunit, a novel splicing variant of the beta1 subunit.** *Eur J Biochem* 2003, **270**:4762-4770.
  51. Oh Y, Waxman SG: **The beta 1 subunit mRNA of the rat brain Na<sup>+</sup> channel is expressed in glial cells.** *Proc Natl Acad Sci U S A* 1994, **91**:9985-9989.
  52. Dib-Hajj SD, Waxman SG: **Genes encoding the beta 1 subunit of voltage-dependent Na<sup>+</sup> channel in rat, mouse and human contain conserved introns.** *FEBS Lett* 1995, **377**:485-488.
  53. Malhotra JD, Kazen-Gillespie K, Hortsch M, Isom LL: **Sodium channel beta subunits mediate homophilic cell adhesion and recruit ankyrin to points of cell-cell contact.** *J Biol Chem* 2000, **275**:11383-11388.
  54. Makita N, Bennett PB Jr., George AL Jr.: **Voltage-gated Na<sup>+</sup> channel beta 1 subunit mRNA expressed in adult human skeletal muscle, heart, and brain is encoded by a single gene.** *J Biol Chem* 1994, **269**:7571-7578.
  55. Qu Y, Isom LL, Westenbroek RE, Rogers JC, Tanada TN, McCormick KA, Scheuer T, Catterall WA: **Modulation of cardiac Na<sup>+</sup> channel expression in Xenopus oocytes by beta 1 subunits.** *J Biol Chem* 1995, **270**:25696-25701.
  56. Nuss HB, Chiamvimonvat N, Perez-Garcia MT, Tomaselli GF, Marban E: **Functional association of the beta 1 subunit with human cardiac (hH1) and rat skeletal muscle (mu 1) sodium channel alpha subunits expressed in Xenopus oocytes.** *J Gen Physiol* 1995, **106**:1171-1191.
  57. Xiao YF, Wright SN, Wang GK, Morgan JP, Leaf A: **Coexpression with beta(1)-subunit modifies the kinetics and fatty acid block of hH1(alpha) Na(+) channels.** *Am J Physiol Heart Circ Physiol* 2000, **279**:H35-H46.
  58. Dehal P, Satou Y, Campbell RK, Chapman J, Degnan B, De TA, Davidson B, Di GA, Gelpke M, Goodstein DM, Harafuji N, Hastings KE, Ho I, Hotta K, Huang W, Kawashima T, Lemaire P, Martinez D, Meinertzhagen IA, Necula S, Nonaka M, Putnam N, Rash S, Saiga H, Satake M, Terry A, Yamada L, Wang HG, Awazu S, Azumi K, Boore J, Branno M, Chin-Bow S, DeSantis R, Doyle S, Francino P, Keys DN, Haga S, Hayashi H, Hino K, Imai KS, Inaba K, Kano S, Kobayashi K, Kobayashi M, Lee BI, Makabe KVV, Manohar C, Matassi G, Medina M, Mochizuki Y, Mount S, Morishita T, Miura S, Nakayama A, Nishizaka S, Nomoto H, Ohta F, Oishi K, Rigoutsos I, Sano M, Sasaki A, Sasakura Y, Shoguchi E, T S, Spagnuolo A, Stainier D, Suzuki MM, Tassy O, Takatori N, Tokuoka M, Yagi K, Yoshizaki F, Wada S, Zhang C, Hyatt PD, Larimer F, Dettler C, Doggett N, Glavina T, Hawkins T, Richardson P, Lucas S, Kohara Y, Levine M, Satoh N, Rokhsar DS: **The draft genome of Ciona intestinalis: insights into chordate and vertebrate origins.** *Science* 2002, **298**:2157-2167.
  59. Canestro C, Bassham S, Postlethwait JH: **Seeing chordate evolution through the Ciona genome sequence.** *Genome Biol* 2003, **4**:208.
  60. Azumi K, De SR, De TA, Rigoutsos I, Yoshizaki F, Pinto MR, Marino R, Shida K, Ikeda M, Ikeda M, Arai M, Inoue Y, Shimizu T, Satoh N, Rokhsar DS, Du PL, Kasahara M, Satake M, Nonaka M: **Genomic analysis of immunity in a Urochordate and the emergence of the vertebrate immune system: "waiting for Godot".** *Immunogenetics* 2003, **55**:570-581.

61. Du PL, Zucchetti I, De SR: **Immunoglobulin superfamily receptors in protochordates: before RAG time.** *Immunol Rev* 2004, **198**:233-248.
62. Sodergren E, Weinstock GM, Davidson EH, Cameron RA, Gibbs RA, Angerer RC, Angerer LM, Arnone MI, Burgess DR, Burke RD, Coffman JA, Dean M, Elphick MR, Etensohn CA, Foltz KR, Hamdoun A, Hynes RO, Klein WH, Marzluff W, McClay DR, Morris RL, Mushagian A, Rast JP, Smith LC, Thorndyke MC, Vacquier VD, Wessel GM, Wray G, Zhang L, Elsik CG, Ermolaeva O, Hlavina W, Hofmann G, Kitts P, Landrum MJ, Mackey AJ, Maglott D, Panopoulou G, Poustka AJ, Pruitt K, Sapojnikov V, Song X, Souvorov A, Solovyev V, Wei Z, Whittaker CA, Worley K, Durbin KJ, Shen Y, Fedrigo O, Garfield D, Haygood R, Primus A, Satija R, Severson T, Gonzalez-Garay ML, Jackson AR, Milosavljevic A, Tong M, Killian CE, Livingston BT, Wilt FH, Adams N, Belle R, Carbonneau S, Cheung R, Cormier P, Cosson B, Croce J, Fernandez-Guerra A, Genevieve AM, Goel M, Kelkar H, Morales J, Mulner-Lorillon O, Robertson AJ, Goldstone JV, Cole B, Epel D, Gold B, Hahn ME, Howard-Ashby M, Scally M, Stegeman JJ, Allgood EL, Cool J, Judkins KM, McCafferty SS, Musante AM, Obar RA, Rawson AP, Rossetti BJ, Gibbons IR, Hoffman MP, Leone A, Istrail S, Materna SC, Samanta MP, Stolc V, Tongprasit W, Tu Q, Bergeron KF, Brandhorst BP, Whittle J, Berney K, Bottjer DJ, Calestani C, Peterson K, Chow E, Yuan QA, Elhaik E, Graur D, Reese JT, Bosdet I, Heesun S, Marra MA, Schein J, Anderson MK, Brockton V, Buckley KM, Cohen AH, Fugmann SD, Hibino T, Loza-Coll M, Majeske AJ, Messier C, Nair SV, Pancer Z, Terwilliger DP, Agca C, Arboleda E, Chen N, Churcher AM, Hallbook F, Humphrey GW, Idris MM, Kiyama T, Liang S, Mellott D, Mu X, Murray G, Olinski RP, Raible F, Rowe M, Taylor JS, Tessmar-Raible K, Wang D, Wilson KH, Yaguchi S, Gaasterland T, Galindo BE, Gunaratne HJ, Juliano C, Kinukawa M, Moy GW, Neill AT, Nomura M, Raisch M, Reade A, Roux MM, Song JL, Su YH, Townley IK, Voronina E, Wong JL, Amore G, Branno M, Brown ER, Cavalieri V, Duboc V, Duloquin L, Flytzanis C, Gache C, Lapraz F, Lepage T, Locascio A, Martinez P, Matassi G, Matranga V, Range R, Rizzo F, Rottinger E, Beane W, Bradham C, Byrum C, Glenn T, Hussain S, Manning G, Miranda E, Thomason R, Walton K, Wikramanayake A, Wu SY, Xu R, Brown CT, Chen L, Gray RF, Lee PY, Nam J, Oliveri P, Smith J, Muzny D, Bell S, Chacko J, Cree A, Curry S, Davis C, Dinh H, Dugan-Rocha S, Fowler J, Gill R, Hamilton C, Hernandez J, Hines S, Hume J, Jackson L, Jolivet A, Kovar C, Lee S, Lewis L, Miner G, Morgan M, Nazareth LV, Okwuonu G, Parker D, Pu LL, Thorn R, Wright R: **The genome of the sea urchin *Strongylocentrotus purpuratus*.** *Science* 2006, **314**:941-952.
63. Jackson FR, Wilson SD, Hall LM: **The tip-E mutation of *Drosophila* decreases saxitoxin binding and interacts with other mutations affecting nerve membrane excitability.** *J Neurogenet* 1986, **3**:1-17.
64. O'Dowd DK, Aldrich RW: **Voltage-clamp analysis of sodium channels in wild-type and mutant *Drosophila* neurons.** *J Neurosci* 1988, **8**:3633-3643.
65. Feng G, Deak P, Chopra M, Hall LM: **Cloning and functional analysis of TipE, a novel membrane protein that enhances *Drosophila para* sodium channel function.** *Cell* 1995, **82**:1001-1011.
66. Warmke JW, Reenan RA, Wang P, Qian S, Arena JP, Wang J, Wunderler D, Liu K, Kaczorowski GJ, Van der Ploeg LH, Ganetzky B, Cohen CJ: **Functional expression of *Drosophila para* sodium channels. Modulation by the membrane protein TipE and toxin pharmacology.** *J Gen Physiol* 1997, **110**:119-133.
67. Hodges DD, Lee D, Preston CF, Boswell K, Hall LM, O'Dowd DK: **tipE regulates Na<sup>+</sup>-dependent repetitive firing in *Drosophila* neurons.** *Mol Cell Neurosci* 2002, **19**:402-416.
68. Derst C, Walther C, Veh RV, Wicher D, Heinemann SH: **Four novel sequences in *Drosophila melanogaster* homologous to the auxiliary Para sodium channel subunit TipE.** *Biochem Biophys Res Commun* 2006, **339**:939-948.
69. Hobert O, Hutter H, Hynes RO: **The immunoglobulin superfamily in *Caenorhabditis elegans* and *Drosophila melanogaster*.** *Development* 2004, **131**:2237-2238.
70. Hartshorne RP, Catterall WA: **Purification of the saxitoxin receptor of the sodium channel from rat brain.** *Proc Natl Acad Sci U S A* 1981, **78**:4620-4624.
71. Hartshorne RP, Catterall WA: **The sodium channel from rat brain. Purification and subunit composition.** *J Biol Chem* 1984, **259**:1667-1675.
72. Hartshorne RP, Keller BU, Talvenheimo JA, Catterall WA, Montal M: **Functional reconstitution of the purified brain sodium channel in planar lipid bilayers.** *Proc Natl Acad Sci U S A* 1985, **82**:240-244.
73. Agnew WS, Levinson SR, Brabson JS, Raftery MA: **Purification of the tetrodotoxin-binding component associated with the voltage-sensitive sodium channel from *Electrophorus electricus* electroplax membranes.** *Proc Natl Acad Sci U S A* 1978, **75**:2606-2610.
74. Miller JA, Agnew WS, Levinson SR: **Principal glycopeptide of the tetrodotoxin/saxitoxin binding protein from *Electrophorus electricus*: isolation and partial chemical and physical characterization.** *Biochemistry* 1983, **22**:462-470.
75. Rosenberg RL, Tomiko SA, Agnew WS: **Single-channel properties of the reconstituted voltage-regulated Na channel isolated from the electroplax of *Electrophorus electricus*.** *Proc Natl Acad Sci U S A* 1984, **81**:5594-5598.
76. Recio-Pinto E, Duch DS, Levinson SR, Urban BW: **Purified and unpurified sodium channels from eel electroplax in planar lipid bilayers.** *J Gen Physiol* 1987, **90**:375-395.
77. Casadei JM, Gordon RD, Barchi RL: **Immunoaffinity isolation of Na<sup>+</sup> channels from rat skeletal muscle. Analysis of subunits.** *J Biol Chem* 1986, **261**:4318-4323.
78. Lombet A, Lazdunski M: **Characterization, solubilization, affinity labeling and purification of the cardiac Na<sup>+</sup> channel using Tityus toxin gamma.** *Eur J Biochem* 1984, **141**:651-660.
79. Zakon HH, Lu Y, Zwickl DJ, Hillis DM: **Sodium channel genes and the evolution of diversity in communication signals of electric fishes: Convergent molecular evolution.** *Proc Natl Acad Sci U S A* 2006, **103**:3675-3680.
80. Abriel H, Kass RS: **Regulation of the voltage-gated cardiac sodium channel Nav1.5 by interacting proteins.** *Trends Cardiovasc Med* 2005, **15**:35-40.
81. Camacho JA, Hensellek S, Rougier JS, Blechschmidt S, Abriel H, Benndorf K, Zimmer T: **Modulation of Nav1.5 channel function by an alternatively spliced sequence in the DII/DIII linker region.** *J Biol Chem* 2006, **281**:9498-9506.
82. Thimmapaya R, Neelands T, Niforatos W, vis-Taber RA, Choi W, Putman CB, Kroeger PE, Packer J, Gopalakrishnan M, Faltynek CR, Surowy CS, Scott VE: **Distribution and functional characterization of human Nav1.3 splice variants.** *Eur J Neurosci* 2005, **22**:1-9.
83. Tan BH, Valdivia CR, Rok BA, Ye B, Ruwaldt KM, Tester DJ, Ackerman MJ, Makielski JC: **Common human SCN5A polymorphisms have altered electrophysiology when expressed in Q1077 splice variants.** *Heart Rhythm* 2005, **2**:741-747.
84. Makielski JC, Ye B, Valdivia CR, Pagel MD, Pu J, Tester DJ, Ackerman MJ: **A ubiquitous splice variant and a common polymorphism affect heterologous expression of recombinant human SCN5A heart sodium channels.** *Circ Res* 2003, **93**:821-828.
85. Zimmer T, Bollensdorff C, Haufe V, Birch-Hirschfeld E, Benndorf K: **Mouse heart Na<sup>+</sup> channels: primary structure and function of two isoforms and alternatively spliced variants.** *Am J Physiol Heart Circ Physiol* 2002, **282**:H1007-H1017.
86. Dietrich PS, McGivern JG, Delgado SG, Koch BD, Eglén RM, Hunter JC, Sangameswaran L: **Functional analysis of a voltage-gated sodium channel and its splice variant from rat dorsal root ganglia.** *J Neurochem* 1998, **70**:2262-2272.
87. O'Dowd DK, Gee JR, Smith MA: **Sodium current density correlates with expression of specific alternatively spliced sodium channel mRNAs in single neurons.** *J Neurosci* 1995, **15**:4005-4012.
88. Meadows L, Malhotra JD, Stetzer A, Isom LL, Ragsdale DS: **The intracellular segment of the sodium channel beta 1 subunit is required for its efficient association with the channel alpha subunit.** *J Neurochem* 2001, **76**:1871-1878.
89. **Simple Modular Architecture Research Tool (SMART)** [<http://smart.embl-heidelberg.de/>]. 2007.
90. **Conserved Domain Database, NCBI** [<http://www.ncbi.nlm.nih.gov/Structure/cdd/wrpsb.cgi>]. 2007.
91. **TM Pred** [<http://www.ch.embnet.org/>]. 2007.
92. Kyte J, Doolittle RF: **A simple method for displaying the hydrophobic character of a protein.** *J Mol Biol* 1982, **157**:105-132.
93. **Prosite** [<http://ca.expasy.org/>]. 2007.
94. **NetNGlyc** [<http://www.cbs.dtu.dk/services/>]. 2007.
95. Saitou N, Nei M: **The neighbor-joining method: a new method for reconstructing phylogenetic trees.** *Mol Biol Evol* 1987, **4**:406-425.

96. Kimura M: *The Neutral Theory of Molecular Evolution* Cambridge, Cambridge University Press; 1983.
97. **Ensembl genome browser** [[www.ensembl.org](http://www.ensembl.org)]. 2007.
98. **National Human Genome Sequencing Center, Baylor College of Medicine** [<http://www.hgsc.bcm.tmc.edu/projects/seaurchin/>]. 2007.
99. Darbar D, Yang T, Churchwell K, Wilde AA, Roden DM: **Unmasking of brugada syndrome by lithium.** *Circulation* 2005, **112**:1527-1531.
100. Eubanks J, Srinivasan J, Dinulos MB, Disteché CM, Catterall WA: **Structure and chromosomal localization of the beta2 subunit of the human brain sodium channel.** *Neuroreport* 1997, **8**:2775-2779.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

