

Mutation in *WNT10A* Is Associated with an Autosomal Recessive Ectodermal Dysplasia: The Odonto-onycho-dermal Dysplasia

Lynn Adaimy, Eliane Chouery, Hala Mégarbané, Salman Mroueh, Valérie Delague, Elsa Nicolas, Hanen Belguith, Philippe de Mazancourt, and André Mégarbané

Odonto-onycho-dermal dysplasia is a rare autosomal recessive syndrome in which the presenting phenotype is dry hair, severe hypodontia, smooth tongue with marked reduction of fungiform and filiform papillae, onychodysplasia, keratoderma and hyperhidrosis of palms and soles, and hyperkeratosis of the skin. We studied three consanguineous Lebanese Muslim Shiite families that included six individuals affected with odonto-onycho-dermal dysplasia. Using a homozygosity-mapping strategy, we assigned the disease locus to an ~9-cM region at chromosome 2q35-q36.2, located between markers *rs16853834* and *D2S353*, with a maximum multipoint LOD score of 5.7. Screening of candidate genes in this region led us to identify the same c.697G→T (p.Glu233X) homozygous nonsense mutation in exon 3 of the *WNT10A* gene in all patients. At the protein level, the mutation is predicted to result in a premature truncated protein of 232 aa instead of 417 aa. This is the first report to our knowledge of a human phenotype resulting from a mutation in *WNT10A*, and it is the first demonstration of an ectodermal dysplasia caused by an altered WNT signaling pathway, expanding the list of WNT-related diseases.

The ectodermal dysplasias (EDs) are a large and complex nosological group of diseases that have in common anomalies of the hair, teeth, nails, and sweat glands, with or without anomalies in other organs and systems. The incidence is estimated to be ~7 in 10,000 births. So far, >200 different pathological clinical conditions have been recognized and defined as EDs, all with Mendelian modes of inheritance.¹ Classification of EDs was first initiated by Freire-Maia and Pinheiro and was based on clinical descriptions and the ectodermal structures involved.²⁻⁴ This classification was followed by another one that introduced biological mechanisms involved in the pathogenesis of EDs.⁵ Two different groups were thus delineated, each likely to result from mutations in genes with similar functions and possibly involved in the same regulation of development mechanisms and/or in pathogenesis. Characterization of the genes' functions and of the different signaling pathways involved in ~30 different types of EDs led to expansion of the classification of EDs into four major functional subgroups that depend on molecular and biochemical criteria: cell-cell communication and signaling, adhesion, transcription regulation, and development.⁶

In the past few years, many publications indicated the importance of the Wnt pathway in hair-follicle morphogenesis, skin and teeth embryogenesis,⁷⁻¹⁰ and tooth renewal.¹¹ The *Wnt* gene family comprises genes showing strong sequence similarities to mouse *Wnt-1* and *Drosophila wingless* genes (Wnt Homepage and Laboratory of

Randall Moon). The *Wnt* genes encode small secreted proteins, ~350–400 residues in length, that are members of a large family of secreted cysteine-rich proteins that regulate cell-to-cell interactions during embryogenesis. They are found in almost all animal genomes (especially vertebrates). At least four Wnt signaling pathways have been elucidated, of which the canonical Wnt/ β -catenin signaling pathway,¹² which activates target genes in the nucleus, is the best understood. In humans, 19 WNT proteins have been identified that share 27%–83% amino acid sequence identity and a conserved pattern of 23 or 24 cysteine residues (reviewed by Cadigan and Nusse¹³ and Wodarz and Nusse¹⁴).

Since mutations in Wnt pathway genes in mice and zebrafish generally lead to developmental defects,¹⁵ severe abnormalities were expected to be due to mutations in WNT pathway genes in humans. However, despite considerable progress in understanding the mechanisms involved in WNT signaling pathways and their roles in development, studies implicating WNT signaling in human diseases and clinical conditions other than cancer have only recently emerged.¹⁶⁻²²

In this report, we show that *WNT10A* is the causative gene for a rare ED, odonto-onycho-dermal dysplasia (MIM 257980),²³⁻²⁷ expanding the list of WNT-related diseases.

Clinical findings.—Three consanguineous Lebanese Shiite Muslim families were included in this study (fig. 1): two of them (families 1 and 3) have already been described elsewhere,^{26,27} and one (family 2) was a branch of the first

From the Unité de Génétique Médicale, Faculté de Médecine, Université Saint-Joseph (L.A.; E.C.; E.N.; H.B.; A.M.), Service de Dermatologie (H.M.), and Department of Pediatrics, American University of Beirut (S.M.), Beirut; INSERM U491, "Génétique Médicale et Développement," Faculté de Médecine de la Timone, Marseille, France (V.D.); and Hôpital Raymond Poincaré, Université Versailles Saint Quentin en Yvelines EA 2493, Garches, France (P.D.M.)

Received April 11, 2007; accepted for publication May 23, 2007; electronically published August 9, 2007.

Address for correspondence and reprints: Dr. André Mégarbané, Unité de Génétique Médicale, Faculté de Médecine, Université Saint-Joseph de Beyrouth, 42 rue de Grenelle, 75007 Paris, France. E-mail: megarbane@usj.edu.lb

Am. J. Hum. Genet. 2007;81:821–828. © 2007 by The American Society of Human Genetics. All rights reserved. 0002-9297/2007/8104-0027\$15.00
DOI: 10.1086/520064






Figure 1. Pedigrees, haplotypes, and *WNT10A* mutation status of family 1 (A), family 2 (B), and family 3 (C). Individuals' numbers are listed below the pedigree symbols. Genotyped markers from the chromosome 2q34-q36.2 region are shown to the left, and individuals' allele numbers for each marker are given next to the bar. Black bars represent the haplotype segregating with the odonto-onycho-dermal gene. *WNT10A* is located between markers *D2S2722* and *rs1109867*. The mutation genotypes of *WNT10A* are shown below haplotypes. + indicates mutation Glu233X; - indicates wild-type allele. A border indicates the common haplotype 1-C-3-3-4-T-1-T between the three families, and the arrow shows the neomutation in patient V:3 of family 3.



of the two families described by Fadhil et al.²³ Clinical findings are summarized in table 1. In brief, patients in family 1 presented with dry hair, pilar keratosis, severe hypodontia, smooth tongue, onychodysplasia, palmar erythema, and keratoderma and hyperhidrosis of palms and soles. Histology of the skin showed orthokeratotic, hyperkeratosis, hypergranulosis, and mild acanthosis in the epidermis. Scanning electron microscopy of the hair showed longitudinal depressions in some hairs.²⁷

In family 2, two patients were examined, a boy and his paternal uncle. The boy (V:3 in fig. 1B and case 1 in family 2 in table 1) was the second son of consanguineous Lebanese parents. Pregnancy was uneventful, and delivery oc-

curred at term. At birth, scalp hair was almost absent, and fingernails were barely visible. Nails progressively started to grow, but very slowly. According to the boy's father, the deciduous teeth erupted normally but were widely spaced and abnormal in shape. Hyperkeratosis of palms and soles appeared at around age 3 years. He never complained of unusual discomfort from heat. Growth and mental development were normal. At age 5 years, his physical measurements were within the normal limits. He had dry skin all over his body, with a diffuse follicular hyperkeratosis, dry and thin hair, sparse eyebrows, widely spaced and malformed deciduous teeth, a smooth tongue with very few fungiform and filiform papillae, dystrophic



fingernails and toenails, and recurrent folliculitis. Palmo-plantar erythema with mild hyperkeratosis and hyperhidrosis of palms and soles was found. His uncle (IV: 4 in fig. 1B and case 2 in family 2 in table 1) was examined at age 46 years. He also presented with dry hair, severe hypodontia, smooth tongue, onychodysplasia, palmar erythema, and mild keratoderma and hyperhidrosis of palms and soles.

In the third family,²⁶ one patient was reinvestigated (V: 3 in fig. 1C). She presented with dry skin, thin and dry hair that never grew normally, almost absent permanent teeth, a smooth tongue with very few fungiform and filiform papillae, thin and concave fingernails, severely dystrophic toenails, and palmar erythema. She complains from time to time of an excessive palmo-plantar sudation. Electronic microscopic examination of the hair, which had already been performed and was reported elsewhere,²⁶ showed thin hair with longitudinal depressions. At the bottom of these depressions, cuticular cells presenting a

longitudinally striated aspect were found, as if the keratin fibrils were observable under the cell membrane.

Molecular genetics.—After informed consent was obtained from the adult subjects and the parents of the children, peripheral blood samples from 18 subjects were collected, and genomic DNA was isolated from lymphocytes by use of standard methods. Because all the patients included in the study were born to consanguineous marriages, we assumed identity by descent and performed homozygosity mapping²⁸ on pooled DNA by a first genomewide screen of family 1.

A set of 382 highly polymorphic fluorescently labeled markers on chromosomes 1–22 (ABI PRISM Linkage Map Set, version 2.0 [PE Biosystems]) were chosen from the Génethon linkage map,²⁹ as described elsewhere.³⁰ The first set of markers was not sufficiently informative, since several loci were homozygous in all affected and unaffected individuals. Genotyping of additional markers around these loci allowed the identification of a locus ho-

Table 1. Review of Clinical Features of Our Patients

Clinical Feature	Family 1			Family 2		Family 3
	Case 1	Case 2	Case 3	Case 1	Case 2	Case 1
Sex	M	M	M	M	M	F
Absence of hair at birth	±	—	+	+	?	—
Congenital absence of nails	±	—	+	+	?	—
Dystrophic fingernails	+	+	+	+	—	+
Dystrophic toenails	+	±	+	+	—	+
Erythematous atrophic patches on the face	—	—	—	—	+	—
Palmoplantar keratoderma	+	+	+	±	+	—
Hyperhidrosis	+	+	+	+	+	+
Oligodontia	+	+	+	+	+	+
Reduced papillae of the tongue and/or smooth tongue	+	+	+	+	+	+
Partial alopecia (male pattern)	—	—	—	—	+	—
Sparse eyelashes and/or eyebrows	—	—	—	+	—	—
Sparse axillary and/or pubic hair	—	—	—	?	—	?
Abnormal hair pattern	±	±	±	—	+	+
Diffuse follicular hyperkeratosis	+	—	+	+	—	±
Dry skin	+	+	+	+	+	+

NOTE.— ± means “more or less.”

mozygous by descent at marker *D2S301* (University of California–Santa Cruz [UCSC] accession number AFM214YE1) at chromosome 2q35. To refine the size of the putative shared homozygous region, 14 additional microsatellite markers (*D2S2319*, *D2S128*, *D2S2361*, *D2S164*, *D2S434*, *D2S2722*, *D2S163*, *D2S344*, *D2S377*, *D2S2372*, *D2S339*, *D2S130*, *D2S2228*, and *D2S353*) chosen from databases (UCSC and Ensembl) and four highly heterozygous SNPs (*rs16853834*, *rs2303400*, *rs4674305*, and *rs1109867*) chosen from dbSNP were genotyped. Haplotype analyses and recombination events allowed the restriction of the homozygous candidate region to a 9-Mb interval between markers *rs16853834* and *D2S353* (UCSC Genome Browser accession number AFM296VH9) at chromosome 2q35-2q36.2 (fig. 1A). In family 2, we found linkage to the same region (fig. 1B). Thus, we performed parametric multipoint linkage analysis with Easylinkage version 5.0, using a recessive model and complete penetrance. Cumulative linkage analyses of the two families showed a significant positive LOD score of 5.74 on chromosome 2q35-36.1 at $\theta = 0.001$.

There were ~70 genes in the homozygous candidate region, several of which were known to be expressed in skin and teguments and could therefore be implicated in the disease (GenAtlas and UCSC Genome Browser). The coding region, as well as exon-intron boundaries and 5' and 3' UTRs of six of those genes (*EPHA4* [GenBank accession number NM_004438], *INHA* [GenBank accession number NM_002191], *DNAJB2* [GenBank accession number NM_006736], *VIL1* [GenBank accession number NM_007127], *WNT6* [GenBank accession number NM_006522], and *WNT10A* [GenBank accession number NM_025216.2]), were amplified by PCR from genomic DNA from one patient per family (families 1 and 2) with the use of intronic primers (sequence of primers and amplification conditions available on request). Primers were designed using Primer3 and OLIGOS v.9.3 and were

checked for specificity by use of BLAST, and DNA sequences were obtained from UCSC or GenBank databases. Mutation screening was performed by fluorescent bidirectional sequencing of genomic amplified PCR products on an ABI3130 automated sequencer (Applied Biosystems).

We identified a homozygous c697G→T transversion in exon 3 of *WNT10A* in each affected patient tested from the two families (fig. 2). The mutation cosegregates with the phenotype in families 1 and 2, comprising a total of five patients, and was not found in 200 control chro-




Figure 2. *WNT10A* mutation associated with odonto-onycho-dermal dysplasia. Electropherograms of *WNT10A* exon 3 in a control individual (top), one parent (middle), and an affected patient (bottom). DNA sequence analysis revealed a homozygous G→T substitution at nucleotide 697 (from the translation start site) in the affected patient, causing the Glu233X mutation. The parent is heterozygous with respect to the Glu233X mutation.

mosomes tested by endonuclease-restriction digestion with the use of *AvaI* (BioLabs), thus making a polymorphism unlikely. At the protein level, the c.697G→T mutation causes the substitution of a negatively charged glutamic acid residue for a stop codon (p.Glu233X), which results in a premature, truncated protein of 232 aa, instead of 417 aa, that lacks the cysteine-rich prosite pattern PS50311, which is highly conserved in the WNT superfamily. Screening of the *WNT10A* gene for mutations in one patient of the third family revealed the same mutation. The three families investigated here denied any relationship with one another. Nevertheless, all affected individuals of the three families showed the identical haplotype of 1-C-3-3-4-T-1 for markers *D2S2361*, *rs2303400*, *D2S301*, *D2S164*, *D2S434*, *rs4674305*, and *D2S2722*, respectively, with a probable neomutation at *D2S434* for patient V:3 of family 3, who is heterozygous 3/4 instead of 4/4 for this marker. These data, in addition to the fact that the three families belong to the same religious community, suggest that there is an ancestral haplotype from marker *D2S2361* to *WNT10A* included, and that the mutation found in *WNT10A* in the patients of the three families comes from a common ancestor.

Studies on mice showed that *Wnt10a* is up-regulated in skin of developing embryos,⁷ in placodes at the onset of follicle morphogenesis, and in postnatal hair follicles beginning a new cycle of hair growth.⁸ Also, *Wnt10a* is expressed in the epithelium when tooth morphogenesis begins.³¹ The involvement of *Wnt10a* through the Wnt/ β -catenin signaling pathway has previously been suggested in the apical ectodermal ridge in chick limb formation³² and in tooth development in mice.¹⁵ In the absence of WNT signaling, β -catenin is phosphorylated and degraded. Since β -catenin is required for localized expression of regulatory genes, initiation of hair- and tooth-follicle placode formation,¹⁰ and tooth renewal as well as tooth aspect,¹¹ its absence results in the disruption of different ectodermal organs. In humans, the present observation shows that *WNT10A* plays a crucial role in the development and/or regulation of the ectodermal derivatives. The mutation found in *WNT10A* might affect β -catenin in the same way as the absence of WNT signaling does, given the fact that the resulting truncated protein might be nonfunctional.

Interestingly, Millar et al.³³ demonstrated that *Wnt3* is expressed in developing and mature hair follicles and that its overexpression in transgenic mouse skin causes a short-hair phenotype, by altered differentiation of hair-shaft precursor cells, and cyclical balding, by hair-shaft structural defects, and is associated with an abnormal profile of protein expression in the hair shaft. Our patients had hair that grew very slowly and never became very long (except for individual IV:4 in fig. 1A). It is likely that the abnormal structure of the hair observed by light and scanning electron microscopy^{27,26} and the short-hair phenotype are caused by defects in the proliferation or differentiation of hair-shaft precursor cells. But, in contrast to

Wnt3, where overexpression led to this phenotype, truncated or loss of functional *WNT10A* altered normal hair growth, suggesting that the two *WNT* genes might use different pathways.

Recently, Liu et al.³⁴ demonstrated that Wnt- β -catenin signaling is activated in developing fungiform placodes and taste-bud cells and plays a central role in the formation of tongue papillae. These authors found that *Wnt10b*, another subtype of the *Wnt10* protein that shares 59% sequence identity with *Wnt10a*,^{7,35} was specifically elevated in developing fungiform placodes. They also found that any alteration of β -catenin leads to an absence or marked reduction in fungiform papilla morphogenesis. In the families we studied, fungiform and filiform papillae were very few. Thus, although there are at present no in vivo data confirming a role of *WNT10A* in the tongue, it is tempting to speculate about the role of the latter in tongue papillae, which could explain the presence of a smooth tongue in our patients.

On the other hand, other studies found that mutations in the genes encoding R-spondin 4 and R-spondin 1, characterized as a small family of growth factors, are responsible for anonychia and palmoplantar hyperkeratosis, respectively.^{36,37} The furin-like repeats encoded by exons 2 and 3 of the R-spondin family of protein genes are believed to be required for activation and stabilization of β -catenin. Thus, it is not surprising that a mutation in the *WNT10A* gene leads to abnormal nails and palmoplantar keratoderma by perturbing keratinocyte proliferation and differentiation, as seen in our patients.

Overall, the phenotypic expression of the *WNT10A* mutation described in the three families was mild when compared with the malformations caused by mutations in other members of WNT family.^{16,18} Moreover, clinical inter- and intrafamilial heterogeneity was noted. Indeed, the patients from family 3 were not given diagnoses of having odonto-onycho-dermal dysplasia because they showed an absence of keratoderma and had normal facial skin without erythema, atrophy, or telangiectasia.²⁶ Nevertheless, the overlapping features with families 1 and 2 led us to sequence *WNT10A*. The same mutation was found. A clinical follow-up with one patient of family 3 showed that, in addition to the details described elsewhere,²⁶ she had a smooth tongue and short hair that she rarely cut. Still, neither follicular hyperkeratosis nor keratoderma was noted. The mild phenotype and the clinical heterogeneity between patients suggest that there are alternate pathways and/or environmental factors that may contribute to compensate for the abnormal *WNT10A* protein.

In conclusion, we report, for the first time to our knowledge, a gene of the WNT family that when mutated causes ED, confirming the importance of the Wnt pathway for the normal formation and regeneration of hair, teeth, skin, and nails in humans. Although the name of the present pathology is odonto-onycho-dermal dysplasia, the abnormal hair indicates that it is more of a tricho-odonto-onycho-dermal dysplasia. The latter could fit in the cell-

cell communication and signaling group of Lamartine's classification,⁶ showing that an altered epithelial-mesenchymal interaction seems to be one of the most important mechanisms in the pathogenesis of EDs. *WNT10A*, but also other *WNT* genes such as *WNT6* that are coexpressed with *WNT10A*³⁵ or *WNT10B* and that are known to be expressed in teeth,^{31,38} could be considered good candidates for conditions similar to the one we report here, such as that described by Fried,³⁹ tricho-odonto-onycho-dermal dysplasia with cysts of the eyelids syndrome (called "Schopf-Schulz-Passarge syndrome" [MIM 224750])⁴⁰; the entity reported by Salamon et al.,⁴¹ characterized by hypodontia, sparse and dry hair, dystrophic nails, dry skin, and hyperkeratotic papules on the hands; and the one described by Pinheiro et al.,⁴² characterized by hypodontia, onychodysplasia, mild palmoplantar keratosis, and sparse hair, eyelashes, and eyebrows. Reports of further families with ED with *WNT* mutations will allow better characterization of the molecular pathways controlling differentiation and proliferation of ectodermal organs.

Acknowledgments

We are grateful to the patients and their parents for their participation in this project. We also thank Drs. Susan Cure, Geraldine Guasch, and Simon Khalifé for their support. This work was supported by grants from the Saint-Joseph University Research Council.

Web Resources

Accession numbers and URLs for data presented herein are as follows:

BLAST, <http://www.ncbi.nlm.nih.gov/BLAST/>
 dbSNP, <http://www.ncbi.nlm.nih.gov/SNP>
 Ensembl, <http://www.ensembl.org/index.html>
 Genatlas, <http://www.genatlas.org>
 GenBank, <http://www.ncbi.nlm.nih.gov/Genbank/> (for *EPHA4* [accession number NM_004438], *INHA* [accession number NM_002191], *DNAJB2* [accession number NM_006736], *VIL1* [accession number NM_007127], *WNT6* [accession number NM_006522], and *WNT10A* [accession number NM_025216.2])
 Laboratory of Randall Moon, <http://faculty.washington.edu/rtmoon>
 Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for odonto-onycho-dermal dysplasia and Schopf-Schulz-Passarge syndrome)
 Primer3, http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi
 UCSC Genome Browser, <http://www.genome.ucsc.edu> (for *D2S301* [accession number AFM214YE1] and *D2S353* [accession number AFM296VH9])
 Wnt Homepage, <http://www-leland.stanford.edu/~rnusse/wntwindow.html>

References

1. Itin PH, Fistarol SK (2004) Ectodermal dysplasias. *Am J Med Genet C Semin Med Genet* 131:45–51
2. Freire-Maia N, Pinheiro M (1984) Ectodermal dysplasias: a

- clinical and genetic study. Alan R. Liss, New York, pp 172–173
3. Freire-Maia N, Pinheiro M (1988) Selected conditions with ectodermal dysplasia. *Birth Defects Orig Artic Ser* 24:109–121
4. Pinheiro M, Freire-Maia N (1994) Ectodermal dysplasias: a clinical classification and a causal review. *Am J Med Genet* 53:153–162
5. Priolo M, Lagana C (2001) Ectodermal dysplasias: a new clinical-genetic classification. *J Med Genet* 38:579–585
6. Lamartine J (2003) Towards a new classification of ectodermal dysplasias. *Clin Exp Dermatol* 28:351–355
7. Wang J, Shackleford GM (1996) Murine Wnt10a and Wnt10b: cloning and expression in developing limbs, face and skin of embryos and in adults. *Oncogene* 13:1537–1544
8. Reddy S, Andl T, Bagasra A, Lu MM, Epstein DJ, Morrisey EE, Millar SE (2001) Characterization of Wnt gene expression in developing and postnatal hair follicles and identification of Wnt5a as a target of Sonic hedgehog in hair follicles morphogenesis. *Mech Dev* 107:69–82
9. Millar SE, Willert K, Salinas PC, Roelink H, Nusse R, Sussman DJ, Barsh GS (1999) WNT signaling in the control of hair growth and structure. *Dev Biol* 207:133–149
10. Andl T, Reddy ST, Gaddapara T, Millar SE (2002) WNT signals are required for the initiation of hair follicle development. *Dev Cell* 2:643–653
11. Jarvinen E, Salazar-Ciudad I, Birchmeier W, Taketo MM, Jernvall J, Thesleff I (2006) Continuous tooth generation in mouse is induced by activated epithelial wnt/beta-catenin signaling. *Proc Natl Acad Sci USA* 103:18627–18632
12. Peifer M, Polakis P (2000) Wnt signaling in oncogenesis and embryogenesis—a look outside the nucleus. *Science* 287:1606–1609
13. Cadigan KM, Nusse R (1997) Wnt signaling: a common theme in animal development. *Genes Dev* 11:3286–3305
14. Wodarz A, Nusse R (1998) Mechanisms of Wnt signaling in development. *Annu Rev Cell Dev Biol* 14:59–88
15. Kratochwil K, Galceran J, Tontsch S, Roth W, Grosschedl R (2002) FGF4, a direct target of LEF1 and Wnt signaling, can rescue the arrest of tooth organogenesis in *Lef1*^{-/-} mice. *Genes Dev* 16:3173–3185
16. Niemann S, Zhao C, Pascu F, Stahl U, Aulepp U, Niswander L, Weber JL, Muller U (2004) Homozygous *WNT3* mutation causes tetra-amelia in a large consanguineous family. *Am J Hum Genet* 74:558–563
17. Jordan BK, Shen JH, Olaso R, Ingraham H A, Vilain E (2003) Wnt4 overexpression disrupts normal testicular vasculature and inhibits testosterone synthesis by repressing steroidogenic factor 1/ β -catenin synergy. *Proc Natl Acad Sci USA* 100:10866–10871
18. Woods CG, Stricker S, Seemann P, Stern R, Cox J, Sherridan E, Roberts E, Springell K, Scott S, Karbani G, et al (2006) Mutations in *WNT7A* cause a range of limb malformations, including Fuhrmann syndrome and Al-Awadi/Raas-Rothschild/Schinzel phocomelia syndrome. *Am J Hum Genet* 79:402–408
19. Robitaille J, MacDonald ML, Kaykas A, Sheldahl LC, Zeisler J, Dube MP, Zhang LH, Singaraja RR, Guernsey DL, Zheng B, et al (2002) Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy. *Nat Genet* 32:326–330
20. Xu Q, Wang Y, Dabdoub A, Smallwood PM, Williams J, Woods C, Kelly MW, Jiang L, Tasman W, Zhang K, et al (2004) Vascular development in the retina and inner ear: control by

- Norrin and Frizzled-4, a high-affinity ligand-receptor pair. *Cell* 116:883–895
21. Lammi L, Arte S, Somer M, Jarvinen H, Lahermo P, Thesleff I, Pirinen S, Nieminen P (2004) Mutations in *AXIN2* cause familial tooth agenesis and predispose to colorectal cancer. *Am J Hum Genet* 74:1043–1050
 22. Biason-Lauber A, Konrad D, Navratil F, Schoenle (2004) A *WNT4* mutation associated with Mullerian-duct regression and virilization in a 46, XX woman. *N Engl J Med* 351:792–798
 23. Fadhil M, Ghabra TA, Deeb M, Der Kaloustian VM (1983) Odontoonychodermal dysplasia: a previously apparently undescribed ectodermal dysplasia. *Am J Med Genet* 14:335–346
 24. Zirbel GM, Ruttum MS, Esterly NB (1995) Odonto-onychodermal dysplasia. *Br J Dermatol* 133:797–800
 25. Arnold WP, Merks MA W, Steijlen PM (1995) Variant of odontoonychodermal dysplasia? *Am J Med Genet* 59:242–244
 26. Mégarbané A, Noujeim Z, Fabre M, Der Kaloustian VM (1998) New form of hidrotic ectodermal dysplasia in a Lebanese family. *Am J Med Genet* 75:196–199
 27. Mégarbané H, Haddad M, Delague V, Renoux J, Bohem N, Megarbane A (2004) Further delineation of the odonto-onychodermal dysplasia syndrome. *Am J Med Genet A* 129:193–197
 28. Lander E, Botstein D (1987) Homozygosity mapping: a way to map human recessive traits with the DNA of inbred children. *Science* 236:1567–1570
 29. Dib C, Faure S, Fizames C, Samson D, Drouot N, Vignal A, Millasseau P, Marc S, Hazan J, Seboun E, et al (1996) A comprehensive genetic map of the human genome based on 5,264 microsatellites. *Nature* 380:152–154
 30. Delague V, Bareil C, Tuffery S, Bouvagnet P, Chouery E, Koussa S, Maisonobe T, Loiselet J, Mégarbané A, Claustres M (2000) Mapping of a new locus for autosomal recessive demyelinating Charcot-Marie-Tooth disease to 19q13.1-13.3 in a large consanguineous Lebanese family: exclusion of *MAG* as a candidate gene. *Am J Hum Genet* 67:236–243
 31. Dassule HR, McMahon AP (1998) Analysis of epithelial-mesenchymal interactions in the initial morphogenesis of the mammalian tooth. *Dev Biol* 202:215–227
 32. Narita T, Sasaoka S, Udagawa K, Ohshima T, Wada N, Nishimatsu S, Takada S, Nohno T (2005) *Wnt10a* is involved in AER formation during chick limb development. *Dev Dyn* 233:282–287
 33. Millar SE, Willert K, Salinas PC, Roelink H, Nusse R, Sussman DJ, Barsh GS (1999) WNT signaling in the control of hair growth and structure. *Dev Biol* 207:133–149
 34. Liu F, Thirumangalathu S, Gallant N, Yang SH, Stoick-Cooper C, Reddy S, Andl T, Taketo M, Dlugosz A, Moon R, et al (2007) Wnt- β -catenin signaling initiates taste papilla development. *Nat Genet* 39:106–112
 35. Kirikoshi H, Sekihara H, Katoh M (2001) WNT10A and WNT6, clustered in human chromosome 2q35 region with head-to-tail manner, are strongly coexpressed in SW480 cells. *Biochem Biophys Res Commun* 283:798–805
 36. Blyden DC, Ishii Y, O'Toole EA, Unsworth HC, Teh MT, Ruschendorf F, Sinclair C, Hopsu-Havu VK, Tidman N, Moss C, et al (2006) The gene encoding R-spondin 4 (*RSPO4*), a secreted protein implicated in Wnt signaling, is mutated in inherited anonychia. *Nat Genet* 38:1245–1247
 37. Parma P, Radi O, Vidal V, Chaboissier MC, Dellambra E, Valentini S, Guerra L, Schedl A, Camerino G (2006) R-spondin1 is essential in sex determination, skin differentiation and malignancy. *Nat Genet* 38:1304–1309
 38. Sarkar L, Sharpe PT (1999) Expression of wnt signalling pathway genes during tooth development. *Mech Dev* 85:197–200
 39. Fried K (1977) Autosomal recessive hydrotic ectodermal dysplasia. *J Med Genet* 14:137–139
 40. Schopf E, Schulz HJ, Passarge E (1971) Syndrome of cystic eyelids, palmo-plantar keratosis, hypodontia and hypotrichosis as a possible autosomal recessive trait. *Birth Defects Orig Artic Ser* 7:219–221
 41. Salamon T, Lazovic O, Stenek S (1972) Hair changes in ectodermal dysplasia associated with multiple anomalies. *Hautarzt* 23:441–445
 42. Pinheiro M, Pereira LC, Freire-Maia N (1981) A previously undescribed condition: tricho-odonto-onychodermal syndrome. A review of the tricho-odonto-onychia subgroup of ectodermal dysplasias. *Br J Dermatol* 105:371–382