



Published in final edited form as:

Insect Biochem Mol Biol. 2007 June ; 37(6): 570–578.

Specific Transcriptional Responses to Juvenile Hormone and Ecdysone in *Drosophila*

Robert B. Beckstead, Geanette Lam, and Carl S. Thummel*

Department of Human Genetics, University of Utah School of Medicine, 15 North 2030 East Room 2100, Salt Lake City, UT 84112-5330, USA

Abstract

Previous studies have shown that ecdysone (E), and its immediate downstream product 20-hydroxyecdysone (20E), can have different biological functions in insects, suggesting that E acts as a distinct hormone. Here we use *Drosophila* larval organ culture in combination with microarray technology to identify genes that are transcriptionally regulated by E, but which show little or no response to 20E. These genes are coordinately expressed for a brief temporal interval at the onset of metamorphosis, suggesting that E acts together with 20E to direct puparium formation. We also show that *E74B*, *pepck*, and *CG14949* can be induced by juvenile hormone III (JH III) in organ culture, and that *CG14949* can be induced by JH independently of protein synthesis. In contrast, *E74A* and *E75A* show no response to JH in this system. These studies demonstrate that larval organ culture can be used to identify *Drosophila* genes that are regulated by hormones other than 20E, and provide a basis for studying crosstalk between multiple hormone signaling pathways.

Keywords

ecdysone; juvenile hormone; metamorphosis; gene regulation

1. Introduction

Like all holometabolous insects, *Drosophila melanogaster* has two primary lipophilic hormones, the steroid hormone 20-hydroxyecdysone (20E) and the sesquiterpenoid juvenile hormone (JH) (Riddiford, 1993). 20E arises from its immediate precursor, ecdysone (E), which is released in bursts from the *Drosophila* prothoracic gland in response to neuropeptide signaling. Pulses of 20E during the first and second larval instars trigger molting of the cuticle while a high titer pulse of 20E at the end of the third instar triggers puparium formation, initiating metamorphosis and the prepupal stage of development. 20E exerts its effects by binding to a heterodimer of two nuclear receptors, the Ecdysone Receptor (EcR) and Ultraspiracle (USP) (Koelle et al., 1991; Thomas et al., 1993; Yao et al., 1993). This activated receptor complex reprograms gene expression in target cells, directing the appropriate stage-specific developmental programs during the fly life cycle.

It has long been known that E is modified by peripheral tissues into 20E, although only recently has the 20-monooxygenase been discovered that catalyzes this reaction (Petryk et al., 2003). E is widely believed to be inactive, with little or no effect on classic EcR-mediated responses

*Corresponding author. Tel.: +801-581-2937; fax: +801-581-5374. E-mail address: carl.thummel@genetics.utah.edu (C.S. Thummel)

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

in *Drosophila* (Richards, 1978; Baker et al., 2000; Hu et al., 2003). Several studies in other insects, however, have invoked developmental functions for E, including effects on DNA synthesis in cultured wing discs of *Galleria mellonella* and molting in *Chironomus tentans* (Oberlander, 1972; Clever et al., 1973). More recently, the high E titer during early *Manduca* metamorphosis has been shown to direct neuroblast proliferation and optic lobe development, establishing a critical role for this hormone that is distinct from functions regulated by 20E (Champlin and Truman, 1998). In addition, *Aedes aegypti* EcR, unlike its *Drosophila* counterpart, can be efficiently activated by both E and 20E (Wang et al., 2000). These observations raise the interesting possibility that E could act as a hormone, much like 20E, although there is no molecular data to support such a role for this ecdysteroid.

Studies in Coleoptera and Lepidoptera have shown that JH defines the nature of the biological responses triggered by 20E during development, leading to the status quo model for JH action. In the presence of high JH titers, pulses of 20E lead to larval-larval molts whereas, in the absence of JH, 20E initiates maturational responses and metamorphosis (reviewed by Riddiford, 1996; Gilbert et al., 2000). In contrast, much less is known about roles for JH in *Drosophila* pre-adult development. Ectopic application of a JH analog during prepupal stages results in lethality with defects in abdominal, nervous system, and muscular development (Riddiford and Ashburner, 1991; Restifo and Wilson, 1998). The defects in abdominal development are accompanied by changes in cuticle gene expression that are mediated by sustained expression of the Broad-Complex transcription factors (Zhou and Riddiford, 2002). No normal developmental functions, however, have been shown for JH during *Drosophila* larval or prepupal stages. Similarly, relatively little is known about its molecular mechanisms of action. Two candidate JH receptors have been proposed in the literature, USP and Methoprene-tolerant (Met) (Jones and Sharp, 1997; Ashok et al., 1998; Miura et al., 2005), although biological evidence for these acting as JH receptors is lacking (Restifo and Wilson, 1998; Gilbert et al., 2000; King-Jones and Thummel, 2005; Palanker et al., 2006). Rather, the identification of several genes that are transcriptionally controlled by JH provides a foundation for understanding its molecular mechanism of action (reviewed by Berger and Dubrovsky, 2005).

In this study we exploit the use of larval organ culture and microarray technology to identify genes regulated by E or JH. We show that E can regulate genes in *Drosophila* that are distinct from those controlled by 20E. These genes are coordinately induced or repressed at puparium formation, suggesting that E contributes to the onset of metamorphosis in parallel with 20E. We also identify three genes that are induced by JH in larval organ culture and characterize their regulation by JH, as well as how this response can be modified by 20E. This study provides a basis for understanding the regulatory functions of E and JH in *Drosophila*.

2. Experimental procedures

2.1 Organ culture

For the E microarray study, third instar larvae at approximately 10 hours prior to puparium formation were staged by the addition of 0.05% bromophenol blue to the food and selecting animals with partial blue guts, as described previously (Andres and Thummel, 1994). Eight animals were dissected in a nine-well glass dish (Corning) and cultured in approximately 0.1 ml oxygenated Schneiders *Drosophila* Medium (Invitrogen) at 25°C for the times indicated below. Each well contains the integument and all internal organs from the dissected animals. Cultures were incubated under a constant flow of oxygen in a styrofoam box. Following an initial incubation of 1 hour, the medium was removed and replaced with either fresh Schneiders *Drosophila* Medium (no hormone) or medium plus 5×10^{-6} M ecdysone (E) (Sigma-Aldrich) for 6 hours at 25°C. This is the same concentration that has been used for studies of 20E responses in cultured larval organs, which is two-fold higher than the amount required for a

50% maximal early puff response in the salivary gland polytene chromosomes and comparable to the peak 20E titers seen *in vivo* (Ashburner, 1973; Karim and Thummel, 1991). The commercial source of E used in our studies was found to be 93% pure by RP-HPLC, as determined by Dr. J. Warren (Univ. of N. Carolina, Chapel Hill). We used highly purified E, kindly provided by Dr. Warren, for all validation northern blot hybridization studies reported here. Similarly, purified (10R)-juvenile hormone III was used for northern blot studies along with (10S)-JH III as a negative control, both kindly provided by Dr. W. Goodman (Univ. of Wisconsin, Madison). Organs were collected and RNA was extracted as described below. Glass culture plates were treated with a 20% solution of PEG 20,000, rinsed at least six times, and autoclaved, to prevent JH binding to the glass. All experiments were done in triplicate and harvested separately for array analysis. For northern blot analysis, third instar larvae were synchronized at the second-to-third instar larval molt and aged for either 28 or 38 hours (approx. -20 or -10 hours relative to puparium formation, respectively). Eight animals were processed as above for organ culture and treated with no hormone, 5×10^{-6} M (10R)-juvenile hormone III, 5×10^{-6} M 20-hydroxyecdysone, 5×10^{-6} M ecdysone, 8.5×10^{-5} M cycloheximide (Sigma-Aldrich), 5×10^{-6} M (10S)-juvenile hormone, 5×10^{-6} M farnesol (Sigma-Aldrich), or 5×10^{-6} M methyl linoleate (Sigma-Aldrich). JH and 20E were added simultaneously in all cases where both hormones were used in combination.

2.2 Microarray and cluster analysis

Total RNA was isolated from organ cultures using TriPure (Roche) and purified on RNeasy columns (Qiagen). All samples were prepared independently in triplicate to facilitate statistical analysis. Probe labeling, hybridization to Affymetrix GeneChip *Drosophila*[®] Genome 1.0 Arrays (Affymetrix), and scanning were performed by the University of Maryland Biotechnology Institute Microarray Core Facility. Raw data was normalized using dChip1.2 and replicate arrays were combined for expression analysis (Li and Wong, 2001). Expression analysis using dChip1.2 was based on the PM-only model, and expression-level differences between control and experimental were analyzed by assigning a fold-change value and filtering for outliers. Data sets were analyzed using Microsoft Access selecting those genes with a change in expression level of ≥ 1.5 -fold. Comparisons made between the E and 20E data sets were performed using Microsoft Access. The 20E gene set for this comparison was derived from both the 20E-regulated genes and 20E primary-response genes described by Beckstead et al. (2005). Cluster analysis was performed using dChip1.2. Comparisons were made between our microarray datasets and previously published microarray datasets using Microsoft Access.

2.3 Northern blot hybridizations

Total RNA was isolated from cultured larval organs using TriPure (Roche), fractionated by formaldehyde gel electrophoresis, transferred to nylon membranes, and probed with radioactively labeled probes, as described (Andres et al., 1993). To facilitate direct comparison, blots were probed, washed, and exposed at the same time. Probes were generated by PCR from genomic DNA using the following pairs of primers: *dro5F* AACCGACAACATGCAGATCA, *dro5R* TTCCGGCAATTTAGAGCAAC, *pepckF* ATGAACGCAAAGTCCTCGAC, *pepckR* GGCTGCTCCAATGAAGACTC, *CG7906F* AGTTGACTCAGCATAACGG, *CG7906-3'* TGGCACAAATGGACAGTG, *CG7924-5'* AGGTTGTGCCAAATGCAG, *CG7924R* TCGGCACACATCCTCATC, *nmdmcF* CTCCGACCAGCTTAAACTGG, *nmdmcR* ACTAGCACACGAGCTCAAGG, *CG11956F* TACATATGGTGGGAGTTCCAGGAG, *CG11956R* CAAGCTTATCCTGCAAGGACCTAC, *CG14949F* ACCAGTTTTGAGAGCCAGGA, *CG14949R* CTAATCCTCGGGTTCCGTCT, *kr-h1F* GTGCGAGAAGACATTCATCCAGTC, *kr-h1R* TTGAAGTGCTCGATCAGCACGGTG, *blackF* CACCAGCACCAGGAACAGTA, *blackR* ACTCGGGCGAGTGTCTGTAG.

3. Results

3.1 Identification of genes regulated by E

Organs were dissected from third instar larvae staged at approximately 10 hours before pupariation, and cultured in the presence of either no hormone or 5×10^{-6} M ecdysone (E), following protocols used for earlier studies of 20E-regulated transcription (Ashburner, 1972; Beckstead et al., 2005). RNA was extracted from these tissues, labeled, and hybridized to Affymetrix *Drosophila* 1.0 Genome Arrays. Data was analyzed using dChip1.2 (Li and Wong, 2001). For statistical comparisons, RNA from three independent organ culture experiments was used for each data point in the microarray study. Comparison of the control and E-treated datasets led to the identification of 267 E-regulated genes that exhibited a change in expression level of 1.5-fold or greater. Given that E can be converted to 20E by *Drosophila* larval organs (Somme-Martin et al., 1988; Petryk et al., 2003), we compared this data set to the 376 genes that exhibit a change in expression level of 1.5-fold or greater when treated by either 20E alone or 20E plus cycloheximide (Beckstead et al., 2005). This resulted in the identification of 55 genes that appear to be responding specifically to E and not to 20E (Fig. 1A; Table 1).

If these 55 genes correspond to part of an E-specific genetic program then we might expect to see them coordinately regulated when E titers peak during development. Such a time occurs late in the third instar, when an increase in E titer precedes the high titer pulse of 20E that triggers puparium formation (Parvy et al., 2005; Warren et al., 2006). To examine the temporal profiles of E-regulated genes at this time, we used data from an earlier microarray study that determined transcript levels in *w¹¹¹⁸* animals staged at -18, -4, 0, 2, 4, 6, 8, 10, and 12 hours relative to pupariation (Beckstead et al., 2005). Interestingly, this study revealed that each of the 55 E-regulated genes is expressed for a brief temporal window at the onset of metamorphosis (Fig. 1B). The majority of genes are induced at puparium formation and down-regulated in 4-6 hr prepupae. A second set of genes are specifically induced in mid-prepupae, immediately preceding the late prepupal 20E peak that occurred at 10 hours after pupariation in these animals (Beckstead et al., 2005). The remaining genes are down-regulated in late third instar larvae. Taken together, these observations suggest that E activates a genetic program that operates in parallel to that of 20E at the onset of metamorphosis.

Northern blot hybridizations were used to validate the microarray results and confirm the specificity of these genes for responding to E as opposed to 20E. Organs were dissected from larvae staged at either 10 or 20 hours prior to puparium formation, and treated with no hormone (control), E, or 20E. RNA was then extracted from these tissues and analyzed on northern blots using probes to detect several E-regulated genes (Fig. 2, lanes 1,4,5,6,9,10). As a control, we examined the expression of two well characterized 20E primary-response transcripts, *E74A* and *E75A*, both of which are induced by 20E at 10 hours before pupariation, as expected (Segraves and Hogness, 1990; Karim and Thummel, 1991) (Fig. 2, lanes 6,9). A similar response is seen with E, albeit somewhat weaker, consistent with the expectation that E can be converted to 20E by the cultured larval organs (Fig. 2, lanes 6,10). Interestingly, this response is only seen in -10 hour animals, indicating that the -20 hour animals have not yet gained competency to respond to 20E, consistent with earlier work (Richards et al., 1999). We next examined the regulation of *CG7924*, *nmdmc*, *dro5*, *CG7906* and *CG12224*, five genes that display high levels of induction by E alone in our microarray study (Table 1). As predicted, these genes are induced by E but not by 20E (Fig. 2), with the exception of *CG12224* which does not respond to either E or 20E (data not shown). The observation that we can validate the E-specific regulation for four of these five genes indicates that most genes identified by our microarray study respond specifically to this hormone. We also examined two genes that show a higher level of induction by E than 20E in the microarray studies, *black* and *CG11956*. These genes show a very weak response to 20E and much more efficient induction by E, suggesting

that at least some of the genes in this class are also likely to represent specific responses to E (Fig. 2 and data not shown). As with *E74* and *E75A*, the E-regulated genes are only induced in organs cultured from -10 hour larvae, suggesting that they also require competence for their response to hormone.

3.2 Many E-regulated genes are also regulated by JH in combination with 20E

In the course of doing this study, we made the unexpected discovery that E-regulated genes can also respond to JH in the presence of 20E. As shown in Figure 2, the northern blots used for validating the E-regulated genes also contain samples of RNA isolated from third instar larval organs cultured with either JH III alone or JH III and 20E. We used JH III in these experiments because it has been shown to have potent effects on both genetic and biological responses in *Drosophila* (reviewed in Riddiford, 1993; Berger and Dubrovsky, 2005). *CG7924*, *nmdmc*, *black* and *CG11956* are all E-inducible genes that are up-regulated in organ cultures that are treated with JH and 20E, but not JH or 20E alone (Fig. 2, lanes 6-9). Moreover, the magnitude of this up-regulation is similar to that seen in response to E (Fig. 2, lane 10). Interestingly, *dro5* shows no significant response to JH and 20E, indicating that not all E-regulated genes are also controlled by JH and 20E. As with the E and 20E response, JH and 20E only has effects in organs that were harvested at 10 hour prior to pupariation (Fig. 2).

Studies in other insects have shown that ecdysone is capable of stimulating JH release from the corpora allata (Gu and Chow, 2003). This raises the possibility that the similar transcriptional responses we observe to treatment with E or 20E plus JH might be due to the production of JH by the ring gland in the larval organs cultured with E. According to this model, E would be converted to 20E in the organ culture system and then either E or 20E might stimulate the release of JH. To test this possibility, we dissected organs from third instar larvae staged at approximately 10 hours before puparium formation and removed the brain and its associated ring gland from one set of organs (Fig. 3, right panels). Given that JH in *Drosophila* arises from the corpora allata in the ring gland, we reasoned that the removal of the brain and ring gland would prevent any endogenous JH response in those tissues. These two sets of organs were cultured with either no hormone, E alone, or JH III and 20E, after which RNA was extracted and analyzed by northern blot hybridization for the expression of E-regulated genes (Fig. 3). *CG7906*, *CG7924*, *nmdmc*, *black*, and *CG11956* are all induced by E or JH plus 20E, as shown previously. We can see some variability in the level of *CG7924* transcription which is typical for this gene, which appears to show more stage-specificity than other E-regulated genes (data not shown). Removal of the brain and ring gland, however, had no significant effect on the response to either E alone or JH and 20E, indicating that the ability of E to upregulate this set of genes is not dependent upon the production of JH by the ring gland.

3.3 Identification of JH-regulated genes

In addition to the response to JH and 20E, we also noted that some genes, such as *E74B*, can be induced by JH III alone in organs dissected from either -20 hour or -10 hour third instar larvae (Fig. 1, lanes 1,2,6,7). This raised the possibility that we might be able to identify JH-regulated genes by microarray analysis of RNA isolated from cultured larval organs treated with JH. Conducting these studies using either methoprene or JH III, however, resulted in a high degree of variation between the JH-treated samples and a low change in expression level between the JH-treated samples and the controls, primarily below 1.5-fold (data not shown). Moreover, most responses could not be validated by northern blot hybridization. Two JH-regulated genes did, however, emerge from these studies: *pepck*, which encodes an enzyme involved in gluconeogenesis, and *CG14949*, which encodes a novel protein of unknown function. To characterize their regulation in more detail, organs were dissected from -20 hour third instar larvae and treated with either the protein synthesis inhibitor cycloheximide, JH III

alone, JH III and 20E, JH III and cycloheximide, or JH III and 20E and cycloheximide. Three negative controls were also included in this study – the biologically inactive JH III 10S isomer, farnesol, and methyl linoleate. RNA was extracted from these organs and analyzed by northern blot hybridization to detect *E74B*, *pepck*, *CG14949*, and *E75A* mRNA (Fig. 4). *E75A* was included in this study because it can be directly induced by JH III in *Drosophila* S2 tissue culture cells and is up-regulated by methoprene in ovaries from *apterous* mutant females (Dubrovsky et al., 2004). *E74B*, *pepck*, and *CG14949* are all up-regulated by treatment with JH alone (Fig. 4, compare lanes 1 and 3). Both *E74B* and *pepck* show approximately the same response to cycloheximide alone and cycloheximide plus JH, although the levels of *pepck* RNA may be increased ~2-fold by JH relative to the control (Fig. 4, lanes 2,5). In contrast, *CG14949* transcription is significantly increased upon treatment with JH and cycloheximide relative to the cycloheximide alone control, suggesting that it is a direct target for JH regulation. Interestingly, the JH induction of *E74B* and *pepck* is suppressed by adding 20E (Fig. 4, lanes 3,4) and, for *pepck*, this down-regulation also occurs in the presence of cycloheximide (Fig. 4, compare lanes 5 and 6). This observation suggests that the response of JH-regulated genes can be suppressed by 20E. Finally, *E75A* is only responsive to 20E, indicating that its JH regulation cannot be recapitulated under the conditions of larval organ culture. All four transcripts failed to respond to any of the negative control compounds, as expected (Fig. 4, lanes 7-9).

4. Discussion

This paper describes the identification of *Drosophila* genes regulated by E or JH using microarrays in combination with RNA isolated from cultured larval organs. We show that E regulates a set of genes that are distinct from those controlled by 20E. In addition, we identify three classes of genes that are regulated by JH: genes that are induced by JH alone, genes induced by JH in combination with 20E, and genes in which their JH induction is blocked by the addition of 20E. This study provides a foundation for future efforts aimed at identifying receptors for these hormones as well as a more complete understanding of the biological and molecular targets of 20E and JH signaling.

We identified 267 genes that change their expression in response to E in cultured larval organs. By comparing this list with those genes shown previously to be regulated by 20E (Beckstead et al., 2005) we identified three sets of genes, (1) those regulated by E but not 20E, (2) those that are regulated by either E or 20E, and (3) those specifically regulated by 20E (Fig. 1A). The set of 55 E-specific genes provides molecular evidence that E can act as a hormone in *Drosophila* independently of 20E. Our effort to validate this list by northern blot hybridization confirmed that four of five genes tested are regulated by E and not 20E – *CG7924*, *nmdmc*, *dro5*, and *CG7906* – suggesting that most of the genes in this list are E-specific. The predicted functions of these 55 genes include transcriptional control, enzymes, and transporters (Table 1). Interestingly, two of the validated E-inducible genes, *CG7906* and *CG7924*, map next to one another on the genome and encode related proteins that carry a Kazal-type serine protease inhibitor motif, although their functions remain undefined. The simple structure of these genes and their close physical association suggest that they are coordinately regulated by E, and provide a good basis for future efforts to identify E-response elements in the genome.

The second set of 212 genes that are regulated by E or 20E most likely include 20E-specific genes that are responding to 20E produced from the E added to the organ cultures. Interestingly, two genes that show a preferential response to E in the microarray studies, *black* and *CG11956*, display a significantly better induction by E than 20E on northern blots (Fig. 2), indicating that these 212 genes include additional E-regulated genes. Finally, 164 20E-regulated genes did not respond to the conversion of E to 20E in organ culture (Fig. 1A). It is possible that these genes require a greater concentration of 20E than was achieved by the conversion of E to 20E during the incubation period.

Upon examining the temporal profiles of the 55 E-regulated genes, we found that most are induced between 4 and 18 hours prior to puparium formation, along with a smaller set of genes that are down-regulated at this time (Fig 1B). The timing of response suggests that these genes may be responding to the increased E titer that precedes the pulse of 20E (Parvy et al., 2005; Warren et al., 2006), and that E and 20E act together to control the appropriate changes in gene expression that direct entry into metamorphosis. Recent genetic studies of the ecdysone biosynthetic pathway in *Drosophila* provide some support for this model. In their characterization of the *neverland* gene, Yoshiyama et al. (2006) reported that 7-dehydrocholesterol rescues the lethality of *neverland* RNAi more effectively than 20E, implying that one or more compounds upstream from 20E are essential for proper development. Similarly, preliminary studies indicate that *phantom* (*phm*), *disembodied* (*dib*), and *shadow* (*sad*) mutants, which block specific steps in E biosynthesis (Warren et al., 2002; Warren et al., 2004), are more effectively rescued by feeding E than 20E (H. Ono and M. O'Connor, personal communication). We are attempting to confirm and extend these studies by constructing transgenic lines that will provide temporal control over *dib* and *shade* (*shd*) inactivation. The *dib* gene encodes the enzyme that catalyzes the conversion of 2,22-dideoxyecdysone to 2-deoxyecdysone, while Shd directs the conversion of E to 20E. By inactivating *dib* or *shd* at specific times during development, we should be able to observe the consequences of removing both E and 20E (through inactivation of *dib*) or just removing 20E (through inactivation of *shd*). These tools should allow us to determine whether E provides functions distinct from those of 20E alone.

The ability of E to regulate a distinct set of genes from 20E raises that question of what receptor might mediate this response. A good candidate is the DHR38 nuclear receptor which can be activated by several ecdysteroids that are distinct from those that activate EcR, including E (Baker et al., 2003). Curiously, E does not manifest this effect through direct binding to the receptor and thus other factors must cooperate with DHR38 to transduce an E signal.

Previous studies have identified several JH-regulated genes in *Drosophila* (for a recent review, see Berger and Dubrovsky, 2005). These include *dRNaseZ*, *minidiscs*, *JHI-26*, and *E75A*, all of which are induced by JH III in *Drosophila* S2 tissue culture cells (Dubrovsky et al., 2000; Dubrovsky et al., 2002; Dubrovsky et al., 2004), as well as the *Drosophila* juvenile hormone esterase gene (Kethidi et al., 2005). In this study, we use cultured larval organs to identify genes regulated by JH and characterize the role of 20E in their transcriptional control. This approach bypasses the need for the hormone to penetrate the cuticle and assays the response of multiple tissues to hormone treatment. Microarray analysis of RNA samples isolated from third instar larval tissues treated with JH III, however, resulted in many false positives that could not be validated by northern blot hybridization. This result, combined with the observation that *E74B*, *pepck*, and *CG14949* can be clearly induced by JH III in cultured larval organs (Fig. 4), suggests that JH III alone may not have widespread effects on gene expression at this stage. This is consistent with biological studies which have failed to reveal critical functions for JH during *Drosophila* larval development (Riddiford, 1993). Alternatively, JH III may be metabolized under our organ culture conditions into less effective or inactive compounds. Another possibility is that a more significant transcriptional effect might be seen in response to JHB₃, which is the major form of JH produced by the *Drosophila* ring gland at larval wandering (Richard et al., 1989). Further studies are required to determine what role, if any, JH might play in regulating normal transcriptional responses during larval stages.

Extensive studies by Riddiford and colleagues have shown that JH modifies the 20E-regulation of a number of key transcription factor-encoding genes in *Manduca*: *EcR*, *USP*, *E74*, *E75*, and the *Broad-Complex* (for a review, see Riddiford et al., 2003). Similar regulatory interactions between JH and 20E have been seen in other insects, providing a molecular context for

understanding the status quo action of JH (Berger and Dubrovsky, 2005). We find that a known *Drosophila* 20E primary-response gene, *E74B*, can be upregulated by JH (Fig. 4), a response opposite that seen in *Manduca* (Stilwell et al., 2003). Another well-characterized 20E primary-response gene, *E75A*, shows no response to JH in cultured larval organs, in contrast to its robust JH induction in S2 tissue culture cells (Dubrovsky et al., 2004). Our result is similar to the lack of an effect of JH or JH mimics on *E75A* transcription in *Manduca* GV1 cells or *Manduca* epidermis (Keshan et al., 2006). Given that *E75A* can be induced by JH in the ovaries of *apterous*⁴ mutants, it is possible that third instar larval organs are not competent to mediate this transcriptional response.

Several genes we identified in this study provide unique insights into the interactions between JH and 20E. We identified five genes that are unresponsive to JH or 20E alone, but specifically induced by these two hormones in combination: *CG7906*, *CG7924*, *nmdmc*, *black*, and *CG11956* (Figs. 2,3). To our knowledge, this is the first such description of genes that require both JH and 20E for their expression. Curiously, these same genes are induced by E but not 20E. The observation that *dro5* can be up-regulated by E, but shows no response to JH and 20E, suggests that these are distinct modes of regulation (Fig. 2). Further studies will be required to determine the basis for this unexpected correlation between transcriptional responses to E and responses to 20E and JH.

Our studies have also shown that 20E can block the JH induction of *E74B* and *pepck* (Fig. 4), providing evidence that 20E can modify the JH regulation of at least some genes. The same effect is seen for *pepck* in the presence of the protein synthesis inhibitor cycloheximide, suggesting that 20E can directly prevent JH induction, possibly through EcR. This observation is the opposite of the well-characterized effects of JH on 20E signaling, where JH modifies responses to 20E, and suggests that the interplay between these two hormones is more complex than previously expected.

The identification of E-regulated genes, JH-regulated genes, and genes regulated by 20E in combination with JH provides a basis for future studies aimed at defining the crosstalk between hormone signaling pathways in insects. For example, studies of the effect of JH on *E74B*, *pepck*, and *CG14949* transcription in organs dissected from *usp* or *met* mutants could provide an opportunity to examine their possible role in mediating JH responses. Similarly, the identification of *CG14949* as a primary target for transcriptional induction by JH may facilitate the identification of JH response elements, providing new insights into the molecular mechanisms of JH action.

Acknowledgements

We thank A. Godinez for assistance with microarray hybridizations and data collection, K. King-Jones for help with microarray analysis, data comparisons and valuable discussions, M. O'Connor for helpful discussions, J. Warren for ecdysone purification, W. Goodman for JH III purification, and W. Goodman and L. Riddiford for helpful comments on the manuscript. R.B.B. was supported by an NIH Multidisciplinary Cancer Research Training Program (T32 CA0093247). This research was supported by the NIH (R01 DK075607).

References

- Andres, A.; Thummel, C. Methods for quantitative analysis of transcription in larvae and prepupae. In: LSB, G.; EA, F., editors. *Drosophila melanogaster: Practical Uses in Cell and Molecular Biology*. Academic Press; New York: 1994. p. 565-573.
- Andres AJ, Fletcher JC, Karim FD, Thummel CS. Molecular analysis of the initiation of insect metamorphosis: a comparative study of *Drosophila* ecdysteroid-regulated transcription. *Dev Biol* 1993;160:388-404. [PubMed: 8253272]

- Ashburner M. Patterns of puffing activity in the salivary gland chromosomes of *Drosophila*. VI. Induction by ecdysone in salivary glands of *D. melanogaster* cultured *in vitro*. *Chromosoma* 1972;38:255–281. [PubMed: 4627363]
- Ashburner M. Sequential gene activation by ecdysone in polytene chromosomes of *Drosophila melanogaster*. I. Dependence upon ecdysone concentration. *Dev Biol* 1973;35:47–61. [PubMed: 4207111]
- Ashok M, Turner C, Wilson TG. Insect juvenile hormone resistance gene homology with the bHLH-PAS family of transcriptional regulators. *Proc Natl Acad Sci USA* 1998;95:2761–2766. [PubMed: 9501163]
- Baker KD, Shewchuk LM, Kozlova T, Makishima M, Hassell A, Wisely B, Caravella JA, Lambert MH, Reinking JL, Krause H, et al. The *Drosophila* orphan nuclear receptor DHR38 mediates an atypical ecdysteroid signaling pathway. *Cell* 2003;113:731–742. [PubMed: 12809604]
- Baker KD, Warren JT, Thummel CS, Gilbert LI, Mangelsdorf DJ. Transcriptional activation of the *Drosophila* ecdysone receptor by insect and plant ecdysteroids. *Insect Biochem Mol Biol* 2000;30:1037–1043. [PubMed: 10989290]
- Beckstead RB, Lam G, Thummel CS. The genomic response to 20-hydroxyecdysone at the onset of *Drosophila* metamorphosis. *Genome Biol* 2005;6:R99. [PubMed: 16356271]
- Berger EM, Dubrovsky EB. Juvenile hormone molecular actions and interactions during development of *Drosophila melanogaster*. *Vitam Horm* 2005;73:175–215. [PubMed: 16399411]
- Champlin DT, Truman JW. Ecdysteroid control of cell proliferation during optic lobe neurogenesis in the moth *Manduca sexta*. *Development* 1998;125:269–277. [PubMed: 9486800]
- Clever U, Clever I, Storbeck I, Young NL. The apparent requirement of two hormones, alpha- and beta-ecdysone, for molting induction in insects. *Dev Biol* 1973;31:47–60. [PubMed: 4787193]
- Dubrovsky EB, Dubrovskaya VA, Berger EM. Juvenile hormone signaling during oogenesis in *Drosophila melanogaster*. *Insect Biochem Mol Biol* 2002;32:1555–1565. [PubMed: 12530223]
- Dubrovsky EB, Dubrovskaya VA, Berger EM. Hormonal regulation and functional role of *Drosophila* E75A orphan nuclear receptor in the juvenile hormone signaling pathway. *Dev Biol* 2004;268:258–270. [PubMed: 15063166]
- Dubrovsky EB, Dubrovskaya VA, Bilderback AL, Berger EM. The isolation of two juvenile hormone-inducible genes in *Drosophila melanogaster*. *Dev Biol* 2000;224:486–495. [PubMed: 10926782]
- Gilbert LI, Granger NA, Roe RM. The juvenile hormones: historical facts and speculations on future research directions. *Insect Biochem Mol Biol* 2000;30:617–644. [PubMed: 10876106]
- Gu S-H, Chow Y-S. Stimulation of juvenile hormone biosynthesis by different ecdysteroids in *Bombyx mori*. *Zool Studies* 2003;42:450–454.
- Hu X, Cherbas L, Cherbas P. Transcription activation by the ecdysone receptor (EcR/USP): identification of activation functions. *Mol Endocrinol* 2003;17:716–731. [PubMed: 12554759]
- Jones G, Sharp PA. Ultraspiracle: an invertebrate nuclear receptor for juvenile hormones. *Proc Natl Acad Sci USA* 1997;94:13499–13503. [PubMed: 9391054]
- Karim FD, Thummel CS. Ecdysone coordinates the timing and amounts of *E74A* and *E74B* transcription in *Drosophila*. *Genes Dev* 1991;5:1067–1079. [PubMed: 2044954]
- Keshan B, Hiruma K, Riddiford LM. Developmental expression and hormonal regulation of different isoforms of the transcription factor E75 in the tobacco hornworm *Manduca sexta*. *Dev Biol* 2006;295:623–632. [PubMed: 16697364]
- Kethidi DR, Xi Z, Palli SR. Developmental and hormonal regulation of juvenile hormone esterase gene in *Drosophila melanogaster*. *J Insect Physiol* 2005;51:393–400. [PubMed: 15890182]
- King-Jones K, Thummel CS. Nuclear receptors—a perspective from *Drosophila*. *Nat Rev Genet* 2005;6:311–323. [PubMed: 15803199]
- Koelle MR, Talbot WS, Segraves WA, Bender MT, Cherbas P, Hogness DS. The *Drosophila* EcR gene encodes an ecdysone receptor, a new member of the steroid receptor superfamily. *Cell* 1991;67:59–77. [PubMed: 1913820]
- Li C, Wong WH. Model-based analysis of oligonucleotide arrays: expression index computation and outlier detection. *Proc Natl Acad Sci USA* 2001;98:31–36. [PubMed: 11134512]

- Miura K, Oda M, Makita S, Chinzei Y. Characterization of the *Drosophila Methoprene-tolerant* gene product. Juvenile hormone binding and ligand-dependent gene regulation. *Febs J* 2005;272:1169–1178. [PubMed: 15720391]
- Oberlander H. •-ecdysone induced DNA synthesis in cultured wing discs of *Galleria mellonella* - Inhibition by 20-hydroxyecdysone and 22-isoecdysone. *J Insect Phys* 1972;18:223–228.
- Palanker L, Necakov AS, Sampson HM, Ni R, Hu C, Thummel CS, Krause HM. Dynamic regulation of *Drosophila* nuclear receptor activity *in vivo*. *Development* 2006;133:3549–3562. [PubMed: 16914501]
- Parvy JP, Blais C, Bernard F, Warren JT, Petryk A, Gilbert LI, O'Connor MB, Dauphin-Villemant C. A role for •FTZ-F1 in regulating ecdysteroid titers during postembryonic development in *Drosophila melanogaster*. *Dev Biol* 2005;282:84–94. [PubMed: 15936331]
- Petryk A, Warren JT, Marques G, Jarcho MP, Gilbert LI, Kahler J, Parvy JP, Li Y, Dauphin-Villemant C, O'Connor MB. Shade is the *Drosophila* P450 enzyme that mediates the hydroxylation of ecdysone to the steroid insect molting hormone 20-hydroxyecdysone. *Proc Natl Acad Sci USA* 2003;100:13773–13778. [PubMed: 14610274]
- Restifo LL, Wilson TG. A juvenile hormone agonist reveals distinct developmental pathways mediated by ecdysone-inducible Broad Complex transcription factors. *Dev Genet* 1998;22:141–159. [PubMed: 9581286]
- Richard DS, Applebaum SW, Sliter TJ, Baker FC, Schooley DA, Reuter CC, Henrich VC, Gilbert LI. Juvenile hormone bisepoxide biosynthesis *in vitro* by the ring gland of *Drosophila melanogaster*: a putative juvenile hormone in the higher Diptera. *Proc Natl Acad Sci U S A* 1989;86:1421–1425. [PubMed: 2493154]
- Richards G. Sequential gene activation by ecdysone in polytene chromosomes of *Drosophila melanogaster*. VI. Inhibition by juvenile hormones. *Dev Biol* 1978;66:32–42. [PubMed: 109334]
- Richards G, Da Lage JL, Huet F, Ruiz C. The acquisition of competence to respond to ecdysone in *Drosophila* is transcript specific. *Mech Dev* 1999;82:131–139. [PubMed: 10354477]
- Riddiford, LM. Hormones and *Drosophila* development. Cold Spring Harbor Press; Cold Spring Harbor, New York: 1993.
- Riddiford, LM. Molecular aspects of juvenile hormone action in insect metamorphosis. In: Gilbert, LI.; Tata, J.; Atkinson, B., editors. *Metamorphosis: Postembryonic Reprogramming of Gene Expression in Amphibian and Insect Cells*. Academic Press; New York: 1996. p. 223-251.
- Riddiford LM, Ashburner M. Effects of juvenile hormone mimics on larval development and metamorphosis of *Drosophila melanogaster*. *Gen Comp Endocrinol* 1991;82:172–183. [PubMed: 1906823]
- Riddiford LM, Hiruma K, Zhou X, Nelson CA. Insights into the molecular basis of the hormonal control of molting and metamorphosis from *Manduca sexta* and *Drosophila melanogaster*. *Insect Biochem Mol Biol* 2003;33:1327–1338. [PubMed: 14599504]
- Segraves WA, Hogness DS. The *E75* ecdysone-inducible gene responsible for the 75B early puff in *Drosophila* encodes two new members of the steroid receptor superfamily. *Genes Dev* 1990;4:204–219. [PubMed: 2110921]
- Somme-Martin G, Colardeau J, Lafont R. Conversion of ecdysone and 20-hydroxyecdysone into 3-dehydroecdysteroids is a major pathway in third instar *Drosophila melanogaster* larvae. *Insect Biochem* 1988;18:729–734.
- Stilwell GE, Nelson CA, Weller J, Cui H, Hiruma K, Truman JW, Riddiford LM. *E74* exhibits stage-specific hormonal regulation in the epidermis of the tobacco hornworm, *manduca sexta*. *Dev Biol* 2003;258:76–90. [PubMed: 12781684]
- Thomas HE, Stunnenberg HG, Stewart AF. Heterodimerization of the *Drosophila* ecdysone receptor with Retinoid X Receptor and Ultraspiracle. *Nature* 1993;362:471–475. [PubMed: 8385270]
- Wang SF, Ayer S, Segraves WA, Williams DR, Raikhel AS. Molecular determinants of differential ligand sensitivities of insect ecdysteroid receptors. *Mol Cell Biol* 2000;20:3870–3879. [PubMed: 10805730]
- Warren JT, Petryk A, Marques G, Jarcho M, Parvy JP, Dauphin-Villemant C, O'Connor MB, Gilbert LI. Molecular and biochemical characterization of two P450 enzymes in the ecdysteroidogenic pathway of *Drosophila melanogaster*. *Proc Natl Acad Sci USA* 2002;99:11043–11048. [PubMed: 12177427]

- Warren JT, Petryk A, Marques G, Parvy JP, Shinoda T, Itoyama K, Kobayashi J, Jarcho M, Li Y, O'Connor MB, et al. Phantom encodes the 25-hydroxylase of *Drosophila melanogaster* and *Bombyx mori*: a P450 enzyme critical in ecdysone biosynthesis. *Insect Biochem Mol Biol* 2004;34:991–1010. [PubMed: 15350618]
- Warren JT, Yerushalmi Y, Shimell MJ, O'Connor MB, Restifo LL, Gilbert LI. Discrete pulses of molting hormone, 20-hydroxyecdysone, during late larval development of *Drosophila melanogaster*: correlations with changes in gene activity. *Dev Dyn* 2006;235:315–326. [PubMed: 16273522]
- Yao TP, Forman BM, Jiang Z, Cherbas L, Chen JD, McKeown M, Cherbas P, Evans RM. Functional ecdysone receptor is the product of *EcR* and *Ultraspiracle* genes. *Nature* 1993;366:476–479. [PubMed: 8247157]
- Yoshiyama T, Namiki T, Mita K, Kataoka H, Niwa R. Neverland is an evolutionally conserved Rieske-domain protein that is essential for ecdysone synthesis and insect growth. *Development* 2006;133:2565–2574. [PubMed: 16763204]
- Zhou X, Riddiford LM. *Broad* specifies pupal development and mediates the 'status quo' action of juvenile hormone on the pupal-adult transformation in *Drosophila* and *Manduca*. *Development* 2002;129:2259–2269. [PubMed: 11959833]




Fig 1. Ecdysone regulates a distinct set of genes from those controlled by 20-hydroxyecdysone (A) Venn diagram depicting the overlap between genes regulated by ecdysone (E) and those regulated by 20E. The set of 376 20E-regulated genes was derived from both 20E-regulated genes and 20E primary-response genes from Beckstead et al. (2005), as described in Experimental Procedures. (B) Cluster analysis of microarray data derived from RNA samples isolated from staged wild type animals (Beckstead et al., 2005). The colors for each time point represent the change in expression level relative to the average expression levels across all time points for that gene, with dark blue indicating the lowest level of expression and red indicating the highest level, as depicted on the bottom. The numbers at the top indicate hours relative to pupariation.




Fig 2. Validation of transcriptional responses to E and JH

Northern blot analysis of RNA samples isolated from cultured third instar larval organs treated with either no hormone (control), (10R)-juvenile hormone III (JH), JH and 20-hydroxyecdysone (JH+20E), 20E alone, or ecdysone (E) for 6 hours. Organ cultures were established from third instar larvae staged at either ~20 hours or ~10 hours prior to pupariation. Lane numbers are shown at the bottom. The blots were hybridized to detect transcripts as shown, with *rp49* mRNA used as a control for loading and transfer.




Fig 3. The brain and ring gland are not required for transcriptional responses to ecdysone

Northern blot analysis of RNA samples isolated from organ cultures derived from -10 hour third instar larvae treated with no hormone (control), (10R)-juvenile hormone III and 20-hydroxyecdysone (JH+20E), or ecdysone alone (E). One set of cultures had the brain and ring gland removed prior to the addition of hormone (- brain). The blots were hybridized to detect transcripts as shown, with *rp49* mRNA used as a control for loading and transfer.




Fig 4. Juvenile hormone regulates gene expression in larval organ culture

Northern blot analysis of RNA samples isolated from organ cultures derived from third instar larvae at ~20 hours prior to pupariation, treated with either no hormone (control), cycloheximide (cyc), (10R)-juvenile hormone III (JH), JH III and 20-hydroxyecdysone (JH, 20E), JH III and cycloheximide (JH,cyc), JH III and 20E and cycloheximide (JH,20E,cyc), (10S)-juvenile hormone III (10S), farnesol (far), and methyl linoleate (ml). The latter three are negative controls for (10R)-JH III. Lane numbers are shown at the bottom. The blots were hybridized to detect transcripts as shown, with *rp49* mRNA used as a control for loading and transfer.

Table 1

E-regulated genes

gene	fold change	molecular function
<i>dro5</i>	5.35	drosomycin
<i>CG7906</i>	3.49	n/a
<i>CG7924</i>	2.41	n/a
<i>CG14957</i>	1.93	n/a
<i>Su(Tpl)</i>	1.87	transcription elongation factor
<i>mob1</i>	1.81	n/a
<i>malvolio</i>	1.69	cation transporter
<i>CG11686</i>	1.69	n/a
<i>nmdmc</i>	1.69	NAD-dependent methylenetetrahydrofolate dehydrogenase
<i>regena</i>	1.68	transcription factor
<i>CG12224</i>	1.67	oxidoreductase activity
<i>smi35A</i>	1.66	serine/threonine kinase
<i>CG14478</i>	1.65	n/a
<i>creg</i>	1.64	transcriptional repressor
<i>snap</i>	1.64	soluble NSF attachment protein
<i>croquemort</i>	1.63	scavenger receptor
<i>CG8054</i>	1.62	n/a
<i>adenosine 3</i>	1.62	phosphoribosylamine-glycine ligase
<i>CG7606</i>	1.6	uncertain
<i>NPC2</i>	1.57	receptor binding
<i>emc</i>	1.57	transcription co-repressor
<i>Vap-33-1</i>	1.57	structural molecule
<i>Hsc70Cb</i>	1.57	chaperone binding
<i>CG15248</i>	1.56	uncertain
<i>egghead</i>	1.56	beta-1,4-mannosyltransferase
<i>CG1965</i>	1.56	transcription regulator
<i>CG7806</i>	1.56	xenobiotic-transporting ATPase
<i>CG9317</i>	1.55	organic cationic transporter
<i>CG1837</i>	1.55	protein disulfide isomerase
<i>CG6145</i>	1.54	n/a
<i>CG11169</i>	1.54	n/a
<i>rab-11</i>	1.53	small GTPase
<i>fu12</i>	1.53	1-acylglycerol-3-phosphate O-acyltransferase
<i>CG2082</i>	1.53	n/a
<i>CG7696</i>	1.53	transcription regulator
<i>smi21F</i>	1.53	n/a
<i>CG5455</i>	1.52	catalytic activity
<i>Stat92E</i>	1.51	transcription factor
<i>CG3823</i>	1.51	vitamin E binding
<i>CG12965</i>	1.51	transcription factor
<i>CG1246</i>	1.51	n/a
<i>rap</i>	1.5	n/a
<i>CG17110</i>	1.5	metallopeptidase
<i>CG12004</i>	1.5	n/a
<i>CG3223</i>	1.5	n/a
<i>CG6169</i>	1.5	n/a
<i>bowl</i>	1.5	transcription factor
<i>CG10114</i>	-1.52	helena Element
<i>new glue 3</i>	-1.55	structural molecule
<i>PGRP-SA</i>	-1.56	peptidoglycan receptor
<i>CG13187</i>	-1.57	protein binding
<i>nemy</i>	-1.58	glutaminase
<i>GST-E1</i>	-1.62	glutathione transferase
<i>acp26Ab</i>	-1.74	hormone
<i>GST-D3</i>	-1.77	glutathione transferase