

Published in final edited form as:


Org Lett. 2006 November 9; 8(23): 5315–5316.

Use of Allylic Strain To Enforce Stereochemistry. Direct Syntheses of 7,8-Dihydroxycalamenene and Mansonone C

George A. Kraus and Insik Jeon

 Department of Chemistry, Iowa State University, Ames, Iowa 50011 gakraus@iastate.edu

Abstract



Direct syntheses of 7,8-dihydroxycalamenene and mansonone C were achieved. The *cis*-stereochemistry required for the synthesis of 7,8-dihydroxycalamenene was introduced by an intramolecular cyclization directed by allylic strain.

Allylic 1,3-strain has been used in acyclic systems to direct the introduction of new stereogenic centers.¹ Notable examples include the work of Kishi,² Adam,³ and Giese.⁴ We are not aware of any application of allylic strain to control the relative stereochemistry in disubstituted tetralins. Tetralins such as **1**, **3**, and **4** have attracted considerable synthetic attention (Figure 1).⁵ 7,8-Dihydroxycalamenene (**1a**) exhibits useful anti-infective activity.⁶ Hydroxycalamenene (**1b**) was isolated from *Hypericum elodeoides*.⁷ Mansonone C (**2**), extracted from the heartwood of *Mansonia altissima*,⁸ was found to possess promising antifungal, larvicidal, and anti-oxidant properties.⁹ Schmalz recorded a synthesis of **1a** using arenechromium complexes to introduce the relative stereochemistry.¹⁰

Many synthetic approaches to these compounds begin with natural products such as menthone wherein the relative stereochemistry has already been established.¹¹

Several researchers have noted that attempts to install stereochemistry by epimerization or by cyclization onto the aromatic ring have led to mixtures.^{12,13} We report herein that a system such as **5**, wherein allylic strain between G and the methyl group forces the methyl group to be axial as the six-membered ring is being formed, affords exclusively the *cis*-stereoisomer **7** (Scheme 1).

In order to evaluate the directing effect, we first synthesized allylic acetate **8** (Scheme 2). Allylic acetate **8** was synthesized starting from 6-methyl-5-hepten-2-one and the anion of 1,4-dimethoxybenzene. Dehydroxylation of the resulting benzylic alcohol using Li/NH₃ followed by allylic oxidation by the method of Sharpless¹⁴ yielded an allylic alcohol which, upon acetylation, afforded **8**. Surprisingly, cyclization of allylic acetate **8** using the conditions of Ma

Correspondence to: George A. Kraus.

Supporting Information Available: Experimental procedures, characterization data, and NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

and Zheng¹⁵ afforded tetralin **10** and naphthalene **11** in 51% and 37% yields, respectively. Compound **9**, the expected product, was not isolated.

We believe that compounds **10** and **11** result from a novel cation-mediated disproportionation reaction. Comparison of the NMR spectrum of **10** with that of the literature compound¹³ showed that the *cis*-stereoisomer was exclusively formed.

In support of this allylic strain assisted stereoselective cyclization, we have found that cyclization of allylic acetate **12**, which does not contain the directing group G, affords tetralin **13** as a 1.3:1 mixture of diastereomers (Scheme 3).

With the stereochemistry of **10** established, we began the synthesis of **1a** by the reaction of 5-methyl-4-hexen-1-ol¹⁶ with the anion of 1,2-bis(methoxymethoxy)-3-methylbenzene (Scheme 4). The resulting alkoxide was in situ acetylated by the addition of acetic anhydride. The displacement of the acetate to the corresponding methyl group was achieved using Me₃Al to afford **15**. In order to enhance the stability for the acid-mediated cyclization, MOM protecting groups were converted to the more stable methyl ether **16**. It was then oxidized and acetylated by the same methods used to generate **8**. Cyclization of **17** using trifluoroacetic acid in acetic acid at 70 °C for 12 h followed by deprotection by the use of BBr₃ generated 7,8-dihydroxycalamenene (**1a**) in two steps (30% yield). Interestingly enough, mansonone C (**2**) was also obtained from the same reaction in two steps (46% yield). The direct synthesis of 7,8-dihydroxycalamenene (**1a**) demonstrates the advantage of employing the concept of allylic strain in organic synthesis.

Acknowledgements

We thank the National Institutes of Health (Grant No. P01 ES12020) and the Office of Dietary Supplements for partial financial support through the Center for Research on Botanical Dietary Supplements at Iowa State University.

References

1. Hoffmann RW. Chem Rev 1989;89:1841. Johnson F. Chem Rev 1968;68:375.
2. Johnson MR, Nakata T, Kishi Y. Tetrahedron Lett 1979:4343.
3. Adam W, Wirth T. Acc Chem Res 1999;32:703.
4. Giese B, Bulliard M, Zeitz HG. Synlett 1991:425.
5. Erogorgiaene (3): Rodriguez AD, Ramirez C. J Nat Prod 2001;64:100. [PubMed: 11170678]
Elisapterosin B (4): Rodriguez AD, Ramirez C, Rodriguez II, Barnes CL. J Org Chem 2000;65:1390. [PubMed: 10814100]
6. Wahyouno S, Hoffmann JJ, Bates RB, McLaughlin SP. Phytochemistry 1991;30:2175.
7. Mathela DK, Mathela CS, Dev V. J Indian Chem Soc 1984;61:792.
8. Marini Bettòlo GB, Casinovi CG, Galeffi C. Tetrahedron Lett 1965:4857.
9. Tiew P, Ioset JR, Kokpol U, Chavasiri W, Hostettmann K. Phytother Res 2003;17:190. [PubMed: 12601687]
10. Schmalz HG, Hollander J, Arnold M, Duerner G. Tetrahedron Lett 1993;34:6259.
11. Serra S, Fuganti C. Tetrahedron Lett 2005;46:4769.
12. Ramaiah P, Rao AS. Synth Commun 1989;19:931.
13. Beckwith PLM, Ghisalberti EL, Jefferies PR, Raston CL, White AH. Aust J Chem 1988;41:351.
14. Umbreit MA, Sharpless KB. J Am Chem Soc 1977;99:5526.
15. Ma S, Zhang J. Tetrahedron 2003;59:6273.
16. Kraus GA, Kim J. Org Lett 2004;6:3115. [PubMed: 15330601]







Figure 1.
Tetralin-derived products and mansonone C.



Scheme 1.



Scheme 2.



Scheme 3.

