

IFN- γ action in the media of the great elastic arteries, a novel immunoprivileged site

Rapid Publication

Albert J. Dal Canto,¹ Paul E. Swanson,¹ Andrew K. O'Guin,¹ Samuel H. Speck,^{1,2} and Herbert W. Virgin^{1,2}

¹Departments of Pathology and Immunology, and

²Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, Missouri, USA

Address correspondence to: Herbert W. Virgin IV, Washington University School of Medicine, 660 South Euclid Avenue, St. Louis, Missouri 63110, USA. Phone: (314) 362-9223; Fax: (314) 362-4096; E-mail: virgin@immunology.wustl.edu.

Received for publication October 12, 2000, and accepted in revised form December 5, 2000.

Infection of medial smooth muscle cells with γ -herpesvirus 68 (γ HV68) causes severe chronic vasculitis that is restricted to the great elastic arteries. We show here that persistence of disease in the great elastic arteries is (a) due to inefficient clearance of viral infection from this site compared with other organs or other vascular sites, and (b) associated with failure of T cells and macrophages to enter the virus-infected elastic media. These findings demonstrate immunoprivilege of the media of the great elastic arteries. We found that IFN- γ acted on somatic cells during acute infection to prevent the establishment of medial infection and on hematopoietic cells to determine the severity of disease in this site. The immunoprivileged elastic media may provide a site for persistence of pathogens or self antigens leading to chronic vascular disease, a process regulated by IFN- γ actions on both somatic and hematopoietic cells. These concepts have significant implications for understanding immune responses contributing to or controlling chronic inflammatory diseases of the great vessels.

J. Clin. Invest. **107**:R15–R22 (2001).

Introduction

We developed a new animal model of chronic viral vasculitis restricted to the great elastic arteries (chronic elastic arteritis; refs. 1 and 2). Severe chronic elastic arteritis persists weeks to months after infection of IFN- γ receptor-deficient (IFN- γ R^{-/-}) mice with doses of the γ -herpesvirus γ HV68 that fail to induce arteritis in wild-type mice. Infection with γ HV68 also increases vascular lesion formation in apoE-deficient animals (3). The role of IFN- γ in regulating γ HV68 arteritis is particularly interesting since IFN- γ is a key regulator of vascular inflammation in multiple systems and can either increase or decrease vascular pathology, depending on the nature of the insult to the vessel (1, 4–6, reviewed in ref. 7).

γ HV68-induced chronic elastic arteritis is characterized by striking inflammation of the adventitia and intima of the affected artery segment in regions containing productively

infected smooth muscle cells (SMCs) in the media (1, 2). Continued productive infection contributes to chronic damage and inflammation (2). Interestingly, while immunodeficiency makes mice more susceptible to chronic elastic arteritis, γ HV68-infected wild-type weanling mice also develop elastic arteritis lasting up to 5 months after infection (1). This raises the question of why infection of the great vessels can persist for prolonged periods in an immunocompetent host.

In this study, we show that the tropism of γ HV68 for the media of the elastic arteries during chronic infection was due to failure of the immune response to effectively clear the virus from the elastic media despite effective clearance of other vascular sites and other organs. Effector T cells and macrophages did not enter the media of the great vessels even when active viral infection was present. We also show that IFN- γ independently regu-

lates the severity and incidence of arteritis by actions on different cell types. These studies defining an immunoprivileged site within the vasculature, and demonstrating the cellular basis for a critical protective role of IFN- γ in this site, have important implications for understanding chronic inflammatory diseases of the great vessels.

Methods

Viruses, mice, and pathologic analysis. γ HV68 was passaged and assayed as described (1). γ HV68 mutants M1.LacZ and v-cyclin.LacZ have been described (8, 9). IFN- γ R^{-/-} mice are on a pure 129Ev/Sv background, and 129Ev/Sv controls were used as controls (10). Mice 5 to 7 weeks old were infected intraperitoneally with γ HV68 in 0.5 ml of complete DMEM for all experiments. Organs were titrated by plaque assay, and prepared and stained for histopathologic examination as described (1). Criterion for lesion scores based on hematoxylin and eosin (H&E) histology were as described (2). Briefly, scores increased from 0 to 5 as the extent of adventitial and intimal thickening, neutrophilic infiltration, and medial necrosis increased. Lesions were scored in a blinded fashion by A.J. Dal Canto and H.W. Virgin, and scores were averaged. Microscopic images were prepared using a Zeiss Axiophot microscope with a Spot camera and Spot software 1.1 (Diagnostics Instruments, Sterling Heights, Michigan, USA) and Northern Eclipse v2.0 software (Empix Imaging Inc., Mississauga, Ontario, Canada).

β-galactosidase staining and immunohistochemistry. Mice were infected intraperitoneally with 5×10^6 PFU M1.LacZ or γ HV68 v-cyclin.LacZ and sacrificed by perfusion with 30 ml 2% paraformaldehyde and 0.2% glutaraldehyde in PBS. The heart, aorta, and visceral organs were removed en bloc, rinsed in PBS, and stained for β -galactosidase (β -gal) activity at 37°C overnight (11), before fixation in 4% paraformaldehyde. Staining was evaluated under a dissecting microscope, and pictures were taken with a Kodak DC120 digital camera. Viral antigen was detected by immunohistochemistry as described (1). For detection of inflammatory cells, frozen sections were fixed in 1:1 methanol/acetone at -20°C. Endogenous peroxidase activity was inhibited (1), slides were blocked (1% BSA, 0.2% powdered skim milk, 0.3% triton in PBS) for 30 minutes, and then biotin blocked (Vector Laboratories, Burlingame, California, USA). Rat mAb's used were: anti-CD4 Ab GK1.5 (12); anti-CD8 Ab H35 (12), anti-B-220 Ab 14.8 (12), and anti-CD11b Ab 5C6 (CRL-1969; American Type Culture Collection, Rockville, Maryland, USA), and were added at 25 ng/ml overnight at 4°C. F4/80 (13) mAb supernatant was used at a 1:2000 dilution. Rat IgG was used as a negative control. Staining was detected using F(ab')₂ biotin-conjugated donkey anti-rat secondary Ab (1.2 μ g/ml; Jackson ImmunoResearch Laboratories Inc., West Grove, Pennsylvania, USA).

Bone marrow reconstitutions and IFN- γ depletion in vivo. Four-week-old mice were lethally γ -irradiated with 1100 R followed by intravenous injection of 1 to 2×10^7 nucleated bone marrow cells from sex-matched 129Ev/Sv or IFN- γ R^{-/-} mice. The percentage of reconstitution was assessed at 6–8 weeks using quantitative Southern blots of peripheral blood lymphocyte DNA (QIAamp blood kit; QIAGEN Inc., Valencia, California, USA) cut with BamHI and probed with a 2-kb fragment corresponding to exon 4 of the IFN- γ R gene (14). To transiently deplete IFN- γ , mice were injected intraperitoneally with 250 μ g of IFN- γ -neutralizing hamster mAb H22 (15) in 0.5 ml sterile saline 1 day before and 6 days after infection. To chronically deplete IFN- γ , mice

were injected weekly until sacrifice (6 weeks). The Ab dose was selected based on previous reports of effective IFN- γ depletion with H22 (15, 16).

Primary cell cultures. Primary mouse aortic cells were prepared based on published protocols (17, 18). Ten thoracic aortas from 4- to 6-week-old male 129Ev/Sv mice were harvested, placed in HBSS with 0.2 mM CaCl (HBSS/Ca), and the adventitia removed by blunt dissection. Digestion with collagenase followed by trypsin yielded an intimal/adventitial culture. Further digestion with collagenase, elastase, and trypsin yielded a medial culture. After 3–4 days, cells were treated with 0.12% trypsin plus 0.5 mM EDTA for 5 minutes at 37°C and replated. Every 2–4 days thereafter, each well was passaged into two new wells. Cells (5 to 50×10^4) were obtained from each culture after the second passage and plated (1 to 3×10^4 cells) into eight-well glass chamber slides (LabTek; Nalge Nunc International, Naperville, Illinois, USA) for use in immunofluorescence studies, 24-well tissue culture (TC) plates for phase-contrast microscopy, or 12-well TC plates for electron microscopy (EM). Two to three days later cultures were treated for 48 hours with or without IFN- γ (250 U/ml for medial cultures and 64 U/ml for intimal/adventitial cultures; Genzyme Pharmaceuticals, Cambridge, Massachusetts, USA), infected at an moi of 5 for 2 hours at 37°C, and then cultured in fresh medium with or without IFN- γ . Fresh media with or without IFN- γ was added every 3 days. Phase-contrast pictures were taken on a Olympus CK2 microscope equipped with a Kodak DC120 digital camera and the Kodak MDS 120 v1.0 acquisition program.

Immunofluorescence and EM. For immunofluorescence, cells were fixed for 20 minutes at -20°C in methanol, washed, and then blocked for 30 minutes at room temperature in 20 μ g/ml BSA, 20% goat serum, 0.01% triton in PBS. Polyclonal anti- γ HV68 antiserum (1), or preimmune serum, was used at a dilution of 1:500 to detect viral antigen. A mouse mAb to human muscle actin (HHF35; DAKO Corp., Carpinteria, California, USA) was used at 0.55 μ g/ml to detect SMCs. An isotype-matched Ab to keyhole limpet hemocyanin was used as a negative control (PharMingen, San

Diego, California, USA). Primary Ab's were incubated overnight at 4°C, and secondary Ab's were added for 1 hour at room temperature. Muscle actin was detected using a Cy3-conjugated goat anti-mouse secondary antibody (10 μ g/ml; Jackson ImmunoResearch Laboratories Inc.), and viral antigen was detected using an Oregon green 488-conjugated goat anti-rabbit secondary antibody (7 μ g/ml; Molecular Probes, Eugene, Oregon, USA). Bisbenzimidazole was added (0.5 μ g/ml) to stain the nuclei blue. For EM, cells were fixed with 3% glutaraldehyde in 0.1 M cacodylate buffer (0.1 M sodium cacodylate in water, pH 7.0, plus 0.54% wt/vol dextrose) for 1 hour, washed in 0.1 M cacodylate buffer, scraped, pelleted, and processed as described (2). SMCs were distinguished by the presence of organized actin filaments, subplasmalemmal attachment plaques, and pinocytosis. The presence of viral capsids was evaluated within nuclei in a blinded fashion by A.J. Dal Canto on high-power photographs ($\times 57,800$).

Statistical analysis. Data were plotted and statistically analyzed using GraphPad Prism (GraphPad Software, San Diego, California, USA). The severity of pathology was statistically analyzed by comparing lesions scores using the Mann-Whitney test. Staining of different regions of the aorta for β -gal and alterations in incidence of arteritis in mice treated with anti-IFN- γ Ab were compared using a χ -square test and Fisher's exact test (which gave comparable results). Data from immunofluorescence and EM studies were analyzed using an unpaired *t* test.

Results

Viral tropism due to anatomic specificity of immune function. We considered two hypotheses to explain the restriction of γ HV68-induced arteritis to the great vessels. γ HV68 might infect only SMCs of the elastic media, leaving other vascular sites uninfected. Alternatively, γ HV68 could infect multiple vascular sites, but be efficiently cleared from all sites other than the elastic media. To distinguish between these hypotheses, we infected IFN- γ R^{-/-} mice with β -galactosidase (β -gal) expressing γ HV68 mutants that induce chronic elastic arteritis (8, 9). At various times, the




Figure 1

Localization of γ HV68 vascular infection over time. IFN γ R $^{-/-}$ mice were infected with either M1.*LacZ* or γ HV68 v-cyclin.*LacZ*. The aorta and its major branches were resected and stained en bloc for β -gal activity. Infection with wild type γ HV68 resulted in no comparable staining.

(a) Staining 13 days after infection of a vessel on the posterior surface of the kidney. (b) Staining 13 days after infection of the descending aorta. Note staining in both the adventitia (Adv) and media (M). (c) Staining 21 days after infection of the descending aorta. Staining is predominately in the media. (d) Staining 21 days after infection at the base of the aorta. Note staining is only in the media. (e) Cross section from the descending aorta 21 days after infection demonstrating inflammation surrounding medial staining. (f) High-power view of boxed region in e. L, lumen; I, intima.

aorta with its major branches and attached heart, liver, and kidneys were dissected out as a unit and stained for β -gal activity to localize regions of viral infection (Figure 1 and 2).


Seven to fifteen days after infection, γ HV68 was present in both the media and adventitia of the great vessels as shown by evaluation under the dissecting microscope (Figure 1a and Figure 2, regions a and b). Together with infection of a peripheral vessel adjacent to the kidney (Figure 1a and Figure 2, region c), this proves that γ HV68

tropism is not due to selective infection of SMCs in the elastic media.

Infection with γ HV68 was efficiently cleared from both the adventitia of the great vessels and a peripheral vessel. Whereas 20 of 28 mice had bilateral staining of a peripheral vessel 7–15 days after infection (Figure 1a and Figure 2, region c), clearance was evident 17–21 days after infection and complete by 42–43 days ($P < 0.0001$ for late times versus days 7–15). Similarly, between 17 and 21 days after infection, adventitial staining was cleared signif-

icantly (Figure 1b compared to Figure 1, c and d, and Figure 2, regions a and b; $P < 0.0001$ for late times versus days 7–15). In contrast, infection of the elastic media at the base of the aorta persisted through days 42–43. In other studies, persistence of medial infection and disease of the great vessels has been seen up to 105 days after infection (1). Clearance of the media of the thoracic aorta also lagged behind clearance of either the adventitia or a peripheral vessel (Figure 1 and 2).

Clearance of adventitial infection was immune mediated because it did not occur in B cell- and T cell-deficient RAG1 $^{-/-}$ mice (not shown). Failure of efficient immune clearance of γ HV68 from the elastic media was associated with a lack of infiltration of CD4 T cells (not shown), CD8 T cells, and macrophages into the media (Fig-



Region	Days 7-15 <i>n</i> = 28	Days 17-21 <i>n</i> = 25	Days 42,43 <i>n</i> = 18
a	Adv	11 ^A	0 ^A
	M	19	15 ^{A,D}
b	Adv	3 ^A	0 ^A
	M	21	2 ^{A,E}
c	40	2 ^A	0 ^A

Figure 2

Quantitation of clearance of γ HV68 from different vascular structures over time. IFN- γ R $^{-/-}$ mice were infected and analyzed as in Figure 1 at days 7, 9, 11, 13, 15, 17, 19, 21, 42, and 43 after infection. *n*, number of mice. The number of aortas with blue staining in the adventitia (Adv) or media (M) is reported for the aortic base (a) or the descending aorta (b), which was representative of the entire aorta. Staining of a peripheral vessel on the posterior surface of both kidneys is also reported (c). For staining at late times compared with staining at days 7–15, ^A $P < 0.0001$; ^B $P > 0.05$; ^C $P = 0.04$. For medial clearance compared with adventitial clearance, ^D $P < 0.0001$; ^E $P = 0.006$; ^F $P = 0.1$.




Figure 3

Localization of inflammatory cells in arteritic lesions. Shown are low- and high-power views of immunohistochemistry on parallel aortic frozen sections from an IFN- γ ^{-/-} mouse infected with γ HV68 4 weeks after infection. Dark-brown staining represents Ab binding. (a and b) CD8 T cells are present in the intima and adventitia, but not the media. Similar staining was observed with an Ab to CD4 (not shown). (c and d) Staining for CD11b shows Mac1-positive cells (macrophages and neutrophils) in all three layers of the aorta. (e and f) Staining for F4/80-positive macrophages is limited to the intima and adventitia. (g and h) Negative control with rat IgG. The area of highest background staining is shown in h for comparison with specific staining in panels b, d, and f. I, intima; M, media; Adv, adventitia.

ure 1, e and f, and Figure 3) despite the presence of active virus infection of medial smooth muscle cells (refs. 1 and 2), data not shown, and Figure 1f). The only inflammatory cells consistently present in the media were CD11b-positive (Mac1-positive) cells with the morphology of neutrophils (Figure 3, c and d).

Failure to clear the elastic media in normal mice. While it was possible that IFN- γ was required for effective immune function in the elastic media, we hypothesized that inefficient clearance of medial infection reflected a fundamental property of the elastic media. This hypothesis makes two predictions: (a) it should be possible to induce chronic elastic arteritis in wild-type mice, which have functional IFN- γ responses, with viral doses high enough to establish medial infection; and (b) transient depletion of IFN- γ in wild-type mice during acute infection should induce chronic elastic arteritis by fostering medial infection.

Consistent with our hypothesis, increasing the dose of γ HV68 in wild-type mice to greater than 5×10^7 PFU γ HV68 resulted in arteritis persisting 5.5 to 10 weeks (Figure 4). In contrast, IFN- γ ^{-/-} mice developed chronic arteritis when infected with 50-fold less virus (10^6 PFU) (1). While arteritis persisted, none of the ten wild-type animals evaluated at 6 weeks had

detectable γ HV68 in spleen, liver, or lung (sensitivity of plaque assay 50 PFU/organ), demonstrating that persistence occurs in the great vessels despite clearance of other organs. Furthermore, transient depletion of IFN- γ increased susceptibility of wild-type mice to chronic arteritis. Seventeen percent (5 of 29) of control mice infected with 5×10^7 PFU of γ HV68 had chronic elastic arteritis 6 weeks after infection. In contrast, 67% (6 of 9) of mice transiently depleted of IFN- γ developed chronic disease (Figure 4a; $P = 0.0043$). At twofold to fivefold lower infectious doses, disease was seen in 50% of transiently depleted mice, but not in controls (Figure 4a; $P = 0.0062$).

We scored the severity of lesions after γ HV68 infection of wild-type mice, wild-type mice transiently or chronically depleted of IFN- γ , and IFN- γ ^{-/-} mice (Figure 4, b–e). All arteritic lesions had viral antigen detectable in the media by immunohistochemistry (not shown and ref. 1 and 2). Wild-type and transiently IFN- γ -depleted wild-type mice had mild lesions with mononuclear infiltrates in the aortic intima and/or adventitia, but no medial neutrophilic infiltrates or necrosis (Figure 4, b and c). In contrast, IFN- γ ^{-/-} and chronically IFN- γ -depleted wild-type mice had severe lesions with intense medial neutrophilic infiltrates and/or medial necrosis (Figure 4, d and e).

Cellular sites of IFN- γ action that determine incidence and severity of arteritic lesions. We performed reciprocal bone marrow transfers to identify the cellular sites of IFN- γ action that determine the incidence and severity of chronic elastic arteritis. Wild-type (129Ev/Sv) and IFN- γ ^{-/-} recipient mice were lethally irradiated and reconstituted with bone marrow cells from either 129Ev/Sv or IFN- γ ^{-/-} donors (donor \rightarrow recipient: 129Ev/Sv \rightarrow 129Ev/Sv; IFN- γ ^{-/-} \rightarrow IFN- γ ^{-/-}; 129Ev/Sv \rightarrow IFN- γ ^{-/-}; IFN- γ ^{-/-} \rightarrow 129Ev/Sv). After 8 weeks, reconstituted mice were infected with 10^7 PFU of γ HV68, and arteritis was evaluated over 12 weeks. Controls demonstrated that the procedure neither protected from, nor predisposed to, arteritis. Thus, 10 of 13 of the IFN- γ ^{-/-} \rightarrow IFN- γ ^{-/-} mice developed severe arteritis, whereas 0 of 11 of the 129Ev/Sv \rightarrow 129Ev/Sv mice developed lesions (Figure 5, groups A and D). The IFN- γ -receptor status of the recipient determined susceptibility to arteritis since 9 of 18 of the 129Ev/Sv \rightarrow IFN- γ ^{-/-} mice, but 0 of 19 of the IFN- γ ^{-/-} \rightarrow 129Ev/Sv mice developed disease (Figure 5, groups B and C). Lesions in IFN- γ ^{-/-} \rightarrow IFN- γ ^{-/-} mice were very severe (Figure 5, group A; average lesions score 4.7 ± 0.3), while lesions in the 129Ev/Sv \rightarrow IFN- γ ^{-/-} mice were much less severe, lacking medial neutrophilic infiltrates and necrosis (Figure

5, group B; average lesion score 1.7 ± 0.2). Thus, the presence of IFN- γ receptor on hematopoietic cells determined lesion severity even when somatic cells lack the IFN- γ receptor.

IFN- γ blocks γ HV68 infection and replication in SMCs and non-SMCs from the aorta. Since γ HV68 replicates in vascular SMCs within arteritic lesions (1, 2) and IFN- γ prevents arteritis, we tested the hypothesis that IFN- γ has antiviral effects in primary aortic cells. SMCs and non-SMCs isolated from mouse aortas were treated with IFN- γ for 48 hours and then infected with γ HV68. IFN- γ reduced virus induced cytopathic effect (Figure 6, a and b). Using dual-label immunofluorescence, we found that IFN- γ decreased the percentage of both non-SMCs and SMCs expressing viral antigen ($P < 0.005$; Figure 6, c, d, and g). IFN- γ treatment significantly decreased the percentage of cells with intranuclear viral capsids as determined by EM (Figure 6, e, f, and g). Thus IFN- γ has antiviral effects in both SMCs and other primary cells derived from the aorta.

Discussion

We found that the immune system functions less efficiently in the elastic media of the great vessels than in small vessels, the adventitia of the great vessels, or visceral organs such as spleen, liver, and lung. For this reason the immune system, rather than viral interactions with cellular receptors, determines tropism of γ HV68 for the great elastic arteries during chronic disease. These findings make the important point that chronic diseases of the great vessels may be due to immunoprivilege of the elastic media, resulting in persistence of pathogens or antigens in this site, in turn leading to chronic inflammation. Lack of effective immune clearance was independent of IFN- γ , although IFN- γ prevented the establishment of medial infection and regulated the severity of disease once infection was established. This essential protective role of IFN- γ contrasts with studies emphasizing promotion of vascular pathology by IFN- γ (reviewed in ref. 7), underscoring the fact that the nature of the vascular insult determines whether specific immune functions are helpful or harmful.

While many mechanisms could explain immunoprivilege within the vasculature, the failure of T cells and macrophages to enter the elastic media suggests one potential mechanism: limited access of effector cells. Lack of immune cell access to the elastic media may be a general contributor to chronic vascular disease and is seen in other

circumstances. For example, injection of mice with splenocytes stimulated in vitro with syngeneic microvascular SMCs resulted in medial and adventitial infiltrates in arterioles and venules, but not in elastic arteries (19). Interestingly, murine cytomegalovirus (MCMV) also induces chronic elastic arteritis with a lack of medial infiltrates




Figure 4

Incidence and severity of chronic elastic arteritis in wild-type mice infected with high doses of γ HV68 with or without transient depletion of IFN- γ . Mice were infected with the indicated doses of γ HV68 with or without depletion of IFN- γ and evaluated for arteritis on H&E-stained sections 5.5 to 10 weeks after infection. (a) The incidence of arteritis in various groups is presented. Seven of 19 mice infected with 17 to 20×10^7 PFU γ HV68 died within 7 days. Twelve of 14 mice infected with 34 to 50×10^7 PFU died within 7 days. Data for transient depletions are from two independent experiments. Data for no depletions are from 4–5 independent experiments. ^A $P = 0.0062$; ^B $P = 0.0043$. (b) Lesion at the aortic base of a 129Ev/Sv mouse sacrificed 10 weeks after infection with 10^8 PFU γ HV68. Lesions in transiently IFN- γ -depleted 129Ev/Sv mice have similar histology (lesion scores for 25 mice from both groups combined = 2.0 ± 0.2 ; $P < 0.0001$ for scores compared with IFN- γ R^{-/-} mice, $P = 0.001$ compared with chronically depleted wild-type mice). (c) High-power view of boxed region in b. (d) Lesion at the aortic base of a chronically IFN- γ -depleted 129Ev/Sv mouse sacrificed 6 weeks after infection with 5×10^7 PFU γ HV68 (lesion scores for four mice = 4.0 ± 0.7). Arteritis in IFN- γ R^{-/-} mice has similar histology (lesion scores for 16 mice = 5.0 ± 0). (e) High-power view of boxed region in d. The black lines show the boundaries of the media. Adv, adventitia; M, media; I, intima; L, lumen; V, aortic valve. ND, not determined.




Figure 5

IFN- γ prevents chronic elastic arteritis by effects on somatic cells but regulates the nature of the pathology by effects on hematopoietic cells. Reciprocal bone marrow reconstitutions were performed between IFN- γ R^{-/-} and 129Ev/Sv mice, and arteritis was evaluated for 12 weeks after γ HV68 infection. Numbers above bars represent the number of mice within a group. Boxed numbers represent average lesion scores for a group. Data are pooled from four independent experiments. For incidence of disease, $P = 0.0001$ comparing group A with D, and $P = 0.0004$ comparing group B with C. For severity of disease, $P < 0.0001$ comparing groups A and B. NA, lesion scores not applicable.

despite the presence of viral antigen in the media (6). Similarly, viral persistence in the kidney may be due to limited entry of CD8 lymphocytes into the kidney despite its rich vascular supply (20, 21). Thus, the failure of immune cells to enter certain anatomic sites may be a significant contributor to immunoprivilege. Perhaps medial elastic laminae, especially in those portions of the media not penetrated by the vasa vasorum, provide a significant barrier to entry of lymphocytes and macrophages. The process of immune cell entry into the media could be regulated also by cytokines (for example IFN- γ) by induction of cell adhesion molecules or proteases critical for tissue penetration (22, 23).

One potential problem with extending observations in the mouse system to the situation in human arteries is the lack of vasa vasorum in mice. However, the inner 29 lamellae of the human aorta do not have vasa vasorum, and the human abdominal aorta lacks vasa vasorum completely (24–26). Consistent with this, there can be striking sparing of the media of the human aorta even when extensive adventitial inflammation, including germinal center formation, is present (27–29). Even when lymphocytes enter the media of

the human great vessels, a segregation of effector cells between layers of the great vessels has been reported, with activated IFN- γ -secreting T cells being excluded from the media in giant cell arteritis (30). Thus, findings in both the mouse and human are consistent with immunoprivilege of the elastic media based on restricted access of immune effector cells.

While intrinsic properties of the elastic media contribute to inefficient clearance of viral infection, viral factors may also play a role. γ HV68 encodes a secreted protein (M3) that binds CC chemokines such as RANTES, MCP, and MIP1 α , as well as the C and CXXXC chemokines lymphotactin and fractalkine, resulting in failure of chemokine-receptor signaling (31, 32). The virus also encodes the secreted γ HV68 regulator of complement activation (RCA) protein that inhibits both classical and alternative complement activation (33). Expression of M3 or the γ HV68 RCA protein in the media could influence mononuclear cell infiltration. Notably, the possible contribution of viral factors to arteritis does not invalidate our conclusion that the elastic media of the great vessels is an immunoprivileged site, since γ HV68 is

cleared from other organs and other sites in the vasculature despite the contribution of such virulence factors.

IFN- γ action during acute and chronic elastic arteritis and cellular sites of IFN- γ action. Together with data presented here for the great vessels, findings in the salivary gland and CNS show that IFN- γ is a key determinant of control of persistent virus infection in multiple pathogen and organ systems, especially when persistent infection of an immunoprivileged site is involved (6, 34–39). Since IFN- γ determined both the incidence and severity of arteritis, we defined the cellular targets for these actions. IFN- γ has effects on somatic cells and, by actions on hematopoietic cells, immune modulatory activities (reviewed in refs. 40 and 41). IFN- γ determined the incidence of arteritis by actions on somatic cells since IFN- γ R^{-/-} recipients of wild-type bone marrow developed arteritis, while wild-type recipients of IFN- γ R^{-/-} bone marrow did not. Since studies using transient depletion of IFN- γ showed that IFN- γ acts early during infection, we conclude that IFN- γ action on somatic cells during acute infection determines the incidence of chronic elastic arteritis. Given this, we tested the hypothesis that IFN- γ has antiviral effects on primary vascular SMCs. IFN- γ significantly protected against γ HV68 infection in both primary SMCs and non-SMCs from mouse aortas. This effect of IFN- γ on somatic cells could well contribute to the role of IFN- γ in controlling the incidence of chronic elastic arteritis.

Interestingly, the presence of 129Ev/Sv derived hematopoietic cells was associated with less severe arteritis in IFN- γ R^{-/-} recipient mice, a finding most consistent with a hematopoietic cell-specific immunoregulatory action of IFN- γ . Similar to the experiments presented here, IFN- γ responsiveness of both somatic and hematopoietic cells is required for resistance to *Toxoplasma gondii*, which, like γ HV68, invades somatic cells as well as macrophages (42). Thus, regulation of infection by the action of IFN- γ on both somatic and hematopoietic cells is a general finding.

Implications. Our results suggest that lesion distribution in chronic vascular disease may result in large part from





Figure 6 IFN- γ inhibits γ HV68 infection of primary aortic cells. IFN- γ -treated and untreated intimal/adventitial and medial cultures were evaluated 3 days after infection. In **c** and **d** nuclei were stained blue, muscle actin was stained red, and viral antigen was stained green. Uninfected cultures and infected cultures stained with control Ab's demonstrated no viral antigen staining. (**a** and **b**) Phase-contrast microscopy of infected intimal/adventitial cultures with or without IFN- γ . (**c**) Representative field of infected medial cultures without IFN- γ treatment. (**d**) Representative field of infected medial cultures treated with IFN- γ . (**e**) EM of a cell with nuclear capsids from an infected, untreated medial culture. (**f**) EM of a cell without nuclear capsids from an infected, IFN- γ -treated medial culture. (**g**) Multiple fields were evaluated by dual immunofluorescence for viral antigen and muscle actin (three experiments) or by electron microscopy (two experiments). IF, immunofluorescence. The numbers above the bars represent the number of cells counted. For comparing results with or without IFN- γ treatment by IF, ^A $P < 0.005$, ^B $P < 0.0002$. For comparing results with or without IFN- γ by EM, ^A $P = 0.0024$, ^B $P = 0.0013$.

research grant from the Monsanto-Searle Biomedical Agreement to SHS and by NIH grants CA43143, CA52004 and CA58524 to SHS, and CA74730, HL60090 and AI39616 to HWV.



properties intrinsic to different vessels and different layers of the great vessels that prevent efficient immune clearance of foreign or self antigens. Given this, it is clear that strategies for blunting the inflammatory response as a way to control chronic vascular disease may well have a price; continued persistence of the inciting agent or antigen. This may be an especially severe problem for herpesvirus infection since these viruses can go latent and reactivate at a later time to cause disease during immunosuppression. This emphasizes the importance of defin-

ing the underlying cause of vascular disease as part of the decision process leading to trials of blockade of immune and inflammatory function in these diseases.

Acknowledgments

We would like to thank Dr. Bob Schreiber, Dr. Wayne Yokoyama, and Dr. Jeff Saffitz for discussions and assistance throughout this project. We also thank members of Dr. David Leib's lab and members of the Speck and Virgin labs for helpful discussions. This work was supported in part by a

- Weck, K.E., et al. 1997. Murine gammaherpesvirus 68 causes large vessel arteritis in mice lacking interferon-gamma responsiveness: a new model for virus induced vascular disease. *Nat. Med.* **3**:1346-1353.
- Dal Canto, A.J., Virgin, H.W., and Speck, S.H. 2000. Ongoing viral replication is required for gammaherpesvirus 68-induced vascular damage. *J. Virol.* **74**:11304-11310.
- Alber, D.G., Powell, K.L., Vallance, P., Goodwin, D.A., and Grahame-Clarke, C. 2000. Herpesvirus infection accelerates atherosclerosis in the apolipoprotein E-deficient mouse. *Circulation.* **102**:779-785.
- Nagano, H., et al. 1997. Interferon-gamma deficiency prevents coronary arteriosclerosis but not myocardial rejection in transplanted mouse hearts. *J. Clin. Invest.* **100**:550-557.
- Gupta, S., et al. 1997. IFN-gamma potentiates atherosclerosis in apoE knock-out mice. *J. Clin. Invest.* **99**:2752-2761.
- Presti, R.M., Pollock, J.L., Dal Canto, A.J., O'Guin, A.K., and Virgin, H.W. 1998. Interferon-gamma regulates acute and latent murine cytomegalovirus infection and chronic disease of the great vessels. *J. Exp. Med.* **188**:577-588.
- Tellides, G., et al. 2000. Interferon-gamma elicits arteriosclerosis in the absence of leukocytes. *Nature.* **403**:207-211.
- Clambey, E.T., Virgin, H.W., and Speck, S.H. 2000. Disruption of the murine gammaherpesvirus 68 M1 open reading frame leads to enhanced reactivation from latency. *J. Virol.* **74**:1973-1984.
- Van Dyk, L.F., Virgin, H.W., and Speck, S.H. 2000. The murine gammaherpesvirus 68 v-cyclin is a

- critical regulator of reactivation from latency. *J. Virol.* **74**:7451-7461.
10. Muller, U., et al. 1994. Functional role of type I and type II interferons in antiviral defense. *Science.* **264**:1918-1921.
 11. Stabell, E.C., and Olivo, P.D. 1992. Isolation of a cell line for rapid and sensitive histochemical assay for the detection of herpes simplex virus. *J. Virol. Methods.* **38**:195-204.
 12. Virgin, H.W., and Tyler, K.L. 1991. Role of immune cells in protection against and control of reovirus infection in neonatal mice. *J. Virol.* **65**:5157-5164.
 13. Austyn, J.M., and Gordon, S. 1981. F4/80, a monoclonal antibody directed specifically against the mouse macrophage. *Eur. J. Immunol.* **11**:805-815.
 14. Huang, S., et al. 1993. Immune response in mice that lack the interferon-gamma receptor. *Science.* **259**:1742-1745.
 15. Buchmeier, N.A., and Schreiber, R.D. 1985. Requirement of endogenous interferon-gamma production for resolution of *Listeria monocytogenes* infection. *Proc. Natl. Acad. Sci. USA.* **82**:7404-7408.
 16. Heise, M.T., and Virgin, H.W. 1995. The T cell independent role of IFN-gamma and TNF-alpha in macrophage activation during murine cytomegalovirus and herpes simplex virus infection. *J. Virol.* **69**:904-909.
 17. Ramos, K.S., and Cox, L.R. 1993. Aortic endothelial and smooth muscle cell cultures. In *Methods in toxicology*. A. Tyson and J.M. Frazier, editors. Academic Press. New York, New York, USA. 159-168.
 18. Gunther, S., Alexander, R.W., Atkinson, W.J., and Gimbrone, M.A.J. 1982. Functional angiotensin II receptors in cultured vascular smooth muscle cells. *J. Cell. Biol.* **92**:289-298.
 19. Hart, M.N., Tassell, S.K., Sadewasser, K.L., Schelper, R.L., and Moore, S.A. 1985. Autoimmune vasculitis resulting from in vitro immunization of lymphocytes to smooth muscle. *Am. J. Pathol.* **119**:448-455.
 20. Ahmed, R., Morrison, L.A., and Knipe, D.M. 1996. Persistence of viruses. In *Fields virology*. B.N. Fields, D.M. Knipe, and P.M. Howley, editors. Lipincott-Raven. Philadelphia, Pennsylvania, USA. 219-249.
 21. Ando, K., Guidotti, L.G., Cerny, A., Ishikawa, T., and Chisari, F.V. 1994. CTL access to tissue antigen is restricted in vivo. *J. Immunol.* **153**:482-488.
 22. Galis, Z.S., Sukhova, G.K., Lark, M.W., and Libby, P. 1994. Increased expression of matrix metalloproteinases and matrix degrading activity in vulnerable regions of human atherosclerotic plaques. *J. Clin. Invest.* **94**:2493-2503.
 23. Galis, Z.S., et al. 1994. Cytokine-stimulated human vascular smooth muscle cells synthesize a complement of enzymes required for extracellular matrix digestion. *Circ. Res.* **75**:181-189.
 24. Wolinsky, H., and Glagov, S. 1967. A lamellar unit of aortic medial structure and function in mammals. *Circ. Res.* **20**:99-111.
 25. Wolinsky, H., and Glagov, S. 1967. Nature of species differences in the medial distribution of aortic vasa vasorum in mammals. *Circ. Res.* **20**:409-421.
 26. Wolinsky, H., and Glagov, S. 1969. Comparison of abdominal and thoracic aortic medial structure in mammals. Deviation of man from the usual pattern. *Circ. Res.* **25**:677-686.
 27. Ramshaw, A.L., and Parums, D.V. 1990. Immunohistochemical characterization of inflammatory cells associated with advanced atherosclerosis. *Histopathology.* **17**:543-552.
 28. Parums, D.V., Dunn, D.C., Dixon, A.K., and Mitchinson, M.J. 1990. Characterization of inflammatory cells in a patient with chronic periaortitis. *Am. J. Cardiovasc. Pathol.* **3**:121-129.
 29. Parums, D.V. 1990. The spectrum of chronic periaortitis. *Histopathology.* **16**:423-431.
 30. Wagner, A.D., Bjornsson, J., Bartley, G.B., Goronzy, J.J., and Weyand, C.M. 1996. Interferon-gamma-producing T cells in giant cell vasculitis represent a minority of tissue-infiltrating cells and are located distant from the site of pathology. *Am. J. Pathol.* **148**:1925-1933.
 31. Parry, B.C., et al. 2000. A broad spectrum secreted chemokine binding protein encoded by a herpesvirus. *J. Exp. Med.* **191**:573-578.
 32. van Berkel, V., et al. 2000. Identification of a gammaherpesvirus selective chemokine binding protein that inhibits chemokine action. *J. Virol.* **74**:6741-6747.
 33. Kapadia, S.B., Molina, H., van Berkel, V., Speck, S.H., and Virgin, H.W. 1999. Murine gammaherpesvirus 68 encodes a functional regulator of complement activation. *J. Virol.* **73**:7658-7670.
 34. Jonjic, S., Pavic, I., Lucin, P., Rukavina, D., and Koszinowski, U.H. 1990. Efficacious control of cytomegalovirus infection after long-term depletion of CD8+ T lymphocytes. *J. Virol.* **64**:5457-5464.
 35. Jonjic, S., Mutter, W., Weiland, F., Reddehase, M.J., and Koszinowski, U.H. 1989. Site-restricted persistent cytomegalovirus infection after selective long-term depletion of CD4+ T lymphocytes. *J. Exp. Med.* **169**:1199-1212.
 36. Lucin, P., Pavic, I., Polic, B., Jonjic, S., and Koszinowski, U.H. 1992. Gamma interferon-dependent clearance of cytomegalovirus infection in salivary glands. *J. Virol.* **66**:1977-1984.
 37. Tishon, A., Lewicki, H., Rall, G., Von Herath, M., and Oldstone, M.B.A. 1995. An essential role for type 1 interferon-gamma in terminating persistent viral infection. *Virology.* **212**:244-250.
 38. von Herrath, M.G., Coon, B., and Oldstone, M.B. 1997. Low-affinity cytotoxic T-lymphocytes require IFN-gamma to clear an acute viral infection. *Virology.* **229**:349-359.
 39. Planz, O., et al. 1997. A critical role for neutralizing-antibody-producing B cells, CD4(+) T cells, and interferons in persistent and acute infections of mice with lymphocytic choriomeningitis virus: implications for adoptive immunotherapy of virus carriers. *Proc. Natl. Acad. Sci. USA.* **94**:6874-6879.
 40. Biron, C. 1994. Cytokines in the generation of immune responses to, and resolution of, virus infection. *Curr. Opin. Immunol.* **6**:530-538.
 41. Stark, G.R., Kerr, I.M., Williams, B.R., Silverman, R.H., and Schreiber, R.D. 1998. How cells respond to interferons. *Annu. Rev. Biochem.* **67**:227-264.
 42. Yap, G.S., and Sher, A. 1999. Effector cells of both nonhemopoietic and hemopoietic origin are required for interferon (IFN)-gamma- and tumor necrosis factor (TNF)-alpha-dependent host resistance to the intracellular pathogen, *Toxoplasma gondii*. *J. Exp. Med.* **189**:1083-1092.