



Published in final edited form as:

Tetrahedron. 2007 August 13; 63(33): 7977–7984.

Neurosteroid Analogues. 13. Synthetic methods for the preparation of 2 β -hydroxygonane derivatives as structural mimics of *ent*-3 α -hydroxysteroid modulators of GABA_A receptors

Cunde Wang^a, Nigam P. Rath^b, and Douglas F. Covey^{a,*}

^a Department of Molecular Biology and Pharmacology, Washington University School of Medicine, Campus Box 8103, 660 S. Euclid Ave., St. Louis, MO 63110, USA

^b Department of Chemistry and Biochemistry, University of Missouri St. Louis, 8001 Natural Bridge Road, St. Louis, MO 63121, USA

Abstract

Many different 3 α -hydroxysteroids in the androstane and pregnane steroid series enhance the actions of γ -aminobutyric acid (GABA) at GABA type-A (GABA_A) receptors in the mammalian central nervous system. Recent studies have shown that (3 α ,5 α)-3-hydroxyandrostane-17-one (androsterone) is less active at these receptors than its enantiomer *ent*-androsterone. Further structure–activity relationship (SAR) studies are needed to explore the structural features of *ent*-androsterone that are important for its enhanced action at these receptors. Molecular modeling shows that 2 β -hydroxysteroids are similar in three-dimensional shape to the enantiomers of 3 α -hydroxysteroids. The development of synthetic methods to gain access to C₁₇-substituted analogues of 2 β -hydroxygonanes for SAR studies is demonstrated with the synthesis of (2 β ,5 α ,13 β ,14 β)-2-hydroxygonane-17-one.

Keywords

2 β -hydroxygonanes; abnormal Beckmann rearrangement; neurosteroids; phenanthrenes

I. Introduction

The steroid (3 α ,5 α)-3-hydroxypregnan-20-one (allopregnanolone, Figure 1) and many analogues of it are known to be potent enhancers of γ -aminobutyric acid (GABA) at GABA Type A (GABA_A) receptors.^{1–4} These neuroactive GABAergic steroids have activity as general anesthetics, anticonvulsants, sedative hypnotics and anxiolytics; and there is considerable current interest in the development of new analogues as pharmaceuticals having these activities.

As part of our ongoing studies of the enantioselectivity of neurosteroid action at GABA_A receptors, we recently investigated the enantioselectivity for (3 α ,5 α)-3-hydroxyandrostane-17-one (androsterone, Figure 1) effects at these receptors.⁵ Androsterone has only weak actions on GABA_A receptor function^{6,7} and, based on our previous results from enantioselectivity

*Corresponding author. Telephone: 314-362-1726, FAX: 314-362-7058, Email: dcovey@wustl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

studies of allopregnanolone,⁸ we expected that *ent*-androsterone (Figure 1) would have even weaker actions than androsterone. Unexpectedly, we found that *ent*-androsterone was more active than androsterone. Moreover, a 17-spiroepoxide derivative of *ent*-androsterone (Figure 1) was shown to have actions comparable to those of allopregnanolone.⁵

Additional steroid analogues are needed for future structure–activity relationship (SAR) studies of *ent*-androgen action at GABA_A receptors. In this regard, we were intrigued by the structural similarity between 2 β -hydroxysteroids and the enantiomers of 3 α -hydroxysteroids. Figure 2A shows the structure of (2 β ,5 α ,13 β ,14 β)-2-hydroxygonan-17-one (**1**) and *ent*-androsterone. Figure 2B shows a three-dimensional overlay of molecular models of steroid **1** and *ent*-androsterone. Both steroids are shown with their β face down because in this orientation the compounds most closely resemble the highly active 3 α -hydroxysteroid allopregnanolone. As presented in Figure 2, each of the four steroid rings of one compound is proximate to the corresponding ring of the other compound. The O₂ and O₃ atoms of the two molecules are 0.68 Å apart and the O₁₇ atoms are 0.54 Å from each other in this overlay.

Two additional design criteria were considered in the decision to select steroid **1** as an initial potential structural mimic of *ent*-androsterone. Previous SAR studies have shown that having methyl groups on the steroid α face decreases the activity of 3 α -hydroxysteroid modulators of GABA_A receptors.^{9,10} When aligned as shown in Figure 2B, the C₁₈ and/or C₁₉ methyl groups of steroids in the androstane, estrane (19-norandrostane) and 18-norandrostane series would be located below the plane of the steroid rings (i.e. they would occupy positions similar to those occupied by methyl groups on the α face of 3 α -hydroxysteroids, and therefore could be expected to negatively affect pharmacological activity). By choosing a steroid in the gonane class as a synthetic target, worries about unfavorable steric effects of these methyl groups is avoided.

The decision to have the 13 β ,14 β *cis* C,D-ring fusion present in steroid **1** was made because this *cis* ring fusion places the C₁₇ carbonyl group further above the plane of the steroid rings than it would be in the corresponding 13 β ,14 α *trans* ring fusion. This was considered to be desirable since previous SAR studies indicate that the D-ring hydrogen bonding groups needs to be above the plane of the steroid rings for high pharmacological activity.^{1,6,7}

2. Results and Discussion

The starting material, (3 β ,5 α)-3-hydroxyestrane-17-one (**3**), was prepared in three steps by known methodology^{11,12} from commercially available 19-nortestosterone (**2**) in 85% total yield (Scheme 1). Steroid **3** was brominated selectively in the C₁₆ position using CuBr₂^{13,14} to give compound **4** in 69% yield (Scheme 2). HBr was eliminated from steroid **4** using LiBr/Li₂CO₃¹⁵ to give compounds **5a** (33%) and **5b** (39%). For characterization purposes, some spectroscopic data was collected on each of these products after their separation by chromatography. For synthetic purposes, products **5a** and **5b** were not separated and the mixture was subjected to catalytic hydrogenation to give compound **6** in 87% yield. The 17-keto group was ketalized in the standard way¹⁶ to afford compound **7** in 85% yield. Oxidation of the 3 β -hydroxyl group of compound **7** using PCC in the presence of NaOAc¹⁷ gave compound **8** (75%).

When treated with 10% ethanolic KOH at room temperature for 24 h, compound **8** underwent an aldol condensation¹⁸ to give compound **9** in 74% yield (Scheme 3). Reduction of the 3-ketone group with NaBH₄ at room temperature led to the 3 β -hydroxysteroid **10** (99%), and acetylation of the hydroxyl group gave the acetate derivative **11** in 90% yield. Compound **11** was treated with ozone in the usual manner¹⁸ to give steroid **12** (80%). Samarium(II) iodide-THF mediated reductive removal of the 3-acetyloxy group from compound **12** gave 2-

ketosteroid **13** (41%), 2 β -hydroxysteroid **14a** (20%) and 2 α -hydroxysteroid **14b** (26%). Reduction of compound **13** with K-Selectride in THF at -78 °C gave additional amounts of compound **14a** (85%).

The hydroxyl group of 2 β -hydroxysteroid **14a** was reacted with acetic anhydride and pyridine in the presence of a catalytic amount of 4-dimethylaminopyridine to afford 2-(acetyloxy)steroid **15** (92%) after 5 min at room temperature. The 17-ketal of compound **15** was readily removed using *p*-TsOH in acetone at room temperature for 12 h to give, in quantitative yield, steroid **16**. The reaction of compound **16** with NH₂OH·HCl and NaOAc gave oxime **17** (97%). Oxime **17** underwent an abnormal Beckmann rearrangement¹⁹ upon treatment with CH(OCH₃)₃/TFA in THF to give carbonitrile **18** (79%). Ozonolysis of compound **18** at -78 °C yielded ketone-nitrile **19** (58%). Treatment of compound **19** with SmI₂ in THF and irradiation with a 500 W lamp²⁰ at 0–10 °C for 8 h gave crude 13-hydroxysteroid **20**. After isolation, crude product **20** was again treated with SmI₂ in THF at room temperature for 10 min to give compound **21** (17%, from compound **19**). Finally, base-catalyzed hydrolysis of the 3-acetyloxy group of compound **21** and column chromatography to remove the presumed 13 α ,14 β stereoisomer gave compound **1** (82%). The structure of steroid **1** was confirmed by single crystal X-ray diffraction analysis (Figure 3).²¹

Using methods reported previously,²² the actions of compound **1** as a modulator of GABA_A receptor function were compared to those of *ent*-androsterone. Whereas *ent*-androsterone allosterically displaced 50% of [³⁵S]-*t*-butylbicyclopophosphorothionate bound to the picrotoxin binding site on GABA_A receptors at a concentration of 0.31 μ M,⁵ no displacement of [³⁵S]-*t*-butylbicyclopophosphorothionate occurred at concentrations up to 30 μ M of compound **1**. *ent*-Androsterone (10 μ M) enhances 2 μ M GABA-mediated chloride currents at rat $\alpha_1\beta_2\gamma_{2L}$ GABA receptors expressed in *Xenopus laevis* oocytes.⁵ Compound **1** at concentrations up to 10 μ M does not enhance 2 μ M GABA-mediated chloride currents at these expressed receptors. Finally, *ent*-androsterone causes 50% loss of righting reflex for tadpoles at a concentration of 3.38 μ M.⁵ By contrast, compound **1** did not cause loss of righting reflex in tadpoles at a concentration as high as 10 μ M. Thus, despite the similarities in the overall shapes of the two steroids, only *ent*-androsterone is effective as a positive modulator of GABA_A receptors at concentrations below 10 μ M.

3. Conclusion

We have described the synthesis and crystal structure of (2 β , 5 α , 13 β , 14 β)-2-hydroxygonan-17-one (**1**) from commercially available 19-nortestosterone. Novel features of the synthetic route developed include the first use of a SmI₂-promoted reaction to remove a 3-acetyloxy group from a 2-keto-3-(acetyloxy)steroid, the first report of the abnormal Beckmann rearrangement on steroids having the 13 β ,14 β *cis* C,D-ring fusion and the first use of SmI₂-promoted reactions to prepare 18-norsteroids having a 13 β ,14 β *cis* C,D-ring fusion. Omission of the part of the reported synthetic sequence that was used to construct the 13 β ,14 β *cis* C,D-ring fusion would also allow other 2 β -hydroxygonane analogues with the 13 β ,14 α *trans* C,D-ring fusion to be prepared using methods reported previously for the synthesis of (13 β ,14 α)-18-norsteroids.^{20, 23}

4. Experimental

4.1 General Methods

Melting points were determined on a Kofler micro hot stage and are uncorrected. NMR spectra were recorded in CDCl₃ at 300 MHz (¹H) or 75 MHz (¹³C). IR spectra were recorded as films on a NaCl plate. Elemental analyses were carried out by M-H-W Laboratories, Phoenix, AZ.

4.1.1 (3 β ,5 α ,16 β β)-16-Bromo-3-hydroxyestr-17-one (4)—A mixture of known compound **3**²⁴ (1.40 g, 5.07 mmol) and copper bromide (3.06 g, 13.7 mmol) in methanol (35 mL) was stirred under reflux for 12 h. Then the solvent was removed under reduced pressure to give a residue and water (25 mL) was added. The product was extracted with CH₂Cl₂ (25 mL), the organic layer was washed with water (2 \times 10 mL), brine (20 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; CH₂Cl₂/EtOAc, 10:1) to give compound **4** (1.24 g, 69%) as white crystals: mp 108–110 °C (EtOAc-hexanes); [α]_D²⁰ = +59.1 (c = 0.30, CHCl₃); ¹H NMR δ 4.56 (m, 1H), 3.56 (brs, 1H), 3.05 (s, 1H), 2.18 (m, 2H), 0.91 (s, 3H); ¹³C NMR δ 13.6, 24.4, 27.7, 29.0, 31.7, 32.5, 33.3, 34.9, 39.2, 40.4, 42.4, 45.4, 46.0, 46.5, 47.2, 47.3, 69.3, 213.0; IR ν_{\max} 3400, 2920, 2853, 1748, 1448, 1023 cm⁻¹. Anal. Calcd. For C₁₈H₂₇BrO₂: C 60.85, H 7.66; Found C 60.78, H 7.87.

4.1.2 (3 β ,5 α)-3-Hydroxyestr-14-en-17-one (5a) and (3 β ,5 α ,14 β β)-3-Hydroxyestr-15-en-17-one (5b)—The mixture of compound **4** (0.65 g, 1.8 mmol), LiBr (0.42 g, 5.0 mmol) and Li₂CO₃ (0.35 g, 5.0 mmol) in DMF (30 mL) was stirred under reflux for 4 h. Afterwards, the reaction mixture was chilled to room temperature and was poured into water (50 mL), the mixture was extracted with EtOAc (2 \times 30 mL), the combined organic extract was washed with saturated NaHCO₃ (10 mL), water (10 mL), brine (10 mL) and dried over Na₂SO₄. The solvent was removed under reduced pressure. The products were purified by column chromatography (silica gel; CH₂Cl₂/EtOAc, 10:1) to give compound **5a** (0.17 g, 33%; R_f = 0.47, CH₂Cl₂/EtOAc, 10:1) and compound **5b** (0.20 g, 39%; R_f = 0.33, CH₂Cl₂/EtOAc, 10:1). Compound **5a** was obtained as an oil: ¹H NMR δ 5.47 (d, *J* = 1.8 Hz, 1H), 3.59 (m, 1H), 2.86 (m, 2H), 2.29 (s, 1H), 1.12 (s, 3H); ¹³C NMR δ 19.7, 25.2, 27.8, 28.0, 32.9, 33.0, 35.4, 40.8, 41.0, 41.3, 43.0, 46.5, 48.6, 50.7, 70.0, 112.3, 152.8, 222.7; IR ν_{\max} 3400, 3059, 2921, 2855, 1744, 1641, 1449, 1046 cm⁻¹.

Compound **5b** was obtained as an oil: ¹H NMR δ 7.64 (dd, *J* = 6.0, 2.4 Hz, 1H), 6.15 (dd, *J* = 6.0, 2.4 Hz, 1H), 3.57 (m, 1H), 2.63 (m, 1H), 1.11 (s, 3H); ¹³C NMR δ 21.3, 24.8, 28.0, 30.1, 31.7, 33.4, 35.5, 39.7, 39.8, 40.8, 42.9, 47.3, 47.5, 54.4, 70.2, 132.2, 163.3, 215.2. IR ν_{\max} 3400, 2921, 2854, 1705, 1585, 1449, 1052 cm⁻¹. Compound **5a** and **5b** were not purified further and were used in the next step directly.

4.1.3 (3 β ,5 α ,14 β)-3-Hydroxyestr-17-one (6)—A mixture of compounds **5a** and **5b** (3.90 g, 14.2 mmol) was dissolved in EtOAc (50 mL) and hydrogenated under H₂ (50 psi) in the presence of 5% Pd-C (400 mg) at room temperature overnight. The reaction mixture was filtered through a pad of Celite 545[®] to remove catalyst and the solvent was removed under reduced pressure. The residue was purified by flash column chromatography (silica gel; EtOAc/hexanes, 1:1) to give compound **6** (3.81 g, 97%) as white crystals: mp 116–118 °C (EtOAc); [α]_D²⁰ = +92.0 (c = 0.27, CHCl₃); ¹H NMR δ 3.56 (m, 1H), 3.27 (brs, 1H), 2.45 (dd, *J* = 19.2, 8.1 Hz, 1H), 2.13 (dd, *J* = 19.2, 9.9 Hz, 1H), 1.07 (s, 3H); ¹³C NMR δ 18.1, 19.3, 24.6, 27.6, 27.7, 30.0, 33.4, 35.1, 35.6, 39.1, 40.0, 40.6, 42.7, 46.2, 47.5, 47.8, 69.6, 222.8; IR ν_{\max} 3401, 2917, 2854, 1732, 1448, 1048 cm⁻¹. Anal. Calcd. For C₁₈H₂₈O₂: C 78.21, H 10.21; Found C 78.42, H 9.97.

4.14 (3 β ,5 α ,14 β)-3-Hydroxyestr-17-one cyclic-(1,2-ethanediyl acetal) (7)—A solution of compound **6** (0.41 g, 1.49 mmol), ethylene glycol (2.2 g, 35 mmol) and *p*-TsOH (60 mg, 0.32 mmol) in benzene (35 mL) was refluxed using a Dean-Stark trap for 12 h. The mixture was cooled to room temperature, diluted with ether (50 mL), washed with saturated NaHCO₃ (2 \times 20 mL) and brine (2 \times 20 mL). The organic phase was dried over Na₂SO₄, and the solvent was removed under reduced pressure to give a crude product which was purified by column chromatography (silica gel; EtOAc/hexanes, 1:1) to yield compound **7** (406 mg, 85%) as an oil: [α]_D²⁰ = +73.9 (c = 0.065, CHCl₃); ¹H NMR δ 3.86 (m, 4H), 3.55 (m, 1H),

0.92 (s, 3H); ^{13}C NMR δ 16.1, 19.7, 25.3, 27.8, 28.9, 30.3, 32.1, 33.6, 35.3, 39.4, 39.7, 40.8, 43.0, 45.1, 46.4, 46.8, 63.7, 65.0, 69.9, 120.4; IR ν_{max} 3368, 2920, 2856, 1447, 1367, 1031 cm^{-1} . Anal. Calcd. For $\text{C}_{20}\text{H}_{32}\text{O}_3$: C 74.96, H 10.06; Found C 75.12, H 9.86.

4.1.5 (5 α ,14 β)-estran-3,17-dione cyclic-17-(1,2-ethanediyl acetal) (8)—To a solution of compound **7** (406 mg, 1.27 mmol) in CH_2Cl_2 (25 mL) was added NaOAc (0.46 g, 5.6 mmol) and PCC (0.72 g, 3.35 mmol) at 0°C . The mixture was stirred at 0°C for 2 h and allowed to stand overnight. Then the reaction mixture was diluted with Et_2O (100 mL) and the mixture was filtered through a pad of silica gel. Removal of solvent gave a residue which was purified by column chromatography (silica gel; hexanes/EtOAc, 5:1) to give compound **8** (302 mg, 75%) as white crystals: mp: 85–86 $^\circ\text{C}$ (hexanes/EtOAc); $[\alpha]_{\text{D}}^{20} = +103.9$ ($c = 0.28$, CHCl_3); ^1H NMR δ 3.90 (m, 4H), 2.29 (m, 4H), 0.94 (s, 3H); ^{13}C NMR δ 16.3, 19.8, 25.7, 29.0, 30.0, 30.1, 32.2, 34.2, 39.4, 39.7, 41.1, 43.4, 45.2, 46.1, 46.9, 48.5, 63.9, 65.2, 120.4, 211.6; IR ν_{max} 2935, 2880, 1714, 1470, 1448 cm^{-1} . Anal. Calcd. For $\text{C}_{20}\text{H}_{30}\text{O}_3$: C 75.43, H 9.50; Found C 75.63, H 9.38.

4.1.6 (5 α ,14 β)-2-(Phenylmethylene)estran-3,17-dione cyclic-17-(1,2-ethanediyl acetal) (9)—A mixture of compound **8** (1.60 g, 5.03 mmol), benzaldehyde (1.75 g, 16.5 mmol) and KOH (400 mg, 7.14 mmol) in EtOH (30 mL) and water (5 mL) was stirred at room temperature in the dark for 24 h. Then the reaction mixture was poured into ice-water (50 mL), and the product was extracted with CH_2Cl_2 (2×25 mL), the organic phase was washed with water (15 mL), brine (15 mL) and dried over Na_2SO_4 . The solvent was removed under reduced pressure to give a residue which was purified by column chromatography (silica gel; hexanes/EtOAc, 6:1) to give compound **9** (1.51 g, 74%) as light yellow crystals: mp: 153–155 $^\circ\text{C}$ (EtOAc); $[\alpha]_{\text{D}}^{20} = +43.0$ ($c = 0.33$, CHCl_3); ^1H NMR δ 7.47 (d, $J = 2.1$ Hz, 1H), 7.40 (m, 5H), 3.89 (m, 4H), 3.36 (dd, $J = 16.2, 2.7$ Hz, 1H), 2.63 (dd, $J = 16.8, 4.2$ Hz, 1H), 0.91 (s, 3H); ^{13}C NMR δ 16.4, 19.9, 25.7, 29.0, 29.9, 32.3, 33.2, 33.7, 39.0, 39.1, 40.8, 44.4, 45.3, 47.0, 47.4, 64.0, 65.3, 120.5, 128.3 ($2 \times \text{C}$), 128.5, 130.3 ($2 \times \text{C}$), 135.4, 135.6, 135.8, 201.5; IR ν_{max} 2918, 2860, 1682, 1594, 1572, 1491, 1085, 735 cm^{-1} . Anal. Calcd. For $\text{C}_{27}\text{H}_{34}\text{O}_3$: C 79.76, H 8.34; Found C 79.66, H 8.35.

4.1.7 (3 β ,5 α ,14 β)-3-Hydroxy-2-(Phenylmethylene)estran-17-one cyclic-(1,2-ethanediyl acetal) (10)—To a solution of compound **9** (1.51 g, 3.72 mmol) in THF (40 mL) and MeOH (25 mL) was added NaBH_4 (70 mg, 1.85 mmol) at room temperature. The mixture was stirred for 30 min. Most of solvent was removed under reduced pressure and EtOAc (35 mL) was added to the residue. The organic phase was washed with water (20 mL), brine (20 mL) and dried over Na_2SO_4 . After the solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; hexanes/EtOAc, 3:1) to give compound **10** (1.50 g, 99%) as a thick oil: $[\alpha]_{\text{D}}^{20} = +67.2$ ($c = 0.49$, CHCl_3); ^1H NMR δ 7.34 (m, 5H), 6.57 (s, 1H), 4.12 (m, 1H), 3.88 (m, 4H), 3.10 (dd, $J = 13.5, 2.4$ Hz, 1H), 2.43 (brs, 1H), 0.91 (s, 3H); ^{13}C NMR δ 16.4, 19.9, 25.3, 29.0, 30.3, 31.2, 32.3, 33.2, 39.3, 40.3, 41.4, 44.7, 45.3, 47.0, 48.5, 63.9, 65.2, 72.4, 118.3, 120.6, 125.9, 128.0 ($2 \times \text{C}$), 128.8 ($2 \times \text{C}$), 137.9, 144.1; IR ν_{max} 3435, 2917, 2859, 1445, 1087 cm^{-1} ; Anal. Calcd. For $\text{C}_{27}\text{H}_{36}\text{O}_3$: C 79.37, H 8.88; Found C 79.54, H 8.96.

4.1.8 (3 β ,5 α ,14 β)-3-(Acetyloxy)-2-(phenylmethylene)estran-17-one cyclic-(1,2-ethanediyl acetal) (11)—A mixture of compound **10** (1.50 g, 3.67 mmol), acetic anhydride (4.5 mL, 480 mmol), pyridine (4.5 mL) and 4-dimethylaminopyridine (10 mg, 0.082 mmol) was stirred at room temperature for 5 min. The reaction mixture was poured into ice-water (50 mL) and was extracted with EtOAc (2×30 mL). The combined organic extract was washed with water (2×20 mL), 10% NaHCO_3 (10 mL), brine (20 mL) and dried over Na_2SO_4 . After solvent removed under reduced pressure, the residue was purified by column chromatography

(silica gel; hexanes/EtOAc, 4:1) to give compound **11** (1.48 g, 90%) as white crystals: mp 141–143 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +105.0$ ($c = 0.20$, CHCl_3); $^1\text{H NMR } \delta$ 7.33 (m, 5H), 6.36 (s, 1H), 5.32 (m, 1H), 3.88 (m, 4H), 3.24 (dd, $J = 13.5, 3.3$ Hz, 1H), 2.16 (s, 3H), 0.91 (s, 3H); $^{13}\text{C NMR } \delta$ 16.4, 19.9, 21.2, 25.3, 29.1, 30.3, 31.5, 32.3, 33.1, 39.5, 40.3, 41.0, 41.2, 45.3, 47.0, 48.3, 64.0, 65.3, 73.8, 118.9, 120.6, 126.2, 128.1 ($2 \times \text{C}$), 128.8 ($2 \times \text{C}$), 137.5, 139.3, 170.1; IR ν_{max} 2931, 2860, 1738, 1599, 1241, 1046 cm^{-1} . Anal. Calcd. For $\text{C}_{29}\text{H}_{38}\text{O}_4$: C 77.30, H 8.50; Found 77.50, H 8.61.

4.1.9 (3 β ,5 α ,14 β)-3-(Acetyloxy)estran-2,17-dione cyclic-17-(1,2-ethanediyl acetal) (12)—A solution of compound **11** (1.48 g, 3.29 mmol) in MeOH (60 mL) and EtOAc (30 mL) was treated with ozone at -78 °C until a blue color persisted (ca 30 min). Oxygen was passed through the solution for 20 min until the blue color disappeared, and Me_2S (10 mL) was added to the solution, the reaction mixture was stirred overnight from -78 °C to room temperature. Solvent was removed under reduced pressure and the residue was purified by column chromatography (silica gel; hexanes/EtOAc, 3:1) to give compound **12** (0.98 g, 80%) as white crystals: mp 188–189 °C (hexanes-EtOAc); $[\alpha]_D^{20} = +125.0$ ($c = 0.45$, CHCl_3); $^1\text{H NMR } \delta$ 5.20 (m, 1H), 3.88 (m, 4H), 2.70 (d, $J = 13.2$ Hz, 1H), 2.13 (s, 3H), 0.91 (s, 3H); $^{13}\text{C NMR } \delta$ 16.1, 19.6, 20.3, 25.0, 28.6, 29.8, 32.0, 32.1, 38.5, 38.7, 39.9, 40.6, 43.3, 45.0, 46.5, 48.0, 63.7, 65.0, 75.5, 120.1, 169.5, 203.8; IR ν_{max} 2933, 2863, 1748, 1728, 1238, 1092 cm^{-1} . Anal. Calcd. For $\text{C}_{22}\text{H}_{32}\text{O}_5$: C 70.18, H 8.57; Found C 69.93, H 8.80.

4.1.10 (5 α ,14 β)-estran-2,17-dione cyclic-17-(1,2-ethanediyl acetal) (13), (2 β ,5 α ,14 β)-2-hydroxyestran-17-one cyclic-(1,2-ethanediyl acetal) (14a) and (2 α ,5 α ,14 β)-2-hydroxyestran-17-one cyclic-(1,2-ethanediyl acetal) (14b)—Iodine (2.06 g, 8.12 mmol) in dried THF (30 mL) was added to samarium filings (1.50 g, 10 mmol) by syringe under Ar. The mixture was stirred at room temperature for 30 min to give a black blue solution. Then compound **12** (0.92 g, 2.45 mmol) in dried THF (15 mL) and MeOH (5 mL) was added to the SmI_2 -THF solution and the reaction mixture was stirred for 15 min. The reaction mixture was poured into 10% Na_2CO_3 (70 mL) and was extracted with EtOAc (3×30 mL), the combined organic extract was washed with water (20 mL), brine (20 mL) and dried over Na_2SO_4 . After the solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; $\text{CH}_2\text{Cl}_2/\text{EtOAc}$, 10:1) to give compound **13** (0.32 g, 41%), compound **14a** (0.16 g, 20%) and compound **14b** (0.20 g, 26%).

Compound **13** was obtained as white crystals: mp 143–145 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +91.4$ ($c = 0.45$, CHCl_3); $^1\text{H NMR } \delta$ 3.89 (m, 4H), 2.59 (dd, $J = 15.6, 2.7$ Hz, 1H), 2.35 (m, 2H), 0.91 (s, 3H); $^{13}\text{C NMR } \delta$ 16.2, 19.8, 24.9, 28.8, 30.0, 32.2, 32.6, 33.3, 38.8, 41.0, 41.1, 41.2, 45.0, 45.1, 46.7, 47.7, 63.9, 65.1, 120.4, 211.7; IR ν_{max} 2948, 2858, 1714, 1455, 1088 cm^{-1} . Anal. Calcd. For $\text{C}_{20}\text{H}_{30}\text{O}_3$: C 75.43, H 9.50; Found C 75.29, H 9.70.

Compound **14a** was obtained as white crystals: mp 136–138 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +105.5$ ($c = 0.12$, CHCl_3); $^1\text{H NMR } \delta$ 4.13 (m, 1H), 3.89 (m, 4H), 0.91 (s, 3H); $^{13}\text{C NMR } \delta$ 16.4, 19.9, 25.1, 27.6, 29.1, 30.5, 32.3, 32.5, 33.8, 36.3, 39.7, 40.1, 40.8, 42.6, 45.2, 47.2, 64.0, 65.2, 66.6, 120.8; IR ν_{max} 3436, 2918, 2860, 1305, 1095 cm^{-1} . Anal. Calcd. For $\text{C}_{20}\text{H}_{32}\text{O}_3$: C 74.96, H 10.06; Found C 74.94, H 9.94.

Compound **14b** was obtained as white crystals: mp 141–143 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +81.2$ ($c = 0.07$, CHCl_3); $^1\text{H NMR } \delta$ 3.89 (m, 4H), 3.55 (m, 1H), 0.91 (s, 3H); $^{13}\text{C NMR } \delta$ 16.4, 20.0, 25.3, 29.2, 30.5, 32.0, 32.4, 33.3, 35.4, 39.2, 39.5, 40.0, 41.9, 45.3, 45.8, 47.1, 64.0, 65.3, 71.2, 120.8; IR ν_{max} 3368, 2920, 2857, 1305, 1097, 1031 cm^{-1} . Anal. Calcd. For $\text{C}_{20}\text{H}_{32}\text{O}_3$: C 74.96, H 10.06; Found C 74.92, H 10.18.

14.1.11 (2 β ,5 α ,14 β)-2-hydroxyestran-17-one cyclic-(1,2-ethanediyl acetal) (Compound 14a)—To a solution of compound **13** (406 mg, 1.28 mmol) in dried THF (35 mL) was added 1M K-Selectride (6.5 mL, 6.5 mmol) in THF at -78 °C under N₂ and the mixture was stirred for 1 h at this temperature. Then the reaction was quenched by adding 10% NaOH (8 mL) and 30% hydrogen peroxide (8 mL), and the reaction was continued for other 30 min. The mixture was extracted with EtOAc (3 \times 30 mL). The combined organic phase was washed with water (10 mL), brine (2 \times 10 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; CH₂Cl₂/EtOAc, 10:1) to give compound **14a** (346 mg, 85%) whose properties were identical to those for this product when isolated directly from the SmI₂ reduction.

14.1.12 (2 β ,5 α ,14 β)-2-(Acetyloxy)estran-17-one cyclic-(1,2-ethanediyl acetal) (15)—A mixture of compound **14** (476 mg, 1.49 mmol), acetic anhydride (2 mL), dry pyridine (2 mL) and 4-dimethylaminopyridine (10 mg) was stirred at room temperature for 5 min. The reaction mixture was poured into ice-water (15 mL) and was extracted with EtOAc (2 \times 15 mL). The organic phase was washed successively with saturated NaHCO₃ (5 mL), water (5 mL), 10% aqueous HCl (5 mL), water (5 mL), brine (5 mL) and dried over Na₂SO₄. The solvent was removed under reduced pressure to give oily compound **15** (496 mg, 92%) that was not purified and was directly converted to compound **16**.

14.1.13 (2 β ,5 α ,14 β)-2-(Acetyloxy)estran-17-one (16)—The mixture of compound **15** (496 mg, 1.37 mmol), *p*-TsOH (50 mg, 0.26 mmol) and water (36 mg, 2 mmol) in acetone (15 mL) was stirred at room temperature overnight. After the solvent was removed under reduced pressure, EtOAc (35 mL) was added to the residue. and the organic phase was washed with saturated NaHCO₃ (10 mL), water (10 mL), brine (10 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was purified by re-crystallization (hexanes/EtOAc) to give compound **16** (430 mg, 99%) as white crystals: mp 144–146 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +92.3$ (*c* = 0.12, CHCl₃); ¹H NMR δ 5.10 (s, 1H), 2.44 (dd, *J* = 18.6, 7.2 Hz, 1H), 2.05 (s, 3H), 1.08 (s, 3H); ¹³C NMR δ 18.2, 19.4, 21.1, 24.3, 27.7, 28.0, 29.4, 30.0, 33.1, 33.4, 35.6, 39.2, 40.2, 41.3, 41.9, 47.4, 48.1, 69.6, 170.1, 222.3; IR ν_{\max} 2910, 2850, 1727, 1250, 1020 cm⁻¹. Anal. Calcd. For C₂₀H₃₀O₃: C 75.43, H 9.50; Found C 75.27, H 9.39.

14.1.14 (2 β ,5 α ,14 β)-2-(Acetyloxy)estran-17-one oxime (17)—The mixture of compound **16** (460 mg, 1.47 mmol), hydroxylamine hydrochloride (1.00 g, 14.4 mmol) and NaOAc (1.15 g, 14.0 mmol) in MeOH (35 mL) was refluxed for 1.5 h. The solvent was removed partially under reduced pressure. Water (20 mL) was added to the residue. Then the mixture was extracted with EtOAc (2 \times 20 mL), the combined organic extract was washed with water (20 mL), brine (20 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was purified by re-crystallization (hexanes/EtOAc) to give compound **17** (476 mg, 97%) as white crystals: mp 186–188 °C (hexanes/EtOAc); $[\alpha]_D^{20} = +76.4$ (*c* = 0.89, CHCl₃); ¹H NMR δ 9.44 (s, 1H), 5.10 (s, 1H), 2.06 (s, 3H), 1.18 (s, 3H); ¹³C NMR δ 20.7, 21.5, 21.8, 25.2, 25.3, 28.2, 29.6, 31.0, 32.0, 33.5, 33.8, 39.1, 40.3, 41.7, 42.2, 43.9, 50.3, 70.1, 173.0, 176.4; IR ν_{\max} 3305, 2918, 2857, 1737, 1369, 1242, 937 cm⁻¹. Anal. Calcd. For C₂₀H₃₁NO₃: C 72.04, H 9.37, N 4.20; Found C 72.18, H 9.12, N 4.10.

14.1.15 (1*R*,4*aR*,4*bS*,6*S*,8*aS*,10*aS*)-6-(Acetyloxy)tetradecahydro-2-methylene-1-phenanthrenepropanenitrile (18)—To a solution of compound **17** (476 mg, 1.43 mmol) and trimethyl orthoformate (2.8 mL, 25.6 mmol) in dried THF (35 mL) was added dropwise trifluoroacetic acid (0.26 mL, 35 mmol) at 60 °C for 10 min, and the reaction was continued for another 30 min. Then saturated Na₂CO₃ (4.5 mL) was added, the solvent was partially removed under reduced pressure and EtOAc (50 mL) was added. The organic phase was

washed with saturated NaHCO₃ (10 mL), water (10 mL), brine (10 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; hexanes/EtOAc, 4:1) to give compound **18** (356 mg, 79%) as white crystals: mp 103–105 °C (EtOAc); [α]_D²⁰ = +21.5 (c = 0.13, CHCl₃); ¹H NMR δ 5.10 (s, 1H), 4.74 (t, *J* = 2.1 Hz, 1H), 4.72 (d, *J* = 9.0 Hz, 1H), 2.03 (s, 3H); ¹³C NMR δ 15.3, 21.5, 22.0, 28.2, 29.7, 29.9, 30.2, 31.7, 33.6, 33.7, 40.2, 42.1, 42.4, 46.7, 47.8, 70.0, 109.7, 112.0, 149.2, 170.6; IR ν_{\max} 2920, 2856, 2244, 1733, 1647, 1246 cm⁻¹. Anal. Calcd. For C₂₀H₂₉NO₂: C 76.15, H 9.27, N 4.44; Found C 76.13, H 9.32, N 4.19.

14.1.16 (1*R*,4*aR*,4*bS*,6*S*,8*aS*,10*aR*)-6-(Acetyloxy)tetradecahydro-2-oxo-1-phenanthrenepropanenitrile (19)—A solution of compound **18** (356 mg, 1.13 mmol) in MeOH (60 mL) and EtOAc (35 mL) was treated with ozone at -78 °C until a blue color persisted (ca 30 min). Oxygen was passed through the solution for 20 min until the blue color disappeared, and Me₂S (4.5 mL) was added at -78 °C and the resultant mixture was stirred overnight while warming to room temperature. Solvent was removed under reduced pressure and the residue was purified by column chromatography (silica gel; hexanes/EtOAc/CH₂Cl₂, 4:1:1) to give compound **19** (206 mg, 58%) as white crystals: mp 157–159 °C (hexanes/EtOAc); [α]_D²⁰ = +24.4 (c = 0.13, CHCl₃); ¹H NMR 5.11 (s, 1H), 2.46 (m, 1H), 2.04 (s, 3H); ¹³C δ NMR δ 15.1, 21.2, 22.6, 27.7, 29.2, 29.4, 30.4, 33.0, 33.6, 37.2, 39.1, 41.5, 41.7, 45.8, 54.5, 69.5, 118.8, 170.3, 213.1; IR ν_{\max} 2922, 2850, 2244, 1727, 1699, 1379, 1241 cm⁻¹. Anal. Calcd. For C₁₉H₂₇NO₃: C 71.89, H 8.57, N 4.41; Found C 71.64, H 8.73, N 4.49.

14.1.17 (2 β ,5 α ,14 β)-2-(Acetyloxy)-13-hydroxy-18-norestran-17-one (20) and (2 β ,5 α ,14 β)-2-(Acetyloxy)gonan-17-one (21)—Iodine (240 mg, 0.95 mmol) in dried THF (30 mL) was added to samarium filings (225 mg, 1.50 mmol) by syringe under Ar. The mixture was stirred at room temperature for 30 min to give a black blue solution. Then a solution of compound **19** (93 mg, 0.30 mmol) and 2-methyl-2-propanol (38 mg, 0.51 mmol) in THF (10 mL) was added to the freshly made SmI₂-THF solution under Ar at 0 °C. The reaction mixture was stirred at 0–10 °C while irradiated with a 500 W tungsten lamp for 8 h and then the reaction mixture was poured into 5% aqueous HCl (10 mL), and extracted with EtOAc (2 \times 15 mL). The combined organic extract was washed with water (10 mL), 10% NaHCO₃ (10 mL), brine (10 mL), and dried over Na₂SO₄. Solvent removal gave crude compound **20**, (46 mg) which was used without purification or characterization.

Under Ar, a solution of crude compound **20** in dried THF (5 mL) and MeOH (0.5 mL) was added to a freshly made SmI₂-THF (0.1 M, 15 mL) solution by syringe. The reaction mixture was stirred at room temperature for 10 min, quenched by adding 5% aqueous HCl (10 mL) and extracted with EtOAc (2 \times 10 mL). The combined organic extract was washed with water (10 mL), 10% NaHCO₃ (10 mL), brine (10 mL) and dried over Na₂SO₄. After solvent was removed under reduced pressure, the residue was purified by column chromatography (silica gel; CH₂Cl₂/EtOAc, 20:1) to give compound **21** (15.6 mg, 17% from compound **19**) as white crystals: mp 118–120 °C (EtOAc); ¹H NMR δ 5.11 (s, 1H), 2.38 (dd, *J* = 18.9, 8.1 Hz, 1H), 2.05 (s, 3H). From the ¹³C NMR data this product was determined to be a 9:1 mixture of epimers at C₁₄. The ¹³C NMR resonances for the major isomer were δ 20.7, 21.5, 22.9, 28.0, 28.3, 29.7, 30.2, 33.5, 33.7, 37.1, 40.6, 41.5, 41.7, 41.8, 42.3, 49.2, 70.0, 170.6, 221.4; IR ν_{\max} 2918, 2855, 1738, 1245 cm⁻¹.

14.1.18 (2 β ,5 α ,13 β ,14 β)-2-Hydroxygonan-17-one (1)—The mixture of compound **21** (15.6 mg, 0.05 mmol), NaOH (2 mg, 0.05 mmol) and water (18 mg, 1 mmol) in MeOH (12 mL) was refluxed for 2 h. The reaction mixture was chilled, and most of the MeOH was removed under reduced pressure to give a residue to which was added EtOAc (20 mL). The organic phase was washed with 10% aqueous HCl (5 mL), water (5 mL), brine (5 mL) and dried over Na₂SO₄. After the solvent was removed under reduced pressure, the residue was

purified by column chromatography (silica gel; hexanes/CH₂Cl₂/EtOAc, 3:1:1) to give compound **1** (11 mg, 82%) as white crystalline needles: mp 123–124 °C (EtOAc/hexanes); [α]_D²⁰ = +140 (c = 0.30, CHCl₃); ¹H NMR δ 4.16 (m, 1H), 2.38 (dd, *J* = 18.9, 8.1 Hz, 1H); ¹³C NMR δ 20.7, 23.0, 27.6, 28.0, 30.3, 32.5, 33.7, 36.3, 37.2, 40.6, 40.8, 41.7, 41.8, 42.6, 49.2, 66.7, 221.5; IR ν_{\max} 3565, 2923, 2851, 1720 cm⁻¹. Anal. Calcd. For C₁₇H₂₆O₂: C 77.82, H 9.99; Found: C 77.68, H 10.02.

Acknowledgements

This work was supported by NIH grant GM47969. Instrumentation for crystallographic studies was made possible by NSF Grant CHE-0420497. The authors thank Dr. Alex S. Evers, Dr. Steven Mennerick, Dr. Charles F. Zorumski, Ann Benz and Brad D. Manion for the biological evaluations of compound **1**.

References and Notes

1. Phillips GH. *J Steroid Biochem* 1975;6:607. [PubMed: 1186246]
2. Covey DF, Evers AS, Mennerick S, Zorumski CF, Purdy RH. *Brain Res Rev* 2001;37:91. [PubMed: 11744077]
3. Hamilton NM. *Curr Top Med Chem* 2002;2:887. [PubMed: 12171578]
4. Belelli D, Lambert JJ. *Nat Rev Neurosci* 2005;6:565. [PubMed: 15959466]
5. Katona BW, Krishnan K, Cai ZY, Manion BD, Benz A, Taylor A, Evers AS, Zorumski CF, Mennerick S, Covey DF. *Eur J Med Chem*. In press
6. Purdy RH, Morrow AL, Blinn JR, Paul SM. *J Med Chem* 1990;33:1572. [PubMed: 2160534]
7. Anderson A, Boyd AC, Clark JK, Fielding L, Gemmell DK, Hamilton NM, Maidment MS, May V, McGuire R, McPhail P, Sansbury FH, Sundaram H, Taylor R. *J Med Chem* 2000;43:4118. [PubMed: 11063608]
8. Wittmer LL, Hu Y, Kalkbrenner M, Evers AE, Zorumski CF, Covey DF. *Mol Pharmacol* 1996;50:1581. [PubMed: 8967980]
9. Han M, Zorumski CF, Covey DF. *J Med Chem* 1996;39:4218. [PubMed: 8863799]
10. Zeng CM, Manion BD, Benz A, Evers AS, Zorumski CF, Mennerick S, Covey DF. *J Med Chem* 2005;48:3051. [PubMed: 15828844]
11. McKinney AR, Ridley, Damon D, Turner P. *Aust J Chem* 2003;56:829.
12. Han M, Hayes BA, Prendergast PT, Gupta S. *J Labelled Compd Radiopharm* 2000;43:1149.
13. Groszek G, Kabat MM, Kurek A, Masnyk M, Wicha J. *Bull Pol Acad Sci Chem* 1986;34:313.
14. Numazawa M, Nagaoka M, Osawa Y. *J Org Chem* 1982;47:4024.
15. Liu Z, Zhang J, Cheng W. *Chinese Chem Lett* 1994;5:39.
16. Leese MP, Newman SP, Purohit A, Reed MJ, Potter BVL. *Bioorg & Med Chem Lett* 2004;14:3135. [PubMed: 15149660]
17. Demir AS, Sabol MR, Jeganathan A, Kurt Dolence E, Watt DS, Moldowan JM. *Org Prep and Proced Int* 1987;19:197.
18. Ohloff G, Maurer B, Winter B, Giersch W. *Helv Chim Acta* 1983;66:192.
19. Wang C, Jiang X, Shi J, Lu J, Hu Y, Hu H. *J Org Chem* 2003;68:4579. [PubMed: 12762777]
20. Jiang X, Wang C, Hu Y, Hu H, Covey DF. *J Org Chem* 2000;65:3555. [PubMed: 10843648]
21. Crystallographic data (excluding structure factors) for the structure in this article have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication number CCDC 634624. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK [fax: +44 1223 330633 or email: deposit@ccdc.cam.ac.uk].
22. Jiang X, Manion BD, Benz A, Rath NP, Evers AS, Zorumski CF, Mennerick S, Covey DF. *J Med Chem* 2003;46:5334. [PubMed: 14640542]
23. Han M, Covey DF. *J Org Chem* 1996;61:7614. [PubMed: 11667699]
24. Counsell RE. *Tetrahedron* 1961;15:202.




Figure 1. Structures of steroid modulators of GABA_A receptor function. Allopregnanolone and *ent*-androsterone 17-spiroepoxide potently enhance the actions of GABA at GABA_A receptors. Androsterone is a weak enhancer of GABA action. *ent*-Androsterone has activity higher than that of androsterone but less than that of the other two steroids.




Figure 2. Panel A: Structures of (2β,5α,13β,14β)-2-hydroxygonan-17-one (**1**) and *ent*-androsterone. Panel B: An overlay of steroid **1** (blue structure) and *ent*-androsterone (orange structure). The structures were overlaid using a RMS fit of atoms C₉, C₁₁, C₁₂, C₁₄, O₃ and O₁₇ in each structure for the alignment. The O₂–O₁₇ (9.37 Å) distance in steroid **1** is shorter than the O₃–O₁₇ distance (9.67 Å) in *ent*-androsterone.







Figure 3. Projection view of one of the two unique molecules of steroid **1** shown with 50% thermal ellipsoids for non-hydrogen atoms.


**Scheme 1.**



Scheme 2.



Scheme 3.



Scheme 4.