

## Transposition of the *mariner* element from *Drosophila mauritiana* in zebrafish

JAMES M. FADOOL<sup>\*†‡</sup>, DANIEL L. HARTL<sup>§</sup>, AND JOHN E. DOWLING<sup>†</sup>

<sup>\*</sup>Department of Zoology and Wildlife Sciences, Auburn University, Auburn, AL 36849; and Departments of <sup>†</sup>Molecular and Cellular Biology and <sup>§</sup>Organismic and Evolutionary Biology, The Biological Laboratories, Harvard University, 16 Divinity Avenue, Cambridge MA 02138

Contributed by John E. Dowling, February 19, 1998

**ABSTRACT** With the increased popularity of zebrafish (*Danio rerio*) for mutagenesis studies, efficient methods for manipulation of its genome are needed. One approach is the use of a transposable element as a vector for gene transfer in this species. We report here the transformation of zebrafish and germ-line transmission of the *mariner* element from *Drosophila mauritiana*. The *mariner* element was selected because its transposition is independent of host-specific factors. One- to two-cell-stage zebrafish embryos were coinjected with a supercoiled plasmid carrying the nonautonomous *mariner* element *peach* and mRNA encoding the transposase. Surviving larvae were reared to adulthood, and the transmission of *peach* to the F<sub>1</sub> generation was tested by PCR. Four of the 12 founders, following plasmid injections on 2 different days, transmitted the element to their progeny. Inheritance of the transgene from the F<sub>1</sub> to the F<sub>2</sub> generation showed a Mendelian pattern. No plasmid sequences were detected by PCR or Southern blot analysis, indicating transposition of *peach* rather than random integration of the plasmid DNA. These data provide evidence of transformation of a vertebrate with a transposable element and support the host-independent mechanism for transposition of the *mariner* element. We suggest this system could be used for insertional mutagenesis or for identifying active regions of the genome in the zebrafish.

Transposons are naturally occurring genetic elements capable of moving from one chromosomal location to another (1). The fundamental components of transposable elements are a gene encoding the enzyme necessary for transposition, a transposase, and flanking sequences required for recognition by the transposase. Autonomous elements encode functional copies of the transposase and possess the recognition sequences to mediate transposition. Nonautonomous elements often carry mutated copies of the transposase but retain the recognition sequences; thus, they can be mobilized by transposase expressed from another source (1, 2).

In many eukaryotes, transposons are powerful tools for genetic research (3–5). This finding is convincingly illustrated by the widespread use of the *P*-element transposon in *Drosophila melanogaster* for mutating genes and for transferring foreign DNA sequences into the genome (3, 6–9). Sequence homology has identified many remnants of transposable elements in vertebrates, although the frequency of active transposition is comparatively low (10), and no functional transposase has been isolated as yet *in vivo* (11). Attempts to use the *P*-element for transformation in vertebrates failed, apparently because of the requirement of additional host-specific factors necessary for transposition (12–14). To overcome these difficulties, we have investigated the utility of the *mariner* element

isolated from *Drosophila mauritiana* as a transformation vector for vertebrates (15).

The original, nonautonomous, *mariner* element was isolated from an unstable eye color allele, *white-peach*, in *D. mauritiana* (15). The *Mos1* factor, an autonomous copy of *mariner*, was identified by its ability to promote high frequency excision of the *mariner* element, *peach*, from its position near the *white* gene (16, 17). The *mariner* element is ≈1.3 kb in length, encodes a single polypeptide chain of 345 amino acids, and is flanked by 28-bp inverted repeats. *mariner*-like elements (MLEs) have been identified in a wide range of host species including humans (18–21). Recent findings that *Mos1* can transform the parasitic protozoan *Leishmania* (22) and a MLE from horn fly can transpose *in vitro* provide evidence that transposition is independent of host-specific factors (23). Thus, *mariner* has been proposed as a vector for transformation of a wide range of eukaryotes (22–24).

In this report, we present evidence for transposition in the zebrafish of *mariner* from *D. mauritiana*. Zebrafish embryos were injected with a plasmid carrying the nonautonomous element, *peach*, and mRNA encoding the *Mos1* transposase. The strategy was to rely upon the host's cellular machinery to translate the mRNA into functional *Mos1* transposase. The transposase should recognize and then transpose the *peach* element from the plasmid vector into the host genome. A transposition event in cells giving rise to the germ line would be detected by inheritance of the *peach* element in the F<sub>1</sub> progeny. Although random integration of plasmid DNA is plausible, distinguishing random integration from transposition is possible by screening for co-inheritance of *peach* and plasmid vector sequences (e.g., the β-lactamase gene and regions adjacent to the multicloning site). We propose that by manipulating the presence of the transposase, a general transformation system for mutagenesis and gene tagging can be developed in a vertebrate model.

### MATERIALS AND METHODS

**Fish Stocks and DNA Constructs.** The Ekwill strain of zebrafish, originally obtained from Ekwill Fish Farm (Gibsonton, FL), was maintained as an inbred population in the Harvard Zebrafish Facility.

The construct pAD31 consists of an immobile version of *Mos1* cloned into the multicloning site of pBluescript II (24). The 1,450-bp insert lacks the 5'-inverted repeat necessary for recognition by the transposase and is flanked by 200 bp of *Drosophila simulans* genomic DNA. pAD31 was linearized 3' of the insert with *XhoI* or *HindIII*, digested with proteinase K, and phenol/chloroform extracted. mRNA was generated by *in vitro* transcription by using T7 RNA polymerase and the mCAP Capping Kit according to the manufacturer's instructions (Stratagene). The mRNA was brought to a concentration

The publication costs of this article were defrayed in part by page charge payment. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. §1734 solely to indicate this fact.

© 1998 by The National Academy of Sciences 0027-8424/98/955182-5\$2.00/0  
PNAS is available online at <http://www.pnas.org>.

Abbreviation: MLE, *mariner*-like element.

<sup>‡</sup>To whom reprint requests should be addressed at: Department of Zoology and Wildlife Sciences, 331 Funchess Hall, Auburn University, Auburn, AL 36849. e-mail: fadooljm@mail.auburn.edu.

of 1  $\mu\text{g}/\mu\text{l}$  with water, and the purity and sizes of the transcript were examined by agarose gel electrophoresis. The pWE1 construct (provided by Daniel De Aguiar, Harvard University) was generated by ligating a PCR amplification product of the original 1.3-kb *peach* element (15) generated with sequence-specific primers flanked by *EcoRI* restriction sites into the unique *EcoRI* site of pBluescript II (Fig. 1N). Prior to injection, plasmid DNA was purified on Wizard miniprep columns (Promega), digested with proteinase K, phenol/chloroform extracted, and precipitated with ethanol. The DNA was dissolved in diethyl-pyrocyanate-treated water at a concentration of 0.1  $\mu\text{g}/\mu\text{l}$ . Gel electrophoresis demonstrated the preparation to consist primarily of supercoiled plasmid.

One- to two-cell-stage zebrafish embryos from timed, pairwise matings were micro-injected just below the cell cytoplasm with an estimated 1–4 nl of a solution containing 45  $\text{ng}/\mu\text{l}$  pWE1 and 50  $\text{ng}/\mu\text{l}$  mRNA or 45  $\text{ng}/\mu\text{l}$  pWE1 plasmid DNA alone. Surviving embryos were reared to breeding stage.

**DNA Isolation and PCR.** Founder fish were mated pairwise or outcrossed to the original Ekwil line. DNA was prepared from pools of 20–30 embryos by digestion at 52°C in 100  $\mu\text{l}$  of 10 mM Tris (pH 8.0) containing 200  $\mu\text{g}/\text{ml}$  proteinase K, 0.2% SDS, and 20 mM NaCl. The reactions were stopped by heating to 95°C. The digests were diluted 1:5 with water and centrifuged to remove suspended material. Two microliters of the digest were used as template in 35 or 50  $\mu\text{l}$  PCRs. Individual embryos were digested in 25  $\mu\text{l}$  digestion buffer and processed as above. Individual zebrafish older than 21 days were screened by isolating DNA from caudal fin clippings. Fish were anesthetized in tricaine (25), and either the dorsal or ventral region of the caudal fin was removed with a razor. The tissue was digested as above for individual embryos.

The PCRs contained 10 mM Tris-HCl (pH 9.0), 50 mM KCl, 0.1% Triton X-100, 2.25 mM  $\text{MgCl}_2$ , 20 mM each dNTP, 25 pmol each primer, and 1 unit *Taq* DNA polymerase (Promega). The reactions were initially denatured at 94°C for 3 min, followed by 35 cycles of 94°C for 30 sec and 60°C for 1 min 30 sec, and a final 72°C elongation for 15 min to 2 hr. For most samples, the presence of suitable template DNA in the preparations was confirmed by amplification in separate reactions of a 250-bp fragment of the zebrafish homeobox gene *ZF21* (26). The primer pairs for the *peach* element were sense primer JMF31 (GAA GTG TCA ACC TTG ACT GTC) and antisense primer JMF36 (CTC ATG TGG CAC CCA TCT AC), which yield an amplification product of 450 bp. These primers demonstrate no known homology to endogenous zebrafish genomic sequences. Primers complementary to plasmid DNA were JMF32 (GCG GCC AAC TTA CTT CTG AC) and JMF33 (CTG ACG CTC AGT GGA ACG AA), which amplify a 650-bp fragment of the  $\beta$ -lactamase gene, and JMF6 (GTT TTC CCA GTC ACG ACG TTG TAA) against M13 forward sequence, which in combination with JMF36 yields a 680-bp product. The primer pair used for amplification of the *ZF21* gene was JMF21 (GCG TTT TGT CCA CAT TAT TCA G)

and JMF22 (GTA GCC AGA CTC ATG CTC TTC AG). Amplification products were identified by agarose gel electrophoresis.

**Southern Blot Analysis.** Adult zebrafish were ground into powder in liquid nitrogen, resuspended in digestion buffer (10 mM Tris, pH 8.0/25 mM EDTA/0.5% SDS/10 mM NaCl/100  $\mu\text{g}/\text{ml}$  proteinase K), and incubated overnight at 50°C. Preparations were repeatedly extracted with phenol, followed by phenol/chloroform, and precipitated with ethanol. Ten to 15  $\mu\text{g}$  of DNA was digested to completion with the *Bam*HI, *Eco*RI, or *Sac*I (Life Technologies, Gaithersburg, MD) in appropriate buffer. Samples were precipitated with ethanol and resuspended in TE prior to electrophoresis on 0.8% agarose gels in TAE. Following electrophoresis, the DNA was blotted to GeneScreenPlus (DuPont/NEN) by capillary transfer and UV cross-linked to the membrane. Prehybridization and hybridization were conducted at 65°C according to the manufacturers recommendations (DuPont/NEN) by using prehybridization solution (Life Technologies). The probe, labeled with [ $^{32}\text{P}$ ]dCTP, was generated from the 1,300-bp *Eco*RI fragment of the pWE1 vector by random priming.

## RESULTS

The plasmid pWE1 carries a single copy of the nonautonomous *mariner* element, *peach*, cloned into a unique *Eco*RI restriction site. mRNA encoding the transposase was transcribed from a plasmid containing the *Mos1* factor, an autonomous copy of *mariner*. On two separate days, 64 1- to 2-cell-stage zebrafish embryos were injected with pWE1 and *Mos1* mRNA. The surviving larvae (13 and 33, respectively) were reared to breeding stage and mated in pairs. DNA was isolated from pools of 20–30 embryos and screened by PCR for the presence of the *peach* element. If a pool tested positive for the transgene, clutches of embryos from the same founding pair were raised to breeding stage. DNA isolated from caudal fin clips was screened to identify transgenic F<sub>1</sub> individuals. The frequency of transmission of the transgene to the F<sub>2</sub> progeny was determined.

Pools of embryos from 4 pairs of founders tested positive for the transgene (Fig. 2). This result represents the identification of a minimum of 4 independent integration events in the 12 founders successfully mated, a frequency of 33% for germ-line transmitting founders. From 3 of these, the transgene was inherited by 1 or 2 (3–14%) of the individual F<sub>1</sub> fish, indicating mosaicism of the germ line. In the fourth case, the transgene was not identified in any of the 22 available F<sub>1</sub> fish. This low frequency of transmission is consistent with the frequency observed in the pooled DNA samples in which possibly less than 2% of the progeny inherited this copy of the transgene.

From four F<sub>1</sub> fish identified in the remaining three independent lines, the frequency of transmission of the transgene to their F<sub>2</sub> progeny was examined. F<sub>1</sub> fish 1060.07, 1061.03, and 1085.26 transmitted the transgene to 50–56% (7/12, 10/18, and 9/18, respectively) of the F<sub>2</sub> generation, consistent with a




FIG. 1. Diagrammatic representation of the *mariner*-containing vector pWE1. A 1,300-bp PCR amplification product derived from the nonautonomous element, *peach*, was ligated into the unique *Eco*RI site of pBluescript II. The locations of the M13 sequences flanking the polylinker site and the ampicillin resistance gene are represented by the hatched and stippled boxes, respectively. The locations of primer pairs used to screen for *mariner* and plasmid sequence in genomic DNA preparations are indicated.




FIG. 2. PCR amplification of the *mariner* element in F<sub>1</sub> progeny. (a) Genomic DNA was isolated from pools of 20–30 F<sub>1</sub> generation larvae from six pairwise matings of founder fish (numbered 1–6) and used as template for amplification of *mariner* sequences. Nine of the pools of larvae from four different crosses tested positive for the transgene as indicated by a product at 450 bp (arrow). (b) Analysis of DNA samples isolated from fin clippings of individual F<sub>1</sub> offspring from mating of founder pair number 5. Genomic DNA from 2 of the 15 individuals demonstrated the presence of *mariner* sequences (arrow) (lane M, 100-bp marker; Promega; lane B, no DNA control; lane C, positive control containing 1 ng pWE1 plasmid).

Mendelian pattern of segregation of a single heterozygous locus. F<sub>1</sub> fish 1061.09 transmitted the transgene to 68% and 70% (11/16 and 7/10) of F<sub>2</sub> progeny in two clutches, respectively (Fig. 3). This result is consistent with the inheritance of two independent insertions. Because the F<sub>1</sub> fish in question was reared from a clutch derived from mating two putative founder fish, the two potential inserts could have been inherited from a single founder or as separate inserts from each parent. Our data cannot rule out either possibility. Overall, the data do demonstrate isolation in the F<sub>1</sub> generation of at least four independent inserts from the original five possible.

DNA preparations from individual F<sub>2</sub> fish from each of the F<sub>1</sub>s were also tested for the presence of plasmid DNA sequences. Two sets of primer pairs were used: primers specific to the  $\beta$ -lactamase gene and a primer pair with one specific to *peach* sequences and one specific to the plasmid M13 sequences flanking the insert (Fig. 1). If the transgenic lines were the result of random integration of plasmid DNA into the fish




FIG. 3. PCR demonstration of inheritance of the *mariner* element without plasmid vector sequences. (a) Genomic DNA isolated from individual F<sub>2</sub> larvae was tested for the presence of the *mariner* element and flanking plasmid sequences. Amplification of *mariner* sequences identified 11 of 16 (68%) larvae as positive for the transgene (arrow) (lane M, 100-bp marker; Promega; lane B, no DNA control). (b) Amplification with one primer specific for *mariner* and one for the M13 reverse sequence flanking the insert in the pWE1 plasmid revealed none possessed this plasmid DNA. The control lane (C), containing 1 ng pWE1 plasmid, demonstrated the expected amplification product of 680 bp (arrowhead).

genome, we would expect to find plasmid sequences inherited with the *peach* sequences. However, in all cases tested, the DNA did not support amplification of either the ampicillin resistance gene or M13 sequences (Fig. 3). Lowering the annealing temperature in PCR conditions gave a similar result. These data suggest that the transgenic lines result from transposition of *peach* into the zebrafish genome.

Genomic DNA isolated from three of the lines was subjected to Southern blot analysis to provide further evidence for transposition. If the transgene had integrated as plasmid DNA, the sequences of the multicloning site of the pWE1 vector should also be inherited. Digestion with *Eco*RI should then liberate the *peach* element from the integrated plasmid, yielding a 1,300-bp band on a Southern blot. The same argument implies that digestion with *Sac*I, which would cut once within the element and once within the multicloning site, would generate one fragment of 800 bp in each line and a second of variable size. However, such results were not observed. Digestion with *Eco*RI resulted in bands of 10, 7, and 9 kb in the three lines tested, and digestion with *Sac*I produced two fragments of high molecular weight in each line (Fig. 4). These data demonstrate that the *peach* element inserted into the zebrafish genome without the flanking plasmid sequences, providing evidence that transgenesis was the result of transposition and not random integration into the host genome. In our study, injection of the plasmid alone yielded no transgenic animals (data not shown) indicating that integration of the transgene was dependent upon the presence of the transposase.




FIG. 4. Evidence for transposition of the *peach* element by Southern blot analysis. Genomic DNA from lines 1085 (a), 1060 (b) and 1061 (c) were digested with *Bam*HI (B), *Eco*RI (E), and *Sac*I (S), and filters were probed with the 1.3-kb *peach* element from pWE1. If the *peach* element had integrated with flanking plasmid sequences, then digestion with *Eco*RI should generate a single 1.3-kb band, and digestion with *Sac*I, which recognizes one site in the *peach* element and one in the multicloning site of pWE1, should produce a fragment of 800 bp and a second fragment with a size dependent upon the site of integration. However, the hybridization patterns for the three samples differ, and the fragments are much larger than expected. These findings indicate that integration of the *peach* element occurred without flanking plasmid sequences and that the genomic sites of integration are different. (Molecular weight standards are shown to the left. The arrow indicates the location of the intact 1.3-kb *peach* element from pWE1 following digestion with *Eco*RI.)



## DISCUSSION

We have demonstrated that the *mariner* element from *D. mauritiana* can be used for generation of stable transgenic lines of zebrafish. This result was achieved by microinjection, into 1- and 2-cell-stage zebrafish embryos, a plasmid carrying a nonautonomous element and mRNA encoding the transposase. This strategy is similar to that originally used for transformation of *D. melanogaster*, which relied upon microinjection of a helper plasmid encoding the transposase with the DNA containing the transposon (6, 7). While this work was under review, transformation of zebrafish with the *Caenorhabditis elegans* transposable element, *Tc3* was achieved (27). *Tc1*-related elements are known to occur in teleosts including zebrafish (10, 11). Although no functional transposase has been isolated *in vivo*, a low frequency of transposition has been demonstrated (10), and a functional transposase based upon these sequences has been engineered *in vitro* (11); thus, it may be that the whole *mariner/Tc1* superfamily of transposons is functional in this genome. These results suggest the possibility of replicating in zebrafish, a vertebrate model, the success observed with transposable elements in *D. melanogaster* (6–9).

Although transformation of zebrafish and germ-line transmission of a transgene was first demonstrated a decade ago (28), the technology is not well developed. The most common method relies on random integration of DNA microinjected into a 1- to 2-cell-stage embryo. The efficiency of transmission through the germ line is usually low (0–20% in different reports), and founder fish demonstrate varying degrees of mosaicism (28–31). Rearrangements and concatomerization of the transgene are common (28–30). Expression of the transgene after passage through the germ line is unpredictable and may decrease in subsequent generations (29, 30, 32).

A major advance in zebrafish transgenics occurred with the application of a pseudotype retrovirus vector for insertional mutagenesis (33–35). First developed as a vector for gene therapy and genetic studies (36), the engineered virus can infect a wide range of organisms and it efficiently integrates into the genome. In the zebrafish, transformation rates are approaching 100%, with most founders transmitting on average 10 proviral inserts to their progeny (37). This vector has been successfully employed for the mutating and cloning of several genes involved in zebrafish embryogenesis (34, 35). Based upon a frequency of isolating 1 embryonic lethal mutation per 70 insertions, it has been proposed that 250–350 mutations can be recovered over a period of several years. However, the application of the retrovirus to modestly sized mutagenesis screens or as a general transformation vector has not been demonstrated.

The *mariner/Tc1* elements could be used as vectors for insertional mutagenesis. One strategy would be similar to that using the pseudotype retrovirus (34–36): generate many transposon inserts, breed these to homozygosity, and screen the larvae for mutant phenotypes. An alternative strategy would be an “enhancer-trap” screen of heterozygous insertions looking for specific patterns of expression of a reporter gene (8, 38–40). Expression of the reporter gene is regulated by host enhancer sequences near the site of insertion. Once a mutant phenotype or interesting pattern of reporter gene expression are identified, the transposon acts as a molecular tag and facilitates cloning of the gene. If large numbers of inserts can be efficiently mobilized to new chromosomal loci in each generation, it would be possible to undertake complementation analysis to clone genes previously identified in chemical mutagenesis screens. A limitation of any such mutant screen is the ability to generate the large number of insertions needed to approach saturation of the potential insertion sites. Steps are underway in our laboratories to improve this transformation frequency.

We thank Stephen M. Beverly for reviewing this manuscript and for helpful comments. We also thank Bill McCarthey and Amy Chin of the Harvard University Fish Facility for maintenance of zebrafish, Daniel De Aguiar for plasmid constructs, and Robert Hagan for assistance in the laboratory. This work was supported by grants awarded by the National Institutes of Health (EY00811 to J.E.D. and EY06554 and EY11774 to J.M.F.).

- Howe, M. & Berg, D., eds. (1989) *Mobile DNA* (Am. Soc. Microbiol., Washington, DC).
- Hartl, D. L., Lozovskaya, E. R. & Lawrence, J. (1992) *Genetica* **86**, 47–53.
- Engels, W. R. (1989) in *Mobile DNA*, eds. Howe, M. & Berg, D. (Am. Soc. Microbiol., Washington, DC), pp. 437–484.
- Moerman, D. G. & Waterston, R. H. (1989) in *Mobile DNA*, eds. Howe, M. & Berg, D. (Am. Soc. Microbiol., Washington, DC), pp. 537–556.
- Osborne, B. I. & Baker, B. (1995) *Curr. Opin. Cell Biol.* **7**, 406–413.
- Spradling, A. C. & Rubin, G. M. (1982) *Science* **218**, 341–347.
- Rubin, G. M. & Spradling, A. C. (1982) *Science* **218**, 348–353.
- O’Kane, C. J. & Gehring, W. J. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 9123–9127.
- Spradling, A. C., Stern, D. M., Kiss, I., Roote, J., Laverty, T. & Rubin, G. M. (1995) *Proc. Natl. Acad. Sci. USA* **92**, 10824–10830.
- Lam, W. L., Lee, T. S. & Gilbert, W. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 10870–10875.
- Ivics, Z., Hackett, P. B., Plasterk, R. H. & Izsvak, Z. (1997) *Cell* **91**, 501–510.
- Laski, F. A., Rio, D. C. & Rubin, G. M. (1986) *Cell* **44**, 7–19.
- Rio, D. C. & Rubin, G. M. (1988) *Proc. Natl. Acad. Sci. USA* **85**, 8929–8933.
- Beall, E. L., Admon, A. & Rio, D. C. (1994) *Proc. Natl. Acad. Sci. USA* **91**, 12681–12685.
- Jacobson, J. W., Medhora, M. M. & Hartl, D. L. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 8684–8688.
- Bryan, G. J., Jacobson, J. W. & Hartl, D. L. (1987) *Science* **235**, 1636–1638.
- Medhora, M. M., MacPeck, A. H. & Hartl, D. L. (1988) *EMBO J.* **7**, 2185–2189.
- Robertson, H. M. (1993) *Nature (London)* **362**, 241–245.
- Auge-Gouillou, C., Bigot, Y., Pollet, N., Hamelin, M. H., Meunier-Rotival, M. & Periquet, G. (1995) *FEBS Lett.* **24**, 541–546.
- Garcia-Fernandez, J., Bayascas-Ramirez, J. R., Marfany, G., Munoz-Marmol, A. M., Casali, A., Baguna, J. & Salo, E. (1995) *Mol. Biol. Evol.* **12**, 421–431.
- Robertson, H. M., Zumpano, K. L., Lohe, A. R. & Hartl, D. L. (1996) *Nat. Genet.* **12**, 360–361.
- Gueiros-Fiho, F. J. & Beverley, S. M. (1997) *Science* **276**, 1716–1719.
- Lampe, D. J., Churchill, M. E. A. & Robertson, H. M. (1996) *EMBO J.* **15**, 5470–5479.
- Lohe, A. R., Lidholm, D.-A. & Hartl, D. L. (1995) *Genetics* **140**, 183–192.
- Westerfield, M. (1995) *The Zebrafish Book* (Univ. of Oregon Press, Eugene), 3rd Ed.
- Njolstad, P. R., Molven, A. & Fjose, A. (1988) *FEBS Lett.* **230**, 25–30.
- Raz, E., van Luenen, H. G. A. M., Schaerringer, B., Plasterk, R. H. A. & Driever, W. (1998) *Curr. Biol.* **8**, 82–88.
- Stuart, G. W., McMurray, J. V. & Westerfield, M. (1988) *Development (Cambridge, U.K.)* **103**, 403–412.
- Stuart, G. W., Vielkind, J. R., McMurray, J. V. & Westerfield, M. (1990) *Development (Cambridge, U.K.)* **109**, 577–584.
- Culp, P., Nusslein-Volhard, C. & Hopkins, N. (1991) *Proc. Natl. Acad. Sci. USA* **88**, 7953–7957.
- Lin, S., Yang, S. & Hopkins, N. (1994) *Dev. Biol.* **161**, 77–83.
- Gibbs, P. D. L., Peek, A. & Thorgaard, G. (1994) *Mol. Mar. Biol. Biotech.* **3**, 307–316.
- Lin, S., Gaiano, N., Culp, P., Burns, J. C., Friedmann, T., Yee, J.-K. & Hopkins, N. (1994) *Science* **265**, 666–669.
- Gaiano, N., Amsterdam, A., Kawakami, K., Allende, M., Becker T. & Hopkins, N. (1996) *Nature (London)* **383**, 829–832.
- Allende, M. L., Amsterdam, A., Becker, T., Kawakami, K., Gaiano, N. & Hopkins, N. (1996) *Genes Dev.* **10**, 3141–3155.

36. Burns, J. C., Friedmann, T., Driever, W., Burrascano, M. & Yee, J.-K. (1993) *Proc. Natl. Acad. Sci. USA* **90**, 8033–8037.
37. Gaiano, N., Allende, M., Amsterdam, A., Kawakami, K., Hopkins, N. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 7777–7783.
38. Allen, N. D., Cran, D. G., Barton, S. C., Hettle, S., Reik, W. & Surani, M. A. (1988) *Nature (London)* **333**, 852–855.
39. Gossler, A., Joyner, A. L., Rossant, J. & Skarnes, W. C. (1989) *Science* **244**, 463–465.
40. Bayer, T. A. & Campos-Ortega, J. A. (1992) *Development (Cambridge, U.K.)* **115**, 421–426.