



Published in final edited form as:

Exp Hematol. 2007 August ; 35(8): 1256–1265.

Unique and redundant roles of $\alpha 4$ and $\beta 2$ integrins in kinetics of recruitment of lymphoid versus myeloid cell subsets to the inflamed peritoneum revealed by studies of genetically deficient mice

Tatiana Ulyanova^{*}, Gregory V. Priestley, Ena Ray Banerjee, and Thalia Papayannopoulou
Department of Medicine/Hematology University of Washington, Seattle, WA

Abstract

OBJECTIVE—Leukocyte recruitment to inflammatory sites is a prominent feature of acute and chronic inflammation. Instrumental in this process is the coordinated upregulation of leukocyte integrins (among which $\alpha 4\beta 1$ and $\beta 2$ integrins are major players) and their cognate receptors in inflamed tissues. To avoid the ambiguity of previous short-term antibody-based studies and to allow for long-term observation, we used genetically deficient mice to compare roles of $\alpha 4$ and $\beta 2$ integrins in leukocyte trafficking.

METHODS—Aseptic peritonitis was induced in $\alpha 4$ or $\beta 2$ integrin-deficient (conditional and conventional knockouts, respectively) and control mice, and recruitment of major leukocyte subsets to the inflamed peritoneum was followed for up to 4 days.

RESULTS—Despite normal chemokine levels in the peritoneum and adequate numbers, optimal recruitment of myeloid cells was impaired in both $\alpha 4$ - and $\beta 2$ -deficient mice. Furthermore, clearance of recruited neutrophils and macrophages was delayed in these mice. Lymphocyte migration to the peritoneum in the absence of $\alpha 4$ integrins was drastically decreased, both at steady state and during inflammation, a finding consistent with impaired lymphocyte *in vitro* adhesion and signaling. By contrast, in the absence of $\beta 2$ integrins, defects in lymphocyte recruitment were only evident when peritonitis was established.

CONCLUSIONS—Our data with concurrent use of genetic models of integrin deficiency reveal non-redundant functions of $\alpha 4$ integrins in lymphocyte migration to the peritoneum and further refine specific roles of $\alpha 4$ and $\beta 2$ integrins concerning trafficking and clearance of other leukocyte subsets at homeostasis and during inflammation.

INTRODUCTION

A prominent feature of acute or chronic inflammation is the recruitment of mature leukocytes to inflammatory sites. For the successful implementation of this process, several highly coordinated adhesion and activation steps need to be accomplished by leukocytes in inflamed tissues [1,2]. Essential molecular players in this multi-step adhesion/migration cascade are $\alpha 4$ and $\beta 2$ integrins. In particular, $\alpha 4\beta 1$ (VLA4) integrin is unique among integrins as it can function in all three steps of the trafficking cascade: rolling/tethering initiated by selectins,

^{*}Corresponding author: Dr. Tatiana Ulyanova, Ph.D., Department of Medicine Division of Hematology, University of Washington, 1705 NE Pacific St., HSB Room K-257, Seattle WA 98195, Phone: 206-543-2368, FAX: 206-543-3560, e-mail: ulyant@u.washington.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

firm adhesion, and transmigration step controlled by activated integrins [3-6]. Expression of $\alpha 4$ integrins is constitutive in all leukocytes except human neutrophils, where it is inducible [7], whereas murine neutrophils constitutively express $\alpha 4\beta 1$ [8]. The $\beta 2$ integrins are expressed exclusively in hematopoietic cells [9]. Both $\alpha 4$ and $\beta 2$ integrins, as well as their cognate receptors are up-regulated by various inflammatory stimuli [1,6].

Function-blocking antibodies and peptides have been extensively used to study the role of $\alpha 4$ and $\beta 2$ integrins in leukocyte trafficking. However, results of antibody studies vary with the animal model used or the route of antibody administration, and off-target effects can not be excluded [10-14]. To avoid the ambiguity of antibody-based studies and to carry out long-term observations, mouse models with genetically modified integrin genes have been generated [15-17]. To circumvent embryonic lethality of $\alpha 4$ knockout mice [18] in studying the role of $\alpha 4$ integrins in vivo, reconstitution of RAG^{-/-} mice with $\alpha 4$ -deficient ES cells was undertaken [19]. In this model though, a profound defect in development of $\alpha 4$ -/- lymphocytes and lymphoid organs was observed in postnatal life, thus precluding the use of this model to study migratory behavior of mature leukocyte populations. A new model of postnatal conditional $\alpha 4$ deficiency with normal development of the immune system was recently established in our laboratory [20].

Using this model, new aspects of the role of $\alpha 4$ integrins in homing and retention of hematopoietic progenitors in the bone marrow at steady state and recovery after hematopoietic stress were revealed, but trafficking patterns of mature leukocytes to the inflammatory sites have not previously been addressed or compared to other integrin-deficient mice.

To uncover unique and overlapping roles of $\alpha 4$ and $\beta 2$ integrins in mature hematopoietic cell trafficking, we analyzed patterns of recruitment of various leukocyte subsets to the peritoneum before and after inflammation in $\alpha 4$ or $\beta 2$ integrin-deficient mice, using a well-studied model of aseptic thioglycolate-induced peritonitis. Our results revealed intrinsic differences of migratory responses in the absence of $\alpha 4$ integrins in lymphoid versus myeloid subsets. Parallel studies using mice with single $\alpha 4$ - or $\beta 2$ -, as well as mice with double ($\alpha 4$ and $\beta 2$) integrin deficiency allowed fine-tuning of the roles of $\alpha 4$ and $\beta 2$ integrins in leukocyte trafficking.

MATERIALS AND METHODS

Mice

Mice used in this study were of C57/Bl6x129 (WT, $\alpha 4\Delta/\Delta$) or C57/Bl6 ($\beta 2$ -/-) background, between 8 and 12 weeks of age. Wild type (WT) animals were purchased from Taconic (Germantown, NY). Beta 2 integrin-deficient mice were obtained from Dr. A. Beudet (Baylor College, Houston, TX) [16]. MxCre⁺ $\alpha 4$ ^{f/f} mice were generated in our laboratory [20]. To induce $\alpha 4$ integrin ablation, these mice were treated neonatally with interferon inducer, poly (I:C) (three injections of 50 μ l of 1mg/ml solution in phosphate-buffered saline (PBS), intraperitoneally (i.p.), 48 hours apart). MMP9^{-/-} mice were kindly provided by Dr R. Senior (Washington University, St. Louis, MO) [21]. All animals were bred and maintained under specific pathogen-free conditions at the University of Washington. All experimental procedures were done in accordance with Institutional Animal Care and Use Committee guidelines on approved protocols.

Antibodies

Anti- $\alpha 4$ integrin antibody, PS/2, was purchased from Southern Biotechnology (Birmingham, AL). PE-Cy5-conjugated F4/80 (Cl:A3-1) was from AbD Serotec Ltd (Raleigh, NC). Gr-1-PE, Gr1-APC, CD45-APC, CD45-FITC, CD3-Cy-chrome, CD4-CyChrome, CD8-FITC, B220-FITC, B220-PE, B220-CyChrome, CD19-biotin, CD18-PE, streptavidin-APC, and

corresponding fluorochrome-conjugated isotype-matched immunoglobulins that served as controls were purchased from BD Biosciences (San Diego, CA)

Peritoneal inflammation

Mice were injected i.p. with 1ml of 3% thioglycolate (TG) and were sacrificed by cervical dislocation 4, 16, 48 and 96 hours post injection. The peritoneal cavity was lavaged with 5 ml of ice-cold phosphate-buffered saline (PBS) containing 5mM EDTA, cells were enumerated and used for further analysis.

Chemokine measurement

Aliquots of frozen peritoneal lavage fluid were sent to Pierce Biotechnology Inc. (Woburn, MA) to measure concentration of an array of chemokines and cytokines by Searchlight Multiplex technology.

FACS analysis

Antibody-labeled cells were analyzed on a FACSCalibur (BD Biosciences, San Jose, CA) using CellQuest software.

In vitro migration

Transwell migration was performed as described elsewhere [22]. In brief, splenocytes were stained with B220-FITC and CD3-PE and 0.5×10^6 cells were transferred into the upper chambers of transwell inserts (pore size $5\mu\text{m}$, Corning Costar, Cambridge, MA). Splenocytes were allowed to migrate through the uncoated inserts (trans-well migration), or transwells coated with bEND3 mouse endothelial cells (trans-endothelial migration) for 4 hours at 37°C towards SDF-1 α (100 ng/ml, Peprotech, Rocky Hill, NJ). Cells collected from the lower chamber were enumerated by FACS. The number of events acquired from this sample was compared to that of input cells (before migration) and the percentage of migration was calculated.

Adhesion to endothelial cells

Splenocytes were isolated and labeled with CarboxyFluorescein Succinimidyl Ester, CFSE (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions, washed and brought to 2×10^6 cells/ml in adhesion buffer (PBS containing 1% BSA, 2mM MgCl_2 , 2mM CaCl_2). Labeled cells were transferred into 24-well plates (1×10^6 cells/well) with a confluent layer of bEND3 cells either untreated or treated overnight with TNF- α (100 ng/ml, Peprotech). After a short spin (2 min., $100\times\text{g}$), adhesion was allowed to occur at 37°C for 40 minutes. Non-adherent cells were washed out with PBS. bEND3 cells and adherent splenocytes were trypsinized with Tryple Select (Invitrogen), washed, and the number of CFSE-positive cells was determined by FACS after 1 min. acquisition. Cell adhesion was expressed as a percent of input cells.

Actin polymerization

Actin polymerization was performed as described [23]. In brief, splenocytes (1×10^6 cells) were stained with CD3-FITC and B220-FITC, stimulated with 100 ng/ml SDF-1 α (Peprotech) for indicated amount of time, fixed, permeabilized, and stained with phalloidin (Invitrogen). After washing with PBS/BSA, cells, gated on lymphocytes (CD3+B220+), were analyzed by FACS.

Ca²⁺ mobilization

Ca²⁺ mobilization experiments were performed as described elsewhere [24]. Splenocytes were isolated and loaded with Indo-1 (Invitrogen), washed and labeled with B220-FITC and CD3-

PE antibodies. Cells were resuspended in Ca^{2+} buffer (PBS, 1% BSA, 1mM CaCl_2 , 1mM MgCl_2) at 4×10^6 cells/ml. Change of the 395/530 fluorescence quotient was monitored over 3 min. by FACS following stimulation with SDF-1 α (200 ng/ml, Peprotech) or Ionomycin (1 μM , Sigma). After analysis using FloJo software, Ca^{2+} mobilization was expressed as a percent of the peak levels after stimulation over baseline.

Statistical analysis

Data shown are means \pm standard error of mean (sem). Statistical analyses were performed using a Student *t* test and P values of < 0.05 were considered significant.

RESULTS

Animal models

In the present study we used genetic models of $\alpha 4$ ($\alpha 4\beta 1$ and $\alpha 4\beta 7$) or $\beta 2$ (CD18) integrin deficiency. Beta 2 integrin deficient (CD18 $^{-/-}$) mice were generated by germ line deletion of $\beta 2$ integrin gene [16]. Alpha4 integrin-deficient mice ($\alpha 4\Delta/\Delta$) were previously generated in our laboratory as conditional knockouts. These were MxCre $^{+}\alpha 4^{f/f}$ mice, in which ablation of $\alpha 4$ integrins occurs after treatment with interferon inducer, poly(I:C) [20]. After ablation of $\alpha 4$ integrin in these mice, percent of $\alpha 4^{+}$ cells was reduced to $3.7\% \pm 0.2\%$ ($n=11$) in the bone marrow and $5.2\% \pm 0.8\%$ ($n=11$) in the peripheral blood. White blood cell (WBC) counts in both $\alpha 4$ integrin-deficient ($19 \pm 2 \times 10^3$ cells/ μl) and $\beta 2$ integrin-deficient ($49 \pm 5 \times 10^3$ cells/ μl) mice were significantly higher than in control mice ($8 \pm 1 \times 10^3$ cells/ μl), as previously described [16,20]. In addition, we recently generated mice with both $\alpha 4$ and $\beta 2$ integrin deficiency by interbreeding single integrin knockouts. Here we report only preliminary results with $\alpha 4\Delta/\Delta \beta 2^{-/-}$ doubly deficient mice, since only a handful of animals were available for experiments due to difficulties in breeding.

To study unique and redundant roles of $\alpha 4$ and $\beta 2$ integrins in leukocyte migration *in vivo*, we employed the aseptic peritonitis model in all mice with integrin deficiencies and monitored various leukocyte subsets by comparing their levels in circulation and in the peritoneal cavity for intervals up to 4 days.

Kinetics of migration of various leukocyte subsets *in vivo*

Neutrophils are the first leukocytes that migrate and accumulate at a site of inflammation. By the 4th hour of peritonitis, a substantial number of neutrophils was detected in the peritoneum of WT and $\alpha 4\Delta/\Delta$ mice and increased further by the 16th hour (Fig. 1A). In $\beta 2^{-/-}$ mice, neutrophils were slower to migrate, but by the 16th hour of peritonitis their numbers were comparable to those in WT animals (Fig. 1A). Defects in optimal neutrophil recruitment became evident when we compared recruitment indices in $\alpha 4$ and $\beta 2$ integrin-deficient versus control mice (Fig. 1B). Recruitment index represents the proportion of cells harvested from the peritoneal cavity measured against the total number of cells available for migration in the circulation, assuming a blood volume of 2 ml in the mouse. The defect was more pronounced in $\beta 2^{-/-}$ mice: recruitment indices at each time point were less than 2% of corresponding control value, whereas in $\alpha 4\Delta/\Delta$ mice at 4 and 16 hours, they comprised 46% and 34%, respectively, of the corresponding control value. This observed defect in $\beta 2^{-/-}$ neutrophil recruitment was even more striking, in view of elevated levels of neutrophil chemoattractant MIP2 in the peritoneal cavity of $\beta 2^{-/-}$ mice (267 ± 63 pg/ml) as compared to controls (74 ± 5 pg/ml, $P < 0.05$). In the $\alpha 4$ -deficient mice, MIP2 levels were similar to controls, as were KC levels in both $\alpha 4^{-/-}$ and $\beta 2^{-/-}$ mice (data not shown).

At later times of peritonitis (96 hours) more neutrophils were harvested from the peritoneum of both $\alpha 4$ ($0.4 \pm 0.03 \times 10^6$ cells, $P < 0.01$) and $\beta 2$ ($3.7 \pm 0.9 \times 10^6$ cells, $P < 0.01$) integrin-deficient mice as compared to controls ($0.1 \pm 0.02 \times 10^6$ cells).

It is worth noting that, in WT mice, all peritoneal neutrophils were $\alpha 4$ -positive, whereas in $\alpha 4$ -deficient mice, all migrated neutrophils were $\alpha 4$ -negative (data not shown), suggesting their $\alpha 4$ -independent migration.

As a general rule, macrophage recruitment follows that of neutrophils during inflammation. We first analyzed inflammatory monocytes, a subset that is recognized by the presence of Gr-1 marker in addition to the monocytic marker, F4/80 [25] (Fig. 2A). Although there was no difference in the numbers of inflammatory monocytes at 4 hours, at the peak of their accumulation (16 hours), significantly fewer inflammatory monocytes were recovered from peritoneum of either $\alpha 4$ or $\beta 2$ -integrin-deficient mice than in controls (Fig. 2B), despite their increased numbers in circulation (Fig. 2C). Calculated recruitment indices further emphasized impairment in recruitment of inflammatory monocytes in integrin-deficient mice (Fig. 2D). The observed defect occurred despite normal levels of MCP1 and MIP1 α , mononuclear cell chemokines, in the peritoneal cavity of these mice (data not shown).

We next studied accumulation of total peritoneal macrophages (F4/80-positive cells). The numbers of F4/80+ cells recovered from the peritoneum were similar in integrin-deficient and control mice, except for a transient but significant decrease in macrophage migration at 16 hours in $\alpha 4$ -deficient mice (Fig. 3A). Of interest, at steady state, 37.61% \pm 3.42% of peritoneal F4/80+ macrophages were $\alpha 4$ -positive and this proportion gradually decreased after TG injection to 32.71% \pm 5.99% at 4 hours, 13.16% \pm 3.98% at 16 hours, and further to 2.78% \pm 1.88% at 96 hours. While the decrease in proportion of $\alpha 4$ -positive macrophages was likely attributable in part to dilution by influxing $\alpha 4$ -negative macrophages, their total numbers also decreased with time (Fig. 3B) in $\alpha 4$ -deficient mice.

To investigate recruitment of lymphocytes, we compared numbers of T and B cells in the peritoneal lavage of naïve mice and 48 hours after TG injection, at which time the peak of lymphocyte recruitment is observed [26]. In $\beta 2^{-/-}$ mice, although the lymphocyte numbers in peritoneal cavity ($2.05 \times 10^6 \pm 0.45 \times 10^6$ cells) were similar to controls ($1.15 \times 10^6 \pm 0.19 \times 10^6$ cells) at steady state, a defect in lymphocyte recruitment became evident after inflammation was established: TG injection resulted in less than 2-fold increase over baseline in $\beta 2^{-/-}$ lymphocyte numbers that is significantly less than a 7-fold increase in controls (Fig. 4A) and the recruitment index was also significantly decreased (Fig. 4B). In non-stimulated $\alpha 4$ -deficient animals, even though only a small number was recovered from the peritoneal cavity, 47.1% \pm 7.6% of T cells and 71.4% \pm 8.1% of B cells were $\alpha 4$ -positive, in contrast to 5.2% \pm 0.8% of $\alpha 4$ -positive cells present in the peripheral blood. These data suggest preferential migration and accumulation over time of $\alpha 4$ -positive cells at steady state. Very few $\alpha 4$ -negative lymphocytes were detected in peritoneal cavities of $\alpha 4$ -deficient mice, and this number did not change after TG injection (Fig. 4A). The recruitment index for $\alpha 4^{-/-}$ lymphocytes was 45-fold less than for control (Fig. 4B), indicating a profound defect in $\alpha 4^{-/-}$ lymphocyte migration and recruitment. Of interest, a decrease in levels of lymphocyte chemoattractant RANTES in the peritoneal cavity was observed in $\alpha 4$ -deficient relative to control mice (6.37 ± 1.42 pg/ml and 14.63 ± 1.65 pg/ml, respectively, $P < 0.05$) whereas measured RANTES levels in $\beta 2^{-/-}$ mice, were similar to controls (data not shown). The majority of migrated lymphocytes in all TG-induced peritonitis mouse models studied was comprised of B cells (data not shown).

In the WT mice, TG injection resulted in transient leucopenia: at 4 and 16 hours of peritonitis circulating WBC numbers dropped to $3.2 \pm 0.5 \times 10^3$ and $4.0 \pm 0.5 \times 10^3$ cells/ μ l blood, respectively, but returned to basal levels by the 96th hour ($7.2 \pm 0.4 \times 10^3$ cells/ μ l blood). In

contrast, in the integrin-deficient mice, no significant changes in WBC counts were detected over the course of peritonitis as compared to baseline (data not shown).

We recently developed animals with both $\alpha 4$ and $\beta 2$ integrin deficiency by interbreeding the single integrin-deficient mice. The baseline WBC counts ($109 \pm 10 \times 10^3$ cells/ μ l, n=14) reached unprecedented levels in these doubly deficient mice, likely reflecting not only persistent inflammatory conditions, but also profound defects in leukocyte emigration to tissues. Indeed, peritoneal leukocyte numbers at 0, 4, 16 and 96 hours after TG injection were $5.6 \pm 0.3 \times 10^6$ (n=4), $4.4 \pm 2.1 \times 10^6$ (n=4), $5.9 \pm 2.6 \times 10^6$ (n=3), and $5.4 \pm 2.4 \times 10^6$ (n=3) cells respectively, indicating abrogation of leukocyte recruitment in the absence of both integrins, despite markedly elevated circulating WBCs (Fig. 5). Comparison of leukocyte recruitment in single integrin- and double-deficient animals indicates that both integrins are absolutely required for leukocyte migration to the peritoneal cavity.

In vitro functional testing of $\alpha 4\Delta/\Delta$ lymphocytes

Considering that the most prominent defect in recruitment to the inflamed peritoneum involved the $\alpha 4\Delta/\Delta$ lymphocytes, we further investigated their migratory behavior in vitro. We compared adhesion of control and $\alpha 4\Delta/\Delta$ lymphocytes to TNF α -stimulated (VCAM-1 expressing) and non-stimulated bEND3 cells, a mouse endothelial cell line. Twice as many control lymphocytes adhered to VCAM-1-expressing bEND3 cells as compared to non-stimulated cells; at the same time, no increase in adhesion was observed with $\alpha 4\Delta/\Delta$ lymphocytes (Fig. 6A). These data suggest the preferential usage of $\alpha 4$ integrins by lymphocytes for adhesion to bEND3 cells, since TNF α not only up-regulates VCAM-1 but also ICAM-1, a $\beta 2$ integrin ligand on endothelial cells [27,28]. To determine whether ability to migrate per se is preserved in $\alpha 4\Delta/\Delta$ lymphocytes, we performed migration through an uncoated Transwell insert (trans-well migration) not requiring engagement of $\alpha 4$ integrins. As a stimulus for migration we used SDF-1 α , a well-characterized lymphocyte chemoattractant agent [29]. In this setting, since equal amounts of chemoattractant are present, differences in migration would reflect cell intrinsic defects. Substantial migration of B- and T-lymphocytes in all genotypes except for $\alpha 4\Delta/\Delta$ B-cells was observed (Fig. 6B and data not shown), suggesting impairment of signaling in these cells. We next performed lymphocyte migration through a Transwell coated with bEND3 cells (trans-endothelial migration). Lymphocytes were allowed to migrate through non-stimulated bEND3 cells or cells stimulated with TNF α upregulating VCAM-1. Control and $\beta 2^{-/-}$ B-cells migrated more efficiently through TNF α -treated endothelium; in contrast, $\alpha 4\Delta/\Delta$ B-cells showed no increase in migration through VCAM1-expressing bEND3 cells (Fig. 6C). Similar results were seen with T-cells (data not shown). Of interest, significantly fewer $\beta 2^{-/-}$ B-cells migrated through the non-stimulated endothelial layer, whereas their migration through VCAM-1-expressing bEND3 cells was similar to control indicating that migration through the TNF α -stimulated endothelial cells is superseded by $\alpha 4$ integrins via VLA4/VCAM-1 interaction. Since trans-well migration was impaired in $\alpha 4\Delta/\Delta$ lymphocytes, we tested early signaling events characteristic of this process. Actin polymerization is a very early event in migration-associated signaling in response to SDF-1 α . No significant differences in actin polymerization, measured by phalloidin staining, were detected in either integrin-deficient lymphocytes or control cells, suggesting preservation of proximal signaling steps leading to actin remodeling (Fig. 7A). At the same time, absence of $\alpha 4$ integrin in lymphocytes resulted in a modest but significant decrease in SDF-1 α -induced Ca²⁺ mobilization (Fig. 7B), likely reflecting impaired signaling. These data suggest impaired response of $\alpha 4$ -deficient lymphocytes to soluble chemoattractants.

DISCUSSION

Although the lack of either $\alpha 4$ or $\beta 2$ integrin resulted in decreased efficiency of leukocyte recruitment to the inflamed peritoneum, the substantial numbers of total leukocytes observed at various times after TG injection emphasize the overlapping roles of $\alpha 4$ and $\beta 2$ integrins in total leukocyte migration. These data are consistent with previous studies showing that only simultaneous functional blockade of $\alpha 4$ and $\beta 2$ integrins (either by treatment of wild type mice with a combination of anti- $\alpha 4$ and anti- $\beta 2$ antibodies, or by treatment of $\beta 2$ -deficient mice with anti- $\alpha 4$ -antibody) prevents leukocyte recruitment to the peritoneum [15,30,31]. Our data with $\alpha 4/\beta 2$ doubly deficient mice support and extend these observations. In addition, they provide direct evidence for an absence of significant compensation by other cytoadhesins [32,33] in these animals.

Our studies of the different leukocyte subsets in genetically deficient animals further delineate distinct versus overlapping contributions of $\alpha 4$ and $\beta 2$ integrins to leukocyte subset-specific migration.

Although the presence of either $\alpha 4$ or $\beta 2$ integrins afforded significant migration of neutrophils to inflamed peritoneum (Fig. 1A), assessment of recruitment indices (Fig. 1B) indicates that optimal recruitment requires the presence of both $\alpha 4$ and $\beta 2$ integrins. The interchangeable usage of $\alpha 4$ or $\beta 2$ integrins by neutrophils may be tissue-specific. While neutrophil migration to the heart and lung tissues is mediated by both $\alpha 4$ and $\beta 2$ integrins [34,35], migration of neutrophils to the skin in non-allergic inflammation depends primarily on $\beta 2$ integrins [17, 36]. Tissue-specific variations in chemokine secretion and expression of adhesion molecules in local inflammatory milieu may explain different patterns of integrin usage by leukocytes during their migration to the various tissues [37].

Of interest are observations regarding the recruitment of monocyte/macrophages in $\alpha 4$ -deficient mice. Macrophage numbers in these mice, although low in early development of peritonitis (Fig. 3A) due to delayed recruitment of inflammatory monocytes to the site of inflammation (Fig. 2), continually increased up to 96 hours (Fig. 3B). As the number of $\alpha 4$ -negative macrophages increased, the number of $\alpha 4$ -positive macrophages progressively decreased during the course of inflammation (Fig. 3B). The decline in total numbers did not simply reflect “dilution” by the incoming $\alpha 4$ -negative cells, but is best explained by their egress from the peritoneal cavity, as was previously shown for normal macrophages [38]. Macrophage egress from the peritoneum observed in control mice (compare the drop in macrophage numbers between 16 and 96 hours of peritonitis, Fig. 3A) was impaired in both $\alpha 4$ and $\beta 2$ integrin-deficient mice, suggesting that the presence of $\alpha 4$ integrins, along with the $\beta 2$ -integrin Mac1 [39], is important in their egress from inflammatory sites.

In contrast to macrophages which migrate from the peritoneal cavity into the draining lymphatics [38], neutrophils undergo apoptosis and are cleared from the peritoneum by macrophages. Thus, increased neutrophil numbers observed in both $\alpha 4\Delta/\Delta$ and $\beta 2^{-/-}$ mice at 96 hours post TG injection may suggest either delay in apoptosis, impaired clearance, or sustained neutrophil recruitment. Involvement of both $\alpha 4$ and $\beta 2$ integrins in regulating apoptosis and/or clearing of apoptotic cells have been demonstrated; deficiency in $\beta 2$ integrins leads to a delay in neutrophil apoptosis and their subsequent accumulation in the peritoneum [40,41], while $\alpha 4$ integrin on apoptotic cells assist in their phagocytosis [42]. Thus, absence of $\alpha 4$ integrins may affect clearance of $\alpha 4\Delta/\Delta$ neutrophils, thereby resulting in increased neutrophil presence in the peritoneum.

Lymphocyte migration to the inflamed peritoneum was greatly impaired only in $\alpha 4$ integrin-deficient mice (Fig. 4). Our results are in agreement with previously reported observations in chimeric mice in which $\alpha 4\Delta/\Delta$ lymphocytes did not migrate into the inflamed peritoneum

[19]. However, the dependence of lymphocyte recruitment on $\alpha 4$ integrin is also a tissue-specific phenomenon; recruitment of lymphocytes to brain tissue is mainly dependent on $\alpha 4$ integrin [43], whereas their migration to inflamed lungs and airways is mediated by both $\alpha 4$ and $\beta 2$ integrins [11,44,45], and migration to inflamed skin depends exclusively upon $\beta 2$ integrin engagement [46].

It is important to emphasize that, beyond the complete and partial impairment in lymphocyte recruitment seen in $\alpha 4$ and $\beta 2$ -deficient mice, respectively, the defects were manifested differently: $\alpha 4\Delta/\Delta$ lymphocytes showed defective migration to the non-inflamed as well as to the inflamed peritoneum; the defect in $\beta 2^{-/-}$ lymphocytes was partial and became apparent only after inflammation was established (Fig. 4A). This finding is in agreement with preferential usage of $\alpha 4$ over $\beta 2$ integrin by lymphocytes for adhesion to the extracellular matrix before and after inflammatory stimulus [47]. However, whether integrin-specific recruitment changes under shear stress is unclear. For example, preferential usage of $\beta 2$ integrin by T-lymphocytes in response to chemoattractive stimuli under shear stress conditions has been reported [48]. In addition, considering pre-existing high systemic levels of inflammatory cytokines in CD18 $^{-/-}$ mice [49], the partial defect seen in $\beta 2^{-/-}$ lymphocytes may also be attributed to an attenuated response to TG of already activated CD18 $^{-/-}$ lymphocytes by pre-existing inflammatory stimuli.

The reasons for impaired $\alpha 4\Delta/\Delta$ lymphocyte recruitment to the inflamed peritoneum are attributable mainly to intrinsic cell defects, rather than being dictated by the environment. This is supported by the fact that $\alpha 4\Delta/\Delta$ lymphocytes showed impaired adhesion and migration to SDF-1 α (both trans-well and trans-endothelial migration) (Fig. 6B, C). Consistent with these results, perturbed $\alpha 4$ integrin/paxillin-dependent signaling also leads to defective lymphocyte migration to the inflamed peritoneum [26]. Beyond the intrinsic migratory defects of $\alpha 4\Delta/\Delta$ lymphocytes, suboptimal stimulation by environmental factors (i.e. decreased levels of RANTES) cannot be excluded. The metalloproteinases (MMP) that are thought to contribute to leukocyte recruitment [50,51] most likely do not play a major role in the model employed, as we detected neither lack of induction of either MMP9 or MMP2 in the peritoneal cavity during peritonitis, nor any difference in trans-endothelial migration of MMP9 $^{-/-}$ lymphocytes compared to controls (data not shown). This is in agreement with the observation of Betsuyaku et. al., that MMP9 is not required for efficient leukocyte migration [52].

In summary, our results demonstrate that, although critically dependent on the expression of both $\alpha 4$ and $\beta 2$ integrins, there is a subset-specific usage of these adhesion molecules by leukocytes in their migration to and egress from the inflamed peritoneum. While the roles of $\alpha 4$ and $\beta 2$ integrins are partially redundant in myeloid cells, they are distinct and non-overlapping in lymphocytes. Taken together, our data with genetically deficient mouse models extend and refine previous knowledge on the kinetics of leukocyte accumulation and their egress from the peritoneum in the aseptic peritonitis model. Whether kinetic changes in inflammatory models of other tissues are different will require further studies.

ACKNOWLEDGMENTS

This work was supported by the NIH grants (HL58734, DK46557) for T.P. Authors thank Ms. Devra Batdorf for expert mouse handling.

REFERENCES

1. Alon R, Feigelson S. From rolling to arrest on blood vessels: leukocyte tap dancing on endothelial integrin ligands and chemokines at sub-second contacts. *Semin Immunol* 2002;14:93–104. [PubMed: 11978081]

2. Yadav R, Larbi KY, Young RE, Nourshargh S. Migration of leukocytes through the vessel wall and beyond. *Thromb Haemost* 2003;90:598–606. [PubMed: 14515179]
3. Chan JR, Hyduk SJ, Cybulsky MI. Chemoattractants induce a rapid and transient upregulation of monocyte alpha4 integrin affinity for vascular cell adhesion molecule 1 which mediates arrest: an early step in the process of emigration. *J Exp Med* 2001;193:1149–1158. [PubMed: 11369786]
4. Kitayama J, Fuhlbrigge RC, Puri KD, Springer TA. P-selectin, L-selectin, and alpha 4 integrin have distinct roles in eosinophil tethering and arrest on vascular endothelial cells under physiological flow conditions. *J Immunol* 1997;159:3929–3939. [PubMed: 9378981]
5. Johnston B, Issekutz TB, Kubes P. The alpha 4-integrin supports leukocyte rolling and adhesion in chronically inflamed postcapillary venules in vivo. *J Exp Med* 1996;183:1995–2006. [PubMed: 8642310]
6. Yusuf-Makagiansar H, Anderson ME, Yakovleva TV, Murray JS, Siahaan TJ. Inhibition of LFA-1/ICAM-1 and VLA-4/VCAM-1 as a therapeutic approach to inflammation and autoimmune diseases. *Med Res Rev* 2002;22:146–167. [PubMed: 11857637]
7. Ibbotson GC, Doig C, Kaur J, et al. Functional alpha4-integrin: a newly identified pathway of neutrophil recruitment in critically ill septic patients. *Nat Med* 2001;7:465–470. [PubMed: 11283674]
8. Pereira S, Zhou M, Mocsai A, Lowell C. Resting murine neutrophils express functional alpha 4 integrins that signal through Src family kinases. *J Immunol* 2001;166:4115–4123. [PubMed: 11238661]
9. Petruzzelli L, Takami M, Humes HD. Structure and function of cell adhesion molecules. *Am J Med* 1999;106:467–476. [PubMed: 10225251]
10. Mulligan MS, Lentsch AB, Miyasaka M, Ward PA. Cytokine and adhesion molecule requirements for neutrophil recruitment during glycogen-induced peritonitis. *Inflamm Res* 1998;47:251–255. [PubMed: 9683032]
11. Gascoigne MH, Holland K, Page CP, et al. The effect of anti-integrin monoclonal antibodies on antigen-induced pulmonary inflammation in allergic rabbits. *Pulm Pharmacol Ther* 2003;16:279–285. [PubMed: 12877819]
12. Henderson WR Jr, Chi EY, Albert RK, et al. Blockade of CD49d (alpha4 integrin) on intrapulmonary but not circulating leukocytes inhibits airway inflammation and hyperresponsiveness in a mouse model of asthma. *J Clin Invest* 1997;100:3083–3092. [PubMed: 9399955]
13. Abraham WM, Sielczak MW, Ahmed A, et al. Alpha 4-integrins mediate antigen-induced late bronchial responses and prolonged airway hyperresponsiveness in sheep. *J Clin Invest* 1994;93:776–787. [PubMed: 8113411]
14. Abraham WM, Gill A, Ahmed A, et al. A small-molecule, tight-binding inhibitor of the integrin alpha (4)beta(1) blocks antigen-induced airway responses and inflammation in experimental asthma in sheep. *Am J Respir Crit Care Med* 2000;162:603–611. [PubMed: 10934094]
15. Henderson RB, Lim LH, Tessier PA, et al. The use of lymphocyte function-associated antigen (LFA)-1-deficient mice to determine the role of LFA-1, Mac-1, and alpha4 integrin in the inflammatory response of neutrophils. *J Exp Med* 2001;194:219–226. [PubMed: 11457896]
16. Scharffetter-Kochanek K, Lu H, Norman K, et al. Spontaneous skin ulceration and defective T cell function in CD18 null mice. *J Exp Med* 1998;188:119–131. [PubMed: 9653089]
17. Mizgerd JP, Kubo H, Kutkoski GJ, et al. Neutrophil emigration in the skin, lungs, and peritoneum: different requirements for CD11/CD18 revealed by CD18-deficient mice. *J Exp Med* 1997;186:1357–1364. [PubMed: 9334375]
18. Yang JT, Rayburn H, Hynes RO. Cell adhesion events mediated by alpha 4 integrins are essential in placental and cardiac development. *Development* 1995;121:549–560. [PubMed: 7539359]
19. Arroyo AG, Taverna D, Whittaker CA, et al. In vivo roles of integrins during leukocyte development and traffic: insights from the analysis of mice chimeric for alpha 5, alpha v, and alpha 4 integrins. *J Immunol* 2000;165:4667–4675. [PubMed: 11035110]
20. Scott LM, Priestley GV, Papayannopoulou T. Deletion of alpha4 integrins from adult hematopoietic cells reveals roles in homeostasis, regeneration, and homing. *Mol Cell Biol* 2003;23:9349–9360. [PubMed: 14645544]
21. Vu TH, Shipley JM, Bergers G, et al. MMP-9/gelatinase B is a key regulator of growth plate angiogenesis and apoptosis of hypertrophic chondrocytes. *Cell* 1998;93:411–422. [PubMed: 9590175]

22. Andrew DP, Spellberg JP, Takimoto H, Schmits R, Mak TW, Zukowski MM. Transendothelial migration and trafficking of leukocytes in LFA-1-deficient mice. *Eur J Immunol* 1998;28:1959–1969. [PubMed: 9645378]
23. Papayannopoulou T, Priestley GV, Bonig H, Nakamoto B. The role of G-protein signaling in hematopoietic stem/progenitor cell mobilization. *Blood* 2003;101:4739–4747. [PubMed: 12595315]
24. Bonig H, Rohmer L, Papayannopoulou T. Long-term functional impairment of hemopoietic progenitor cells engineered to express the S1 catalytic subunit of pertussis toxin. *Exp Hematol* 2005;33:689–698. [PubMed: 15911093]
25. Geissmann F, Jung S, Littman DR. Blood monocytes consist of two principal subsets with distinct migratory properties. *Immunity* 2003;19:71–82. [PubMed: 12871640]
26. Feral CC, Rose DM, Han J, et al. Blocking the alpha 4 integrin-paxillin interaction selectively impairs mononuclear leukocyte recruitment to an inflammatory site. *J Clin Invest* 2006;116:715–723. [PubMed: 16470243]
27. Kuldo JM, Westra J, Asgeirsdottir SA, et al. Differential effects of NF- κ B and p38 MAPK inhibitors and combinations thereof on TNF- α - and IL-1 β -induced proinflammatory status of endothelial cells in vitro. *Am J Physiol Cell Physiol* 2005;289:C1229–C1239. [PubMed: 15972838]
28. Min JK, Kim YM, Kim SW, et al. TNF-related activation-induced cytokine enhances leukocyte adhesiveness: induction of ICAM-1 and VCAM-1 via TNF receptor-associated factor and protein kinase C-dependent NF- κ B activation in endothelial cells. *J Immunol* 2005;175:531–540. [PubMed: 15972689]
29. Ding Z, Xiong K, Issekutz TB. Chemokines stimulate human T lymphocyte transendothelial migration to utilize VLA-4 in addition to LFA-1. *J Leukoc Biol* 2001;69:458–466. [PubMed: 11261794]
30. Winn RK, Harlan JM. CD18-independent neutrophil and mononuclear leukocyte emigration into the peritoneum of rabbits. *J Clin Invest* 1993;92:1168–1173. [PubMed: 8104195]
31. Henderson RB, Hobbs JA, Mathies M, Hogg N. Rapid recruitment of inflammatory monocytes is independent of neutrophil migration. *Blood* 2003;102:328–335. [PubMed: 12623845]
32. Katayama Y, Hidalgo A, Chang J, Peired A, Frenette PS. CD44 is a physiological E-selectin ligand on neutrophils. *J Exp Med* 2005;201:1183–1189. [PubMed: 15824084]
33. Lenter M, Levinovitz A, Isenmann S, Vestweber D. Monospecific and common glycoprotein ligands for E- and P-selectin on myeloid cells. *J Cell Biol* 1994;125:471–481. [PubMed: 7512971]
34. Bowden RA, Ding ZM, Donnachie EM, et al. Role of alpha4 integrin and VCAM-1 in CD18-independent neutrophil migration across mouse cardiac endothelium. *Circ Res* 2002;90:562–569. [PubMed: 11909820]
35. Burns JA, Issekutz TB, Yagita H, Issekutz AC. The alpha 4 beta 1 (very late antigen (VLA)-4, CD49d/CD29) and alpha 5 beta 1 (VLA-5, CD49e/CD29) integrins mediate beta 2 (CD11/CD18) integrin-independent neutrophil recruitment to endotoxin-induced lung inflammation. *J Immunol* 2001;166:4644–4649. [PubMed: 11254723]
36. Mizgerd JP, Bullard DC, Hicks MJ, Beaudet AL, Doerschuk CM. Chronic inflammatory disease alters adhesion molecule requirements for acute neutrophil emigration in mouse skin. *J Immunol* 1999;162:5444–5448. [PubMed: 10228023]
37. Hillyer P, Mordelet E, Flynn G, Male D. Chemokines, chemokine receptors and adhesion molecules on different human endothelia: discriminating the tissue-specific functions that affect leukocyte migration. *Clin Exp Immunol* 2003;134:431–441. [PubMed: 14632748]
38. Bellingan GJ, Xu P, Cooksley H, et al. Adhesion molecule-dependent mechanisms regulate the rate of macrophage clearance during the resolution of peritoneal inflammation. *J Exp Med* 2002;196:1515–1521. [PubMed: 12461086]
39. Cao C, Lawrence DA, Strickland DK, Zhang L. A specific role of integrin Mac-1 in accelerated macrophage efflux to the lymphatics. *Blood* 2005;106:3234–3241. [PubMed: 16002427]
40. Weinmann P, Scharffetter-Kochanek K, Forlow SB, Peters T, Walzog B. A role for apoptosis in the control of neutrophil homeostasis in the circulation: insights from CD18-deficient mice. *Blood* 2003;101:739–746. [PubMed: 12393639]

41. Coxon A, Rieu P, Barkalow FJ, et al. A novel role for the beta 2 integrin CD11b/CD18 in neutrophil apoptosis: a homeostatic mechanism in inflammation. *Immunity* 1996;5:653–666. [PubMed: 8986723]
42. Johnson JD, Hess KL, CookMills JM. CD44, alpha(4) integrin, and fucoidin receptor-mediated phagocytosis of apoptotic leukocytes. *J Leukoc Biol* 2003;74:810–820. [PubMed: 12960273]
43. Deloire MS, Touil T, Brochet B, Dousset V, Caille JM, Petry KG. Macrophage brain infiltration in experimental autoimmune encephalomyelitis is not completely compromised by suppressed T-cell invasion: in vivo magnetic resonance imaging illustration in effective anti-VLA-4 antibody treatment. *Mult Scler* 2004;10:540–548. [PubMed: 15471371]
44. Xu B, Wagner N, Pham LN, et al. Lymphocyte homing to bronchus-associated lymphoid tissue (BALT) is mediated by L-selectin/PNAd, alpha4beta1 integrin/VCAM-1, and LFA-1 adhesion pathways. *J Exp Med* 2003;197:1255–1267. [PubMed: 12756264]
45. Banerjee ER, Jiang Y, Henderson WR, Scott LM, Papayannopoulou T. a4 and b2 integrins have nonredundant roles for asthma development, but for optimal allergen sensitization only a4 is critical. *Exp Hematol* 2007;35:605–617. [PubMed: 17379071]
46. Grabbe S, Varga G, Beissert S, et al. Beta2 integrins are required for skin homing of primed T cells but not for priming naive T cells. *J Clin Invest* 2002;109:183–192. [PubMed: 11805130]
47. Solpov A, Shenkman B, Vitkovsky Y, et al. Platelets enhance CD4+ lymphocyte adhesion to extracellular matrix under flow conditions: role of platelet aggregation, integrins, and non-integrin receptors. *Thromb Haemost* 2006;95:815–821. [PubMed: 16676073]
48. Schreiber TH, Shinder V, Cain DW, Alon R, Sackstein R. Shear flow-dependent integration of apical and subendothelial chemokines in T-cell transmigration: implications for locomotion and the multistep paradigm. *Blood* 2007;109:1381–1386. [PubMed: 17038526]
49. Wu H, Prince JE, Brayton CF, et al. Host resistance of CD18 knockout mice against systemic infection with *Listeria monocytogenes*. *Infect Immun* 2003;71:5986–5993. [PubMed: 14500519]
50. Kolaczowska E, Chadzinska M, Scislowska-Czarnecka A, Plytycz B, Opendakker G, Arnold B. Gelatinase B/matrix metalloproteinase-9 contributes to cellular infiltration in a murine model of zymosan peritonitis. *Immunobiology* 2006;211:137–148. [PubMed: 16530081]
51. Renckens R, Roelofs JJ, Florquin S, et al. Matrix metalloproteinase-9 deficiency impairs host defense against abdominal sepsis. *J Immunol* 2006;176:3735–3741. [PubMed: 16517742]
52. Betsuyaku T, Shipley JM, Liu Z, Senior RM. Neutrophil emigration in the lungs, peritoneum, and skin does not require gelatinase B. *Am J Respir Cell Mol Biol* 1999;20:1303–1309. [PubMed: 10340950]




Figure 1. Both $\alpha 4$ and $\beta 2$ integrins are required for efficient neutrophil migration to the peritoneum Mice were injected with TG, peritoneal leukocytes were harvested 4 and 16 hours post injection and stained with various leukocyte markers (see Materials and Methods). Neutrophil (Gr-1 +, F4/80-) numbers (A) and recruitment indices (B) were calculated. Note that although sufficient numbers of neutrophils are present in the peritoneal cavity of $\alpha 4\Delta/\Delta$ and $\beta 2^{-/-}$ mice, there is a significant decrease in recruitment index at each time point. Asterisk (*) indicates significant difference compared to controls, $P < 0.05$. Mice used per group: WT, n=5; $\beta 2^{-/-}$, n=4; $\alpha 4\Delta/\Delta$, n=4.




Figure 2. Recruitment of inflammatory monocytes to the peritoneum is impaired in $\alpha 4\Delta/\Delta$ and $\beta 2^{-/-}$ mice

Aseptic peritonitis was induced in mice by TG injection and 4 and 16 hours later, numbers of inflammatory monocytes were counted. (A) Typical FACS profile of peritoneal exudate of a WT mouse 4 hours (left panel) and 16 hours (right panel) after TG injection. Neutrophils are Gr-1+F4/80- (R1), inflammatory monocytes are Gr-1+F4/80+ (R2), and non-inflammatory monocytes/macrophages are Gr-1-F4/80+ (R3); Numbers of inflammatory monocytes in WT (n=5), $\beta 2^{-/-}$ (n=4), and $\alpha 4\Delta/\Delta$ (n=4) mice were determined in the peritoneal cavity (B) and in peripheral blood (C). Recruitment indices (D) were calculated to assess the efficiency of migration from the circulation. Asterisk (*) indicates significant difference over control (P<0.05).




Figure 3. Macrophages use $\alpha 4$ and $\beta 2$ integrins interchangeably to maintain sufficient numbers in peritoneum during inflammation

(A) Peritonitis was induced in WT (n=5), $\beta 2^{-/-}$ (n=4), and $\alpha 4^{\Delta/\Delta}$ (n=4) mice and 4, 16, and 96 hours later numbers of macrophages (F4/80+) and (B) numbers of $\alpha 4$ -positive and -negative macrophages in $\alpha 4^{\Delta/\Delta}$ peritoneal lavage were determined by FACS. Asterisk (*) indicates significant difference over control (P<0.05); M ϕ denotes macrophages.




Figure 4. Lymphocyte migration to the inflamed peritoneum is absent in $\alpha 4\Delta/\Delta$ and decreased in $\beta 2^{-/-}$ mice

(A) Aseptic peritonitis was induced in WT (n=5), $\beta 2^{-/-}$ (n=5) and $\alpha 4\Delta/\Delta$ (n=5) mice.

Lymphocyte numbers were determined by FACS, before and 48 hours after TG injection. (B) Efficiency of migration, as indicated by recruitment index, was calculated at 48 hours after TG injection. Asterisk (*) indicates significant difference from control mice at the corresponding hour of peritonitis ($P < 0.05$).




Figure 5. Alpha4 and beta2 integrins are necessary and sufficient for leukocyte migration to the inflamed peritoneum

Total leukocyte numbers in peritoneum of WT (n=5), $\beta 2^{-/-}$ (n=5), $\alpha 4\Delta/\Delta$ (n=5), and $\alpha 4\Delta/\Delta \beta 2^{-/-}$ double deficient (n=4) mice at various times after induction of aseptic peritonitis. Note that, most of the time leukocyte numbers in mice deficient in a single integrin are similar to that of controls, whereas no recruitment of double deficient leukocytes occurs. Asterisk (*) indicates significant difference from control (P<0.05)




Figure 6. Alpha-4 integrins are essential for lymphocyte adhesion and migration in vitro
 (A) Adhesion of control (n=5) and $\alpha 4\Delta\Delta$ (n=3) splenocytes to endothelial cells. Splenocytes were labeled with CFSE and allowed to adhere to either non-stimulated or TNF α -stimulated bEND3 monolayer. Fold increase of adhesion to stimulated over non-stimulated endothelium is displayed. (B) Trans-well and (C) trans-endothelial migration of B-lymphocytes towards SDF-1 α . Note that migration of $\alpha 4\Delta\Delta$ B cells (and T cells, not shown) through VCAM-1-expressing endothelium (TNF α) did not differ from that of non-stimulated endothelium, in contrast to control and $\beta 2^{-/-}$ cells. Asterisk (*) indicates significant difference over controls; ‡ indicates significant difference over non-stimulated endothelium (P<0.05). Three mice per group were used.




Figure 7. Lymphocyte early responses to chemoattractant SDF-1α

(A) Actin polymerization in WT, $\alpha 4\Delta/\Delta$ and $\beta 2^{-/-}$ lymphocytes. Splenocytes were stained for lymphocyte markers (CD3 and B220), stimulated with SDF-1 α for indicated amount of time and phalloidin staining was performed. Fold increase over non-stimulated cells was calculated. (B) SDF-1 α -induced Ca²⁺ mobilization. Splenocytes were loaded with Indo-1 and stained for lymphocyte markers. Ca²⁺ flux was assessed following stimulation with SDF-1 α or ionomycin. Results are displayed as percent of the peak stimulation levels over baseline. Asterisk (*) indicates significant difference over control ($P < 0.05$). Three mice per group were used.