
Gene silencing mechanisms mediated by Aubergine–piRNA complexes in *Drosophila* male gonad

KAZUMICHI M. NISHIDA, KUNIAKI SAITO, TOMOKO MORI, YOSHINORI KAWAMURA, TOMOKO NAGAMI-OKADA, SACHI INAGAKI, HARUHIKO SIOMI, and MIKIKO C. SIOMI

Institute for Genome Research, University of Tokushima, Tokushima 770-8503, Japan

ABSTRACT

Genetic studies have shown that *Aubergine* (*Aub*), one of the Piwi subfamily of *Argonautes* in *Drosophila*, is essential for germ cell formation and maintaining fertility. *aub* mutations lead to the accumulation of retrotransposons in ovaries and testes, and *Stellate* transcripts in testes. *Aub* in ovaries associates with a variety of Piwi-interacting RNAs (piRNAs) derived from repetitive intergenic elements including retrotransposons. Here we found that *Aub* in testes also associates with various kinds of piRNAs. Although in ovaries *Aub*-associated piRNA populations are quite diverse, piRNAs with *Aub* in testes show a strong bias. The most abundant piRNAs were those corresponding to antisense transcripts of *Suppressor of Stellate* [*Su(Ste)*] genes known to be involved in *Stellate* gene silencing. The second most abundant class was made up of those from chromosome X and showed strong complementarity to *vasa* transcripts. Immunopurified *Aub*–piRNA complexes from testes displayed activity in cleaving target RNA containing sequences complementary to *Stellate* and *vasa* transcripts. These results provide the first biochemical insights into gene silencing mechanisms mediated by *Aub* and piRNAs in fly testes.

Keywords: Aubergine; piRNA; *Stellate*; RNA silencing; germlines

INTRODUCTION

RNA silencing is an evolutionarily conserved physiological process involved in regulating gene expression, defending against viruses, and maintaining genome stability (Tomari and Zamore 2005; Zaratiegui et al. 2007). RNA interference (RNAi) (Fire et al. 1998) is the most typical of such processes and causes gene silencing by cleaving (slicing) the target mRNAs in a sequence-dependent manner (Tomari and Zamore 2005). A molecule rendering sequence specificity is a 21–23 nucleotide (nt) long RNA, termed short interfering RNA (siRNA) (Tomari and Zamore 2005).

microRNAs (miRNAs) are a large group of endogenous small RNAs (21–23 nt in length) encoded in the genome of a variety of organisms (Ambros 2004; Du and Zamore 2005; Kloosterman and Plasterk 2006). miRNA also functions in RNA silencing, as does siRNA, a process through which the expression levels of genes involved in various developmental and metabolic processes are modulated. Besides their functional commonality, siRNA and miRNA

show strong similarities even in maturation events. Both are processed from a longer precursor by an RNase III enzyme (Dicer), unwound, and then loaded onto a member of the Argonaute family of proteins (Carmell et al. 2002; Parker and Barford 2006). After maturation and association with Argonautes, they ultimately exert their effects in RNA silencing as guide molecules to specify the RNA targets to be silenced.

Members of the Argonaute family are defined by the presence of the PAZ and PIWI domains (Carmell et al. 2002). In *Drosophila*, five genes—*AGO1*, *AGO2*, *AGO3*, *Piwi*, and *Aubergine* (*Aub*)—exist in the genome as members of the *Argonaute* family. *AGO1* and *AGO2* are expressed ubiquitously through development, whereas the other three are only expressed in germline cells (Williams and Rubin 2002). The latter are grouped as the Piwi subfamily of *Argonautes* because they show strong sequence similarities to each other.

AGO1 and *AGO2* in *Drosophila* function in gene silencing through binding with miRNA and siRNA, respectively (Okamura et al. 2004; Tomari et al. 2004; Miyoshi et al. 2005). Recent studies have shown that the structure of a double-stranded small RNA intermediate strongly influences its partitioning between *AGO1*- and *AGO2*-RISC (RNA-induced silencing complex) (Forstemann et al. 2007;

Reprint requests to: Mikiko C. Siomi, Institute for Genome Research, University of Tokushima, Tokushima 770-8503, Japan; e-mail: siomim@genome.tokushima-u.ac.jp; fax: 81-88-6339451.

Article published online ahead of print. Article and publication date are at <http://www.rnajournal.org/cgi/doi/10.1261/rna.744307>.

Tomari et al. 2007). siRNA-loaded AGO2 functions in RNAi as Slicer and is directly responsible for cleaving a target completely complementary to siRNA (Miyoshi et al. 2005). AGO2 is also involved in RISC formation by slicing the passenger strand within the siRNA duplex (Matranga et al. 2005; Miyoshi et al. 2005; Rand et al. 2005; Kim et al. 2006; Leuschner et al. 2006). Although AGO1 is thought to function in miRNA-mediated translational repression, like AGO2 it also possesses Slicer activity (Miyoshi et al. 2005).

Piwi, one of the Piwi subfamily proteins in *Drosophila*, is an essential factor in germline stem cell (GSC) self-renewal in both males and females (Cox et al. 1998, 2000; Szakmary et al. 2005). *piwi* mutations impact transposon mobility (Sarot et al. 2004; Kalmykova et al. 2005); this most likely results from the deleterious effects of transposon activation. Indeed, recent studies have shown that Piwi in ovaries is associated with repeat-associated siRNAs (rasiRNAs) derived from a variety of repetitive intergenic elements such as retrotransposons (Saito et al. 2006; Vagin et al. 2006; Brennecke et al. 2007). Their longer length distinguishes rasiRNAs (24–30 nt) from siRNAs and miRNAs (Aravin et al. 2003). Recombinant Piwi protein produced in *Escherichia coli* is able to exhibit Slicer activity in vitro (Saito et al. 2006), and Piwi is localized in the nucleus (Cox et al. 2000; Saito et al. 2006; Brennecke et al. 2007). Taken together, it can be postulated that Piwi functions, at least in part, like Slicer in silencing repetitive/transposable genes through its association with rasiRNAs in the nucleus, and thus protects the genome from invasive elements. rasiRNAs in *Drosophila* are also called piRNAs (Brennecke et al. 2007; Zamore 2007); we use the latter term hereinafter.

Genetic studies have revealed that *Aub* is required for pole cell formation (Harris and Macdonald 2001) and for activating RNAi during *Drosophila* oocyte maturation (Kennerdell et al. 2002). *Aub* is also involved in silencing retrotransposons in the germline (Vagin et al. 2004, 2006; Savitsky et al. 2006) and *Stellate* genes in testis through the homologous *Suppressor of Stellate* [*Su(Ste)*] repeats on chromosome Y from which *Su(Ste)* piRNAs are derived (Aravin et al. 2004; Vagin et al. 2006). *Aub* is required for accumulation of *Su(Ste)* piRNAs (Aravin et al. 2004; Vagin et al. 2006). Mutations in *Aub* cause male sterility, which is directly attributable to the failure in silencing the repetitive *Stellate* locus. *Aub* in ovaries is associated with piRNAs originating mainly from retrotransposon antisense transcripts (Vagin et al. 2006; Gunawardane et al. 2007). A large-scale study identifying small RNAs associated with *Aub* in ovary has recently been reported (Brennecke et al. 2007). However, questions for which answers have yet to be elucidated include:

1. What are the differences between piRNAs associated with *Aub* and Piwi in male and female gonads?
2. Is *Aub* in testes physically associated with *Su(Ste)* piRNAs?

3. If so, how is the *Aub*–piRNA complex involved in gene silencing at the molecular level?

Here we closely investigated the profiles of piRNAs associated with Piwi and *Aub* in ovaries and testes. Although cellular localizations of Piwi and *Aub* in ovaries differ from each other, they bind to a similar set of piRNAs. In contrast to this, we found that piRNAs associated with *Aub* in testes were rather unique. Of these piRNAs, the most abundant (~46% of the total) were those corresponding to *Su(Ste)* antisense transcripts. *Aub*–piRNA complex immunopurified from testes was able to cleave target RNAs containing sequences of *Su(Ste)* and *Stellate*. The second largest class of piRNAs associated with *Aub* in testes contained only two kinds of sequences, both of which were mapped to tiny, repetitive regions on chromosome X. Interestingly, these two, termed AT-chX-1 and -2, showed strong complementarity to *vasa (vas)* mRNA, a germline-specific transcript involved in oocyte differentiation and cyst development (Lasko and Ashburner 1988; Styhler et al. 1998). Indeed, the protein levels of VAS in *aub* mutant testes were increased by about twofold compared to those in wild type (wt). The *Aub*–piRNA complex from testes even showed activity in cleaving target RNAs containing part *vas* mRNA. Immunofluorescence showed that *Aub* is localized in the cytoplasm, as is AGO2 (Findley et al. 2003); however, 21-nt siRNA incubated in ovary lysate was only loaded onto AGO2, not onto *Aub*. Taken together, we propose that both in ovaries and testes, *Aub* functions in cytoplasmic RNA silencing only through its association with 24–30-nt piRNAs. Our studies provide the first biochemical insights into gene silencing mechanisms mediated by *Aub* in accord with piRNAs in fly testes.

RESULTS

Expression of *Aub* in germline cells

To biochemically elucidate the functional involvement of *Aub* in the presumptive RNA silencing process in fly gonads, we produced monoclonal antibodies against *Aub*. Western blotting on ovary and S2 cell lysates using the anti-*Aub* antibody revealed that *Aub* is strongly expressed in ovaries, but little in S2 cells (Fig. 1A). AGO1 and AGO2 expressed in S2 cells (Miyoshi et al. 2005) were not cross-reacted by anti-*Aub* (Fig. 1A). Western blotting on Piwi and AGO3 recombinant proteins produced in *E. coli* confirmed that the anti-*Aub* cross-reacts to neither of those (Supplemental Fig. 1A), indicating that the antibody is specific to *Aub*. In early embryos, *Aub* is strongly expressed, but its protein levels gradually diminish throughout development (Supplemental Fig. 1B).

To examine the expression patterns of endogenous *Aub* in ovaries and embryos, immunofluorescent staining was carried out. In the germarium, *Aub* was found in the




FIGURE 1. Aub expression. (A) Aub expression is detected in *Drosophila* ovaries, but not in S2 cells. Western blotting was performed on lysates of fly ovaries and S2 cells using an anti-Aub antibody produced in our laboratory. Anti-tubulin was utilized as a loading control. (B) Fluorescent image of a germarium region stained with anti-Aub, which clearly indicates that Aub is localized in the cytoplasm of germline stem cells (GSC) and germline cyst cells (GCC), but Aub appears not to be expressed in somatic cells in the germarium. (C) Fluorescent image of Aub in an egg chamber at stage 6. Aub is found both in oocytes (Oo) and nurse cells. In the latter, Aub accumulation at the peripheral region of the nuclei and in the cytoplasmic particles is observed (nuage). (D) Fluorescent image of Aub in an egg chamber at stage 10A. Around this stage, Aub accumulation to the posterior pole (PP) in oocytes is clearly found. (E) Fluorescent image of an embryo (stage 14) stained with anti-Aub. Pole cells are indicated as Pc. (F) The DIC image of the embryo shown in E and the immunostaining in E were merged. (G) This immunostaining image clearly shows that Aub is localized in the cytoplasm of pole cells in the interior of embryos. (H–M) Testes were double-stained with anti-Aub and anti-Piwi antibodies (Saito et al. 2006). Fluorescent images of (H,K) Aub and (I,L) Piwi. Merged images are shown in J and M, which clearly indicate that the expressions of Aub and Piwi are mutually exclusive in testes. Aub expression is restricted to GSC, gonialblast cells (Gb), and spermatogonia cells (Sg), while Piwi is only expressed in hub (Hub) and supporting somatic cells (SSC). All images shown in this figure represent a single confocal section.

cytoplasm of germline stem cells (GSC) and germline cyst cells (GCC) (Fig. 1B). There are strong accumulations of Aub at the cytoplasmic side of perinuclear regions, called the nuage, a germline-specific body conserved in species (Eddy 1975). In egg chambers at stages 6 and 10A, Aub was clearly detected in the cytoplasm of both nurse cells and oocytes (Fig. 1C,D). Nuage staining was observed only in nurse cells, not in oocytes at stage 6 (Fig. 1C), which is consistent with the notion that nuage is known to be lost from oocytes (Mahowald and Strassheim 1970). Clear, sharp accumulation of Aub was observed at the posterior of oocytes at stage 10A (Fig. 1D). Aub signals were under the detection limit in somatic cells such as terminal filament cells, cap cells, and follicle cells (Fig. 1B,C), where strong Piwi expression has been observed (Saito et al. 2006). These staining patterns rather resembled those previously reported with GFP-tagged Aub (GFP-Aub) overexpressed in fly ovaries (Harris and Macdonald 2001) and with polyclonal antibodies against Aub (Brennecke et al. 2007). In embryos, endogenous Aub was clearly accumulated in the cytoplasm of pole cells, as expected (Fig. 1E–G).

Testis staining with anti-Aub revealed Aub in GSC, gonialblast, and spermatogonia cells (Fig. 1H,K). However, Aub expression was undetected in somatic cells surrounding gonialblast and spermatogonia cells, and in hub (Fig. 1H,K), where the strong expression of Piwi was observed (Fig. 1I,L; Cox et al. 2000; Saito et al. 2006). Hub is a tiny cluster of post-mitotic somatic cells localized at the apical tip of the testis and functions in the maintenance of GSC identity and influences its behavior (Ohlstein et al. 2004; Snee and Macdonald 2004). Double-staining of testes with anti-Aub and anti-Piwi confirmed that Aub signals were only detected in germ cells, whereas Piwi signals were restricted to somatic cells in testes (Fig. 1J,M). Namely, expression of these two proteins was mutually exclusive in respect to both cellular localization and cell type.

Comparison of piRNAs associated with Aub and Piwi in fly ovaries

We previously identified small RNAs associated with Aub immunoprecipitated from ovary lysate with the anti-Aub antibody (Gunawardane et al. 2007). Silver staining patterns of protein and RNA components in the immunoprecipitate are shown in Supplemental Figure 2. Western blotting analyses on the immunoprecipitate using anti-AGO3 and anti-Piwi antibodies confirmed that the Aub complex does not contain either protein (Supplemental Fig. 2C). To gain insight into the piRNAs associated with Aub and Piwi in ovaries, we reexamined the compositions of those clones, which are listed in Supplemental Tables 1, 2, and 3. Northern blotting analyses demonstrated that piRNAs associated with Aub seemed also to be associated with Piwi (Fig. 2A–C). A size difference was clearly observed between piRNAs with Piwi and Aub (Fig. 2A–C). This agrees with previous data showing that ^{32}P -end-labeled piRNAs associated with Piwi in fly ovaries contained molecules migrating slightly slower than those associated with Aub on denaturing acrylamide gels (Brennecke et al. 2007; Gunawardane et al. 2007). piRNAs associated with Aub and Piwi most likely contain an equal amount of monophosphate at the 5'-ends (data not shown) and are methylated at their 3'-ends (Saito et al. 2007). Thus, piRNAs with Aub might simply be shorter than those associated with Piwi.

We previously reported that miRNAs were not present in Piwi complex immunopurified from ovaries and that siRNAs pre-programmed in ovary lysate were not loaded on Piwi (Saito et al. 2006). We speculated that neither siRNAs nor miRNAs were loaded onto Aub, as was the case for Piwi. To examine this possibility, small RNAs isolated from Aub complex in ovaries were probed with an oligo that specifically recognizes miR-310, one of the miRNAs expressed in fly ovaries. A positive band was detected in just the AGO1 lane (Fig. 2D). Another oligo




FIGURE 2. Detection of Aub association with piRNAs. (A) Northern blotting with probes for 17.6 and GATE retrotransposons shows that piRNAs derived from the transposable elements are involved in Aub and Piwi complexes immunopurified from ovary lysate. It should be noted that the Piwi complex contains piRNAs that are apparently a little longer than those within the Aub complex. (B) The same set of samples used in A was probed with a *roo* retrotransposon sequence. As in A, the fraction of piRNAs with Piwi seems to contain longer molecules compared with those with Aub. (C) An oligo recognizing #4 *roo* piRNA, originally found to exist in fly ovaries (Vagin et al. 2006), was used for Northern blotting analysis. The anti-Piwi lane clearly contains bands migrating slower than those observed in the total and anti-Aub lanes (an asterisk shows one such prominent band). (D) microRNAs are not loaded onto Aub in ovaries. RNA molecules within the AGO1 and Aub complexes immunopurified from ovary lysate were probed with a DNA oligo to detect miR-310. The band corresponding to miR-310 is only found in the anti-AGO1 lane. It should be noted that no miRNAs were obtained through the experiment identifying the small RNAs within Aub complex in ovaries, indicating that it is most likely that no miRNAs are loaded onto Aub in ovaries. (E) *luc* siRNA duplex was first added to ovary lysate. After incubation, AGO2 and Aub were immunopurified from the reaction mixture using specific antibodies. RNA molecules were isolated from the complexes and run on a denaturing acrylamide gel. In the anti-AGO2 lane, the *luc* guide siRNA appears, but not in the anti-Aub and (n.i.) non-immune lanes, indicating that *luc* guide siRNA is specifically loaded onto AGO2 in the ovary lysate.

specifically recognizing miR-186 showed the same result (data not shown). This strongly suggested that Aub does not function together with miRNAs. Indeed, only a trace of miRNAs was obtained (nine of a total 925 *Drosophila* small RNAs; 0.97%) as Aub-associating small RNAs (Supplemental Table 1). We then assessed if the siRNA duplex of 21 nt in length, a triggering molecule of RNAi mediated by AGO2, was loaded onto Aub as well as AGO2 upon addition to ovary lysate. *luc* guide siRNA was efficiently loaded onto AGO2 but not onto Aub (Fig. 2E). These data suggested that the 21 nt siRNA loading machinery composed with Dicer-2 and R2D2 (Liu et al. 2003) was unable to load the small RNA onto both Piwi (Saito et al. 2006) and Aub (this study) proteins.

Su(Ste) piRNAs are the most abundant associated with Aub in testes

Immunoprecipitation of Aub from testes was also performed using the anti-Aub antibody. As in ovaries, small RNAs 24–30 nt long were observed (Supplemental Fig. 3). A small RNA library was constructed, and 458 clones were sequenced (Table 1). The list of piRNAs obtained is shown in Supplemental Table 4. Although piRNAs associated with Aub in testes also contain those derived from LINE and LTR-type transposons (Table 1) and show a strong preference for U at their 5'-ends (~87%) (Supplemental Fig. 4), the most abundant class of piRNAs identified was made up of those derived from *Su(Ste)* antisense transcripts (195 of 425 piRNAs; ~46%) (Fig. 3A; Table 1; Supplemental Table 4). Consistent with these findings, previously genetic studies have shown that both *aub* and *Su(Ste)* mutations cause an accumulation of *Stellate* mRNA in testes (Bozzetti et al. 1995, Aravin et al. 2004; Vagin et al. 2006). *Su(Ste)* piRNAs in a sense orientation were found only once in this screening. This bias was apparently not a consequence of any loading bias to Aub because it has been reported that even in total RNAs, *Su(Ste)* piRNAs in sense are hardly detected in testes (Vagin et al. 2006). piRNAs generated from *Stellate* sense transcript were also obtained in this experiment (two of 425) (Fig. 3A; Table 1; Supplemental Table 4). Notably, *Su(Ste)* piRNAs seemed not evenly produced from the precursor molecules (Fig. 3A). Particularly, one *Su(Ste)* piRNA, termed Su(Ste)-4, was so abundant that it appeared 120 times [including closely related ones such as Su(Ste)-4a shown in Supplemental Table 4] in all the clones sequenced. The association of Su(Ste)-4 with Aub in testes was further confirmed by Northern blotting analysis using a DNA oligo specific for the small RNA (Fig. 3B).

A second large class of piRNAs associated with Aub in testes was made up of those derived from two tiny regions (A, 274 nt; and B, 173 nt) on chromosome X (Fig. 4A). The A and B regions are both located between retrotransposons (Fig. 4A). Interestingly, one of these piRNAs, termed AT-

TABLE 1. (A) Profiles of small RNAs in Aub and Piwi complexes in testes. (B) 195 *Su(Ste)* piRNAs were found in the Aub complex from testes, but only one piRNA originated from the *Su(Ste)* sense transcript

A. Profiles of small RNAs in Aub and Piwi complexes in testes			
Species	Type	Aub in testes	Piwi in testes
D.m.	rRNA	1	14
	tRNA	0	1
	miRNA	0	0
	piRNA	425	246
	mRNA	9	80
	snoRNA	0	6
	snRNA	0	0
	mito.genome.	0	0
	7SL	0	0
	D.m.	Total	435
Others		23	62
	Total	458	409

B. 195 *Su(Ste)* piRNAs were found in the Aub complex from testes, but only one piRNA originated from the *Su(Ste)* sense transcript

piRNAs	Aub in testes	Piwi in testes
LINE	31	42
LTR	68	169
DNA	14	3
Satellite	1	0
Repeat	1	3
Su(Ste)	195	4
AT-chX	104	1
Stellate	2	0
Others	9	24
Total	425	246

Notably, we found no *Su(Ste)* piRNAs in the Aub complex from ovaries. 104 AT-chX piRNAs (including both -1 and -2) (Fig. 4) appeared as piRNAs associated with Aub in testes (24.5%). AT-chX-2 was found in just one of these. The appearance of AT-chX in piRNAs associated with Aub in ovary was just 0.44%.

chX-1, whose sequence appeared 103 times in this screening (~24% of the total piRNAs), showed strong complementarity to *vas* mRNA (Fig. 4B), a germline-specific transcript involved in oogenesis (Lasko and Ashburner 1988; Styhler et al. 1998). Northern blotting using an oligo specific for AT-chX-1 verified its association with Aub in testes (Fig. 4C). AT-chX-2, another piRNA found in this subgroup, appeared only once in this experiment but also showed high complementarity to *vas* mRNA (Fig. 4B). Western blotting revealed that the protein levels of VAS in *aub*⁻¹/*aub*⁻² mutant fly testes were increased by about twofold compared with those of wild-type (wt) and *aub*^{-1/+} (Fig. 4D; Supplemental Fig. 5A). Mislocalization or ectopic expression of VAS was not observed in *aub* mutant testes (data not shown). We also found that VAS expression was not increased in *piwi* mutant testes (Supplemental Fig. 5B). Although it has been shown that the phosphorylation of VAS in ovaries is altered in mutants

A

```

+1
Ste  CATATTTCTGTGAACAGTGAACCTGGCAACA***CGAGCTCGTAAGTAACCTAGGTTTTT
Su(Ste) CATATTTCTGTGAACAGTGAACCTGGCAACA***TTGAGCCGTAAGTAACCTAGGTTTTT

Ste  TCTATAGAAAATTATAGCAAGTCCACAGTAAATCTTCTAGCCAGAACAACAAC-----
Su(Ste) CCTATAGAAAATTATAGCAAGTCCACAGTAAATCTTTCACCCAGAACAACAACAACA

Ste  -----AGCAGCTGGATCGATTGGTTCCCTCGGATCAAGGGCAACGAGTTCCTC
Su(Ste) ACAACAACAACAGCAGCTGGATCAATTGGTTCCCTCGGATCAAGGGCAACGAGTTCCTC

Ste  TGCCCGTGCCCAACCAGTACCTGCAGGATACGTTCAACCAGATGGGCTTGGAGTATTT
Su(Ste) TGCCACGTGCCCCAACACTACTTTCAGGACACGTTCAACCAGATGGGCTTGGAGTACTT

Ste  CAGCGAGACTGGACGTGATCCTGAAGCCGGTATCGACAATTCCTCTGGCTTGTGT
Su(Ste) CAGCCAGCTACTGGACGTGATCCTGAAGCCGGTATCGACAATTCCTCTGGCTTGTGT
Su(Ste)-4 CAGCCAGCTACTGGACGTGATCCTGAAGCCGGTATCGACAATTCCTCTGGCTTGTGT

Ste  ACGCGATGAAAAAAGTGGTACGGCATGATTCACGCCGATACATCAAGTCAGAGCGT
Su(Ste) ACGACGATGAAAAAAGTGGTACGGCATGATTCACGCCGATACATCAAGTCAGAGCGT

Ste  GGCCTGATTCGTATGCACCGAAAATATATGCAGGAGATTTTGAATCGTGTCCAAATAT
Su(Ste) GGCCTGATTCGTATGCACCGAAAATATATGCAGGAGATTTTGAATCGTGTCCAAATAT

Ste  CTCCTGTGATAGGCAGAACACCTGCCAGTGGGCCACAGCGATGTATGGGGCAGTCAA
Su(Ste) CTCCTGTGATAGGCAGAACACCTGCCAGTGGGCCACAGCGATGTATGGGGCAGTCAA

Ste  CGGTCAAGATCTACTGCCACCGGTGTA AAAAGAACTTTCATCCGAAGTCTGATACACAG
Su(Ste) CGGTCAAGATCTACTGCCACCGGTGTA AAAAGAACTTTCATCCGAAGTCTGATACACAG

Ste  CTGGACGGAGCGATGTTCCGGCCACGCTCCCGGACATCTCTTCGCTGTGCGCGAA
Su(Ste) CC-----ATGTTCCGGCCACGCTCCCGGACATCTT--AACCGACGCTGCGCGAA

Ste  CTTGAGATCGCCCTGGACACCCACGGTAAATAATTCCTCGAATATAGTCTGGTTGT
Su(Ste) CTTGAGATCGCCCTGGACACCCACGGTAAATAATTCCTCGAATATAGTCTGGTTGT

Ste  TTTCTAAACAAAGCGCTTGCACCTGCAGTACCTAGGCTTTCGGTTGCACCTGAAAGCCT
Su(Ste) TTTCTAAACAAAGCGCTTAACTTGCATTCCTATGCTTTCGGTTGCACCTGAAAGCCT

Ste  TGATGCACCTCAATTCGCCCAAAATTC
Su(Ste) TGATGCACCTCAATTCGCCCAAAATTC
    
```

B




FIGURE 3. Analyses of *Stellate* and *Su(Ste)* piRNAs associated with Aub in testes. (A) The nucleotide sequence of *Stellate* pre-mRNA (accession number: X15899) (Balakireva et al. 1992) was aligned with that of *Su(Ste)* (accession number: Z11734) (Balakireva et al. 1992). (+1) The transcription start site, (***) the translation start site, and (+++) the stop codon of the *Stellate* transcript. (Boxes in yellow) Nucleotides unmatched between *Su(Ste)* and *Stellate* sequences. (Shaded areas with gray boxes) The intron sequences. *Stellate* and *Su(Ste)* piRNAs identified as being in the Aub complex are underlined in blue and red, respectively. (A thick red underline) *Su(Ste)*-4 piRNA, the most abundantly found through the experiment. (A black dot line) A particular region in *Su(Ste)* from which various piRNAs (29 of 425 piRNAs) were originated (Supplemental Table 4). Note that all the *Su(Ste)* piRNAs obtained in this experiment contained sequences of *Su(Ste)* antisense transcript. *Stellate* piRNAs were from the sense mRNA. (B) The Northern blot confirms that the Aub complex immunoprecipitated from testes contains *Su(Ste)*-4 piRNA (*).

defective in piRNA production and/or function (Klattenhoff et al. 2007), we found no change in the electrophoretic mobility of VAS in *aub* mutant and wt testes (Fig. 4D; Supplemental Fig. 5A; data not shown), suggesting that the

observed twofold changes in VAS abundance do not reflect an effect on its phosphorylation. In contrast, in ovaries, where little AT-chX-1 and -2 was found in the Aub complex (three of 689), we did not observe such an alteration of VAS levels (Fig. 4D) as reported (Wilson et al. 1996). Taken together, it is suggested that *vas* expression in testes might possibly be down-regulated by the Aub complex containing AT-chX-1 and -2 through RNA silencing mechanisms (see below).

Piwi and piRNA association in testes

We also carried out a parallel examination of small RNAs residing within the Piwi complex in testes (Table 1; Supplemental Fig. 3; Supplemental Table 5). We detected only a few *Su(Ste)* piRNAs (four of 246 piRNAs) and no AT-chX-1 and -2 as Piwi-associating small RNAs (Table 1; Supplemental Table 5), which suggested that Piwi hardly contributes to the repression of *Stellate* and *vas* gene expression. It is, however, worth noting that the loss of Piwi in testes increases the production of *Su(Ste)* piRNAs (Vagin et al. 2006). The majority of the small RNAs associated with Piwi in testes were piRNAs derived from transposable elements and other repetitive sequences in the genome, as observed in ovaries (Saito et al. 2006). Interestingly, pieces of protein-coding sense mRNAs were conspicuous in this screening (80 of 347 small RNAs present in the Piwi complex in testes) (Table 1; Supplemental Table 5) as has been reported in the small RNA profiling study done for MIWI, a Piwi homolog expressed in mouse (Girard et al. 2006). However, the significance of the association of Piwi/MIWI in testes with small RNAs originating from sense mRNAs remains unclear.

Immunoprecipitated Aub-piRNA complexes from ovaries and testes show Slicer activity

Previously, we demonstrated that *Drosophila* AGO1 and AGO2 exhibit Slicer activity in vitro (Miyoshi et al. 2005). Moreover, we recently showed that recombinant Piwi, Aub, and AGO3 produced in *E. coli* also exhibit Slicer activity (Saito et al. 2006; Gunawardane et al. 2007). We then assessed if the Aub-piRNA complexes immunopurified from ovaries and testes could, indeed, show the target RNA cleavage activity in vitro. A target RNA containing a sequence perfectly complementary to #4 *roo* piRNA, one of the piRNAs residing in the Aub complex in ovaries (Fig. 2C), was produced and used in the assays. As shown in Figure 5A, the #4 *roo* target RNA was efficiently cleaved with Aub complex immunopurified from ovaries, indicating that endogenous Aub residing in fly gonads also has Slicer activity. Similar experiments were carried out using Aub complex immunopurified from testes. An RNA target containing a sequence completely matching *Su(Ste)*-4 piRNA [Su(Ste)-4 target] or part of the *Stellate* sense mRNA (*Stellate* target), which should be recognized by




FIGURE 5. The Aub-piRNA complexes show Slicer activity. (A) In vitro target RNA cleavage assay using a target containing a sequence complementary to #4 *roo* piRNA (Vagin et al. 2006) (#4 *roo* target). #4 *roo* target was radiolabeled with ^{32}P at the 5'-end. The Aub complex immunopurified from ovaries was incubated with #4 *roo* target, and after the reaction RNAs were prepared and analyzed on a gel. The Aub complex shows its ability to cleave the target, whereas the control (n.i.) shows no such activity. (B) In vitro target RNA cleavage assay was carried out using the Aub complex immunopurified from testes and ovaries. An RNA target, *Su(Ste)-4* target, which has a sequence completely complementary to *Su(Ste)-4* piRNA was produced and used. An expected cleavage product (*) is observed in the Aub complex from testes, but not in that from ovaries, indicating that the Aub complex from ovaries does not have the ability to cleave the particular target RNA. (C) An RNA target that has a partial sequence of *Stellate* mRNA was used in the cleavage assay. The Aub complex used was immunopurified from testes. (D) An RNA target that has a partial sequence of *vas* mRNA was also cleaved with the Aub complex from testes.

Su(Ste) transcripts in *aub* mutants (Aravin et al. 2001; Vagin et al. 2006), this study is the first to show that Aub in testes is physically associated with *Su(Ste)* piRNAs and that the Aub-piRNA complex from testes is able to cleave *Stellate* mRNA. The Aub-piRNA complex was also able to cleave a target RNA, showing complete complementarity to one of the *Su(Ste)* piRNAs, *Su(Ste)-4*, demonstrating that the Aub-piRNA complex can act in RNA silencing

both on the locus piRNA is derived from and loci that show similarity to the piRNA. We found that two piRNAs, AT-chX-1 and -2, originating from tiny repetitive regions located between retrotransposons on chromosome X, showed considerable complementarity to *vas* mRNA. Cleavage on the target RNA sequence in the RNAi pathway has been shown to occur between the +10 and +11 positions of the guide siRNA (the 5'-end of the guide siRNA is assigned as +1) (Elbashir et al. 2001; Schwarz et al. 2004). Base-pairings between siRNA and the target RNA at the +10 and +11 positions is required for the cleavage. If these roles were also applicable to piRNAs, we speculated that AT-chX-1 (and -2) would possibly guide cleavage of *vas* mRNA. Indeed, the *vas* target was cleaved by the Aub complex purified from testes (Fig. 5D), although the cleavage efficiency was relatively low compared with that for *Stellate* mRNA. In agreement with this observation, the levels of *vas* mRNA were moderately increased by loss of *Aub* expression, whereas those of *Stellate* mRNA were severely diminished under the same conditions (Supplemental Fig. 5C). The Aub-piRNA complex may also repress the translation of *vas* mRNAs, just as Argonaute-miRNA complexes repress their mRNAs in various circumstances (Ambros 2004; Du and Zamore 2005; Kloosterman and Plasterk 2006). It has been reported that *spindle-E* (*spnE*) mutations cause piRNAs not to be accumulated in gonads, as in the case of *aub* mutations (Savitsky et al. 2006; Vagin et al. 2006). It should be noted that similar effects on derepression of *vas* gene expression were also observed in testes of *spnE* mutants (data not shown), in which the amounts of AT-chX-1 are markedly reduced (data not shown). As well, we found that *vas* overexpression (EGFP-VAS transgenic flies) (Sano et al. 2002) results in oversized apices in testes, where spermatogonial cells and spermatocytes are located (Supplemental Fig. 5D). Interestingly, *aub* mutants also showed similarly oversized apices in the testes. These results indicate a connection between such phenotypes and increased levels of VAS proteins. At present, we do not know what are the biological consequences of *vas* gene silencing in testis, but our data support a model in which *vas* gene expression is regulated in male germ cells by Aub in concert with the piRNAs.

Notably, >70% of Aub-associated piRNAs in testes were either *Su(Ste)* piRNAs or AT-chXs. In contrast, Piwi-associated piRNAs in testes are mostly derived from these piRNAs. At present, we do not know where this strong bias comes from. Piwi and AGO2 are not required for *Stellate* silencing (Vagin et al. 2006). This study showed that siRNAs and miRNAs were not loaded onto Aub (Fig. 2D,E). These findings support the view that a distinct pathway exists for loading of piRNAs onto Aub in testes. This strong bias could simply be due to a lack of Piwi expression in primary spermatocytes where *Su(Ste)* is expressed (Aravin et al. 2004). Given the model of piRNA biogenesis (Brennecke et al. 2007; Gunawardane et al.

2007), identification of AGO3-associated piRNAs in testes will shed light on the mechanism of the preferential association of *Su(Ste)* piRNAs and AT-chXs with Aub.

In this study, we fortuitously encountered a type of piRNA that originated from non-annotated, intergenic regions and was likely involved in triggering the silencing of protein-coding mRNA targets in collaboration with Aub. More comprehensive biochemical characterization of piRNAs associated with Aub, Piwi, and/or AGO3 in *Drosophila* should shed light on the biological processes involved in the production of the gametes necessary for development of new individuals and thus species perpetuation.

MATERIALS AND METHODS

Drosophila strains

The *yw* strain was used as a wild type. The strains bearing *aub* mutations, *aub^{HN2}* and *aub^{QC42}*, were provided by P. Zamore (University of Massachusetts). Females of *aub^{HN2}* and males of *aub^{QC42}* were crossed to yield *aub* heterozygote flies, *aub^{QC42}/aub^{HN2}*.

Western blot analysis

Two hundred amino acids at the N terminus of Aub fused with GST were used as the antigen to immunize mice. Anti-Aub monoclonal antibodies were produced essentially as described previously (Ishizuka et al. 2002). Western blotting was performed as described previously (Miyoshi et al. 2005). Ten micrograms of proteins from each sample were loaded on gels (Figs. 1A, 4D). Anti-tubulin was from the Developmental Studies Hybridoma Bank (1:1000 dilution). Anti-VAS was provided by A. Nakamura (Kobe-RIKEN CDB) and used at 1:2500 dilution.

Immunofluorescence

Testes and ovaries were dissected manually from adult flies in 1× PBS. Embryos were collected and dechorionated. Immunostaining was performed following standard procedures. Anti-Aub was purified from culture supernatants of hybridoma cells using Thiophilic-Superflow Resion (BD Biosciences) and directly labeled using a HiLyte Fluor 555 Labeling Kit-NH₂ (Dojindo Molecular Technologies). Culture supernatants of anti-Piwi hybridoma cells were used without dilution. Alexa 488-conjugated anti-mouse IgG (Molecular Probe) was used as the secondary antibody to detect the anti-Piwi antibody. All images were collected using a Zeiss LSM510 laser scanning microscope.

Small RNA cloning and sequence analysis

Cloning of small RNAs associated with Aub in ovaries and testes and Piwi in testes was carried out essentially as described (Saito et al. 2006) with minor modifications. Immunoprecipitation was performed essentially as described previously (Miyoshi et al. 2005). Immunoprecipitation buffer contained 30 mM HEPES-KOH (pH 7.3), 150 mM KOAc, 2 mM MgOAc, 5 mM DTT, 0.1% NP-40, 2 μg/mL Pepstatin, 2 μg/mL Leupeptin, and 0.5% Aprotinin. About 300 ovaries or about 1000 testes were used per immunoprecipitation. After immunoprecipitation, total RNAs

were isolated from the immunopurified complexes with phenol-chloroform and precipitated with ethanol. RNAs were dephosphorylated with CIP (NEB) and labeled with [γ -³²P]ATP with T4 polynucleotide kinase (TaKaRa) for visualization. For cloning of small RNAs isolated from Aub complex in ovaries, we used adapters and primers (i.e., RT and PCR primers) as described by Saito et al. (2006). Adapters (MI-5' Linker and MI-3' Linker) and primers used for small RNAs associated with Aub and Piwi in testes were included in the DynaExpress miRNA Cloning Kit (BioDynamics Laboratory). The sequences of each oligo were as follows:

MI-3' Linker: 5'-pCTGTAACCTCGGGTCAATddC-3' (DNA);
MI-5' Linker: 5'-AUCGUCUCGGGAUGAAA-3' (RNA);
3' RT primer: 5'-ATTGACCCGAGTTACAG-3' (DNA);
5' Primer: 5'-ATCGTCTCGGGATGAAA-3' (DNA).

First-strand cDNA synthesis was performed with Stratascript RT (Stratagene) or Reverse Transcriptase (BioDynamics Laboratory). KOD plus (TOYOBO) or Ex Taq polymerase (TaKaRa) was used as the polymerase. PCR products were cloned into the EcoRV site of the pBS SK+ vector (for small RNAs associated with Aub in ovaries) or pTAC-1 (for small RNAs associated with Aub or Piwi in testes) (BioDynamics Laboratory) and sequenced. RNA sequences between 5' and 3' adapters were analyzed by undertaking searches in annotated genomic databases (UCSC: <http://genome.ucsc.edu/cgi-bin/hgBlat>; NCBI: <http://www.ncbi.nlm.nih.gov/blast/>; and FlyBase: <http://flybase.bio.indiana.edu/blast/>) to determine whether the cloned RNAs map to the genome. We only used the best matches up to a maximum of two differences (mismatch, insertion, or deletion) for each small RNA sequence. For repeat annotation, we used results from the RepeatMasker program (UCSC BLAT), NCBI BLAST, or FlyBase BLAST.

Northern blot analysis

Total RNA either from fly ovaries/testes or immunoprecipitated complexes was isolated using ISOGEN (Nippon Gene). Five micrograms of total RNA were used in Figures 2, C and D, 3B, and 4C. Northern blotting was performed as reported previously (Saito et al. 2006). Probes used for *roo* piRNA and miR-310 were as described previously (Saito et al. 2006). Probes used for detecting #4 *roo* piRNA, *Su(Ste)*-4 and AT-chX-1 were as follows:

For #4 *roo* piRNA: 5'-TCGACTCAGTGGCACAATAAAT-3';
for *Su(Ste)*-4: 5'-TCGGGCTTGTCTACGACGATGAGA-3';
for AT-chX-1: 5'-GCCCGAGCCGTCTAACGATGAAACA-3'.

DNA fragments for detecting piRNAs that originated from 17.6 (accession number: XD1472) and GATE (accession number: AJ010298) retrotransposons were obtained by PCR on the *Drosophila melanogaster* genomic DNA and cloned into a pBS SK+ vector. The primers used are as follows:

For 17.6: 5'-ATTCTTGTAACAAATCTTA-3' and 5'-CAGCCTCTCACAATTCAAT-3';
for GATE: 5'-TCGAGCAGGGGCACTTCTCAGTC-3' and 5'-AC TATGGCATGCGCGTCGCCA-3'.

PCR was again performed using primers for the T7 and T3 promoter sequences on each construct, and the PCR products

were used as templates for in vitro transcription using MAXIscript T7 and T3 kits (Ambion) in the presence of ^{32}P -UTP. RNAs transcribed were extracted with phenol:chloroform, precipitated with ethanol, and partially hydrolyzed as described previously (Saito et al. 2006).

siRNA loading assay

Three hundred ovaries were homogenized in a hypotonic buffer (30 mM HEPES-KOH at pH 7.4, 2 mM MgOAc, 5 mM DTT, and 1 mg/mL Pefablock SC) to prepare ovary lysate. *luc* siRNA duplex (Miyoshi et al. 2005) was incubated in the lysate for 1 h at 26°C. Of the *luc* siRNA duplex, only guide siRNA was labeled with [γ - ^{32}P]ATP for visualization. Aub and AGO2 complexes were then immunopurified from the mixture using the specific antibodies. Sodium chloride was added to the lysates to 800 mM just before immunoprecipitation was started. Reaction mixtures were rocked for 1 h at 4°C, and the beads were washed extensively with a washing buffer (30 mM HEPES-KOH at pH 7.4, 800 mM NaCl, 2 mM MgOAc, 5 mM DTT, 1 mg/mL Pefablock SC). RNAs were isolated from the immunoprecipitates, separated on a denaturing polyacrylamide gel, and visualized on BAS-2500 (Fuji).

In vitro target RNA cleavage assay

Target RNA cleavage assays using immunoprecipitates were performed in a reaction buffer containing 30 mM HEPES-KOH (pH 7.4), 100 mM KOAc, 2 mM MgOAc, 5 mM DTT, 10 mM creatine phosphate, 0.5 mM ATP, 30 $\mu\text{g}/\text{mL}$ creatine kinase (Roche), 0.1 U/mL RNasin (Promega), and 0.5 μg of yeast RNA (Ambion). After reaction for 3 h at 26°C, RNAs were isolated from the whole reaction mixtures, run on gels, and visualized on the BAS 2500. To make a target harboring a sequence completely complementary to miR-310, a short DNA fragment produced by annealing a set of oligo DNAs (5'-AGCTTAAAGGCCGGGAA GTGTGCAATAA-3' and 5'-AGCTTTATTGCACACTTCCCGGC CTTTA-3') was inserted in a pBS SKII+ vector digested with HindIII (pBS310). To make a target harboring a sequence completely complementary to #4 *roo* piRNA (#4 *roo* piRNA target), a short DNA fragment produced by annealing a set of oligo DNAs (5'-GATCCATAGTCTGACTCAGTGCCACAATAAA TAAAGGC-3' and 5'-GGCCGCTTTATTTATTGTGCCACTGA GTCGACTATG-3') was inserted in a pBS310 vector digested with BamHI and NotI. To make a target harboring a sequence completely complementary to Su(Ste)-4 piRNA [Su(Ste)-4 target], a short DNA fragment produced by annealing a set of oligo DNAs (5'-GATCTCGGGCTTGTCTACGACGATGAGA-3' and 5'-GGCCTCTCATCGTCGTAGAACAAAGCCCGA-3') was inserted in a pBS310 vector digested with BamHI and NotI. To make a target harboring a partial sequence of *Stellate* sense mRNA (*Stellate* target), a short DNA fragment produced by annealing a set of oligo DNAs (5'-AATTCTCTGGCTTGTGTACGGCGATG AAAG-3' and 5'-GATCCTTTCATCGCCGTACAACAAGCCAGA G-3') was inserted into a pBS SK+ vector digested with BamHI and EcoRI. To make a target harboring a partial sequence of *vas* mRNA (*vas* target; as a target for AT-chX-1), a short DNA fragment produced by annealing a set of oligo DNAs (5'-TTC CTCCCAGCCGTCCAACGATGCAATTGAGA-3' and 5'-TCT CAATTGCATCGTTGGACGGCTCGGGAGGAA-3') was inserted in a pBS SKII+ vector that was digested with BamHI and NotI and

blunted. PCR was again performed using primers for the T7 and T3 promoter sequences, and the PCR products were then used as templates for in vitro transcription using a MEGAscript T7 kit (Ambion). The resultant RNAs were radiolabeled at the 5'-G cap by guanylyltransferase (Ambion) as previously described (Okamura et al. 2004) or with [γ - ^{32}P]ATP using T4 polynucleotide kinase and gel-purified. *vasa* target (Fig. 5D) was labeled with pCp at the 3'-end and gel-purified before use.

Accession numbers

The accession numbers are: Aub-associated piRNAs in ovary, AB296387-AB297014; Aub-associated piRNAs in testis, AB297015-AB297190; and Piwi-associated piRNAs in testis, AB297191-AB297373.

SUPPLEMENTAL DATA

Supplemental materials are available upon request by sending an e-mail message containing the keyword "Aub-supplement" to siomim@genome.tokushima-u.ac.jp.

ACKNOWLEDGMENTS

We thank P. Zamore for providing the *aub* mutant stocks *aub*^{HN2} and *aub*^{QC42} and A. Nakamura for the anti-VAS antibody and technical advice. We also thank H. Kose, T. Suzuki, and V. Vagin for technical advice. We thank members of the Siomi laboratory for discussion and comments on the manuscript. K.S. is a postdoctoral fellow of the 21st Century COE Program from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT). This study was supported in part by grants to M.C.S. and H.S. from MEXT and the New Energy and Industrial Technology Development Organization (NEDO).

Received July 19, 2007; accepted August 15, 2007.

REFERENCES

- Ambros, V. 2004. The functions of animal microRNAs. *Nature* **431**: 350–355.
- Aravin, A.A., Naumova, N.M., Tulin, A.V., Vagin, V.V., Rozovsky, Y.M., and Gvozdev, V.A. 2001. Double-stranded RNA-mediated silencing of genomic tandem repeats and transposable elements in the *D. melanogaster* germline. *Curr. Biol.* **11**: 1017–1027.
- Aravin, A.A., Lagos-Quintana, M., Yalcin, A., Zavolan, M., Marks, D., Snyder, B., Gaasterland, T., Meyer, J., and Tuschl, T. 2003. The small RNA profile during *Drosophila melanogaster* development. *Dev. Cell* **5**: 337–350.
- Aravin, A.A., Klenov, M.S., Vagin, V.V., Bantignies, F., Cavalli, G., and Gvozdev, V.A. 2004. Dissection of a natural RNA silencing process in the *Drosophila melanogaster* germ line. *Mol. Cell. Biol.* **24**: 6742–6750.
- Balakireva, M.D., Shevelyov, Y.Y., Nurminsky, D.I., Livak, K.J., and Gvozdev, V.A. 1992. Structural organization and diversification of Y-linked sequences comprising *Su(Ste)* genes in *Drosophila melanogaster*. *Nucleic Acids Res.* **20**: 3731–3736. doi: 10.1093/nar/20.14.3731.
- Bozzetti, M.P., Massari, S., Finelli, P., Meggio, F., Pinna, L.A., Boldyreff, B., Issinger, O.G., Palumbo, G., Ciriaco, C.,

- Bonaccorsi, S., et al. 1995. The Ste locus, a component of the parasitic cry-Ste system of *Drosophila melanogaster*, encodes a protein that forms crystals in primary spermatocytes and mimics properties of the β subunit of casein kinase 2. *Proc. Natl. Acad. Sci.* **92**: 6067–6071.
- Brennecke, J., Aravin, A.A., Stark, A., Dus, M., Kellis, M., Sachidanandam, R., and Hannon, G.J. 2007. Discrete small RNA-generating loci as master regulators of transposon activity in *Drosophila*. *Cell* **128**: 1–15.
- Carmell, M.A., Xuan, Z., Zhang, M.Q., and Hannon, G.J. 2002. The Argonaute family: Tentacles that reach into RNAi, developmental control, stem cell maintenance, and tumorigenesis. *Genes & Dev.* **16**: 2733–2742.
- Cox, D.N., Chao, A., Baker, J., Chang, L., Qiao, D., and Lin, H. 1998. A novel class of evolutionarily conserved genes defined by piwi are essential for stem cell self-renewal. *Genes & Dev.* **12**: 3715–3727.
- Cox, D.N., Chao, A., and Lin, H. 2000. piwi encodes a nucleoplasmic factor whose activity modulates the number and division rate of germline stem cells. *Development* **127**: 503–514.
- Du, T. and Zamore, P.D. 2005. microPrimer: The biogenesis and function of microRNA. *Development* **132**: 4645–4652.
- Eddy, E.M. 1975. Germ plasm and the differentiation of the germ cell line. *Int. Rev. Cytol.* **43**: 229–280.
- Elbashir, S.M., Lendeckel, W., and Tuschl, T. 2001. RNA interference is mediated by 21- and 22-nucleotide RNAs. *Genes & Dev.* **20**: 1993–1997.
- Findley, S.D., Tamanaha, M., Clegg, N.J., and Ruohola-Baker, H. 2003. Maelstrom, a *Drosophila* spindle-class gene, encodes a protein that colocalizes with Vasa and RDE1/AGO1 homolog, Aubergine, in nuage. *Development* **130**: 859–871.
- Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E., and Mello, C.C. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811.
- Forstemann, K., Horwich, M.D., Wee, L., Tomari, Y., and Zamore, P.D. 2007. *Drosophila* microRNAs are sorted into functionally distinct Argonaute complexes after production by Dicer-1. *Cell* **130**: 287–297.
- Girard, A., Sachidanandam, R., Hannon, G.J., and Carmell, M.A. 2006. A germline-specific class of small RNAs binds mammalian Piwi proteins. *Nature* **442**: 199–202.
- Gunawardane, L.S., Saito, K., Nishida, K.M., Miyoshi, K., Kawamura, Y., Nagami, T., Siomi, H., and Siomi, M.C. 2007. A Slicer-mediated mechanism for repeat-associated siRNA 5' end formation in *Drosophila*. *Science* **315**: 1587–1590.
- Harris, A.N. and Macdonald, P.M. 2001. Aubergine encodes a *Drosophila* polar granule component required for pole cell formation and related to eIF2C. *Development* **128**: 2823–2832.
- Ishizuka, A., Siomi, M.C., and Siomi, H. 2002. A *Drosophila* fragile X protein interacts with components of RNAi and ribosomal proteins. *Genes & Dev.* **16**: 2497–2508.
- Kalmykova, A.I., Klenov, M.S., and Gvozdev, V.A. 2005. Argonaute protein PIWI controls mobilization of retrotransposons in the *Drosophila* male germline. *Nucleic Acids Res.* **33**: 2052–2059. doi: 10.1093/nar/gki323.
- Kennerdell, J.R., Yamaguchi, S., and Carthew, R.W. 2002. RNAi is activated during *Drosophila* oocyte maturation in a manner dependent on aubergine and spindle-E. *Genes & Dev.* **16**: 1884–1889.
- Kim, K., Lee, Y.S., and Carthew, R.W. 2006. Conversion of pre-RISC to holo-RISC by Ago2 during assembly of RNAi complexes. *RNA* **13**: 22–29.
- Klattenhoff, C., Bratu, D.P., McGinnis-Schultz, N., Koppetsch, B.S., Cook, H.A., and Theurkauf, W.E. 2007. *Drosophila* rasiRNA pathway mutations disrupt embryonic axis specification through activation of an ATR/Chk2 DNA damage response. *Dev. Cell* **12**: 45–55.
- Kloosterman, W.P. and Plasterk, R.H.A. 2006. The diverse functions of microRNAs in animal development and disease. *Dev. Cell* **11**: 441–450.
- Lasko, P.F. and Ashburner, M. 1988. The product of the *Drosophila* gene vasa is very similar to eukaryotic initiation factor-4. *Nature* **335**: 611–617.
- Leuschner, P.J., Ameres, S.L., Kueng, S., and Martinez, J. 2006. Cleavage of the siRNA passenger strand during RISC assembly in human cells. *EMBO Rep.* **7**: 314–320.
- Lin, H. and Spradling, A.C. 1997. A novel group of pumilio mutations affects the asymmetric division of germline stem cells in the *Drosophila* ovary. *Development* **124**: 2463–2476.
- Liu, Q., Rand, T.A., Kalidas, S., Du, F., Kim, H.E., Smith, D.P., and Wang, X. 2003. R2D2, a bridge between the initiation and effector steps of the *Drosophila* RNAi pathway. *Science* **301**: 1921–1925.
- Mahowald, A.P. and Strassheim, J.M. 1970. Intercellular migration of centrioles in the germlarium of *Drosophila melanogaster*. An electron microscopic study. *J. Cell Biol.* **45**: 306–320.
- Matranga, C., Tomari, Y., Shin, C., Bartel, D.P., and Zamore, P.D. 2005. Passenger-strand cleavage facilitates assembly of siRNA into Ago2-containing RNAi enzyme complexes. *Cell* **123**: 607–620.
- Miyoshi, K., Tsukumo, H., Nagami, T., Siomi, H., and Siomi, M.C. 2005. Slicer function of *Drosophila* Argonautes and its involvement in RISC formation. *Genes & Dev.* **19**: 2837–2848.
- Ohlstein, B., Kai, T., Decotto, E., and Spradling, A. 2004. The stem cell niche: Theme and variations. *Curr. Opin. Cell Biol.* **16**: 693–699.
- Okamura, K., Ishizuka, A., Siomi, H., and Siomi, M.C. 2004. Distinct roles for Argonaute proteins in small RNA-directed RNA cleavage pathways. *Genes & Dev.* **18**: 1655–1666.
- Parker, J.S. and Barford, D. 2006. Argonaute: A scaffold for the function of short regulatory RNAs. *Trends Biochem. Sci.* **31**: 622–630.
- Rand, T.A., Peterson, S., Du, F., and Wang, X. 2005. Argonaute2 cleaves the anti-guide strand of siRNA during RISC activation. *Cell* **123**: 621–629.
- Saito, K., Nishida, K.M., Mori, T., Kawamura, Y., Miyoshi, K., Nagami, T., Siomi, H., and Siomi, M.C. 2006. Specific association of Piwi with rasiRNAs derived from retrotransposon and heterochromatic regions in the *Drosophila* genome. *Genes & Dev.* **20**: 2214–2222.
- Saito, K., Sakaguchi, Y., Suzuki, T., Suzuki, T., Siomi, H., and Siomi, M.C. 2007. Pimet, the *Drosophila* homolog of HEN1, mediates 2'-O-methylation of Piwi-interacting RNAs at their 3' ends. *Genes & Dev.* **21**: 1603–1608.
- Sano, H., Nakamura, A., and Kobayashi, S. 2002. Identification of a transcriptional regulatory region for germline-specific expression of vasa gene in *Drosophila melanogaster*. *Mech. Dev.* **112**: 129–139.
- Sarot, E., Payen-Groschene, G., Bucheton, A., and Pelisson, A. 2004. Evidence for a piwi-dependent RNA silencing of the gypsy endogenous retrovirus by the *Drosophila melanogaster flamenco* gene. *Genetics* **166**: 1313–1321.
- Savitsky, M., Kwon, D., Georgiev, P., Kalmykova, A., and Gvozdev, V. 2006. Telomere elongation is under the control of the RNAi-based mechanism in the *Drosophila* germline. *Genes & Dev.* **20**: 345–354.
- Schmidt, A., Palumbo, G., Bozzetti, M.P., Tritto, P., Pimpinelli, S., and Schafer, U. 1999. Genetic and molecular characterization of sting, a gene involved in crystal formation and meiotic drive in the male germ line of *Drosophila melanogaster*. *Genetics* **151**: 749–760.
- Schwarz, D.S., Tomari, Y., and Zamore, P.D. 2004. The RNA-induced silencing complex is a Mg²⁺-dependent endonuclease. *Curr. Biol.* **14**: 787–791.
- Snee, M.J. and Macdonald, P.M. 2004. Live imaging of nuage and polar granules: Evidence against a precursor-product relationship and a novel role for Oskar in stabilization of polar granule components. *J. Cell Sci.* **117**: 2109–2120.
- Styhler, S., Nakamura, A., Swan, A., Suter, B., and Lasko, P. 1998. vasa is required for GURKEN accumulation in the oocyte, and is involved in oocyte differentiation and germline cyst development. *Development* **125**: 1569–1578.
- Szakmary, A., Cox, D.N., Wang, Z., and Lin, H. 2005. Regulatory relationship among piwi, pumilio, and bag-of-marbles in *Drosophila*

- germline stem cell self-renewal and differentiation. *Curr. Biol.* **15**: 171–178.
- Tomari, Y. and Zamore, P.D. 2005. Machines for RNAi. *Genes & Dev.* **19**: 517–529.
- Tomari, Y., Matranga, C., Haley, B., Martinez, N., and Zamore, P.D. 2004. A protein sensor for siRNA asymmetry. *Science* **306**: 1377–1380.
- Tomari, Y., Du, T., and Zamore, P.D. 2007. Sorting of *Drosophila* small silencing RNAs. *Cell* **130**: 299–308.
- Vagin, V.V., Klenov, M.S., Kalmykova, A.I., Stolyarenko, A.D., Kotelnikov, R.N., and Gvozdev, V.A. 2004. The RNA interference proteins and Vasa locus are involved in the silencing of retrotransposons in the female germline of *Drosophila melanogaster*. *RNA Biol.* **1**: 54–58.
- Vagin, V.V., Sigova, A., Li, C., Seitz, H., Gvozdev, V., and Zamore, P.D. 2006. A distinct small RNA pathway silences selfish genetic elements in the germline. *Science* **313**: 320–324.
- Williams, R.W. and Rubin, G.M. 2002. ARGONAUTE1 is required for efficient RNA interference in *Drosophila* embryos. *Proc. Natl. Acad. Sci.* **99**: 6889–6894.
- Wilson, J.E., Connell, J.E., and Macdonald, P.M. 1996. *aubergine* enhances *oskar* translation in the *Drosophila* ovary. *Development* **122**: 1631–1639.
- Zamore, P.D. 2007. RNA silencing: Genomic defence with a slice of pi. *Nature* **19**: 864–865.
- Zaratiegui, M., Irvine, D.V., and Martienssen, R.A. 2007. Noncoding RNAs and gene silencing. *Cell* **23**: 763–776.