

Rhizobacterial Volatiles Affect the Growth of Fungi and *Arabidopsis thaliana*[∇]

Anja Vespermann, Marco Kai, and Birgit Piechulla*

University of Rostock, Department of Biological Sciences, Albert-Einstein-Str. 3, 18059 Rostock, Germany

Received 15 May 2007/Accepted 24 June 2007

Volatiles of *Stenotrophomonas*, *Serratia*, and *Bacillus* species inhibited mycelial growth of many fungi and *Arabidopsis thaliana* (40 to 98%), and volatiles of *Pseudomonas* species and *Burkholderia cepacia* retarded the growth to lesser extents. *Aspergillus niger* and *Fusarium* species were resistant, and *B. cepacia* and *Staphylococcus epidermidis* promoted the growth of *Rhizoctonia solani* and *A. thaliana*. Bacterial volatiles provide a new source of compounds with antibiotic and growth-promoting features.

The rhizosphere of plants is the habitat of a community comprising many different organisms. Soil bacteria often possess traits that enable them to act as antagonists by suppressing soilborne plant diseases, for example, by excreting antifungal metabolites that directly or indirectly support plant growth (7, 8, 9, 19). Many of these specialized compounds, such as antibiotics, are either liquid or solid at room temperature, and little is known about volatiles (with molecular masses less than 300 Da, low polarity, and a high vapor pressure) that can act as antibiotics and cause growth inhibition or have more deleterious effects on organisms. The microbial world synthesizes and emits many volatile compounds (1, 3, 4, 6, 16, 17). We previously showed that rhizobacterial isolates of *Serratia plymuthica*, *Serratia odorifera*, *Stenotrophomonas maltophilia*, *Stenotrophomonas rhizophila*, *Pseudomonas fluorescens*, and *Pseudomonas trivialis* emit complex blends of volatiles that inhibit the growth of *Rhizoctonia solani* (12). These rhizobacterial isolates and one human pathogen isolate were used to investigate their biological effects on 14 fungi, as well as on the model plant *Arabidopsis thaliana*.

Fungi were cocultivated with rhizobacteria in divided petri dishes; the fungal medium contained 20 g liter⁻¹ peptone, 10 g liter⁻¹ glucose, and 20 g liter⁻¹ agar-agar (pH 6.8) at 20°C, and the bacterial medium contained 5 g liter⁻¹ peptone from casein, 2.5 g liter⁻¹ peptone from meat, 2.5 g liter⁻¹ peptone from gelatin, 1.5 g liter⁻¹ yeast extract, 1.5 g liter⁻¹ NaCl, and 15 g liter⁻¹ agar-agar (pH 7.2) at 20°C. Two representative examples are shown in Fig. 1. Growth of the mycelium was recorded from the first to the seventh or ninth day after inoculation using a digital camera (C-3030 Zoom Camedia; Olympus) and the Image Gauge software of the image analyzer LAS-1000 (Fujifilm, Tokyo, Japan). Fungal mycelial growth and plant growth were visible in the absence of bacteria (Fig. 1, left panels), but mycelial development of *Paecilomyces carneus* and *R. solani* was strongly inhibited by adjacent growth of *S. maltophilia* R3089 and *A. thaliana* development was strongly inhibited by adjacent growth of *S. odorifera* 4Rx13 (Fig. 1, middle panels). To prove that bacterial volatiles were the prin-

cipal component impeding mycelial and plant growth, charcoal was added to the test system to trap the volatiles. These cocultivations showed that growth inhibition was partly abrogated in the charcoal test system (Fig. 1, right panels). In the presence of charcoal the fungal mycelial size reached 50% or more of the mycelial size of the control, and 80% of *A. thaliana* plants exhibited normal growth.

The strength of fungal growth inhibition depends on the rhizobacterial isolate. Growth retardation was often visible after the second day of incubation, and in most cases this inhibition accelerated after the fourth day. To allow comparisons, growth alterations (expressed in percentages) were calculated for all fungus-




FIG. 1. Cocultivation of rhizobacteria with fungi or *A. thaliana* in the presence or absence of charcoal. Nutrient broth (left panels) or overnight rhizobacterial cultures (20 or 50 μ l) (middle and right panels) were plated in one compartment of a tripartite petri dish. After 2 days of incubation, a fungal mycelium plug (*P. carneus* [A] or *R. solani* [B]) or 10 *A. thaliana* seedlings with or without charcoal (right and middle panels, respectively) were placed in the other compartments, and incubation was continued for 4 days for the fungi or for 14 days for *A. thaliana*.

* Corresponding author. Mailing address: University of Rostock, Institute of Biological Sciences, Albert-Einstein-Str. 3, 18059 Rostock, Germany. Phone and fax: 49 381 498 6130. E-mail: birgit.piechulla@uni-rostock.de.

[∇] Published ahead of print on 29 June 2007.




FIG. 2. Determination of growth inhibition. Mycelial growth inhibition or promotion was calculated after 4 days of cocultivation. The fresh weight of *A. thaliana* was determined after 14 days of cocultivation. Experiments were repeated three times, and each experiment comprised three to seven replicates with each bacterial isolate. The error bars indicate standard errors. The bacterial isolates used were *B. subtilis* B2g (*B.s.*), *B. cepacia* 1S18 (*B.c.*), *P. fluorescens* L13-6-12 (*P.f.*), *P. trivialis* 3Re2-7 (*P.t.*), *S. odorifera* 4Rx13 (*S.o.*), *S. plymuthica* 3Re4-18 (*S.p.*), *S. plymuthica* HRO C48 (*S.p.H.*), *S. epidermidis* 2P13-18 (*S.e.*), *S. maltophilia* R3089 (*S.m.*), and *S. rhizophila* P69 (*S.r.*). The soilborne fungi used were *F. culmorum* PR 19-12-11, *F. solani*, *M. bolleyi* PR 5-11-6, *P. carneus* PR 16-10-1, *Penicillium* sp. strain 2-1-20, *Penicillium waksmanii* PR 17-11-8, *P. betae*, *Phoma eupyrena* PC 17-12-10, *R. solani* AG3, *S. sclerotiorum*, *Trichoderma strictipile* PC26-12-6, and *V. dahliae* V25. In addition, *A. niger* and *N. crassa* wild-type 1202A were used.

bacterium combinations on the fourth day (Fig. 2). *Aspergillus niger* and *Fusarium culmorum* were inhibited by approximately 20% or less or mycelial growth was promoted (*Fusarium solani* and *A. niger*, 3 to 10%) by all or many bacteria. All tested fungi showed individual inhibition patterns. *Microdochium bolleyi*, *P. carneus*, *Phoma betae*, and *Sclerotinia sclerotiorum* were strongly inhibited, by 40% or more. The inhibition of *Verticillium dahliae*, *R. solani*, *Penicillium* sp., and *Neurospora crassa* was moderate in comparison to the inhibition of the other fungi tested. At present, the biologically active volatiles causing the inhibition are not known because many volatiles have not been detected or identified (12). Antifungal effects of organic volatiles were previously shown to inhibit germination or mycelial growth of *S. sclerotiorum* (6, 11), and unidentified compounds from *Bacillus subtilis* cause structural deformation of pathogenic fungi (2). Inorganic volatiles also control growth or inhibit hyphal formation and extension (10, 13). Bacterial volatiles can also promote fungal growth (11, 18).

Bacteria and plants were incubated together for 14 days before the fresh weight of the plants was determined. The following conditions were used for plant growth: surface-sterilized vernalized seeds, half-strength Murashige-Skoog medium, a cycle consisting of 16 h of light and 8 h of darkness, a light intensity of 84 $\mu\text{mol}/\text{m}^2/\text{s}$, and 20°C. Drastic growth inhibition (>80%) leading to small, pale yellow, and dead *A. thaliana* plants was obtained with *P. fluorescens*, *P. trivialis*, both *S. plymuthica* isolates, *S. odorifera*, *S. rhizophila*, and *S. maltophilia* (Fig. 1 and 2). In comparison to controls (nutrient broth), the fresh weight of *A. thaliana* increased by 20 to 30% under the influence of *B. cepacia* and *S. epidermidis*, and cocultivation with *B. subtilis* had no significant effect on plant development. The latter result seems to contradict observations made by Ryu et al. (14, 15) and Farag et al. (5), who showed that *B. subtilis* GB03 and *Bacillus amyloliquefaciens* IN937a promote growth and trigger induced systemic resistance in *A. thaliana*. The use of different bacillus species or isolates and variations in the growth conditions and test systems may account for the different results. Growth promotion of *A. thaliana* and *R. solani*, however, was observed when these organisms were cocultivated with *B. cepacia* or *S. epidermidis*.

Concluding remarks. The experiments described here clearly demonstrate that bacterial volatiles add another component to the action profile of relevant growth-promoting or -inhibiting strategies of rhizobacteria. Volatiles can be an advantageous tool for rhizobacteria because they are small molecules that can easily diffuse through the porous structure of the soil and over great distances in the atmosphere. Many situations can be anticipated where this communication among plants, fungi, and rhizobacteria may be advantageous to at least one of the parties involved and has consequences for the organisms living in a community. The growth-inhibiting effects of rhizobacterial volatiles on phytopathogenic fungi, which cause many economically relevant crop diseases, have potential for agronomical applications, for example, by using the volatiles or rhizobacteria as biological control agents. Most of

the investigated fungi also cause diseases in humans, such as opportunistic infections, mycoses, and allergic reactions (e.g., *Aspergillus*, *Fusarium*, *Paecilomyces*, or *Penicillium* species). Therefore, medicinal applications for rhizobacterial volatiles can also be anticipated.

We thank G. Berg (University of Graz, Austria) and Hella Goshcke (University of Rostock, Germany) for the bacterial and fungal isolates, T. Rönneberg (University of Munich, Germany) for *N. crassa*, and A. Podbielski (University of Rostock, Germany) for *A. niger*.

This research was financially supported by a grant from the University of Rostock to B.P.

REFERENCES

- Alstrom, S. 2001. Characteristics of bacteria from oil seed rape in relation to their biocontrol activity against *Verticillium dahliae*. *J. Phytopathol.* **149**:57–64.
- Chaurasia, B., A. Pandey, L. M. S. Palni, P. Trivedi, B. Kumar, and N. Colvin. 2005. Diffusible and volatile compounds produced by an antagonistic *Bacillus subtilis* strain cause structural deformations in pathogenic fungi in vitro. *Microbiol. Res.* **160**:75–81.
- Dickschat, J. S., S. C. Wenzel, H. B. Bode, R. Müller, and S. Schulz. 2004. Biosynthesis of volatiles by the Myxobacterium *Myxococcus xanthus*. *Chem. Biol. Chem.* **5**:778–787.
- Dickschat, J. S., R. Martens, T. Brinkhoff, M. Simon, and S. Schulz. 2005. Volatiles released by a *Streptomyces* species isolated from the North Sea. *Chem. Biodivers.* **2**:837–865.
- Farag, M. A., C. M. Ryu, L. W. Summer, and P. W. Pare. 2006. GC-MS SPME profiling of rhizobacterial volatiles reveals prospective inducers of growth promotion and induced systemic resistance in plants. *Phytochemistry* **67**:2262–2268.
- Fernando, W. G., R. Ramarathnam, A. S. Krishnamoorthy, and S. C. Savchuk. 2005. Identification and use of potential bacterial organic antifungal volatiles in biocontrol. *Soil Biol. Biochem.* **37**:955–964.
- Gupta, A. M., K. V. B. Gopal, and R. Tilak. 2000. Mechanism of plant growth promotion by rhizobacteria. *Indian J. Exp. Biol.* **38**:856–862.
- Haas, D., and G. Defago. 2005. Biological control of soil-borne pathogens by fluorescent pseudomonads. *Nat. Rev. Microbiol.* **3**:307–319.
- Handelsman, J., and E. V. Stabb. 1996. Biocontrol of soilborne plant pathogens. *Plant Cell* **8**:1855–1869.
- Howell, C. R., R. C. Beier, and R. D. Stipanovic. 1988. Reproduction of ammonia by *Enterobacter cloacae* and its possible role in the biological control of *Pythium* preemergence damping-off by the bacterium. *Phytopathology* **78**:1075–1078.
- Huang, H., J. W. Huang, G. Saidon, and R. Erickson. 1997. Effect of allyl alcohol and fermented agricultural wastes on carpogenic germination of sclerotia of *Sclerotinia sclerotiorum* and colonization of *Trichoderma* spp. *Can. J. Plant Pathol.* **9**:43–46.
- Kai, M., U. Efmert, G. Berg, and B. Piechulla. 2007. Volatiles of bacterial antagonists inhibit mycelial growth of the plant pathogen *Rhizoctonia solani*. *Arch. Microbiol.* **187**:351–360.
- Robinson, P. M., N. D. McKee, L. A. A. Thompson, D. B. Harper, and J. T. G. Hamilton. 1989. Autoinhibition of germination and growth in *Geotrichum candidum*. *Mycol. Res.* **93**:214–222.
- Ryu, C. M., M. A. Farag, C. H. Hu, M. S. Reddy, H. X. Wie, P. W. Pare, and J. W. Kloepper. 2003. Bacterial volatiles promote growth in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **100**:4927–4932.
- Ryu, C. M., M. A. Farag, C. H. Hu, M. S. Reddy, J. W. Kloepper, and P. W. Pare. 2004. Bacterial volatiles induce systemic resistance in *Arabidopsis*. *Plant Physiol.* **134**:1017–1026.
- Schöller, C. E. G., H. Gürtler, R. Petersen, S. Molin, and K. Wilkins. 2002. Volatile metabolites from Actinomycetes. *J. Agric. Food Chem.* **50**:2615–2621.
- Stotzky, G., and S. Schenk. 1976. Volatile organic compounds and microorganisms. *Crit. Rev. Microbiol.* **4**:333–382.
- Wheatley, R. E. 2002. The consequences of volatile organic compound mediated bacterial and fungal interactions. *Antonie Leeuwenhoek* **81**:357–364.
- Whipps, J. M. 2001. Microbial interaction and biocontrol in the rhizosphere. *J. Exp. Bot.* **52**:487–511.