



Published in final edited form as:

Toxicol Appl Pharmacol. 2007 November 1; 224(3): 274–283.

Resveratrol: A Review of Pre-clinical Studies for Human Cancer Prevention

Mohammad Athar¹, Jung Ho Back¹, Xiuwei Tang¹, Kwang Ho Kim², Levy Kopelovich³, David R. Bickers¹, and Arianna L. Kim^{1,*}

¹Department of Dermatology, College of Physicians and Surgeons, Columbia University, 630 West 168th Street VC15-204, New York, NY

²Department of Dermatology, Hallym University College of Medicine, Korea

³Division of Cancer Prevention, National Cancer Institute, Bethesda, MD

Abstract

The search for novel and effective cancer chemopreventive agents has led to the identification of various naturally occurring compounds one of which is resveratrol (*trans*-3, 4', 5-trihydroxystilbene), a phytoalexin derived from the skin of grapes and other fruits. Resveratrol is known to have potent anti-inflammatory and anti-oxidant effects and to inhibit platelet aggregation and the growth of a variety of cancer cells. Its potential chemopreventive and chemotherapeutic activities have been demonstrated in all three stages of carcinogenesis (initiation, promotion, and progression), in both chemically and UVB-induced skin carcinogenesis in mice, as well as in various murine models of human cancers. Evidence from numerous *in vitro* and *in vivo* studies has confirmed its ability to modulate various targets and signaling pathways. This review discusses the current preclinical and mechanistic data available and assesses resveratrol's anti-cancer effects to support its potential as an anticancer agent in human populations.

Keywords

Resveratrol; human cancer; skin cancer; breast cancer; lung cancer; gastric; colorectal cancer; hepatoma; neuroblastoma; pancreatic cancer; leukemia

Introduction

There is growing interest in using naturally occurring compounds as potential cancer chemopreventive agents in human populations. In this regard, a significant correlation between dietary intake and many types of cancer has been shown in epidemiological data generated throughout the world, and in animal experiments, many dietary substances have been documented to have anti-cancer properties. These include green tea catechins, lycopene, soy isoflavones, pomegranate phenolics, selenium, vitamins E and D, curcumin, silibinin, and resveratrol.

* To whom correspondence should be addressed. Departments of Dermatology, Columbia University College of Physicians & Surgeons, 630 West 168th Street VC15-204, New York, NY 10032, USA, Phone: 212-851-4542, Fax: 212-305-7391, Email: ak309@columbia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Resveratrol, *trans*-3,5,4'-trihydroxy-*trans*-stilbene is a phytoalexin produced by plants, and the skin of red grapes is particularly rich in resveratrol which affects the processes underlying all three stages of carcinogenesis; namely, tumor initiation, promotion and progression. It has also been shown to suppress angiogenesis and metastasis. Extensive data in human cell cultures indicate that resveratrol can modulate multiple pathways involved in cell growth, apoptosis, and inflammation. The anti-carcinogenic effects of resveratrol appear to be closely associated with its antioxidant activity, and it has been shown to inhibit cyclooxygenase, hydroperoxidase, protein kinase C, Bcl-2 phosphorylation, Akt, focal adhesion kinase, NF- κ B, matrix metalloprotease-9, and cell cycle regulators. These and other *in vitro* and *in vivo* studies provide a rationale in support of the use of resveratrol in human cancer chemoprevention, in a combinatorial approach with either chemotherapeutic drugs or cytotoxic factors for the highly efficient treatment of drug refractory tumor cells (Seve *et al.*, 2005). This review discusses the current pre-clinical and mechanistic data available to support its potential use in diminishing risk for a variety of human cancers.

Bioavailability

Resveratrol is produced by plants in response to infection by the pathogen *Botrytis cinerea* (Delmas *et al.*, 2006). It is also induced in response to a variety of stress conditions, such as vicissitudes in climate, exposure to ozone, sunlight and heavy metals (Bavaresco, 2003). It has been detected in more than 70 plant species, including grapes, peanuts, berries, and pines. Fresh grape skin contains about 50 to 100 μ g of resveratrol per gram wet weight (Baliga *et al.*, 2005), which contributes to a relatively high concentration of resveratrol in red wine and grape juice.

trans-Resveratrol, also referred to as 3,5,4'-tri-hydroxystilbene, belongs to the stilbene class of polyphenolic compounds. It exists in both *cis* and *trans* isomeric forms. In plants, it mostly exists in glycosylated piceid forms (3-O-B-D-glucosides). Other minor conjugated forms containing 1–2 methyl groups (pterostilbene), a sulfate group (*trans*-resveratrol-3-sulfate) or a fatty acid have also been identified. Glycosylation is known to protect resveratrol from oxidative degradation, and glycosylated resveratrol is more stable and more soluble and readily absorbed in the human gastrointestinal tract (Regev-Shoshani *et al.*, 2003). In humans, following its absorption, it is readily metabolized in the liver by Phase-2 drug-metabolizing enzymes to water-soluble *trans*-resveratrol-3-O-glucuronide and *trans*-resveratrol-3-O-sulfate, accounting for its predominant urine excretion (Walle *et al.*, 2004). Compared to resveratrol, which has a plasma half-life of 8–14 min, the metabolites have a plasma half-life of about 9.2 hours (Walle *et al.*, 2004). However, the bioavailability and efficacy of these resveratrol metabolites is unknown (Baur and Sinclair, 2006). Compared to other known polyphenols, such as quercetin and catechin, *trans*-resveratrol is well absorbed much more efficiently following oral administration to humans (Soleas *et al.*, 2001).

Numerous studies exist that have utilized a wide range of concentrations of resveratrol, suggesting that its biological effects may vary depending on cell and tissue types. For example, concentrations between 32 nM and 100 μ M have been used to study various effects of resveratrol *in vitro*, and 100 ng-1,500 mg per kg (body weight) for animal studies (Baur and Sinclair, 2006). Preclinical studies in rats, using HPLC methods, have suggested that intragastric administration of 20 mg/kg *trans*-resveratrol generated peak values of 1.2 μ M in plasma (Asensi *et al.*, 2002). In a separate study, male rats treated with 300, 1000, and 3000 mg/kg body weight per day were reported to achieve plasma concentrations of 576, 991, and 2728 ng/ml, respectively, and whereas in females, it was 333, 704, and 1137 ng/ml (Crowell *et al.*, 2004). A plasma concentration of approximately 1.1 μ g/ml was determined to be approximately 5 μ M (Crowell *et al.*, 2004). A single oral administration of ¹⁴C-*trans*-resveratrol to male Balb/c mice showed preferential binding of radio-labeled resveratrol in the

stomach, liver, kidney, intestine, bile, and urine, and penetrated the tissues of the liver and kidney (Vitrac *et al.*, 2003). Both the parent compound and the Phase-II metabolites were also detected in these tissues (Vitrac *et al.*, 2003). In humans, 24.6% of the oral dose administered appeared in the urine, including metabolites (Soleas *et al.*, 2001), whereas after intragastric administration to rodents, only 1.5% of resveratrol reached the plasma compartment (Asensi *et al.*, 2002). This may reflect, in part, polymorphisms related to intestinal absorption and/or rates of metabolism in the liver and intestine among the species.

Skin cancer

Non-melanoma skin cancer

Skin cancer is the most common type of human malignancy. Each year, more than one million new cases of non-melanoma skin cancer (NMSC) are diagnosed in the United States alone. Topical resveratrol has been tested for its efficacy against the development of several cutaneous disorders, including skin cancer (Afaq *et al.*, 2002; Baliga and Katiyar, 2006; Bode and Dong, 2000). A single topical application of resveratrol (25 μmol) to SKH-1 hairless mice significantly inhibited ultraviolet B (UVB) (180 mJ/cm^2)-mediated phototoxicity including enhancement of bifold skin thickness and skin edema (Afaq *et al.*, 2003). Skin hyperplasia induced by multiple exposures to UVB radiation (180 mJ/cm^2 ; 7 exposures on alternate days) was also inhibited when resveratrol was applied topically at a dose of 10 $\mu\text{mol}/\text{animal}$, 30 min prior to each UVB exposure. The anti-proliferative effects of resveratrol were shown to be modulated by cell cycle regulatory proteins (Reagan-Shaw *et al.*, 2004). Resveratrol decreased the expression of cyclins D1 and D2, Cdk 2, 4 and 6, and proliferating cell nuclear antigen (PCNA) whereas p21WAF1/CIP1 was increased. Furthermore there was inhibited expression of anti-apoptotic proteins, such as survivin, and markers of tumor promotion, cyclooxygenase (COX)-2, and ornithine decarboxylase (ODC) were observed (Aziz *et al.*, 2005).

The chemopreventive effects of resveratrol were assessed, employing UVB-mediated skin tumorigenesis in the SKH-1 hairless mouse model. Topical application of resveratrol either pre- or post-UVB significantly inhibited tumor incidence and delayed the onset of tumorigenesis. Similarly in a standard chemical carcinogenesis mouse tumor protocol using a two-stage, DMBA-initiated and TPA-promoted murine skin cancer model showed a 98% reduction in skin tumors (Jang *et al.*, 1997). Soleas *et al.* also found a 60% reduction in papillomas with topically applied resveratrol (Soleas *et al.*, 2002), which could be related to its cytotoxic and free radical scavenging activities (Kapadia *et al.*, 2002). Orally administered resveratrol was also shown to inhibit DMBA/croton oil-induced mouse skin papillomas, correlated with prolonging the latent period of tumor occurrence and inhibiting croton oil-induced enhancement of epidermal ODC activities (Fu *et al.*, 2004). *In vitro* data generated in human squamous cell carcinoma cells demonstrate that resveratrol induces G1-phase cell cycle arrest, accompanied by p21WAF1/CIP1 induction, and it decreases the cell cycle regulators, cyclins D1/D2/E and Cdks, hyperphosphorylated pRb proteins, MEK1 > ERK1/2, and AP-1 signaling (Adhami *et al.*, 2001; Ahmad *et al.*, 2001; Kim *et al.*, 2006).

Much of the work regarding skin tumorigenesis has focused on the effects of UVB radiation, whereas much less is known about UVA-induced signaling pathways and their role in tumor promotion. UVA comprises more than 90% of incident solar radiation that reaches the surface of the earth and has been linked to the development of cutaneous squamous cell carcinomas (SCCs) (Bachelor and Bowden, 2004). Seve *et al.* showed that resveratrol potentiates generation of 8-oxo-7,8-dihydro-2'-deoxyguanosine in UVA-irradiated genomic DNA in immortalized HaCat human keratinocyte cells. In the presence of resveratrol, UVA was shown to significantly enhance the induction of DNA strand breaks and cell death (Seve *et al.*, 2005). These *in vitro* data suggest the differential effects and/or mechanism of resveratrol in the context of the UV spectrum.

Melanoma

Resveratrol can inhibit growth and induce apoptosis in melanoma cell lines (Hsieh *et al.*, 2005; Niles *et al.*, 2003). However, oral resveratrol (20 mg/kg twice per day or in the drinking water at 23 mg/l) did not inhibit the growth of B16M melanoma cells inoculated into the footpads of mice but, interestingly, it decreased the hepatic metastatic invasion of B16M cells inoculated intrasplenically (Asensi *et al.*, 2002). The intraperitoneal administration of resveratrol at the time of i.m. injection of B16-BL6 cells into syngeneic mice resulted in a dose-dependent delay of tumor growth, without toxicity. In this study, resveratrol, was found to be less effective when compared with other polyphenols tested, such as EGCG, apigenin, and quercetin (Caltagirone *et al.*, 2000). In a human melanoma xenograft model, Niles *et al.* reported that resveratrol did not have a statistically significant effect on melanoma growth, and it might even stimulate tumor growth at higher dose levels (0.006% in food or 100 mg in slow-release pellets). In addition, piceatannol, a major resveratrol metabolite, did not affect the *in vitro* growth of a murine melanoma cell line, but significantly stimulated the number of lung metastases (Niles *et al.*, 2006). The resveratrol content in the skin of these mice, measured 5 min after a bolus of 75 mg/kg introduced, was found to be 21 nmol/g and 4.67 nmol/g in glucuronide-conjugate forms. A measurable amount of resveratrol was found in the tumors, although it was less than the amount found in the skin. The *in vivo* studies appear to show that resveratrol is not an effective chemotherapeutic agent in inhibiting melanoma growth in animals, although more pre-clinical studies would be needed for confirmation.

Breast cancer

Resveratrol is considered to be a phytoestrogen, based on its structural similarity to diethylstilbestrol, a synthetic estrogen. It can bind to both alpha- and beta-estrogen receptors, and activates estrogen receptor-dependent transcription in human breast cancer cells. Despite a number of studies performed using both hormone-sensitive and hormone-resistant breast cancer cells, the estrogen-modulatory effects of resveratrol remain controversial (Le Corre *et al.*, 2005). In some cell types, such as estrogen receptor (ER) positive MCF-7 and T47D cells, resveratrol acts as a superagonist, whereas in others, it produces activation equal to or less than that of estradiol (Gehm *et al.*, 1997). In other studies, Clement *et al.* reported Fas/Fas ligand-mediated growth inhibition of T47D by resveratrol (Clement *et al.*, 1998). Resveratrol (> 1 μ M) inhibited the growth of MCF-7 cells by antagonizing the growth-stimulatory effect of 17-beta-estradiol (E2) in a dose-dependent manner (Lu and Serrero, 1999), as well as the highly invasive MDA-MB-435 cells (Hsieh *et al.*, 1999). In the absence of E2, resveratrol was observed to carry out mixed estrogen agonist/antagonist activities in some mammary cancer cell lines, but in the presence of E2, resveratrol (1 nM) was shown to function as an anti-estrogen (Bhat *et al.*, 2001). Resveratrol was also shown to inhibit the proliferation of the estrogen-receptor negative human breast carcinoma cell line, MDA-MB-468, by inhibiting the levels of autocrine growth stimulators, transforming growth factor-alpha (TGF- α), PC cell-derived growth factor, and insulin-like growth factor I receptor, and increasing the growth inhibitor TGF- β 2 (Serrero and Lu, 2001).

One implication of these varied *in vitro* results is that the chemopreventive effects of resveratrol likely to be very complex. In fact, in addition to its antioxidant scavenging of free radicals and modulating ER activity (Magee and Rowland, 2004), resveratrol can interfere with an ER α -associated PI3K pathway, following a process that could be independent of the nuclear functions of the ER α (Pozo-Guisado *et al.*, 2004), and acts as an agonist for the cAMP/kinase-A system (El-Mowafy and Alkhalaf, 2003). It also promotes the accumulation of growth inhibitory/pro-apoptotic ceramide (Scarlatti *et al.*, 2003) and induction of quinone reductase (QR, a phase II detoxification enzyme) (Bianco *et al.*, 2005), and induces caspase-independent apoptosis through Bcl-2 downregulation (Pozo-Guisado *et al.*, 2005). It has been shown to

suppress Src tyrosine kinase activity (Kotha *et al.*, 2006a), nitric oxide generation, and the NF κ B pathway (Bhat and Pezzuto, 2002).

In adult rats, resveratrol inhibited the formation of estrogen-dependent preneoplastic ductal lesions induced by DMBA in these mammary glands (IC₅₀ = 3.2 μ M) and reduced MNU-induced mammary tumorigenesis when administered by gavage, suggesting the potential beneficial effects of resveratrol against mammary tumorigenesis (Bhat *et al.*, 2001). Furthermore, dietary resveratrol (10 ppm) in Sprague Dawley rats produced striking reductions in the incidence (45%; $p < 0.05$), multiplicity (55%; $P < 0.001$), and extended the latent period of tumor induction by DMBA in a mammary carcinogenesis protocol. This was associated with decreased COX-2 and matrix metalloprotease-9 expression and suppression of NF κ B activation (Banerjee *et al.*, 2002). Recently, lifetime administration of resveratrol in the diet (1 g/kg) to rats reportedly reduced susceptibility to mammary cancer in DMBA-induced mammary carcinogenesis, which correlated with a significant reduction in proliferative cells and an increase in apoptotic cells in mammary terminal ductal structures, and more differentiated lobular structures (Whitsett *et al.*, 2006).

Resveratrol supplementation (1 mg/l; 4 μ g/mouse) delayed spontaneous mammary tumor development and reduced metastasizing capacity in HER-2/new overexpressing transgenic mice, which develop multiple mammary tumors at an early age. This anti-cancer effect was associated with the downregulation of HER-2/neu (Provinciali *et al.*, 2005). In human breast cancer xenografts, lower tumor growth, decreased angiogenesis, and an increased apoptotic index was seen in MDA-MB-231 tumors in resveratrol-treated (25 mg/kg/day) nude mice (Garvin *et al.*, 2006). In one study, however, no growth-inhibitory effect was observed in highly metastatic 4T1 murine mammary cancer cells when resveratrol was administered intraperitoneally (1–5 mg/kg daily for 23 days, starting at the time of tumor inoculation), although the growth of these cells was inhibited in *in vitro* studies (Bove *et al.*, 2002). The differences in these xenograft experiments remain largely unexplained, but it may be possibly be due to inadequate dosing.

The available experimental evidence regarding breast cancer risk and consumption of estrogenic chemicals during critical periods of development indicates that the timing and level of exposure to estrogenic chemicals are likely to be important risk factors. A short treatment of pre-pubertal female Sprague-Dawley rats with high-dose resveratrol (100 mg/kg) enhanced MNU-induced mammary carcinogenesis in an estrogen-free environment, as reflected by significant increases in the incidence and multiplicity of mammary tumors (Sato *et al.*, 2003). These data suggest that pre-pubertal resveratrol exposure affects endocrine function. Although the precise mechanism by which resveratrol accelerates the occurrence of mammary cancer in these mice requires further investigation, an increase in the number of TEBs (terminal end buds) and reduction of their differentiation into alveolar buds has been shown to play a critical role in increasing the risk of breast cancer (Russo and Russo, 1987; Sato *et al.*, 2003). It is unclear whether a high dose of resveratrol mediates this process. Although the dose of 100 mg/kg/day of resveratrol represents 5,000 times the amount that is consumed by a person drinking one glass of red wine a day (Juan *et al.*, 2002), these studies point to the need for further investigation to determine whether the potential use of resveratrol and/or other dietary consumption of phytoestrogens as a chemopreventive agent for breast cancer is safe in younger, high-risk populations.

Gastric and colorectal cancer

Because the primary etiological determinants for gastric cancer are thought to be exposure to chemical carcinogens and/or chronic infection with *Helicobacter pylori*, and resveratrol was found to be effective in inhibiting the replication of *Helicobacter pylori* (Mahady and Pendland,

2000), this provides a reason for the intervention studies using resveratrol for combating gastric cancer (Atten *et al.*, 2005). A number of cell types respond to resveratrol treatment by manifesting cell cycle arrest and apoptosis that, in part, is mediated through nitric oxide formation, which then interferes with endogenously produced reactive oxygen (Holian *et al.*, 2002). The signaling through PKC alpha and delta activity also seems to be affected by resveratrol treatment (Atten *et al.*, 2005). The intracellular apoptotic signals engaged by resveratrol may be cell type-dependent, and may be related to differentiation status in various gastric adenocarcinoma cell lines (Riles *et al.*, 2006). In colon cancer cells, resveratrol activates various caspases and triggers apoptosis, which involves the accumulation of the pro-apoptotic proteins Bax and Bak and redistribution of the Fas receptor in membrane rafts (Delmas *et al.*, 2003). Relatively high concentrations also substantially downregulate telomerase activity (Fuggetta *et al.*, 2006).

The *in vivo* efficacy of resveratrol has been tested in two animal models of colorectal cancer, dimethylhydrazine-induced AOM and mutant *Min* mice. AOM-induced tumors share many histopathologic similarities with human tumors, and they often carry mutations in *K-ras* and β -catenin genes but, unlike human tumors, the *Apc* gene (15%) is less frequently mutated. The *Min* mice harbor a mutated *Apc* gene similar to that found in patients with familial adenomatous polyposis, and in many sporadic cancers (Corpet and Pierre, 2003). Administered orally at 200 μ g/kg/day in the drinking water, resveratrol significantly reduced the number of AOM-induced aberrant crypt foci (ACF) associated with changes in Bax and p21 expression (Tessitore *et al.*, 2000). *Min* mice receiving resveratrol (0.01% in the drinking water for 7 weeks) showed a 70% reduction in the formation of small intestinal tumors and prevented colon tumor development. Resveratrol treatment led to the downregulation of genes that are directly involved in cell cycle progression or cell proliferation (cyclins D1 and D2, DP-1 transcription factor, and Y-box binding protein) and the upregulation of genes that are involved in the recruitment and activation of immune cells (cytotoxic T lymphocyte Ag-4, leukemia inhibitory factor receptor, and monocyte chemotactic protein 3) and in the inhibition of the carcinogenic process and tumor expansion (tumor susceptibility protein TSG101, transforming growth factor-beta, inhibin-beta A subunit, and desmocollin 2), suggesting the multiplicity of the molecular targets and signaling cascades (Schneider *et al.*, 2001). An inhibitory effect was also observed in the experiments conducted in the xenograft gastric tumor model, which utilized high doses of resveratrol (500–1,500 mg/kg), directly injected beside the tumor body 6 times, at an interval of 2 d (Zhou *et al.*, 2005). The synthetic resveratrol analog 3,4,5,4'-tetramethoxystilbene (DMU-212) also inhibited the development of adenomas in the *Apc*(*Min*+) mouse (Sale *et al.*, 2005). In addition, the efficacy of resveratrol was replicated in other rodent models of colon chemical carcinogenesis. Resveratrol (8 mg/kg body weight, administered every day for 30 weeks) markedly reduced tumor incidence and the occurrence of histological lesions, as well as the size of tumors in 1,2-dimethylhydrazine (DMH)-induced colon carcinogenesis in Wistar rats (Sengottuvelan *et al.*, 2006). However, the data generated in *Min* mice are conflicting. Although resveratrol administered in the drinking water strongly reduced the formation of colon and small intestinal tumors (Schneider *et al.*, 2001), the doses used in this study when consumed ad libitum in the diet, up to 90 mg/kg for 7 weeks, were not effective (Ziegler *et al.*, 2004). The reason for this discrepancy is not understood (resveratrol in diet vs. in drinking water).

Lung Cancer

A lower risk of lung cancer among consumers of wine compared with consumers of other beverages has been observed, which may be partly attributed to the high resveratrol content, particularly in red wine. Resveratrol alters the expression of PAH (polycyclic aromatic hydrocarbons)-metabolizing genes, such as the cytochrome P450 1A1 (CYP1A1) and 1B1 (CYP1B1), microsomal epoxide hydrolase (mEH), and glutathione S-transferase P1 (GSTP1)

genes, resulting in the altered formation of carcinogenic benzo[a]pyrene (BaP) metabolites in human bronchial epithelial cells (Mollerup *et al.*, 2001). It inhibits the expression of CYP1A1 and CYP1B1 and the generation of reactive diol-epoxides that can bind to DNA forming covalent adducts that cause structural alterations with mutations (Berge *et al.*, 2004a). BaP is ubiquitous environmental pollutant and is also present in cigarette smoke. There are numerous carcinogens in cigarette smoke that are likely involved in the pathogenesis of this type of tumor. Interestingly, BaP metabolism requires the induction of cytochrome P-450 1A1 (CYP1A1) through the activation of the AhR. Balb-C mice receiving resveratrol (50 mg/kg/week for 5 weeks) were found to have significantly fewer BPDE-DNA adducts (Revel *et al.*, 2003). Doses of 2.5 and 10 mg/kg resveratrol were also found to significantly reduce the tumor volume (42%), tumor weight (44%) and metastatic potential (56%) in mice bearing highly metastatic Lewis lung carcinomas (LLCs), through the inhibition of DNA synthesis and LLC-induced neovascularization and tube formation of HUVEC (Kimura and Okuda, 2001).

However, dietary resveratrol (68 mg/kg) that did not alter CYP1A1 and CYP1B1 gene expression showed no effect on B[a]P-induced lung tumorigenesis in A/J mice (Berge *et al.*, 2004b). Similarly, no effect on lung tumor multiplicity was shown in A/J mice fed a diet supplemented with resveratrol (500 ppm) from 1 week after B[a]P and 4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone (NNK) treatment until termination (Hecht *et al.*, 1999). No resveratrol or resveratrol conjugates were detectable by HPLC in the lung tissue of animals receiving a resveratrol-supplemented diet. It remains to be defined whether resveratrol given in the diet reaches the lung tissue in sufficient concentrations, or in a biologically active form.

Esophageal Tumorigenesis

Esophageal cancer is common worldwide, and generally has a poor prognosis because the diagnosis is often delayed. Smoking, including exposure to polyaromatic hydrocarbons like benzo[a]pyrene (BaP), is known to be a major risk factor. Resveratrol was shown to suppress the growth of an esophageal cancer cell line EC-9706 (Zhou *et al.*, 2003). The *in vivo* anti-cancer effects of resveratrol were evaluated at concentrations between 1 and 2 mg/kg body weight in N-nitrosomethylbenzylamine (NMBA)-induced esophageal tumorigenesis in rats. Resveratrol suppressed both the number and size of NMBA-induced esophageal tumors per rat, by targeting COXs and PGE(2) (Li *et al.*, 2002).

Prostate Cancer

The growth inhibitory effect of resveratrol has been demonstrated in various cultured prostate cancer cells, both hormone-sensitive and hormone-refractory, which mimic the initial or advanced stages of prostate carcinoma, respectively. These studies have shown that resveratrol substantially modulates the growth of these cells and alters the expression of more than one set of functionally related molecular targets. Resveratrol can repress different classes of androgen-responsive genes, including prostate-specific antigen (PSA), human glandular kallikrein-2, AR (androgen receptor)-specific coactivator ARA70, and p21WAF1/CIP1 in hormone-responsive cells (Mitchell *et al.*, 1999); activate p53-responsive genes such as PIG7, p300/CBP and Apaf-1 (Narayanan *et al.*, 2003); inhibit PI3K/AKT activation; and increase Bax, Bak, Bid, and Bad (Aziz *et al.*, 2006). Resveratrol downregulates PSA by a mechanism independent of changes in AR (Hsieh and Wu, 1999; Hsieh and Wu, 2000). No or minimal interaction of resveratrol with AR was detected (Kampa *et al.*, 2000). Moreover, resveratrol was shown to modulate NO production (Kampa *et al.*, 2000) and prevent the increase in reactive oxygen species (ROS) (Sgambato *et al.*, 2001). Resveratrol treatment of various prostate cells also accompanied the activation of MAPK signaling and an increase in cellular p53 content, likely due to stabilization by serine-15 phosphorylation (Gao *et al.*, 2004; Lin *et al.*, 2002),

ceramide-associated growth inhibition (Sala *et al.*, 2003), and the blockade of Stat3-mediated dysregulation of growth and survival pathways (Kotha *et al.*, 2006a). Currently no pre-clinical studies have been reported on prostate carcinogenesis.

Hepatoma

Oxidative stress has been implicated in the pathogenesis of liver cancer and, therefore, the use of antioxidants as a therapeutic or preventive agent has been recommended. *In vitro*, resveratrol was shown to block the ROS-potentiated invasion of the hepatoma cells (Daiki *et al.*, 2004) and induce apoptosis (Michels *et al.*, 2006). The pre-treatment of the mice with resveratrol at a dose of 2.5 mg/kg body weight for 2 weeks was also shown to block N-nitrosodiethylamine (NDEA)-induced ODC and COX activities in hepatic tissue (Khanduja *et al.*, 2004). In the transplantable murine hepatoma22 model, resveratrol could induce the S phase arrest of H22 cells after the tumor-bearing mice were treated with 10 mg/kg or 15 mg/kg resveratrol for ten days (Wu *et al.*, 2004). It also enhanced the anti-tumor effect of 5-FU, suggesting that resveratrol could be a biochemical modulator to enhance the therapeutic effects of 5-FU for hepatocellular carcinoma. Resveratrol caused a significant decrease (25%) in tumor cell content by inducing apoptosis. In this study, it was shown that resveratrol treatment also induces an accumulation of cells in the G2/M phase in rats inoculated with a fast-growing tumor (the Yoshida AH-130 ascites hepatoma) (Carbo *et al.*, 1999). The inhibition of the cell cycle progression was further shown to involve decreases in the expressions of cyclin B1 and p34cdc2 in murine transplantable liver tumors, following the administration of resveratrol at 10 or 15 mg/kg bodyweight for 10 days (Yu *et al.*, 2003). Dietary resveratrol (10 or 50 ppm) is also shown to have anti-tumor growth effects and anti-metastasis effects in Donryu rats subcutaneously implanted with an ascites hepatoma cell line of AH109A (Miura *et al.*, 2003). In another study, daily i.p. injections of resveratrol at 1 mg/kg for 7 days were shown to be effective in inhibiting ascites hepatoma in rats (Carbo *et al.*, 1999).

Neuroblastoma

In neuronal-like cells, such as the human neuroblastoma SH-SY5Y resveratrol was shown to inhibit caspase 7 activation, as well as degradation of poly-(ADP-ribose)-polymerase, which occur in cells exposed to paclitaxel, an anti-cancer drug (Nicolini *et al.*, 2001). The neuroprotective action of resveratrol was suggested to occur through modulating the signal pathways that commit these neuronal-like cells to apoptosis. Resveratrol was shown to induce S-phase arrest, preventing SH-SY5Y from entering mitosis, the phase of the cell cycle in which paclitaxel exerts its activity (Rigolio *et al.*, 2005). Furthermore, phosphorylation of Bcl-2 and JNK/SAPK, which specifically occurs after paclitaxel exposure, was reversed by resveratrol (Nicolini *et al.*, 2003). The long-term treatment of neuroblastoma cells with resveratrol is also reported to enhance the differentiation state of the cells (Melzig and Escher, 2002). In stage 4 MYCN-amplified neuroblastoma cell lines, resveratrol was shown to decrease cell viability and induce cell cycle arrest and apoptosis, which accompanied transiently up-regulated p53 expression and nuclear translocation of p53, followed by induction of p21(WAF-1/CIP-1) and Bax expression (Liontas and Yeger, 2004). In mice, intraperitoneal injections of 40 mg/kg body weight resveratrol daily for 28 days suppressed the growth rate of subcutaneous neuroblastomas, resulting in 70% long-term survival (Chen *et al.*, 2004).

Fibrosarcoma

Resveratrol was shown to inhibit FGF2-induced angiogenesis and significantly inhibited platelet/fibrin clot-promoted human colon and fibrosarcoma tumor growth in the CAM (chick chorioallantoic membrane) tumor model (Mousa *et al.*, 2005). Oral administration of resveratrol (1 mg/kg per day) inhibited the growth of a murine T241 fibrosarcoma implanted in C57B16/J mice (Brakenhielm *et al.*, 2001).

Pancreatic Cancer

Several *in vitro* studies reported that trans-resveratrol enhanced apoptosis in pancreatic cancer cells, which is associated with mitochondrial depolarization and cytochrome c release followed by caspase-3 activation (Mouria *et al.*, 2002). Resveratrol (100 μ M) inhibited the proliferation of human pancreatic cancer cell lines, PANC-1 and AsPC-1, in a concentration- and time-dependent manner, and increased the fraction of sub-G0/G1 cells (Ding and Adrian, 2002). Furthermore, treatment with resveratrol was shown to sensitize MiaPaCa2 cells for tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced apoptosis through p53-independent induction of p21 and p21-mediated cell cycle arrest associated with survivin depletion (Fulda and Debatin, 2004). Resveratrol can inhibit Src tyrosine kinase activity and thereby blocks the constitutive signal transducer and activator of transcription 3 (Stat3) protein activation, suggesting Src-Stat3 signaling is a target for resveratrol (Kotha *et al.*, 2006b). However, resveratrol given in the diet at a concentration of 10 ppm failed to show significant effects on BOP (N-nitrosobis(2-oxopropyl)amine)-initiation of hamster pancreatic carcinogenesis (Kuroiwa *et al.*, 2006). Additional studies are required for further pre-clinical evaluation of resveratrol efficacy on pancreatic cancer.

Leukemia

A number of *in vitro* studies have demonstrated the anti-proliferative effects of resveratrol in various leukemic cell lines (U937, HL-60) (Gautam *et al.*, 2000; Surh *et al.*, 1999), inducing apoptosis through the CD95-CD95 pathway (Clement *et al.*, 1998). Resveratrol was also shown to induce apoptosis in B-lineage leukemic cells (acute lymphoblastic leukemias) that are resistant to CD95 signaling, through the CD95-independent, mitochondria/caspase-9-specific pathway (Dorrie *et al.*, 2001). Furthermore, involvement of the Cdc42/apoptosis signal-regulating kinase 1/c-Jun N-terminal kinase/FasL signaling cascade was observed in HL-60 cells treated with resveratrol (Su *et al.*, 2005). Downregulation of the two anti-apoptotic proteins, iNOS and Bcl-2, was observed in leukemic B cells from chronic leukaemia treated with resveratrol (Roman *et al.*, 2002). Interleukin 1beta (IL-1 β) plays a key role in the proliferation of acute myeloid leukemia (AML) cells, and in the AML cell lines OCIM2 and OCI/AML3, resveratrol was shown to inhibit proliferation, arresting the cells at the S phase. Resveratrol significantly reduced the production of IL-1 β and suppressed the IL-1 β -induced activation of NF- κ B (Estrov *et al.*, 2003). Interestingly, resveratrol at low concentrations (4–8 μ M) was shown to inhibit apoptosis in human leukemia cells via NADPH oxidase-dependent elevation of intracellular superoxide that blocks mitochondrial hydrogen peroxide production, thus resulting in an cellular environment not susceptible to apoptosis induction (Ahmad *et al.*, 2004). The combined effects of resveratrol have been tested with Ara-C or tiazofurin, both antimetabolites, and showed synergistic growth inhibition and apoptosis induction in HL-60 cells (Horvath *et al.*, 2005). *In vivo*, however, only weak potential anti-leukemic resveratrol activity was suggested at a dose of 80 mg/kg body daily for 60 days in mice implanted with a mouse myeloid leukemia cell line, 32Dp210, despite the strong anti-proliferative and pro-apoptotic activities of resveratrol against these cells *in vitro* (Gao *et al.*, 2002).

Conclusion

The attractiveness of naturally occurring compounds for cancer chemoprevention has escalated in recent years. An ideal chemopreventive/therapeutic agent would restore normal growth control to preneoplastic or cancerous cells by modulating aberrant signaling pathways and/or inducing apoptosis. It should target the multiple biochemical and physiological pathways involved in tumor development, while minimizing toxicity in normal tissues (Manson *et al.*, 2005; Mukhtar and Ahmad, 1999a; Mukhtar and Ahmad, 1999b; Yance and Sagar, 2006). Resveratrol has been shown to be an effective chemopreventive agent in multiple murine

models of human cancers. It has the capacity to interact with multiple molecular targets and appears to be relatively non-toxic at least at the doses tested in these models. However, much work needs to be done: 1) to improve the bioavailability and pharmacologic properties in the different target tissues, and 2) to better understand its exact mechanisms of action in order to predict its efficacy.

The results of these investigations depended critically not only on the agent evaluated, but also on the target population studied and the specific endpoints evaluated (1999). Several phase-I clinical trials are currently underway to evaluate the pharmacokinetics and safety of resveratrol, utilizing various dosages of resveratrol (50 mg–1000 mg) in HIV seronegative, healthy subjects (the Institute of Human Virology) (Baur and Sinclair, 2006). Clinical studies with appropriate end points and surrogate markers for anticancer response could help evaluate resveratrol for larger, more intensive clinical trials. A phase-I clinical trial in colon cancer patients (University of California, Irvine) is intended to examine the effects of resveratrol treatment on colon cancer progression and colonic mucosa in colon cancer patients, as well as its effects in modulating the Wnt signaling pathway. Moreover, given the multiple effects of resveratrol and the relatively low dose of resveratrol obtained from red wine or other dietary sources, its health benefit may lie in synergistic combinations with other agents. Resveratrol is shown to have a synergistic effect *in vitro* with both quercetin and ellagic acid for apoptosis (Baur and Sinclair, 2006). Researchers have started to explore resveratrol treatment in combination with other agents in some pre-clinical studies. Taken together, these studies are expected to provide data on resveratrol's mechanisms of action and provide a foundation for future prevention trials and therapeutic clinical research, utilizing this interesting agent.

Acknowledgements

This work was partially supported by the NIH Grants K01-AR048582-03 to ALK and R01 CA-097249-01 to DRB/MA.

References

- Prevention of Cancer in the Next Millennium: Report of the Chemoprevention Working Group to the American Association for Cancer Research. 1999;59:4743–4758.
- Adhami VM, Afaq F, Ahmad N. Involvement of the Retinoblastoma (pRb)-E2F/DP Pathway during Antiproliferative Effects of Resveratrol in Human Epidermoid Carcinoma (A431) Cells. *Biochemical and Biophysical Research Communications* 2001;288:579–585. [PubMed: 11676482]
- Afaq F, Adhami VM, Ahmad N. Prevention of short-term ultraviolet B radiation-mediated damages by resveratrol in SKH-1 hairless mice. *Toxicology and Applied Pharmacology* 2003;186:28–37. [PubMed: 12583990]
- Afaq F, Adhami VM, Ahmad N, Mukhtar H. Botanical antioxidants for chemoprevention of photocarcinogenesis. *Front Biosci* 2002;7:d784–92. [PubMed: 11897547]
- Ahmad KA, Clement MV, Hanif IM, Pervaiz S. Resveratrol inhibits drug-induced apoptosis in human leukemia cells by creating an intracellular milieu nonpermissive for death execution. *Cancer Research* 2004;64:1452–9. [PubMed: 14973069]
- Ahmad, N.; Adhami, VM.; Afaq, F.; Feyes, DK.; Mukhtar, H. *Cells*. 7. 2001. Resveratrol Causes WAF-1/p21-mediated G1-phase Arrest of Cell Cycle and Induction of Apoptosis in Human Epidermoid Carcinoma A431; p. 1466-1473.
- Asensi M, Medina I, Ortega A, Carretero J, Bano MC, Obrador E, Estrela JM. Inhibition of cancer growth by resveratrol is related to its low bioavailability. *Free Radical Biology and Medicine* 2002;33:387–398. [PubMed: 12126761]
- Atten MJ, Godoy-Romero E, Attar BM, Milson T, Zopel M, Holian O. Resveratrol regulates cellular PKC alpha and delta to inhibit growth and induce apoptosis in gastric cancer cells. *Investigational New Drugs* 2005;23:111–9. [PubMed: 15744586]

- Aziz MH, Afaq F, Ahmad N. Prevention of ultraviolet-B radiation damage by resveratrol in mouse skin is mediated via modulation in survivin. *Photochemistry and Photobiology* 2005;81:25–31. [PubMed: 15469386]
- Aziz MH, Nihal M, Fu VX, Jarrard DF, Ahmad N. Resveratrol-caused apoptosis of human prostate carcinoma LNCaP cells is mediated via modulation of phosphatidylinositol 3'-kinase/Akt pathway and Bcl-2 family proteins. 2006;5:1335–1341.
- Bachelor MA, Bowden GT. UVA-mediated activation of signaling pathways involved in skin tumor promotion and progression. *Seminars in Cancer Biology* 2004;14:131–138. [PubMed: 15018897]
- Baliga MS, Katiyar SK. Chemoprevention of photocarcinogenesis by selected dietary botanicals. *Photochem Photobiol Sci* 2006;5:243–53. [PubMed: 16465310]
- Baliga MS, Meleth S, Katiyar SK. Growth inhibitory and antimetastatic effect of green tea polyphenols on metastasis-specific mouse mammary carcinoma 4T1 cells in vitro and in vivo systems. *Clinical Cancer Research* 2005;11:1918–27. [PubMed: 15756018]
- Banerjee S, Bueso-Ramos C, Aggarwal BB. Suppression of 7,12-Dimethylbenz(a)anthracene-induced Mammary Carcinogenesis in Rats by Resveratrol: Role of Nuclear Factor- κ B, Cyclooxygenase 2, and Matrix Metalloprotease 9. 2002;62:4945–4954.
- Baur JA, Sinclair DA. Therapeutic potential of resveratrol: the in vivo evidence. *Nat Rev Drug Discov* 2006;5:493–506. [PubMed: 16732220]
- Bavaresco L. Role of viticultural factors on stilbene concentrations of grapes and wine. *Drugs under Experimental and Clinical Research* 2003;29:181–7. [PubMed: 15134373]
- Berge G, Ovrebø S, Botnen IV, Hewer A, Phillips DH, Haugen A, Møllerup S. Resveratrol inhibits benzo[a]pyrene-DNA adduct formation in human bronchial epithelial cells. *British Journal of Cancer* 2004a;91:333–8. [PubMed: 15162144]
- Berge G, Ovrebø S, Eilertsen E, Haugen A, Møllerup S. Analysis of resveratrol as a lung cancer chemopreventive agent in A/J mice exposed to benzo[a]pyrene. *British Journal of Cancer* 2004b;91:1380–3. [PubMed: 15316568]
- Bhat KPL, Lantvit D, Christov K, Mehta RG, Moon RC, Pezzuto JM. Estrogenic and Antiestrogenic Properties of Resveratrol in Mammary Tumor Models. 2001;61:7456–7463.
- Bhat KPL, Pezzuto JM. Cancer Chemopreventive Activity of Resveratrol. 2002;957:210–229.
- Bianco NR, Chaplin LJ, Montano MM. Differential induction of quinone reductase by phytoestrogens and protection against oestrogen-induced DNA damage. *Biochem J* 2005;385:279–87. [PubMed: 15456407]
- Bode AM, Dong Z. Signal transduction pathways: targets for chemoprevention of skin cancer. *The Lancet Oncology* 2000;1:181–8. [PubMed: 11905657]
- Bove K, Lincoln DW, Tsan MF. Effect of resveratrol on growth of 4T1 breast cancer cells in vitro and in vivo. *Biochemical and Biophysical Research Communications* 2002;291:1001–5. [PubMed: 11866465]
- Brakenhielm E, Cao R, Cao Y. Suppression of angiogenesis, tumor growth, and wound healing by resveratrol, a natural compound in red wine and grapes. 2001;15:1798–1800.
- Caltagirone S, Rossi C, Poggi A, et al. Flavonoids apigenin and quercetin inhibit melanoma growth and metastatic potential. *International Journal of Cancer* 2000;87:595–600.
- Carbo N, Costelli P, Baccino FM, Lopez-Soriano FJ, Argiles JM. Resveratrol, a natural product present in wine, decreases tumour growth in a rat tumour model. *Biochemical and Biophysical Research Communications* 1999;254:739–43. [PubMed: 9920811]
- Chen Y, Tseng SH, Lai HS, Chen WJ. Resveratrol-induced cellular apoptosis and cell cycle arrest in neuroblastoma cells and antitumor effects on neuroblastoma in mice. *Surgery* 2004;136:57–66. [PubMed: 15232540]
- Clement MV, Hirpara JL, Chawdhury SH, Pervaiz S. Chemopreventive agent resveratrol, a natural product derived from grapes, triggers CD95 signaling-dependent apoptosis in human tumor cells. *Blood* 1998;92:996–1002. [PubMed: 9680369]
- Corpet DE, Pierre F. Point: From Animal Models to Prevention of Colon Cancer. *Systematic Review of Chemoprevention in Min Mice and Choice of the Model System* 2003;12:391–400.
- Crowell JA, Korytko PJ, Morrissey RL, Booth TD, Levine BS. Resveratrol-associated renal toxicity. *Toxicol Sci* 2004;82:614–9. [PubMed: 15329443]

- Daiki M, Yutaka M, Kazumi Y. Resveratrol inhibits hepatoma cell invasion by suppressing gene expression of hepatocyte growth factor via its reactive oxygen species-scavenging property. *Clinical and Experimental Metastasis* 2004;V21:445–451.
- Delmas D, Lancon A, Colin D, Jannin B, Latruffe N. Resveratrol as a Chemopreventive Agent: A Promising Molecule for Fighting Cancer. *Current Drug Targets* 2006;7:423–442. [PubMed: 16611030]
- Delmas D, Rebe C, Lacour S, et al. Resveratrol-induced apoptosis is associated with Fas redistribution in the rafts and the formation of a death-inducing signaling complex in colon cancer cells. *The Journal of Biological Chemistry* 2003;278:41482–90. [PubMed: 12902349]
- Ding XZ, Adrian TE. Resveratrol inhibits proliferation and induces apoptosis in human pancreatic cancer cells. *Pancreas* 2002;25:e71–6. [PubMed: 12409844]
- Dorrie J, Gerauer H, Wachter Y, Zunino SJ. Resveratrol induces extensive apoptosis by depolarizing mitochondrial membranes and activating caspase-9 in acute lymphoblastic leukemia cells. *Cancer Research* 2001;61:4731–9. [PubMed: 11406544]
- El-Mowafy AM, Alkhalaf M. Resveratrol activates adenylyl-cyclase in human breast cancer cells: a novel, estrogen receptor-independent cytostatic mechanism. 2003;24:869–873.
- Estrov Z, Shishodia S, Faderl S, et al. Resveratrol blocks interleukin-1beta-induced activation of the nuclear transcription factor NF-kappaB, inhibits proliferation, causes S-phase arrest, and induces apoptosis of acute myeloid leukemia cells. *Blood* 2003;102:987–95. [PubMed: 12689943]
- Fu ZD, Cao Y, Wang KF, Xu SF, Han R. [Chemopreventive effect of resveratrol to cancer]. *Ai zheng = Aizheng = Chinese Journal of Cancer* 2004;23:869–73. [PubMed: 15301705]
- Fuggetta MP, Lanzilli G, Tricarico M, Cottarelli A, Falchetti R, Ravagnan G, Bonmassar E. Effect of resveratrol on proliferation and telomerase activity of human colon cancer cells in vitro. *Journal of Experimental Clinical Cancer Research* 2006;25:189–93. [PubMed: 16918129]
- Fulda S, Debatin K-M. Sensitization for Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Induced Apoptosis by the Chemopreventive Agent Resveratrol. 2004;64:337–346.
- Gao S, Liu GZ, Wang Z. Modulation of androgen receptor-dependent transcription by resveratrol and genistein in prostate cancer cells. *The Prostate* 2004;59:214–25. [PubMed: 15042621]
- Gao X, Xu YX, Divine G, Janakiraman N, Chapman RA, Gautam SC. Disparate in vitro and in vivo antileukemic effects of resveratrol, a natural polyphenolic compound found in grapes. *The Journal of Nutrition* 2002;132:2076–81. [PubMed: 12097696]
- Garvin S, Ollinger K, Dabrosin C. Resveratrol induces apoptosis and inhibits angiogenesis in human breast cancer xenografts in vivo. *Cancer Letters* 2006;231:113–122. [PubMed: 16356836]
- Gautam SC, Xu YX, Dumaguin M, Janakiraman N, Chapman RA. Resveratrol selectively inhibits leukemia cells: a prospective agent for ex vivo bone marrow purging. *Bone Marrow Transplantation* 2000;25:639–45. [PubMed: 10734298]
- Gehm BD, McAndrews JM, Chien P-Y, Jameson JL. Resveratrol, a polyphenolic compound found in grapes and wine, is an agonist for the estrogen receptor. 1997;94:14138–14143.
- Hecht SS, Kenney PM, Wang M, Trushin N, Agarwal S, Rao AV, Upadhyaya P. Evaluation of butylated hydroxyanisole, myo-inositol, curcumin, esculetin, resveratrol and lycopene as inhibitors of benzo [a]pyrene plus 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced lung tumorigenesis in A/J mice. *Cancer Letters* 1999;137:123–30. [PubMed: 10374832]
- Holian O, Wahid S, Atten MJ, Attar BM. Inhibition of gastric cancer cell proliferation by resveratrol: role of nitric oxide. *American Journal of Physiology* 2002;282:G809–16. [PubMed: 11960777]
- Horvath Z, Saiko P, Illmer C, et al. Synergistic action of resveratrol, an ingredient of wine, with Ara-C and tiazofurin in HL-60 human promyelocytic leukemia cells. *Experimental Hematology* 2005;33:329–35. [PubMed: 15730856]
- Hsieh, T-c; Wang, Z.; Hamby, CV.; Wu, JM. Inhibition of melanoma cell proliferation by resveratrol is correlated with upregulation of quinone reductase 2 and p53. *Biochemical and Biophysical Research Communications* 2005;334:223–230. [PubMed: 15993843]
- Hsieh, T-c; Wu, JM. Differential Effects on Growth, Cell Cycle Arrest, and Induction of Apoptosis by Resveratrol in Human Prostate Cancer Cell Lines. *Experimental Cell Research* 1999;249:109–115. [PubMed: 10328958]

- Hsieh TC, Burfeind P, Laud K, Backer JM, Traganos F, Darzynkiewicz Z, Wu JM. Cell cycle effects and control of gene expression by resveratrol in human breast carcinoma cell lines with different metastatic potentials. *International Journal of Oncology* 1999;15:245–52. [PubMed: 10402233]
- Hsieh TC, Wu JM. Grape-derived chemopreventive agent resveratrol decreases prostate-specific antigen (PSA) expression in LNCaP cells by an androgen receptor (AR)-independent mechanism. *Anticancer Research* 2000;20:225–8. [PubMed: 10769659]
- Jang M, Cai L, Udeani GO, et al. Cancer Chemopreventive Activity of Resveratrol, a Natural Product Derived from Grapes. *1997;275:218–220.*
- Juan ME, Vinardell MP, Planas JM. The daily oral administration of high doses of trans-resveratrol to rats for 28 days is not harmful. *Journal of Nutrition* 2002;132:257–260. [PubMed: 11823587]
- Kampa M, Hatzoglou A, Notas G, et al. Wine antioxidant polyphenols inhibit the proliferation of human prostate cancer cell lines. *Nutrition and Cancer* 2000;37:223–33. [PubMed: 11142097]
- Kapadia GJ, Azuine MA, Tokuda H, Takasaki M, Mukainaka T, Konoshima T, Nishino H. Chemopreventive effect of resveratrol, sesamol, sesame oil and sunflower oil in the epstein-barr virus early antigen activation assay and the mouse skin two-stage carcinogenesis. *Pharmacological Research* 2002;45:499–505. [PubMed: 12162952]
- Khanduja KL, Bhardwaj A, Kaushik G. Resveratrol inhibits N-nitrosodiethylamine-induced ornithine decarboxylase and cyclooxygenase in mice. *Journal of Nutritional Science and Vitaminology* 2004;50:61–5. [PubMed: 15228220]
- Kim AL, Zhu Y, Zhu H, Han L, Kopelovich L, Bickers DR, Athar M. Resveratrol inhibits proliferation of human epidermoid carcinoma A431 cells by modulating MEK1 and AP-1 signalling pathways. *2006;15:538–546.*
- Kimura Y, Okuda H. Resveratrol isolated from *Polygonum cuspidatum* root prevents tumor growth and metastasis to lung and tumor-induced neovascularization in Lewis lung carcinoma-bearing mice. *The Journal of Nutrition* 2001;131:1844–9. [PubMed: 11385077]
- Kotha A, Sekharam M, Cilenti L, et al. Resveratrol inhibits Src and Stat3 signaling and induces the apoptosis of malignant cells containing activated Stat3 protein. *2006a;5:621–629.*
- Kotha A, Sekharam M, Cilenti L, et al. Resveratrol inhibits Src and Stat3 signaling and induces the apoptosis of malignant cells containing activated Stat3 protein. *Molecular Cancer Therapeutics* 2006b;5:621–9. [PubMed: 16546976]
- Kuroiwa Y, Nishikawa A, Kitamura Y, Kanki K, Ishii Y, Umemura T, Hirose M. Protective effects of benzyl isothiocyanate and sulforaphane but not resveratrol against initiation of pancreatic carcinogenesis in hamsters. *Cancer Letters* 2006;241:275–80. [PubMed: 16386831]
- Le Corre L, Chalabi N, Delort L, Bignon YJ, Bernard-Gallon DJ. Resveratrol and breast cancer chemoprevention: molecular mechanisms. *Molecular Nutrition & Food Research* 2005;49:462–71. [PubMed: 15786518]
- Li ZG, Hong T, Shimada Y, et al. Suppression of N-nitrosomethylbenzylamine (NMBA)-induced esophageal tumorigenesis in F344 rats by resveratrol. *Carcinogenesis* 2002;23:1531–6. [PubMed: 12189197]
- Lin HY, Shih A, Davis FB, Tang HY, Martino LJ, Bennett JA, Davis PJ. Resveratrol induced serine phosphorylation of p53 causes apoptosis in a mutant p53 prostate cancer cell line. *The Journal of Urology* 2002;168:748–55. [PubMed: 12131363]
- Liontas A, Yeger H. Curcumin and resveratrol induce apoptosis and nuclear translocation and activation of p53 in human neuroblastoma. *Anticancer Research* 2004;24:987–98. [PubMed: 15161054]
- Lu R, Serrero G. Resveratrol, a natural product derived from grape, exhibits antiestrogenic activity and inhibits the growth of human breast cancer cells. *Journal of Cellular Physiology* 1999;179:297–304. [PubMed: 10228948]
- Magee PJ, Rowland IR. Phyto-oestrogens, their mechanism of action: current evidence for a role in breast and prostate cancer. *The British Journal of Nutrition* 2004;91:513–31. [PubMed: 15035679]
- Mahady GB, Pendland SL. Resveratrol inhibits the growth of *Helicobacter pylori* in vitro. *The American Journal of Gastroenterology* 2000;95:1849. [PubMed: 10926010]
- Manson MM, Farmer PB, Gescher A, Steward WP. Innovative agents in cancer prevention. *Recent Results in Cancer Research Fortschritte der Krebsforschung* 2005;166:257–75.

- Melzig MF, Escher F. Induction of neutral endopeptidase and angiotensin-converting enzyme activity of SK-N-SH cells in vitro by quercetin and resveratrol. *Die Pharmazie* 2002;57:556–8. [PubMed: 12227197]
- Michels G, Watjen W, Weber N, et al. Resveratrol induces apoptotic cell death in rat H4IIE hepatoma cells but necrosis in C6 glioma cells. *Toxicology* 2006;225:173–182. [PubMed: 16843582]
- Mitchell SH, Zhu W, Young CYF. Resveratrol Inhibits the Expression and Function of the Androgen Receptor in LNCaP Prostate Cancer Cells. 1999;59:5892–5895.
- Miura D, Miura Y, Yagasaki K. Hypolipidemic action of dietary resveratrol, a phytoalexin in grapes and red wine, in hepatoma-bearing rats. *Life Sciences* 2003;73:1393–400. [PubMed: 12850500]
- Mollerup S, Ovrebo S, Haugen A. Lung carcinogenesis: resveratrol modulates the expression of genes involved in the metabolism of PAH in human bronchial epithelial cells. *International Journal of Cancer* 2001;92:18–25.
- Mouria M, Gukovskaya AS, Jung Y, Buechler P, Hines OJ, Reber HA, Pandol SJ. Food-derived polyphenols inhibit pancreatic cancer growth through mitochondrial cytochrome C release and apoptosis. *International Journal of Cancer* 2002;98:761–9.
- Mousa SS, Mousa SS, Mousa SA. Effect of resveratrol on angiogenesis and platelet/fibrin-accelerated tumor growth in the chick chorioallantoic membrane model. *Nutrition and Cancer* 2005;52:59–65. [PubMed: 16091005]
- Mukhtar H, Ahmad N. Cancer chemoprevention: future holds in multiple agents. *Toxicology and Applied Pharmacology* 1999a;158:207–10. [PubMed: 10438653]
- Mukhtar H, Ahmad N. Green tea in chemoprevention of cancer. *Toxicol Sci* 1999b;52:111–7. [PubMed: 10630599]
- Narayanan BA, Narayanan NK, Re GG, Nixon DW. Differential expression of genes induced by resveratrol in LNCaP cells: P53-mediated molecular targets. *International Journal of Cancer* 2003;104:204–12.
- Nicolini G, Rigolio R, Miloso M, Bertelli AA, Tredici G. Anti-apoptotic effect of trans-resveratrol on paclitaxel-induced apoptosis in the human neuroblastoma SH-SY5Y cell line. *Neuroscience Letters* 2001;302:41–4. [PubMed: 11278107]
- Nicolini G, Rigolio R, Scuteri A, Miloso M, Saccomanno D, Cavaletti G, Tredici G. Effect of trans-resveratrol on signal transduction pathways involved in paclitaxel-induced apoptosis in human neuroblastoma SH-SY5Y cells. *Neurochemistry International* 2003;42:419–29. [PubMed: 12510025]
- Niles RM, Cook CP, Meadows GG, Fu Y-M, McLaughlin JL, Rankin GO. Resveratrol Is Rapidly Metabolized in Athymic (Nu/Nu) Mice and Does Not Inhibit Human Melanoma Xenograft Tumor Growth. 2006;136:2542–2546.
- Niles RM, McFarland M, Weimer MB, Redkar A, Fu Y-M, Meadows GG. Resveratrol is a potent inducer of apoptosis in human melanoma cells. *Cancer Letters* 2003;190:157–163. [PubMed: 12565170]
- Pozo-Guisado E, Lorenzo-Benayas MJ, Fernandez-Salguero PM. Resveratrol modulates the phosphoinositide 3-kinase pathway through an estrogen receptor alpha-dependent mechanism: relevance in cell proliferation. *International Journal of Cancer* 2004;109:167–73.
- Pozo-Guisado E, Merino JM, Mulero-Navarro S, Lorenzo-Benayas MJ, Centeno F, Alvarez-Barrientos A, Fernandez-Salguero PM. Resveratrol-induced apoptosis in MCF-7 human breast cancer cells involves a caspase-independent mechanism with downregulation of Bcl-2 and NF-kappaB. *International Journal of Cancer* 2005;115:74–84.
- Provinciali M, Re F, Donnini A, Orlando F, Bartozzi B, Di Stasio G, Smorlesi A. Effect of resveratrol on the development of spontaneous mammary tumors in HER-2/neu transgenic mice. *International Journal of Cancer* 2005;115:36–45.
- Reagan-Shaw S, Afaq F, Aziz MH, Ahmad N. Modulations of critical cell cycle regulatory events during chemoprevention of ultraviolet B-mediated responses by resveratrol in SKH-1 hairless mouse skin. *Oncogene* 2004;23:5151–60. [PubMed: 15122319]
- Regev-Shoshani G, Shoseyov O, Bilkis I, Kerem Z. Glycosylation of resveratrol protects it from enzymic oxidation. *The Biochemical Journal* 2003;374:157–163. [PubMed: 12697026]

- Revel A, Raanani H, Younglai E, et al. Resveratrol, a natural aryl hydrocarbon receptor antagonist, protects lung from DNA damage and apoptosis caused by benzo[a]pyrene. *Journal of Applied Toxicology* 2003;23:255–61. [PubMed: 12884409]
- Rigolio R, Miloso M, Nicolini G, Villa D, Scuteri A, Simone M, Tredici G. Resveratrol interference with the cell cycle protects human neuroblastoma SH-SY5Y cell from paclitaxel-induced apoptosis. *Neurochemistry International* 2005;46:205–11. [PubMed: 15670636]
- Riles WL, Erickson J, Nayyar S, Atten MJ, Attar BM, Holian O. Resveratrol engages selective apoptotic signals in gastric adenocarcinoma cells. *World Journal of Gastroenterology* 2006;12:5628–34. [PubMed: 17007014]
- Roman V, Billard C, Kern C, et al. Analysis of resveratrol-induced apoptosis in human B-cell chronic leukaemia. *British Journal of Haematology* 2002;117:842–51. [PubMed: 12060119]
- Russo J, Russo IH. Biological and molecular bases of mammary carcinogenesis. *Laboratory Investigation: A Journal of Technical Methods and Pathology* 1987;57:112–137. [PubMed: 3302534]
- Sala G, Minutolo F, Macchia M, Sacchi N, Ghidoni R. Resveratrol structure and ceramide-associated growth inhibition in prostate cancer cells. *Drugs Under Experimental and Clinical Research* 2003;29:263–9. [PubMed: 15134383]
- Sale S, Tunstall RG, Ruparella KC, Potter GA, Steward WP, Gescher AJ. Comparison of the effects of the chemopreventive agent resveratrol and its synthetic analog trans 3,4,5,4'-tetramethoxystilbene (DMU-212) on adenoma development in the Apc(Min+) mouse and cyclooxygenase-2 in human-derived colon cancer cells. *International Journal of Cancer* 2005;115:194–201.
- Sato M, Pei RJ, Yuri T, Danbara N, Nakane Y, Tsubura A. Prepubertal resveratrol exposure accelerates N-methyl-N-nitrosourea-induced mammary carcinoma in female Sprague-Dawley rats. *Cancer Letters* 2003;202:137–45. [PubMed: 14643443]
- Scarlatti F, Sala G, Somenzi G, Signorelli P, Sacchi N, Ghidoni R. Resveratrol induces growth inhibition and apoptosis in metastatic breast cancer cells via de novo ceramide signaling. *03-0292fje*. 2003
- Schneider Y, Duranton B, Gosse F, Schleiffer R, Seiler N, Raul F. Resveratrol inhibits intestinal tumorigenesis and modulates host-defense-related gene expression in an animal model of human familial adenomatous polyposis. *Nutrition and Cancer* 2001;39:102–7. [PubMed: 11588890]
- Sengottuvelan M, Viswanathan P, Nalini N. Chemopreventive effect of trans-resveratrol--a phytoalexin against colonic aberrant crypt foci and cell proliferation in 1,2-dimethylhydrazine induced colon carcinogenesis. *Carcinogenesis* 2006;27:1038–46. [PubMed: 16338953]
- Serrero G, Lu R. Effect of Resveratrol on the Expression of Autocrine Growth Modulators in Human Breast Cancer Cells. 2001;3:969–979.
- Seve M, Chimienti F, Devergnas S, et al. Resveratrol enhances UVA-induced DNA damage in HaCaT human keratinocytes. *Medicinal Chemistry (Shariqah, United Arab Emirates)* 2005;1:629–33.
- Sgambato A, Ardito R, Faraglia B, Boninsegna A, Wolf FI, Cittadini A. Resveratrol, a natural phenolic compound, inhibits cell proliferation and prevents oxidative DNA damage. *Mutation Research/ Genetic Toxicology and Environmental Mutagenesis* 2001;496:171–180.
- Soleas GJ, Grass L, Josephy PD, Goldberg DM, Diamandis EP. A comparison of the anticarcinogenic properties of four red wine polyphenols. *Clinical Biochemistry* 2002;35:119–124. [PubMed: 11983346]
- Soleas GJ, Yan J, Goldberg DM. Ultrasensitive assay for three polyphenols (catechin, quercetin and resveratrol) and their conjugates in biological fluids utilizing gas chromatography with mass selective detection. *Journal of Chromatography B: Biomedical Sciences and Applications* 2001;757:161–172.
- Su JL, Lin MT, Hong CC, et al. Resveratrol induces FasL-related apoptosis through Cdc42 activation of ASK1/JNK-dependent signaling pathway in human leukemia HL-60 cells. *Carcinogenesis* 2005;26:1–10. [PubMed: 15217905]
- Surh YJ, Hurh YJ, Kang JY, Lee E, Kong G, Lee SJ. Resveratrol, an antioxidant present in red wine, induces apoptosis in human promyelocytic leukemia (HL-60) cells. *Cancer Letters* 1999;140:1–10. [PubMed: 10403535]
- Tessitore L, Davit A, Sarotto I, Caderni G. Resveratrol depresses the growth of colorectal aberrant crypt foci by affecting bax and p21(CIP) expression. *Carcinogenesis* 2000;21:1619–22. [PubMed: 10910967]

- Vitrac X, Desmouliere A, Brouillaud B, et al. Distribution of [14C]-trans-resveratrol, a cancer chemopreventive polyphenol, in mouse tissues after oral administration. *Life Sciences* 2003;72:2219–33. [PubMed: 12628442]
- Walle T, Hsieh F, DeLegge MH, Oatis JE Jr, Walle UK. High Absorption buy Very Low Bioavailability of Oral Resveratrol in Humans. 2004;32:1377–1382.
- Whitsett T, Carpenter M, Lamartiniere C. Resveratrol, but not EGCG, in the diet suppresses DMBA-induced mammary cancer in rats. 2006;5:15.
- Wu SL, Sun ZJ, Yu L, Meng KW, Qin XL, Pan CE. Effect of resveratrol and in combination with 5-FU on murine liver cancer. *World Journal of Gastroenterology* 2004;10:3048–52. [PubMed: 15378791]
- Yance DR Jr, Sagar SM. Targeting Angiogenesis With Integrative Cancer Therapies. 2006;5:9–29.
- Yu L, Sun ZJ, Wu SL, Pan CE. Effect of resveratrol on cell cycle proteins in murine transplantable liver cancer. *World Journal of Gastroenterology* 2003;9:2341–3. [PubMed: 14562407]
- Zhou HB, Chen JJ, Wang WX, Cai JT, Du Q. Anticancer activity of resveratrol on implanted human primary gastric carcinoma cells in nude mice. *World Journal of Gastroenterology* 2005;11:280–4. [PubMed: 15633232]
- Zhou HB, Yan Y, Sun YN, Zhu JR. Resveratrol induces apoptosis in human esophageal carcinoma cells. *World Journal of Gastroenterology* 2003;9:408–11. [PubMed: 12632486]
- Ziegler CC, Rainwater L, Whelan J, McEntee MF. Dietary resveratrol does not affect intestinal tumorigenesis in Apc(Min/+) mice. *The Journal of Nutrition* 2004;134:5–10. [PubMed: 14704285]

Abbreviations

NF-κB	Nuclear factor kappa B
NMSC	non-melanoma skin cancer
COX-2	cyclooxygenase-2
ODC	ornithine decarboxylase
DMBA	9,10-dimethylbenz[<i>a</i>]anthracene
TPA	12- <i>O</i> -tetradecanoylphorbol-13-acetate
UVB	ultraviolet B
UVA	ultraviolet A
SCC	squamous cell carcinoma
i.m	intramuscular
EGCG	(-)-Epigallocatechin gallate
ER	estrogen receptor

TGF-α	transforming growth factor-alpha
PI3K	Phosphoinositide-3 kinase
i.p	intraperitoneal
TEBs	terminal end buds
PKC	protein kinase C
MNU	N-methyl-N-nitrosourea
AOM	azoxymethane
DMH	1,2-dimethylhydrazine
ACF	aberrant crypt foci
ROS	reactive oxygen species
NDEA	N-nitrosodiethylamine
5-FU	5-Fluorouracil
BaP	benzo[a]pyrene
LLC	Lewis lung carcinoma
HUVEC	Human Umbilical Vein Endothelial cells
NNK	4-(methyl-nitrosamino)-1-(3-pyridyl)-1-butanone