

Evolution of Chemotactic-Signal Transducers in Enteric Bacteria

MICHAEL K. DAHL,¹ WINFRIED BOOS,¹ AND MICHAEL D. MANSON^{2*}

Universitaet Konstanz, Department of Biology, D-7750 Konstanz, Federal Republic of Germany,¹ and Department of Biology, Texas A&M University, College Station, Texas 77483²

Received 4 November 1988/Accepted 2 February 1989

The methyl-accepting chemotactic-signal transducers of the enteric bacteria are transmembrane proteins that consist of a periplasmic receptor domain and a cytoplasmic signaling domain. To study their evolution, transducer genes from *Enterobacter aerogenes* and *Klebsiella pneumoniae* were compared with transducer genes from *Escherichia coli* and *Salmonella typhimurium*. There are at least two functional transducer genes in the nonmotile species *K. pneumoniae*, one of which complements the defect in serine taxis of an *E. coli* *tsr* mutant. The *tse* (taxis to serine) gene of *E. aerogenes* also complements an *E. coli* *tsr* mutant; the *tas* (taxis to aspartate) gene of *E. aerogenes* complements the defect in aspartate taxis, but not the defect in maltose taxis, of an *E. coli* *tar* mutant. The sequence was determined for 5 kilobases of *E. aerogenes* DNA containing a 3' fragment of the *cheA* gene, *cheW*, *tse*, *tas*, and a 5' fragment of the *cheR* gene. The *tse* and *tas* genes are in one operon, unlike *tsr* and *tar*. The cytoplasmic domains of Tse and Tas are very similar to those of *E. coli* and *S. typhimurium* transducers. The periplasmic domain of Tse is homologous to that of Tsr, but Tas and Tar are much less similar in this region. However, several short sequences are conserved in the periplasmic domains of Tsr, Tar, Tse, and Tas but not of Tap and Trg, transducers that do not bind amino acids. These conserved regions include residues implicated in amino-acid binding.

The methyl-accepting chemotaxis proteins are the best-characterized signal transducers of the chemosensory apparatus in *Escherichia coli* and *Salmonella typhimurium*. Homologous proteins are found in a wide range of bacterial species (35).

Four transducers have been characterized in *E. coli*, and the genes encoding them have been sequenced (5, 6, 24). The Tsr transducer (taxis to serine and away from some repellents) is the receptor for the attractant L-serine and related amino acids and is responsible for chemotaxis away from a wide range of repellents, including leucine, indole, and weak acids (41, 44). Tsr also is involved in thermotaxis (26). The Tar transducer (taxis to aspartate and away from some repellents) is the receptor for L-aspartate and related amino and dicarboxylic acids; Tar also mediates taxis to the attractant maltose via an interaction with the periplasmic maltose-binding protein (14) and taxis away from the repellents Co²⁺ and Ni²⁺ (41). The Trg transducer (taxis to ribose and galactose) interacts with the periplasmic ribose- and galactose-binding proteins to accomplish chemotaxis toward those two sugars (15, 21). The Tap transducer (taxis-associated protein) (7) mediates taxis toward dipeptides via an interaction with the periplasmic dipeptide-binding protein (28; V. Blank, diploma thesis, University of Konstanz, Federal Republic of Germany, 1987).

All transducers characterized to date are composed of the same structural domains (24). The amino terminus resembles the signal peptide of exported proteins, but it is not removed from the mature protein and serves as the first membrane-spanning region. The next ca. 160 amino acids extend into the periplasm, where they form the receptor domain (23). After a second membrane-spanning region, which seems to act as a stop transfer signal (27), the final 300 or more amino acids are localized in the cytoplasm. This portion of the protein generates the intracellular signals to the flagella (32, 36) and contains the glutamic acid residues that are methyl-

ated and demethylated during adaptation to chemotactic stimuli (19, 42). The amino acid sequence in regions containing the methylation sites and signaling domain is highly conserved among the four transducers (5, 24).

The genes encoding the signal transducers are located at different points on the *E. coli* chromosome (1); *tsr* at 99 min, *tar* and *tap* together in the *meche* operon at 41 min, and *trg* at 31 min. The genes are part of the flagellar regulon (20), and their promoters contain the consensus sequence present in flagellar operons that require an alternate σ factor for expression (2, 16, 17).

The *tar* gene of *S. typhimurium* also has been sequenced (37). *S. typhimurium* Tar mediates aspartate but not maltose taxis (9, 31). *S. typhimurium* also lacks the Tap transducer and therefore shows no taxis toward dipeptides (28). *S. typhimurium* does respond to serine, ribose, and galactose, so it probably has transducers equivalent to Tsr and Trg. *Enterobacter aerogenes* and *Klebsiella pneumoniae* are somewhat more distantly related to *E. coli* and *S. typhimurium* than the last two species are to each other (38). *E. aerogenes* exhibits chemotaxis to aspartate, maltose, and serine (9; M. Dahl, diploma thesis, University of Konstanz, Federal Republic of Germany, 1985), whereas *K. pneumoniae* is nonmotile.

This work was initiated on the premise that comparison of the transducers from the enteric bacteria will contribute to the study of these proteins. Genes encoding functional transducers were identified in phage λ libraries of chromosomal DNA from *E. aerogenes* and *K. pneumoniae*. We sequenced 5 kilobases (kb) of DNA that encodes serine and aspartate transducers and adjacent chemotaxis-related genes. Analysis of this DNA sequence and of the predicted amino acid sequences derived from it provides information about the function and evolution of proteins involved in chemosensing.

MATERIALS AND METHODS

Reagents. Restriction endonucleases, T4-DNA ligase, Klenow enzyme (DNA polymerase I large fragment), the kit for

* Corresponding author.

TABLE 1. Strains and plasmids

Strain	Known markers and properties	Source or reference
Bacterial strains		
<i>E. coli</i>		
LE392	F ⁻ <i>supF supE hsdR galK trpR metB tonA</i>	39
LE392.23	LE392 $\Delta(\textit{argF-lac})U169$	39
RM41	Q358 F ⁻ (r ⁻ m ⁺) <i>supE</i> $\phi 80^f$	18
RM42	Q359 P2 lysogen	18
TG1	<i>pro thi hsd</i> (r ⁻) <i>lacF'</i> <i>lacI^a</i> <i>lacZ</i> Δ M15	8
VB12	<i>ara-14</i> $\Delta(\textit{argF-lac})U169$ <i>his-4 leuB6 metF159_{am} mtl-1 rpsL136</i> $\Delta(\textit{tar-tap})5201$ <i>thi-1 tonA31</i> $\Delta\textit{tsr7021 xyl-15}$	28
<i>E. aerogenes</i> ATCC 13048	Wild type	13
<i>K. pneumoniae</i> KAY2026	Wild type	40
Phage		
M13mp10	M13 sequencing vector	29
M13mp18	M13 sequencing vector	34
M13mp19	M13 sequencing vector	34
λ gt4- <i>lac5</i>	<i>cI857</i> (Ts), <i>lacZ</i> ⁺ <i>lacY</i> ⁺ , helper phage for lysogenizing with λ SE6	10
λ SE6	Kan ^r , low-copy-number phasmid	11
λ SE6-M1	λ SE6 carrying <i>E. aerogenes</i> DNA containing ' <i>cheA</i> , <i>cheW</i> , <i>tse</i> , <i>tas</i> , and <i>cheR</i> '	This study
Plasmids		
pBR322	Ap ^r Tc ^r	4
pJFG5	pBR322 carrying <i>E. coli tsr</i>	12
pRK41	pBR322 with 2.4-kb <i>Clal</i> fragment from <i>S. typhimurium</i> carrying <i>tar</i> , pBR322 with 3.3-kb <i>EcoRI-AvaI</i> fragment with <i>E. coli</i> ' <i>cheA</i> , <i>cheW</i> , <i>tar</i> , and <i>tap</i> '	37
pMK1	pBR322 with 5.0-kb <i>NruI</i> fragment from λ SE6-M1 carrying ' <i>cheA</i> , <i>cheW</i> , <i>tse</i> , <i>tas</i> , and <i>cheR</i> ' from <i>E. aerogenes</i>	22
pMD5	pBR322 with the 3.5-kb <i>NdeI</i> fragment from pRK41 and the 1.8-kb <i>NdeI</i> fragment from pMK1	This study
pMD6	pBR322 with the 3.2-kb <i>NdeI</i> fragment from pRK41 and the 4.5-kb <i>NdeI</i> fragment from pMK1	This study
pMD7	pBR322 with the 2.3-kb <i>EcoRI-StuI</i> fragment from pMD5 carrying ' <i>cheW</i> , <i>tse</i> , and <i>tas</i> '	This study
pMD8		

in vitro packaging of phage λ DNA, and reagents for the M13 dideoxy nucleotide sequencing system were purchased from Boehringer Mannheim Biochemicals, Indianapolis, Ind. Isopropyl-1-thio- β -D-galactopyranoside was purchased from Serva. The *lacZ* oligonucleotide primer (17-mer) was obtained from Bethesda Research Laboratories, Inc., Gaithersburg, Md., or Boehringer. Deoxy-adenosine-5'- α -[³⁵S]thiotriphosphate was purchased from Amersham Corp., Arlington Heights, Illinois. All other chemicals were of reagent grade.

Strains and plasmids. The bacterial and phage strains and plasmids used in this work are listed in Table 1. *E. coli* VB12 ($\Delta\textit{tsr7021}$ $\Delta\textit{tar-tap5201}$) is defective in serine, aspartate, maltose, and dipeptide taxis because of the deletion of the three transducer genes. This strain does not form normal swarms in tryptone soft agar (41) because its steady-state run-tumble behavior is strongly skewed toward running due to the loss of signal input from the missing transducers.

Media and swarm plates. Cells were grown in Luria broth (LB) or minimal medium A supplemented with 0.2% (wt/vol) carbon source, required amino acids at 100 μ g/ml, and ampicillin or kanamycin at 50 μ g/ml as needed. Media were prepared by the method of Miller (30). Tryptone swarm plates contained 0.3% Bacto-Agar (Difco Laboratories, Detroit, Mich.), 1% Bacto-Tryptone (Difco), and 0.8% NaCl. Minimal swarm plates contained 0.3% Bacto-Agar, motility salts [10 mM potassium phosphate (pH 7.0), 1 mM

(NH₄)₂SO₄, 1 mM MgSO₄, 0.5% (wt/vol) NaCl, and 0.5 μ g of FeCl₃ per ml], and 100 μ M maltose, L-aspartate, or L-serine. Aspartate and serine swarm plates also contained 1 mM glycerol. Ampicillin or kanamycin was added to swarm plates to a final concentration of 50 μ g/ml. The plates were scored after 12 to 20 h of incubation at 32°C.

Cloning of genes coding for chemotactic-signal transducers from *E. aerogenes* and *K. pneumoniae*. Chromosomal DNA from *E. aerogenes* ATCC 13048 and *K. pneumoniae* KAY2026 was prepared as described previously (39). Genomic libraries were constructed in phage λ SE6 by the method of Elledge and Walker (11), using the *E. coli* strains RM41 and RM42. These libraries were used to infect strain VB12 ($\Delta\textit{tsr}$ $\Delta\textit{tar-tap}$) lysogenized with the helper phage λ gt4-*lac5*, and a 100- μ l sample of washed, infected cells (ca. 5×10^8 cells) was applied in a 5-cm-long trough on swarm plates containing kanamycin. With the *E. aerogenes* library, minimal-aspartate swarm plates were used. With the *K. pneumoniae* library, tryptone swarm plates were used. Within 24 h of incubation at 32°C in minimal agar or 16 h in tryptone agar, chemotactic swarms spreading away from the trough appeared on some of the plates. Single colonies were isolated from these swarms on LB-kanamycin plates, and phage lysates were prepared from well-aerated cultures of these cells growing in LB medium. Phage production was induced by a 15-min shift from 32°C to 42°C, and lysis of the cells proceeded during continued incubation of the cultures




FIG. 1. Construction of *E. coli*-*S. typhimurium* hybrid *tar* genes. Plasmid pMD6 was made by ligating the 3.5-kb *Nde*I fragment of pRK41 that contains the 5' end of *S. typhimurium tar* with the 1.8-kb *Nde*I fragment from plasmid pMK1 that contains the 3' end of *E. coli tar*. Plasmid pMD7 was made by ligating the 4.5-kb *Nde*I fragment of pMK1 that contains the 5' end of *E. coli tar* with the 3.2-kb *Nde*I fragment from pRK41 that contains the 3' end of *S. typhimurium tar*. *E. coli* DNA is indicated by black bars, *S. typhimurium* DNA is indicated by hatched bars, and pBR322 DNA is indicated by white bars. The genes carried by each plasmid are indicated below the corresponding restriction map.

at 38°C. The λ SE6 and helper phage were separated by plating dilutions of these lysates with *E. coli* LE392.23 on agar containing 5-bromo-4-chloro-3-indolyl- β -D-galactoside; λ SE6 forms white plaques under these conditions, whereas λ gt4-*lac5* forms blue plaques.

From the *E. aerogenes* library, 10 independently isolated, plaque-purified phages were retested to confirm that they conferred an aspartate-chemotaxis-positive phenotype upon infecting strain VB12. One of these phages, λ SE6-M1, was chosen for further analysis. From the *K. pneumoniae* library, 20 plaque-purified isolates were tested for their ability to complement mutations in *E. coli* transducer genes. Phage λ lysates and DNA were prepared by the methods of Silhavy et al. (39).

Determination of nucleotide sequence. DNA sequencing was performed by the modified dideoxy-chain termination method of Biggin et al. (3) with deoxyadenosine-5'- α -[35 S]thiotriphosphate. DNA restriction fragments for sequencing were produced by digesting the purified 5-kb *Nru*I fragment of plasmid pMD5 with the enzymes *Bam*HI, *Cla*I, *Eco*RI, *Eco*RV, *Hae*III, *Hinc*II, *Hind*III, *Pvu*II, *Sau*3A, *Sma*I, and *Stu*I. These fragments were ligated into the polycloning sites of DNA from phage M13mp10, M13mp18, and M13mp19 digested with appropriate enzymes. M13 phage were propagated on *E. coli* TG1.

Computer analysis. The management of sequence information and the subsequent analysis of compiled data were done with the Macintosh DNA Inspector II program (Microsoft




FIG. 2. Restriction map of the plasmids pMD5 and pMD8. The pBR322 DNA is indicated by the white bar, and *E. aerogenes* DNA is shown as a black bar. The genes present on the plasmids are shown between the two restriction maps.

Basic Runtime, version 2.11; Microsoft Corp.). The DNA Translate program (Mac 1.25) was kindly provided by J. S. Parkinson (Biology Department, University of Utah).

RESULTS

Maltose taxis in cells making hybrid Tar proteins. The difference in the abilities of *E. coli* and *S. typhimurium* to sense maltose is logically attributed to differences in the periplasmic domains of Tar from the two species (9, 31), but this suspicion had not been confirmed experimentally. We decided to utilize chimeric transducers, which were previously used to localize functional domains within the proteins (23), to test this inference.

Construction of the hybrid genes (Fig. 1) took advantage of the *NdeI* site in the *tar* genes from *E. coli* and *S. typhimurium*. Plasmid pMD6 codes for a protein containing the N-terminal 256 residues from *S. typhimurium* Tar and the C-terminal 297 residues from *E. coli* Tar, whereas the protein coded by plasmid pMD7 has the N-terminal 256 residues from *E. coli* Tar and the C-terminal 296 residues from *S. typhimurium*. These two plasmids were transformed into *E. coli* VB12 ($\Delta tsr7021 \Delta tar-tap5201$), and the transformants were tested on minimal-aspartate and minimal-maltose swarm plates containing ampicillin.

Cells containing pMD6 formed normal swarm rings only with aspartate, whereas strains containing pMD7 formed swarm rings on both types of plates. Thus, the inability of *S. typhimurium* to respond to maltose is a property of the first 256 amino acids of Tar, which encompass the periplasmic domain.

Transducer genes from *K. pneumoniae*. The differences in transducer identity and function between *E. coli* and *S. typhimurium* raise a question about the transducer complement of other enteric bacteria. *K. pneumoniae* presents an especially intriguing case, since cells of this species are nonmotile. Twenty phage isolated from a *K. pneumoniae* genomic library restored the ability of *E. coli* strain VB12 to form swarm rings in tryptone soft agar. Lysogens of strain VB12 containing these phage were then screened on serine, aspartate, and maltose swarm plates (data not shown). Five




FIG. 3. Chemotactic swarms formed by strain VB12 containing different plasmids. The plasmid in each strain is indicated by the number next to the swarm as follows: 1, pBR322; 2, pMK1; 3, pRK41; 4, pJFG5; 5, pMD5; 6, pMD8. The swarm plates used were tryptone (TB), minimal-serine (Ser), minimal-aspartate (Asp), and minimal-maltose (Mal). All plates contained 50 μ g of ampicillin per ml.

phage presumably carry a serine transducer gene, since they confer the ability to form swarm rings in serine soft agar but not in aspartate or maltose soft agar. Fifteen phage did not restore swarm rings with these three attractants. These phage must carry genes for transducers mediating responses to other compounds present in the tryptone extract.

Transducer genes from *E. aerogenes*. Unlike *K. pneumoniae*, *E. aerogenes* is motile. It also responds to many of the same attractants as *E. coli*, suggesting that these two species should have at least some similar transducers. To compare the structure and function of a related transducer from the two bacteria, we set out to clone the *tar* gene equivalent from *E. aerogenes*.

Phage λ SE6-M1, isolated from the *E. aerogenes* genomic library, is able to complement the defect in aspartate taxis of strain VB12. DNA prepared from this phage was digested with *Nru*I, and a 5-kb fragment was ligated into the *Nru*I site of plasmid pBR322 to yield pMD5 (Fig. 2). This plasmid and the reference plasmids pBR322, pJFG5 (*E. coli tsr*⁺), pMK1 (*E. coli tar*⁺), and pRK41 (*S. typhimurium tar*⁺) were transformed into *E. coli* VB12, and the chemotactic phenotypes of the transformants were tested on swarm plates (Fig. 3). Plasmid pMD5 restored the ability to form swarm rings with both aspartate and serine, but not with maltose. None of the plasmids restored the ability to form swarm rings with L-prolyl-L-leucine, a good attractant for dipeptide chemotaxis (data not shown).

DNA sequence analysis (described below) revealed that the 5-kb *Nru*I fragment in pMD5 contained two genes that encode transducers. One of these genes, including its promoter, was located on a 2.3-kb *Eco*RI-*Stu*I restriction fragment that was inserted into the *Eco*RI and *Nru*I sites of pBR322 to generate pMD8 (Fig. 2). This plasmid was transformed into strain VB12, where it restored the ability to form swarm rings with serine but not with aspartate (Fig. 3). The transducer gene on plasmid pMD8 was named *tse* (taxis to serine), because it complemented the defect in serine taxis caused by a *tsr* mutation. The transducer gene present on pMD5 but not on pMD8 was named *tas* (taxis to aspartate) because it was required to complement the defect in aspartate taxis caused by a *tar* mutation.

Nucleotide sequence of chemotaxis-related genes from *E. aerogenes*. The nucleotide sequence of the 5-kb *Nru*I restriction fragment from plasmid pMD5 was determined from both strands between two and five times, and the amino acid sequences of the five longest open reading frames were predicted (Fig. 4). Based on the similarity of these sequences to their homologs in *E. coli* (33), three of the open reading frames correspond to *cheW* and fragments of the '*cheA* and *cheR*' genes. The open reading frames corresponding to *tse* and *tas* were identified by their resemblance to transducer genes from *E. coli* and *S. typhimurium*.

Comparison of Che proteins from *E. aerogenes* and *E. coli*. The open reading frame for the 3'-terminal portion of *cheA* from *E. aerogenes* (Fig. 4) codes for 226 amino acids, of which 176 (78%) are identical to residues within the last 226 amino acids of CheA from *E. coli* (33). One stretch of 19 amino acids has only one match, but we do not believe our sequence is in error because neither alternate reading frame in this region gives a better match. Without these 19 residues the percent identity rises to 85% (175 of 207 amino acids).

The 5' region of *cheR* in our sequence contains an open reading frame coding for 100 amino acids starting with the ATG codon at nucleotides 4715 through 4717 (Fig. 4). This codon is preceded by the sequence GAGCT at nucleotides 4707 through 4711, which may serve as a ribosome-binding

site. (The C does not match the Shine-Dalgarno consensus sequence.) Alternatively, translation could start at the ATG codon at nucleotides 4769 through 4771. Significant homology between the predicted CheR amino acid sequences of *E. aerogenes* and *E. coli* begins at the glutamine preceding the second methionine residue. Of the next 82 residues, 68 (83%) are identical in the two species. The sequences preceding this glutamine bear little similarity in the two species.

The predicted amino acid sequence of the CheW protein of *E. aerogenes* is 165 residues long and begins with the methionine coded by the ATG triplet at nucleotides 715 through 717 (Fig. 4). This protein is 77% identical to CheW from *E. coli* and 79% identical to CheW from *S. typhimurium* (Fig. 5). The CheW proteins of the latter two species are 92% identical.

Comparison of Tse and Tas with signal transducers from *E. coli* and *S. typhimurium*. The predicted amino acid sequences of Tse and Tas can be aligned with the sequences of four other signal transducers (Fig. 6). The amino acid sequences of the periplasmic domains of the six transducers are compared in Table 2. Tse and Tsr are more like each other than any of the other transducers, being almost as similar as the *E. coli* and *S. typhimurium* Tar proteins. Surprisingly, the periplasmic portion of Tas has a low percent identity with all of the other transducers, exhibiting similarities only in a few, limited regions. The periplasmic portion of Tap diverges even more widely.

DISCUSSION

The properties of *E. coli*-*S. typhimurium* Tar hybrids suggest that differences in the N-terminal halves of the two proteins (presumably in their periplasmic receptor domains) determine whether they function in maltose taxis. Since *E. aerogenes* carries out both aspartate and maltose taxis (19), the periplasmic domain of its Tar transducer might share structural features with *E. coli* Tar that contribute to its interaction with maltose-binding protein. These features could be recognizable as amino acid sequences present in *E. coli* and *E. aerogenes* Tar but not in *S. typhimurium* Tar. Our interest in making this comparison led us to isolate transducer genes from *E. aerogenes*. As the work progressed, the broader goal emerged of using a comparative approach to study the evolution of the signal transducers.

The nonmotile bacterium *K. pneumoniae* contains at least two transducer genes that can function in *E. coli*. One of them codes for a serine transducer. No defined substrates have been identified for the other *K. pneumoniae* transducers, which overcome the general chemotactic defect of strain VB12.

Why have transducer genes been retained in a nonmotile species? In *E. coli*, the transducer genes are expressed late in the regulatory cascade of the flagellar regulon (2, 20), and their expression is blocked by *fla* mutations that inhibit flagellar synthesis. If comparable mutations have occurred in *K. pneumoniae*, then the transducer genes would not be expressed in that species, and the selective pressure to delete them would be relatively low.

Two transducer genes were found on a 5-kb DNA fragment from *E. aerogenes*. The *tse* gene encodes a serine transducer, since it complements an *E. coli tsr* mutation. The *tas* gene is required to restore aspartate taxis in an *E. coli tar* mutant. Since we do not have a construct in which *tas* is expressed in the absence of *tse*, we have not demonstrated that Tas alone functions as an aspartate transducer, although that is the simplest assumption.

cheA¹ 27 54 81 108
 GAA AAA ATT CTC GCC AAA GCG GCG GCG CAG GGG CTG GCG GTC ACG GAC ACG ATG AGT GAT GAA GAG GTC GGA ATG CTT ATT TTT GCG CCG GGC TTT TCA ACC GCG GAA
 Glu Lys Ile Leu Ala Lys Ala Ala Ala Gln Gly Leu Ala Val Thr Asp Thr Met Ser Asp Glu Glu Val Gly Met Leu Ile Phe Ala Pro Gly Phe Ser Thr Ala Glu
 135 162 189 216
 CAG GTG ACC GAC GTC TCT GCG GCG GGC GTC GGC ATG GAC GTC GTC AAA CCG AAT ATT CAG GAG ATG GGC GGT CAC GTA GAA ATC CAT TCC CGT GCG GGC AAA GGG ACC
 Gln Val Thr Asp Val Ser Gly Arg Gly Val Gly Met Asp Val Val Lys Arg Asn Ile Gln Glu Met Gly Gly His Val Glu Ile His Ser Arg Ala Gly Lys Gly Thr
 243 270 297 324
 TCG ATT CGT ATT TTG TTG CCG CTA ACG CTC GCC ATC CTC GAC GGC ATG TCG GTC AAG GTC AAT GAA GAG GTC TTT ATT CTG CCG CTC AAC GCG GTG ATG GAA TCG CTG
 Ser Ile Arg Ile Leu Leu Pro Leu Thr Leu Ala Ile Leu Asp Gly Met Ser Val Lys Val Asn Glu Glu Val Phe Ile Leu Pro Leu Asn Ala Val Met Glu Ser Leu
 351 378 405 432
 CAG CCG CAG CCG GAA GAC CTG CAT CCA ATG GCC GCG GCG GAG CCG ATG CTG CAG GTT CCG GGC GAG TAT CTA CCG CTG GTG GAG CTC TAC CCG GTG TTT GAA TGT GCC
 Gln Pro Gln Arg Glu Asp Leu His Pro Met Ala Gly Gly Glu Arg Met Leu Gln Val Arg Gly Glu Tyr Leu Pro Leu Val Glu Leu Tyr Arg Val Phe Glu Cys Ala
 459 486 513 540
 GGG GCG AAA ACC GAG GCC ACT CAG GGC ATC ATG GTG ATT CTG CAA AGC GCC GGC CCG CGT AAT GCG CTG CTG GTG GAT CAA CTG ATC GGC CAG CAC CAG TGT GTG AAA
 Gly Ala Lys Thr Glu Ala Thr Gln Gly Ile Val Val Ile Leu Gln Ser Ala Gly Arg Arg Asn Ala Leu Leu Val Asp Gln Leu Ile Gly Gln His Gln Cys Val Lys
 567 594 621 648
 AAC CTG GAA ACG AAT TAC CCG AAA GTG CTG CCG GGA ATT TCC GCG GCG ACG ATC CTC GGC GAC GGC AGC GTG GCG CTG ATC GTC GAC GTG TCG CCG CTG CAA ATG CTC AAT
 Asn Leu Glu Thr Asn Tyr Arg Lys Val Pro Gly Ile Ser Ala Ala Thr Ile Leu Gly Asp Gly Ser Val Ala Leu Ile Val Asp Val Ser Ala Leu Gln Met Leu Asn
 675 702 729 756
 CCG GAA AAG CTG CTG AGC GCA GCG GGC GCA TAA CGA CTC ATC TCA TCA AAT TAA CTG GTG CAG ACC ATG GCA GGA TTA GCA ACC GTC AGC AAA TTG GCT GGC GAA ACG
Arg Glu Lys Leu Leu Ser Ala Ala Ala * MET Ala Gly Leu Ala Thr Val Ser Lys Leu Ala Gly Glu Thr
 783 810 837 864
 GTA GGT CAG GCG TTT TTA ATC TTT ACC CTC GGC AAT GAA GAA TAC GGC ATC GAT ATC CTG AAA GTG CAG AAG ATC CCG GGC TAT GAC CAG GTG ACG CCG ATC GCC AAC
 Val Gly Gln Ala Phe Leu Ile Phe Thr Leu Gly Asn Glu Glu Tyr Gly Ile Asp Ile Leu Lys Val Gln Lys Ile Arg Gly Tyr Asp Gln Val Thr Arg Ile Ala Asn
 891 918 945 972
 ACC CCG GAT TTC ATC AAA GGC CTC ACC AAT CTG CCG GGG GTG ATC GTG CCG ATT ATC GAC CTG CCG GTA AAA TAT GCC CAG CAG GGC GTC TCT TAT GAT GAA AAC ACG
 Thr Pro Asp Phe Ile Lys Gly Val Thr Asn Leu Arg Gly Val Ile Val Pro Ile Ile Asp Leu Arg Val Lys Tyr Ala Gln Gln Gly Val Ser Tyr Asp Glu Asn Thr
 999 1026 1053 1080
 GTG GTT ATC GTG CTT AAC TTC GGC CAG CCG GTG GTG GGG ATT GTG GTC GAC GGG GTC TCC GAC GTG TTG TCT CTT ACC GCC GAA CAG ATC CCG CCG GCG GAA TTC
 Val Val Ile Val Leu Asn Phe Gly Gln Arg Val Val Gly Ile Val Val Asp Gly Val Ser Asp Val Leu Ser Leu Thr Ala Glu Gln Ile Arg Pro Ala Pro Glu Phe
 1107 1134 1161 1188
 GCA GTG ACG ATG GCG ACC GAA TAT CTC ACC GGT CTT GGC GCG CTC GGA GCG CTG TTG ATC CTT GTG GAT ATC GAA AAG CTG CTC AGC ACG GAA GAG ATG GCG CTG GTC
 Ala Val Thr Met Ala Thr Gly Tyr Leu Thr Gly Leu Gly Ala Leu Gly Ala Leu Leu Ile Leu Val Asp Ile Glu Lys Leu Leu Ser Thr Glu Glu Met Ala Leu Val
 1215 1242 1269 1296
 GAT AAC GTC GCC AAA AGC CAC TAA GCA ATC GGG CCG GCC GGT AAA AAT AGT CCC CCG CTG GCT AAA GTT CCC CTC CCG TAC GCG GAT AAC CCT TTC AGT CAC ATA CGT
 Asp Asn Val Ala Lys Ser His * flagellar operon consensus
 1324 1352 1379 1406
 AAA GCC TGG CCG TTC AGG TTC CAGGAGGGGA AAT ATG TTT AAT CGT ATT AAG GTC GTC ACC AGT CTG CTC TTA TTA GTG CTG GTG CTA TTT GGC GCA TTG CAG CTG ATT TCA
 SD MET Phe Asn Arg Ile Lys Val Val Thr Ser Leu Leu Leu Val Leu Val Leu Phe Gly Ala Leu Gln Leu Ile Ser
 1433 1460 1487 1514
 GGC GGT CTG TTT TTT TCG TCG CTG AAA GGC GAT AAA GAG AAC TTT ACC GTC CTG CAA ACC ATC CGT CAG CAG CAG TTG CTG CTG AGT GAA AGT CCG GTC GAT CTG CTG
 Gly Gly Leu Phe Phe Ser Ser Leu Lys Gly Asp Lys Glu Asn Phe Thr Val Leu Gln Thr Ile Arg Gln Gln Gln Leu Leu Leu Ser Glu Ser Arg Val Asp Leu Leu
 1541 1568 1595 1622
 CAG GCG CGT AAC TCC CTG AAC CCG GCA GGG ATC CCG TAC ATG ATG GAT ACC AAC AAA ATC GGC AGC GGC GCG ACT ATC GAC GAG CTG CTG GCG AAA GCG GAA AAA GAA
 Gln Ala Arg Asn Ser Leu Asn Arg Ala Gly Ile Arg Tyr Met Met Asp Thr Asn Lys Ile Gly Ser Gly Ala Thr Ile Asp Glu Leu Leu Ala Lys Ala Glu Lys Glu
 1649 1676 1703 1730
 AAG CTG GCG GCG GCC GAG CCG AAC TAC ACC GCC TAT GAA AAA ATC CCG CAG GAC CCG CGT CAG GAT CCT CAG GCG ACG GAA AAG CTT AAG CAG CAG TAT GGC ATC CTG
 Lys Leu Ala Arg Ala Glu Arg Asn Tyr Thr Ala Tyr Glu Lys Ile Pro Gln Asp Pro Arg Gln Asp Pro Gln Ala Thr Glu Lys Leu Lys Gln Gln Tyr Gly Ile Leu

FIG. 4. Nucleotide sequence of the '*cheA-cheW-tse-tas-cheR*' region from *E. aerogenes*. The predicted amino-acid sequences for the longest open reading frames are given below. The amino acids underlined in the '*cheA*' and '*cheR*' gene fragments are identical with residues in the corresponding genes of *E. coli*. The inverted triangles in '*cheA*' indicate single amino acid deletions relative to *E. coli*, and the bracket denotes an inserted amino acid. The underlined sequence designated "flagellar operon consensus" corresponds to the consensus sequence found in the promoters of flagellar operons in *E. coli* (2, 16). The underlined sequences designated SD refer to regions surrounding the predicted Shine-Dalgarno sequences for the *tse* and *tas* genes.

Since *E. aerogenes* responds chemotactically to maltose and synthesizes maltose-binding protein (9), it should have a maltose transducer. Our attempt to isolate this transducer was based on the assumption that it should also be an aspartate receptor. In *E. aerogenes*, however, maltose taxis could be mediated by a minor transducer (like Trg or Tap in *E. coli*) that is unable to restore normal chemotactic behavior to the smooth-swimming VB12 strain. Thus, a search for the gene encoding the maltose transducer of *E. aerogenes* should be repeated with a *tsr*⁺ *tar* *E. coli* strain, which shows normal swimming behavior but no aspartate or maltose taxis.

The genes on the 5-kb DNA fragment from *E. aerogenes* appear in the order 5' *cheA-cheW*-promoter-*tse-tas-cheR* 3' (Fig. 4). This segment spans the 3' and 5' ends of the *E. aerogenes* equivalents of the *E. coli* *mocha* and *meche* operons (33). In *E. coli* the first two genes in the *meche* operon are *tar* and *tap*, whereas in *S. typhimurium* *tar* is the sole transducer gene in the *meche* operon. The *che* genes are in the same relative location in all three species.

The 36 nucleotides between *cheA* and *cheW* in *E. aerogenes* show no obvious similarity to the corresponding 23-nucleotide regions in *E. coli* or *S. typhimurium*. The untranslated region between *cheW* and the start codon of the

first gene in the *meche* operon also varies considerably among the three species. This region contains 122 nucleotides in *E. aerogenes*, 147 nucleotides in *E. coli*, and 243 nucleotides in *S. typhimurium*. The proposed rho-independent site for transcription termination distal to *cheW* in *E. coli* is missing in *E. aerogenes*, as is the perfect 13-base inverted repeat found 3' to the tandem TAA stop codons of *cheW* in *S. typhimurium*.

The region (underlined in Fig. 4) around the Shine-Dalgarno sequence of the first gene in the *meche* operon is conserved in all three species. So is the consensus sequence (also underlined in Fig. 4) identified in promoters of flagellar operons of *E. coli* (2, 16). Conservation of these regulatory sequences is consistent with the observation that the *tse* and *tas* genes are expressed in *E. coli*. No significant homology was detected between the *tse-tas* (85 base pairs) and *tar-tap* (48 base pairs) or the *tas-cheR* (91 base pairs) and *tap-cheR* (16 base pairs) intergenic regions.

Some regions of the predicted amino acid sequences of the '*cheA*' and '*cheR*' polypeptide fragments of *E. aerogenes* are more similar to their counterparts in *E. coli* than others (Fig. 4). The highly conserved sequences presumably represent regions of special structural and functional importance. The CheW proteins from *E. aerogenes*, *E. coli*, and *S. typhimu-*

1757 1784 1811 1838
TAC GGC GCG CTG TCG GAG CTG ATC CAG CTG CTG GGC GGA GGG TGT AAA ATC AAC GCC TTC TTC GAC CAG CCG ACG CAA AAA TAC CAG GAC GAT TTC GAG CAG ACC TAT
Tyr Gly Ala Leu Ser Glu Leu Ile Gln Leu Leu Gly Gly Gly Cys Lys Ala Val Asp Ala Ser Asn Ser Ser Tyr Ser Ser Ala Ile Trp Thr Leu Ile Val Val Ile Ile Val Val

1865 1892 1919 1946
AAC GCC TAT CTG CAG CAA AAC GGC AAG CTG TAT CAA ATC GCC GTT GAC GCT AGC AAC AGC TCG TAT AGC TCG GCG ATC TGG ACC TTA ATT GTG GTC ATC ATC GTG GTG
Asn Ala Tyr Leu Gln Gln Asn Gly Lys Leu Tyr Gln Ile Ala Val Asp Ala Ser Asn Ser Ser Tyr Ser Ser Ala Ile Trp Thr Leu Ile Val Val Ile Ile Val Val

1973 2000 2027 2054
CTG GCG GCG ATC GTC GGC GTG TGG ATG GGT ATC CAC CAT ATC CTG GTG GCG CCG CTG AAC CGC ATG ATA GAA CAC ATC AAA CCG ATC GCG TCC GGC GAT CTG ACC CAG
Leu Ala Ala Ile Val Gly Val Trp Met Gly Ile His His Ile Leu Val Arg Pro Leu Asn Arg Met Ile Glu His Ile Lys Arg Ile Ala Ser Gly Asp Leu Thr Gln

2081 2108 2135 2162
CCT ATT CCG GTG ACC AGC CGT AAT GAA ATC GGC GTG CTG GCC GCC AGC CTC AAG CAC ATG CAG AAC GAG CTT ATC GAA ACG GTA AGC GGC GTG CGT CAG GGC GCG GAT
Pro Ile Pro Val Thr Ser Arg Asn Glu Ile Gly Val Leu Ala Ala Ser Leu Lys His Met Gln Asn Glu Leu Ile Glu Thr Val Ser Gly Val Arg Gln Gly Ala Asp

2189 2216 2243 2270
CGA TCT ATT GAC GGC GCA TCG GAA AGC GCC GCG GGC AAT AAC GAT CTC TCT TCG GCG ACC GAA CAG CAG GCG GCA CCG CTG GAA GAG ACC GCC GCC AGT ATG GAA CAG
Arg Ser Ile Asp Gly Ala Ser Glu Ser Ala Ala Gly Asn Asn Asp Leu Ser Ser Ser Arg Thr Glu Gln Gln Ala Ala Ala Leu Glu Glu Thr Ala Ala Ser Met Glu Gln

2297 2324 2351 2378
CTG ACA GCG ACG GTG AAA CAG AAC GCC GAA AAC GCG CGT CAG GCC AGC CAG CTG GCG CTG AGC GCC TCC GAA ACT GCG CAG AAG GGC GGC AAA GTG GTG GCT AAC GTG
Leu Thr Ala Thr Val Lys Gln Asn Ala Glu Asn Ala Arg Gln Ala Ser Gln Leu Ala Leu Ser Ala Ser Glu Thr Ala Gln Lys Gly Gly Lys Val Val Ala Asn Val

2405 2432 2459 2486
GTC GAA ACC ATG CAC GAC ATC GCC AGC AGC TCG CAG AAA ATT GCC ATC ACC GGC GTT ATC GAC GTA ATC GCC TTC CAG ACC AAC ATC CTG GCG CTT AAC GCC GGC
Val Glu Thr Met His Cys Asp Ile Ala Ser Ser Ser Gln Lys Ile Ala Asp Ile Thr Gly Val Ile Asp Val Ile Ala Phe Gln Thr Ser Ala Leu Ala Leu Asn Ala Ala

2513 2540 2567 2594
GTA GAA GCG GCG CGT GCG GGT GAA CAA GGT CCG GCG TTC GCG GTC GTG GCG GGC GAA GTC CCG AAC CTG GCG CAG CCG AGC GCC CAG GCG GCC AAA GAG ATC AAA CCG
Val Glu Ala Ala Arg Ala Gly Glu Gln Gly Arg Ala Phe Ala Val Val Ala Gly Glu Val Arg Asn Leu Ala Gln Arg Ser Ala Gln Ala Ala Lys Glu Ile Lys Ala

2621 2648 2675 2702
CTG ATC GAA GAC TCG GTC AAC CGT GTG GAT ATG GGC TCG GTA CTG GTG GAA AGC GCC GGC GAC ACC ATG GGC GAT ATC GTC AAT GCG GTC ACC GCG GTG ACC GAC ATC
Leu Ile Glu Asp Ser Val Asn Arg Val Asp Met Gly Ser Val Leu Val Glu Ser Ala Gly Asp Thr Met Gly Asp Ile Val Asn Ala Val Thr Arg Val Thr Asp Ile

2729 2756 2783 2810
ATG GGT GAA ATC GCC TCT GCT TCC GAT GAA CAG AGC CCG GGT ATC GAC CAG GTC GGC CAG GCG GTA CCG GAA ATG GAT GCG GTC ACC CAG CAG AAC GCC TCG CTG GTG
Met Gly Glu Ile Ala Ser Ala Ser Asp Glu Gln Ser Arg Gly Ile Asp Gln Val Gly Gln Ala Val Arg Glu Met Asp Arg Val Thr Gln Gln Asn Ala Ser Leu Val

2837 2864 2891 2918
GAG GAG TCG GCC TCG GCC GCC GCG CTG GAA GAG CAG GCC AGC CTG CTG ACG CAG TCG GGT GCC GTC TTC CCG CTG AAG TCG GAA GGG CAG GAA GAG TAT AAA CCG CCG
Glu Glu Ser Ala Ser Ala Ala Ala Leu Glu Leu Gln Ala Ser Leu Thr Gln Ser Val Ala Val Phe Arg Leu Lys Ser Glu Gln Glu Glu Tyr Lys Ala Phe

2945 2972 2999 3026
GTC AGC AAT AAA ACC GCG CCT GCC GGC ATC GCT ACG CAT AAA AAA ACC AGC GCC AGC GAC TAC CAG GAT AAC TGG GAG ACG CTG TAA CCG GTA CCG GCC GCG ATA CCC
Val Ser Asn Lys Thr Ala Ala Ile Ala Thr His Lys Lys Thr Ser Ala Ser Asp Tyr Gln Asp Asn Trp Phe *

3053 3080 3105 3132
TAC ACG CCG CCG CCT CTC TTG CGT TGT AAG CCG GCC GAG ACC GGC TAC CAA GGA GAT TGC T ~~Met~~ ^{Lys} ~~Met~~ ^{Met} TTT AAC CGA ATC CGT ATC TCT ACC AGT CTT TTC CTG CTG CTG
SD MET Phe Asn Arg Ile Arg Ile Ser Thr Ser Leu Phe Leu Leu Leu

3159 3186 3213 3240
ATC TCC TTT TGC ATC ATG CAA CTG ATC AGC ACC GGG CTC TCT TAC GTC CCG TAC CCG CCA ACC ACA ATC TTG AAC GTA TTA CCG TCA GTA GCC AGC ACG GAT GCG
Ile Ser Phe Cys Ile Met Gln Leu Ile Ser Thr Gly Leu Ser Tyr Val Arg Tyr Ala Pro Thr Thr Thr Ile Leu Asn Val Leu Pro Ser Val Ala Ser Thr Asp Ala

3267 3294 3321 3348
CTT AGC CTG AGC TGG GTA TCG CTG TTA CAG GCG AGA AAT ACC CTT AAC CCG GCT ACC CCG GCG GCT GAA GGT GCC GCA GGA GCA GGT GGA AGT GCG CTG ATC GCG
Leu Ser Leu Ser Trp Val Ser Leu Leu Gln Ala Arg Asn Thr Leu Asn Arg Ala Gly Thr Arg Gly Ala Glu Gly Ala Ala Gly Ala Gly Ser Ala Leu Met Ala

3375 3402 3429 3456
GCG CCC GCA GCT CCG TGC AAA AAG CCG ATC TCT ATT TTA ACC AGT TCC TCG ACA CCG CTC GCG CCG ATG AGC AGG AAC AGC AGT Trp CCG ATG CCA CCG GTG ATA GCT
Ala Glu Ser Ile Leu Ile Gly Lys Lys Pro Ile Ser Ile Leu Ser Thr Ser Arg Leu Ala Leu Ser Thr Ser Arg Asn Ser Ser Arg Met Pro Arg Val Ile Ala

3483 3510 3537 3564
ACG AGA ATC TAC GCG CTG GCG GAA CTG ATC GTG TTC CTC GAA AAT CGC AAC CTG CAG GCC TTT ATG GAC CAG CCG ACG CAG AAA ATA CAG GAC CCG TTT GAG GCC GAC
Thr Arg Ile Tyr Ala Leu Arg Glu Leu Ile Val Phe Leu Glu Ile Val Phe Leu Glu Leu Ala Phe Met Asp Gln Pro Thr Gln Lys Ile Gln Asp Arg Phe Glu Ala Asp

3591 3618 3645 3672
TTT GTG CAA TAT CTG CAG CTG GCG AAG CCG ACG ACC GAT GAA GCC AGC GCC TCC AGC CAG CAG GCC TAC GCG TTG GTT CCA ATC TGG TTG GTT GCC GGG GCG GTG CTG
Phe Val Gln Tyr Leu Gln Leu Ala Lys Ala Thr Thr Asp Glu Ala Ser Ala Ser Ser Gln Gln Ala Tyr Ala Leu Val Pro Ile Trp Leu Val Ala Gly Ala Val Leu

3699 3726 3753 3780
ATG CTC TTG GTA GTC ACC CTC AGC GCG ATG TGG TGG CTG CGA ACC ATG CTG GTG CAG CCG CTG AAT ATT ATT CCG GGC CAT TTC GAA CGT ATC GCC AGC GGC GAT CTT
Met Leu Leu Val Val Thr Leu Ser Ala Met Trp Trp Leu Arg Thr Met Leu Val Gln Pro Leu Leu Asn Ile Ile Arg Gly His Phe Glu Arg Ile Ala Ser Gly Asp Leu

3807 3834 3861 3888
TCC GCA CCG ATT GAG GTC TAT GGC CGT AAC GAA ATC AGC CAG CTA TTT GCC AGC CTG CAG CCG ATG CAA CAG TCG CTG ATT GGT ACC GTC GGC GCG GTG CCG GAC GGC
Ser Ala Pro Ile Glu Val Tyr Gly Arg Asn Glu Ile Ser Gln Leu Phe Ala Ser Leu Gln Arg Met Gln Gln Ser Leu Ile Gly Thr Val Gly Ala Val Arg Asp Gly

3915 3942 3969 3996
GCG GAG TCG ATT CTT ATC GGC CTG CAG GAG ATT GCC GAA GGC AAC AAC GAT CTC TCC TCG CCG ACC GAA CAA CAG GAC CGT TCG CTG GAA GAG ACC GCC GCC AGT ATG
Ala Glu Ser Ile Leu Ile Gly Lys Gln Asn Ala Asp Asn Ala Ser Ser Ser Arg Thr Glu Gln Gln Asp Arg Ser Thr Leu Thr Gln Ala Leu Thr Ala Leu Ser Met

4023 4050 4077 4104
GAG CAG CTA ACG GCG ACG GTA AAA CAG AAT GCC GAT AAC GCC CCG CAG GCA TCG CAG CTG GCG CGT GAT GCC TCC TCC ACC GCC AAA CCG ACG CTG GCG GAT GAC
Glu Gln Leu Ser Ile Leu Ile Gly Lys Gln Asn Ala Asp Asn Ala Ser Ser Ser Thr Ala Ala Lys Arg Thr Leu Ala Asp Asp

4131 4158 4185 4212
GTG GTC ACC ACG ATG CAC GAT ATC GCC AAC AGT TCG CAA AAA ATC GGC GCG ATT ACC AGC GTC ATC GAC GGC ATC GCT TTC CAG ACC AAT ATT CTG GCG CTT AAC GCG
Val Val Thr Thr Met His Asp Ile Ala Asn Ser Ser Gln Lys Ile Gly Ala Ile Thr Ser Val Ile Asp Gly Ile Ala Phe Gln Thr Asn Ile Leu Ala Leu Asn Ala

4239 4266 4293 4320
GCG GTG GAG GCG GCG CCG GCC GGT GAA CAG GGG CCG GCG TTT GCG GTG GTC GCG GGT GAA GTG CCG AAC CTG GCC AGC CCG CAG CCG GCA AAA GAA ATC AAG
Ala Val Glu Ala Ala Arg Ala Gly Glu Gln Gly Arg Gly Phe Ala Val Val Ala Gly Glu Val Arg Asn Leu Ala Ser Arg Ser Ala Gln Ala Ala Lys Glu Ile Lys

4347 4374 4401 4428
CTG CTG ATC GAT GAA TCC GTC AGC CCG GTC AAA CAC GGC TCG GTG CTG GTG GAA AAT TCC GGC GCC ACC ATG CAG GAC ATC GTG CCG TCG GTC ACC CCG GTC ACC GAC
Leu Leu Ile Asp Glu Ser Val Ser Arg Val Lys His Gly Ser Val Leu Val Glu Asn Ser Gly Ala Thr Met Gln Asp Ile Val Arg Ser Val Thr Arg Val Thr Asp

4455 4482 4509 4536
ATC ATG GGC GAA ATC GCC TCC GCC TCC GAT GAG CAG AGC CCG GGC ATC GAG CAG GTG ACG CAG GCG TTA CCC AGA TAT CAG GTC ACG CAA CAG AAC GCC GCC CTG GTG
Ile Met Gly Glu Ile Ala Ser Ala Ser Asp Glu Gln Ser Arg Gly Ile Thr Gln Ala Leu Pro Arg Tyr Gln Val Thr Gln Gln Asn Ala Leu Ala Leu Val

4563 4590 4617 4644
GTG GAA TCG GCT TCC GCC GCT GCG CTG GAA GAG CAG GCG ATT ACC CTC GGC GAT GCA CTG GCG GTA TTC CCG CTG GCG GAC GAC TAA TTG TCG CCG ATA TAC
Val Glu Ser Ala Ser Ala Ala Ala Leu Glu Glu Gln Ala Ile Thr Leu Ala Asp Ala Val Ala Val Phe Arg Leu Ala Asp Asp *

4671 4698 4723 4750
CTA TCC AGG TAA AGG AGA CCC GGT GGA TTG TCA AAC CCG ATT AAA CCG TCG TAT TTG CCG GTG AGC TGG C ^{cheR'} ATG AAG CAA GAC GAC ATC AAC CCG CCG CGT GAA AGC
MET Lys Gln Asp Asp Ile Asn Ala Ala Arg Glu Ser

4777 4804 4831 4858
GGA TCG GCC GTG GCG CAG ATG GCC CAG CGT CTG CCG CTC TCC GAC GCG CAT TTT CCG CCG ATC AGC CAG CTT ATC TAT CAG CCG GCC GCG ATC GTG CTG GCG CCG CAT
Gly Ser Ala Val Ala Gln Met Ala Gln Arg Leu Pro Leu Ser Asp Ala His Phe Arg Arg Ile Ser Gln Leu Ile Tyr Gln Arg Ala Gly Ile Val Leu Ala Pro His

4885 4912 4939 4966
AAG CCG GAG ATG GTG TAC AAC CCG CTG GTG CCG CGT TTA CGT CTG CTG GCG ATT CAT GAT TTC GCG GAC TAC CTG CCG CTG CTG GAA AGC GAC CCG CAC AGC GCC GAG
Lys Arg Glu Met Val Tyr Asn Arg Leu Val Arg Arg Leu Arg Leu Leu Gly Ile His Asp Phe Gly Asp Tyr Leu Ala Leu Leu Glu Ser Asp Pro His Ser Ala Glu

4993 5015
TGG CAG GCG TTT ATC AAT GCG CTG ACC ACC AAC CTG ACC GCC TTT TTT C
Trp Gln Ala Phe Ile Asn Ala Leu Thr Thr Asn Leu Thr Ala Phe Phe

FIG. 4—Continued




FIG. 5. Comparison of the predicted amino acid sequences of CheW from *E. coli* (33), *S. typhimurium* (43), and *E. aerogenes*. Identical residues are boxed. The β - α - β structural motif (45) extends over residues 128 through 148 and 155 through 160.

rium also have a pattern of conserved and variable regions (Fig. 5). CheW has been proposed to contain a nucleotide-binding site (43). An 11-residue consensus sequence that determines the β - α - β motif of the binding site has been described (45). All three species have the identical amino acid at each of the 11 positions, and 10 of the 11 residues conform to the consensus. Other highly conserved segments of CheW may correspond to sites of interaction between CheW and the signaling domains of the transducers (P. Ames, J. Chen, C. Wolff, and J. S. Parkinson, Cold Spring Harbor Symp. Quant. Biol., in press).

Tse and Tas have the typical structural features of signal transducers (Fig. 6): a positively charged N terminus (residues 1 through 6) followed by a hydrophobic membrane-spanning region (residues 7 through 30), a hydrophilic periplasmic domain (residues 31 through 190), a second hydrophobic membrane-spanning region (residues 191 through 210), and a hydrophilic cytoplasmic domain (residue 210 through the C terminus). Of 187 residues that are identical in the six transducers in Fig. 6, 168 are within the cytoplasmic domain and 145 are within a stretch of 236 amino acids (residues 280 through 516 in Tsr).

The K1 and R1 tryptic peptides of Tsr (residues 295 through 317 and 483 through 507) contain the sites of covalent methylation (19). Within the region corresponding to K1, Tse and Tsr are identical at 22 of 23 residues and Tas and Tsr are identical at 21 of 23 residues. The same three methylation sites, which are Glu-Glx sequences, are present in Tse and Tas as well as in Tsr and Tar. Within R1, Tse and Tsr are identical at 21 of 25 residues and the two methylation sites are present. Tas and Tsr are identical at 20 of 25 residues, but one of the methylation sites may be absent in Tas due to the replacement of the Glu-Glu sequence at residues 492 and 493 of Tsr with Val-Glu in Tas.

Another highly conserved stretch runs from residues 360 through 407 of Tsr. The same 48 residues are present in Tar, and there is only one difference in Tap. Tse and Tas share 44

of the 48 residues with Tsr. In Trg, only 37 of these 48 residues are identical to Tsr (5), underscoring the proposed early divergence of Trg from the other transducers. The strong conservation of this region and of the K1 and R1 peptides emphasizes their importance in signaling and adaptation, respectively (Ames et al., in press).

The Tse and Tsr proteins are 60% identical in their periplasmic domains (Fig. 6 and Table 2). Even their cytoplasmic domains resemble one another more closely than they do those of other transducers. The periplasmic domain of Tas bears little resemblance to that of any other known transducer, including Tap, which like Tas is the product of the second gene in a *meche* operon, or Tar, which is a functional analog of Tas. Given the dissimilarity of Tar and Tas, it is not surprising that Tas does not participate in maltose taxis.

The amino acid transducers Tsr, Tse, Tas, and Tar share three sites that are partially or wholly absent in Tap and Trg: (i) Arg at residues 64, 69, and 73 (Arg-64 is present in Trg); (ii) Glu-Leu-Ile at residues 138 through 140; (iii) Gln-Pro-Thr-Gln at residues 154 through 157 (Pro-155 and Gln-157 are present in Tap). Genetic studies indicate that some of these residues are important in chemoreception. *E. coli* mutants in which Arg-64, Arg-69, or Arg-73 of Tar is replaced by other residues have a drastically reduced ability to carry out aspartate taxis, and the apparent affinity for aspartate is reduced by a factor of 10^2 or more (46). Furthermore, a mutation affecting the Arg-64 residue of *E. coli* Tsr greatly reduces the ability of cells to sense serine (25). The three Arg residues may form a positively charged pocket that interacts with a common element of amino acids, such as their α -carboxyl group (25, 46; Ames et al., in press). Other data suggest that Thr-156 of Tsr and Tar is important for serine (25) and aspartate (C. Wolff, personal communication) sensing.

Tse (with 557 residues) and Tas (with 510 residues) are the largest and smallest transducers, respectively, known in the




FIG. 6. Comparison of the predicted amino acid sequences of signal transducer proteins Tsr (*E. coli*), Tse (*E. aerogenes*), Tar (*E. aerogenes*); Tar_E (*E. coli*), Tar_S (*S. typhimurium*), and Tap (*E. coli*). The sequences are aligned to give the maximal number of residue identities among the six transducers, using the corrected sequence of Tsr (12, 25, 27). This alignment introduces some gaps, most of which can reasonably be explained by the deletion or insertion of codons in particular lineages. The alignment for a sequence (covering residues 194 through 212 of Tsr) that includes most of the second membrane-spanning region contains the most gaps and has the lowest qualitative confidence level. Residues identical in all six transducers, or in all except Tap, are boxed. Symbols: *, positions at which residues are identical in Tsr and Tse and at least one other transducer; O, positions at which residues are identical in Tse and Tsr but different in the other transducers. For all other positions, residues that are the same in two or more transducers are printed boldface and underlined. The residue count for each transducer is indicated within parentheses at the end of each line.

enteric bacteria. Most of this variation in length occurs at the extreme C termini of the proteins, which are among the most highly variable segments (Fig. 6). It is striking that the C termini of Tse and Tar are of about the same length and that the last five residues are identical. These features may be correlated with the location of *tse* and *tar* at the 5' end of the *meche* operon. Tsr, encoded by a gene located elsewhere on the chromosome, has a very different C terminus, although the Tsr protein is otherwise homologous with Tse.

Within the enteric bacteria there appear to be at least five evolutionary lines among the transducers. Trg, which looks least like the others, may represent the fusion of the N terminus of a membrane transport protein with a C-terminal domain common to all transducers (5). The remaining four groups (Tsr-Tse, Tar, Tas, and Tap) are similar enough that they probably arose by mutational divergence of intact

genes. Their progenitor may have been an amino acid receptor of rather low specificity. Crucial features of the binding site may be conserved in the three Arg residues and the sequences at positions 138 through 140 and 154 through 157. Subsequent gene duplication would have allowed individual transducers to evolve a higher affinity for certain substrates. The interaction with substrate-binding proteins (maltose-binding protein for Tar, dipeptide-binding protein for Tap) could have elaborated later.

E. coli, *S. typhimurium*, and *E. aerogenes* all have one or two transducer genes in their *meche* operons, but the genes occur in three different combinations. It may be that all transducers other than Trg evolved at this locus and that some of the genes were then deleted or moved to new sites by chromosome rearrangements. Alternatively, duplicated genes may have been redistributed on the chromosome

TABLE 2. Sequence identities in the periplasmic domain of chemotactic signal transducers^a

Amino acid	No. (%) of identical residues in:				
	Tse	Tas	Tar _E	Tar _S	Tap
Tsr	100 (60)	43 (26)	53 (32)	51 (31)	32 (19)
Tse		50 (30)	52 (31)	50 (30)	27 (16)
Tas			35 (21)	33 (20)	29 (17)
Tar _E				113 (68)	35 (21)
Tar _S					35 (21)

^a The amino acid sequences for the six transducers, aligned as in Fig. 6, were compared for the region encompassing residues 31 to 193 of Tsr. The region contains 167 residue positions, although no single transducer has an amino acid at each position. The Table presents the number of residue identities over these 167 positions in pairwise combinations of transducers. Since the deletion or insertion of a codon can occur during gene evolution, the absence of an amino acid at the same position in two or more transducers was also scored as an identity. The percent identity is given in parentheses; it would be altered only slightly if shared gaps were not counted as identities.

before their evolutionary divergence. Genes evolved elsewhere could then have reinvaded the *meche* operon by recombining with the resident transducer genes. The study of transducer evolution will be enhanced by information from a wide taxonomic spectrum of bacteria.

ACKNOWLEDGMENTS

We thank J. S. Parkinson for communication of results before publication and Karin Eigelmeier and Joerg Vreemann for many helpful tips in computer editing.

This work was supported by Sonderforschungsbereich SFB 156 of the Deutsche Forschungsgemeinschaft and by funds supplied by Texas A&M University, including an institutional biomedical research support grant from the National Institutes of Health administered by the Office of University Research Services. M. K. Dahl is a Boehringer Ingelheim Fonds fellow.

LITERATURE CITED

- Bachmann, B. J. 1987. Linkage map of *Escherichia coli* K12, edition 7. In J. L. Ingraham, K. B. Low, B. Magasanik, M. Schaechter, and H. E. Umbarger (ed.), *Escherichia coli* and *Salmonella typhimurium*: cellular and molecular biology. American Society for Microbiology, Washington, D.C.
- Bartlett, D. H., B. B. Frantz, and P. Matsumura. 1988. Flagellar transcriptional activators FliB and FliA: gene sequences and 5' consensus sequences of operons under FliB and FliA control. *J. Bacteriol.* **170**:1575-1581.
- Biggin, M. D., T. J. Gibson, and G. F. Hong. 1983. Buffer gradient gels and ³⁵S-label as an aid to rapid DNA sequence determination. *Proc. Natl. Acad. Sci. USA* **80**:3963-3965.
- Bolivar, F., R. L. Rodriguez, P. J. Green, M. C. Betlach, H. L. Heyneker, and H. W. Boyer. 1977. Construction of new cloning vehicles. II. A multipurpose cloning system. *Gene* **2**:95-113.
- Bollinger, J., C. Park, S. Harayama, and G. L. Hazelbauer. 1984. Structure of the Trg protein: homologies with and differences from other sensory transducers of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **81**:3287-3291.
- Boyd, A., K. Kendall, and M. I. Simon. 1983. Structure of the serine chemoreceptor in *Escherichia coli*. *Nature (London)* **301**:623-626.
- Boyd, A., A. Krikos, and M. I. Simon. 1981. Sensory transducers of *Escherichia coli* are encoded by homologous genes. *Cell* **26**:333-343.
- Carter, P., H. Bedouelle, and G. Winter. 1985. Improved oligonucleotide site directed mutagenesis using M13 vectors. *Nucleic Acids Res.* **13**:4431-4443.
- Dahl, M. K., and M. D. Manson. 1985. Interspecific reconstitution of maltose transport and chemotaxis in *Escherichia coli* with maltose-binding protein from various enteric bacteria. *J. Bacteriol.* **164**:1057-1063.
- Davis, R. W., D. Botstein, and J. R. Roth. 1980. Advanced bacterial genetics, p. 231, 239, and 240. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Elledge, S. J., and G. C. Walker. 1985. Phasmid vectors for identification of genes by complementation of *Escherichia coli* mutants. *J. Bacteriol.* **162**:777-783.
- Gebert, J. F., B. Overhoff, M. D. Manson, and W. Boos. 1988. The Tsr chemosensory transducer of *Escherichia coli* assembles into the cytoplasmic membrane via a *secA*-dependent process. *J. Biol. Chem.* **263**:16652-16660.
- German Collection of Microorganisms. 1983. Catalogue of strains, 3rd ed. Goettingen, Federal Republic of Germany.
- Hazelbauer, G. L. 1975. The maltose chemoreceptor of *Escherichia coli*. *J. Bacteriol.* **122**:206-214.
- Hazelbauer, G. L., and S. Harayama. 1979. Mutants in transmission of chemotactic signals from two independent receptors of *E. coli*. *Cell* **16**:617-625.
- Helmann, J. D., and M. J. Chamberlin. 1987. DNA sequence analysis suggests that expression of flagellar and chemotaxis genes in *Escherichia coli* and *Salmonella typhimurium* is controlled by an alternative σ factor. *Proc. Natl. Acad. Sci. USA* **84**:6422-6424.
- Helmann, J. D., L. M. Marquez, and M. J. Chamberlin. 1988. Isolation and characterization of the *Bacillus subtilis* σ^{28} factor. *J. Bacteriol.* **170**:1568-1574.
- Karn, J., S. Brenner, L. Barnett, and G. Cesareni. 1980. Novel bacteriophage lambda cloning vehicle. *Proc. Natl. Acad. Sci. USA* **77**:5172-5176.
- Kehry, M. R., and F. W. Dahlquist. 1982. The methyl-accepting chemotaxis proteins in *Escherichia coli*. Identification of multiple methylation sites on methyl-accepting chemotaxis protein I. *J. Biol. Chem.* **257**:10378-10386.
- Komeda, Y. 1986. Transcriptional control of flagellar genes in *Escherichia coli* K-12. *J. Bacteriol.* **168**:1315-1318.
- Kondoh, H., C. B. Ball, and J. Adler. 1979. Identification of a methyl accepting chemotaxis protein for ribose and galactose chemoreceptors of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **76**:260-264.
- Kossmann, M., C. Wolff, and M. D. Manson. 1988. The maltose chemoreceptor of *Escherichia coli*: interaction of maltose-binding protein and the Tar signal transducer. *J. Bacteriol.* **170**:4516-4521.
- Krikos, A., M. P. Conley, A. Boyd, H. C. Berg, and M. I. Simon. 1985. Chimeric chemosensory transducers of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* **82**:1326-1330.
- Krikos, A., N. Mutoh, A. Boyd, and M. I. Simon. 1983. Sensory transducers of *E. coli* are composed of discrete structural and functional domains. *Cell* **33**:615-622.
- Lee, L., T. Mizuno, and Y. Imae. 1988. Thermosensing properties of *Escherichia coli* mutants defective in serine chemoreception. *J. Bacteriol.* **170**:4769-4774.
- Maeda, K., and Y. Imae. 1979. Thermosensory transduction in *Escherichia coli*: inhibition of the thermoresponse by L-serine. *Proc. Natl. Acad. Sci. USA* **76**:91-95.
- Manoil, C., and J. Beckwith. 1986. A genetic approach to analyzing membrane protein topology. *Science* **233**:1403-1408.
- Manson, M. D., V. Blank, G. Brade, and C. F. Higgins. 1986. Peptide chemotaxis in *Escherichia coli* involves the Tap signal transducer and the dipeptide permease. *Nature (London)* **321**:253-256.
- Messing, J. 1983. New M13 vectors for cloning. *Methods Enzymol.* **101**:20-78.
- Miller, J. H. 1972. Experiments in molecular genetics. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
- Mizuno, T., N. Mutoh, S. M. Panasenko, and Y. Imae. 1986. Acquisition of maltose chemotaxis in *Salmonella typhimurium* by the introduction of the *Escherichia coli* chemosensory transducer gene. *J. Bacteriol.* **165**:890-895.
- Mutoh, N., K. Oosawa, and M. I. Simon. 1986. Characterization of *Escherichia coli* chemotaxis receptor mutants with null phenotypes. *J. Bacteriol.* **167**:992-998.
- Mutoh, N., and M. I. Simon. 1986. Nucleotide sequence corre-

- sponding to five chemotaxis genes in *Escherichia coli*. *J. Bacteriol.* **165**:161-166.
34. **Norlander, J., T. Kempe, and J. Messing.** 1983. Construction of improved M13 vectors using oligodeoxynucleotide-directed mutagenesis. *Gene* **26**:101-106.
 35. **Nowlin, D. M., D. O. Nettleton, G. W. Ordal, and G. L. Hazelbauer.** 1985. Chemotactic transducer proteins of *Escherichia coli* exhibit homology with methyl-accepting proteins from distantly related bacteria. *J. Bacteriol.* **163**:262-266.
 36. **Oosawa, K., N. Mutoh, and M. I. Simon.** 1988. Cloning of the C-terminal cytoplasmic fragment of the Tar protein and effects of the fragment on chemotaxis of *Escherichia coli*. *J. Bacteriol.* **170**:2521-2526.
 37. **Russo, A. F., and D. E. Koshland, Jr.** 1983. Separation of signal transduction and adaptation functions of the aspartate receptor in bacterial sensing. *Science* **220**:1016-1020.
 38. **Sanderson, K. E.** 1977. Genetic relatedness in the family *Enterobacteriaceae*. *Annu. Rev. Microbiol.* **30**:327-349.
 39. **Silhavy, T. J., M. L. Berman, and L. W. Enquist.** 1984. Experiments with gene fusions. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.
 40. **Sprenger, G. A., and J. W. Lengeler.** 1984. L-Sorbose metabolism in *Klebsiella pneumoniae* and Sor⁺ derivatives of *Escherichia coli* K-12 and chemotaxis toward sorbose. *J. Bacteriol.* **157**:39-45.
 41. **Springer, M. S., M. F. Goy, and J. Adler.** 1977. Sensory transduction in *Escherichia coli*: two complementary pathways of information processing that involve methylated proteins. *Proc. Natl. Acad. Sci. USA* **74**:3312-3316.
 42. **Springer, M. S., M. F. Goy, and J. Adler.** 1979. Protein methylation in behavioral control mechanisms and in signal transduction. *Nature (London)* **280**:279-284.
 43. **Stock, A., J. Mottonen, T. Chen, and J. Stock.** 1987. Identification of a possible nucleotide binding site in CheW, a protein required for sensory transduction in bacterial chemotaxis. *J. Biol. Chem.* **262**:535-537.
 44. **Tso, W. W., and J. Adler.** 1974. Negative chemotaxis in *Escherichia coli*. *J. Bacteriol.* **118**:560-576.
 45. **Wierenga, R. K., P. Terpstra, and W. G. J. Hol.** 1986. Prediction of the occurrence of the ADP-binding β - α - β fold in proteins, using an amino acid sequence fingerprint. *J. Mol. Biol.* **187**:101-107.
 46. **Wolff, C., and J. S. Parkinson.** 1988. Aspartate taxis mutants of the *Escherichia coli* Tar receptor. *J. Bacteriol.* **170**:4509-4515.