

ANOMALOUS STRUCTURES IN THE CYTOPLASM OF HeLa  
CELLS CULTURED IN THE PRESENCE OF 5-BROMODEOXYURIDINE

WILLIAM R. HENDEE, WILLIAM ZEBRUN, and FREDERICK J. BONTE. From the Department of Radiology and Department of Anatomy, The University of Texas Southwestern Medical School, Dallas, Texas. Dr. Hendee's present address is the Department of Physics and Astronomy, Millsaps College, Jackson, Mississippi

INTRODUCTION

The presence of 5-bromodeoxyuridine, an analogue of the pyrimidine nucleoside thymidine, within the culture medium of tissue culture cells

results in incorporation of this analogue into cellular DNA together with suppression of *de novo* thymidine synthesis (1-5). It has been suggested that incorporation of this analogue into the DNA




FIGURE 1

Section through a HeLa cell which has been incubated in BUDR-containing medium for 4 days. Round or oval anomalous bodies (*DB*) of uniform density are present in appreciable numbers. The internal structure of these bodies varies from an essentially homogeneous, granular matrix to a complex network of disorganized membranes. Normal-appearing structures are present adjacent to the dense bodies, although there may be some evidence of increased vacuolization of the Golgi region. Four-day incubation in  $5 \times 10^{-5}$  M BUDR-containing medium.  $\times 18,000$ .

strand results in altered base pairing and base sequence (6-8), actions which may contribute to the decreased viability and increased radiosensitivity observed by different investigators in various analogue-labeled biological organisms (9-12). However, since the possibility exists that analogue action on other cell components may be at least partially responsible for these observations, an investigation of the influence of 5-bromodeoxyuridine on cell parameters other than radiosensitivity was undertaken. The report presented here is confined to structural anomalies observed in HeLa cells grown in medium supplemented with 5-bromodeoxyuridine.

#### MATERIALS AND METHODS

Stock HeLa parental cells (13) were maintained in modified Eagle's growth medium (14) as monolayer

cultures grown on the surface of glass prescription flasks. Uniformity of the growth sheet and absence of heavily clumped areas were criteria for selecting cells to be subjected to experimental conditions. Selected cultures were trypsinized and placed in suspension by vigorously pipetting them into a fresh supply of growth medium supplemented with the appropriate concentration of 5-bromodeoxyuridine (BUDR). Analogue concentration ranged from  $10^{-7}$  M to  $10^{-4}$  M. Aliquot samples of this suspension were allowed to attach to glass coverslips. At all times, appropriate non-labeled control cells were maintained.

At various time intervals between 1 and 7 days following introduction to the analogue, the cells were fixed directly on the coverslips in cold 2 per cent osmium tetroxide buffered with Dalton's potassium dichromate (15). The cells were dehydrated, carefully removed from the coverslips in a monomeric mixture of butyl and methyl methacrylate, and em-



FIGURE 2

Section through another cell incubated in BUDR-containing medium, demonstrating a large number of homogeneously dense, unidentified bodies (*DB*). Although these anomalous bodies are of uniform density and resemble somewhat the dense bodies found in control cells, they are present in much greater numbers in cells which have been incubated in BUDR-containing growth medium. Four-day incubation in  $5 \times 10^{-5}$  M BUDR-containing medium.  $\times 19,000$ .

bedded in gelatin capsules containing the same mixture. Cells were observed and photographed with an RCA EMU 3-F electron microscope. Each group of BUDR-labeled cells was compared with its own control cell group to account for morphological variations among cell samples caused by factors other than those purposefully introduced.

#### OBSERVATIONS AND DISCUSSION

The cytoarchitecture of representative control HeLa cells has been described by other investigators (16-20) and will not be included here. A large number of the HeLa cells cultured in the presence of higher concentrations of BUDR ( $10^{-5}$  to  $10^{-4}$  M) are observed to contain aggregates of round or oval dense cytoplasmic bodies possessing varying degrees of internal structure (Figs. 1 to 3). The variation in internal structure ranges from an essentially homogeneous, granular matrix

to a complete network of disorganized membranes. Frequently the dense bodies contain an accumulation of small, round vesicles and/or microcrystals similar to those described by Dourmashkin and Dougherty (21). The anomalous structures are usually seen adjacent to other organelles of apparently normal configuration, although there may be some evidence of increased vacuolization of the Golgi complex (Fig. 1). The dense bodies are clearly demonstrable following 72 hours of incubation in BUDR-containing medium.

At higher magnification (Fig. 3), the internal structure of these bodies is observed to consist of a complex network of membranes within a dense matrix and bound by a single, double, or multi-layered membrane. However, in some cells the dense bodies show little evidence of internal structure other than a homogeneous matrix.

Although the anomalous bodies bear a structural resemblance to the "globoid bodies" (22), "lysosomal granules" (18), or "lamellar bodies" (23) observed by different investigators in control HeLa cells, the increased size and much greater abundance of the homogeneously granular bodies in BU DR-labeled cells suggest that their formation is a reaction of the cell to cultivation in BU DR-containing medium. The bodies also structurally resemble to some degree the microbodies of rat liver which Rouillier and Bernhard (24) interpret as either degenerating or regenerating mitochondria. Since the dense bodies observed in this study on HeLa cells are often found adjacent to mitochondria of normal configuration, a different interpretation is suggested in this case, namely, that these dense bodies may represent structures synthesized *de novo* as a response to a substance foreign to the cell. The intense osmophilia of the anomalous bodies suggests the presence of quantities of lipid or lipid-like substance, possibly arising from an alteration in the phospholipid or lipid metabolism of the cell.

#### SUMMARY


Cultivation of HeLa cells in the presence of the thymidine analogue 5-bromodeoxyuridine for periods of 72 hours or longer results in the accumulation in the cytoplasm of aggregates of round or oval, dense, homogeneously granular bodies. Although the origin of these bodies is unknown, it is suggested that they may arise as a result of altered phospholipid or lipid metabolism.

#### REFERENCES

- CHEONG, L., RICH, M. A., and EIDINOFF, M. L., Introduction of the 5-halogenated uracil moiety into deoxyribonucleic acid of mammalian cells in culture, *J. Biol. Chem.*, 1960, **235**, 1441.
- DJORDJEVIC, B., and SZYBALSKI, W., Genetics of human cell lines: III. incorporation of 5-bromo and 5-iododeoxyuridine into the deoxyribonucleic acid of human cells and its effect on radiation sensitivity, *J. Exp. Med.*, 1960, **112**, 509.
- HAKALA, M. T., Mode of action of 5-bromodeoxyuridine on mammalian cells in culture, *J. Biol. Chem.*, 1959, **234**, 3072.
- KRISS, J. P., and RÉVÉSZ, L., Quantitative studies of incorporation of exogenous thymidine and 5-BU DR into DNA of mammalian cells *in vitro*, *Cancer Research*, 1961, **21**, 1141.
- SIMON, E. H., Transfer of DNA from parent to progeny in a tissue culture line of human carcinoma of the cervix (strain HeLa), *J. Mol. Biol.*, 1961, **3**, 101.
- SZYBALSKI, W., Ultraviolet light sensitivity and other biological and physico-chemical properties of halogenated DNA, *Progress in Photobiology*, New York, Elsevier Publishing Co., 1960, 542.
- SHAPIRO, H. S., and CHARGAFF, E., Severe distortion by 5-BU of the sequence characteristics of a bacterial DNA, *Nature*, 1960, **188**, 62.
- SINSHEIMER, R. L., Nucleic acids of the bacterial viruses, *The Nucleic Acids*, (E. Chargaff and J. Davidson, editors), New York, Academic Press, Inc., 1960, **3**, 223.
- BERRY, R. J., and ANDREWS, J. R., Modification of the radiation effect on the reproductive capacity of tumor cells *in vivo* with pharmacologic agents, *Rad. Research*, 1962, **16**, 82.
- ERIKSON, R. L., and SZYBALSKI, W., Molecular radiobiology of human cell lines. I. Comparative sensitivity to x-rays and ultraviolet light of cells containing halogen-substituted DNA, *Biochem. and Biophysic. Res. Comm.*, 1961, **4**, 258.
- KAPLAN, H. S., SMITH, K. C., and TOMLIN, P. A., Effect of halogenated pyrimidines on radiosensitivity of *E. coli*, *Rad. Research*, 1962, **16**, 98.
- KOZINSKI, W., and SZYBALSKI, W., Dispersive transfer of parental DNA molecule to the progeny of phage  $\Phi$ X-174, *Virology*, 1959, **9**, 260.
- GEY, G. O., COFFMAN, W. D., and KUBICEK, M. T., Tissue culture studies of the proliferative capacity of cervical carcinoma and normal epithelium, *Cancer Research*, 1952, **12**, 264.
- EAGLE, H., Nutrition needs of mammalian cells in tissue culture, *Science*, 1955, **122**, 501.
- DALTON, A. J., A chrome-osmium fixative for electron microscopy, *Anat. Rec.*, 1955, **121**, 281.

#### FIGURE 3

The internal structure of the unidentified dense bodies seen at higher magnification. Membranous and vesicular elements are embedded within the uniformly dense matrix. Three-day incubation in  $5 \times 10^{-5}$  M BU DR-containing medium.  $\times 42,000$ .



16. EPSTEIN, M. A., Some unusual features of fine structure observed in HeLa cells, *J. Biophysic. and Biochem. Cytol.*, 1961, **10**, 153.
17. HARFORD, C. G., HAMLIN, A., and PARKER, E., Electron microscopy of HeLa cells after ingestion of colloidal gold, *J. Biophysic. and Biochem. Cytol.*, 1957, **3**, 749.
18. JOURNEY, L. J., and GOLDSTEIN, M. N., Electron microscope studies on HeLa cell lines sensitive and resistant to actinomycin D, *Cancer Research*, 1961, **21**, 929.
19. MORGAN, C., GODMAN, G. C., BREITENFELD, P. M., and ROSE, H. M., A correlative study by electron and light microscopy of the development of type 5 adenovirus, *J. Exp. Med.*, 1960, **112**, 373.
20. STOKES, M. G. P., SMITH, K. M., and ROSS, R. W., Electron microscope studies of HeLa cells infected with herpes virus, *J. Gen. Microbiol.*, 1958, **19**, 244.
21. DOURMASHKIN, R., and DOUGHERTY, R., Phagocytosis of crystalline particles by cells grown in tissue culture, *Exp. Cell Research*, 1961, **25**, 480.
22. HARFORD, C. G., HAMLIN, A., PARKER, E., and VON RAVENSWAAY, T., Globoid structures in the cytoplasm of rapidly growing HeLa cells, *J. Biophysic. and Biochem. Cytol.*, 1956, **2**, No. 4, 347.
23. KOJIMA, K., and KOZUKA, S., Structural changes in HeLa cells cultivated in serum-free medium, *J. Cell. Biol.*, 1962, **14**, 141.
24. ROULLER, C., and BERNHARD, W., Microbodies and the problem of mitochondrial regeneration, *J. Biophysic. and Biochem. Cytol.*, 1956, **2**, No. 4, 355.