

The Phosphatidylinositol Transfer Protein Domain of *Drosophila* Retinal Degeneration B Protein Is Essential for Photoreceptor Cell Survival and Recovery from Light Stimulation

Scott C. Milligan,* James G. Alb, Jr.,[‡] Raya B. Elagina,* Vytas A. Bankaitis,[‡] and David R. Hyde*

*Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556; and [‡]Department of Cell Biology, University of Alabama at Birmingham, Birmingham, Alabama 35294

Abstract. The *Drosophila* retinal degeneration B (*rdgB*) gene encodes an integral membrane protein involved in phototransduction and prevention of retinal degeneration. RdgB represents a nonclassical phosphatidylinositol transfer protein (PITP) as all other known PITPs are soluble polypeptides. Our data demonstrate roles for RdgB in proper termination of the phototransduction light response and dark recovery of the photoreceptor cells. Expression of RdgB's PITP domain as a soluble protein (RdgB-PITP) in *rdgB*² mutant flies is sufficient to completely restore the wild-type electrophysiological light response and prevent the degeneration. However, introduction of the T59E mutation, which does not affect RdgB-PITP's phosphatidylinositol (PI) and phosphatidylcholine (PC) transfer in vitro, into the soluble (RdgB-PITP-T59E) or full-length (RdgB-T59E) proteins eliminated rescue of retinal degeneration in *rdgB*² flies, while the light response was partially maintained. Substitution of the rat

brain PITP α , a classical PI transfer protein, for RdgB's PITP domain (PITP α or PITP α -RdgB chimeric protein) neither restored the light response nor maintained retinal integrity when expressed in *rdgB*² flies. Therefore, the complete repertoire of essential RdgB functions resides in RdgB's PITP domain, but other PITPs possessing PI and/or PC transfer activity in vitro cannot supplant RdgB function in vivo. Expression of either RdgB-T59E or PITP α -RdgB in *rdgB*⁺ flies produced a dominant retinal degeneration phenotype. Whereas RdgB-T59E functioned in a dominant manner to significantly reduce steady-state levels of rhodopsin, PITP α -RdgB was defective in the ability to recover from prolonged light stimulation and caused photoreceptor degeneration through an unknown mechanism. This in vivo analysis of PITP function in a metazoan system provides further insights into the links between PITP dysfunction and an inherited disease in a higher eukaryote.

THE *Drosophila* retinal degeneration B protein (RdgB)¹ plays a critical role in the fly photoreceptor cell. The *rdgB* mutant phenotype is characterized by retinal degeneration whose onset, while discernible in dark-reared flies, is greatly accelerated by raising the flies in light (Harris and Stark, 1977; Stark et al., 1983). Typically, *rdgB* mutant flies begin to exhibit the morphological hallmarks of photoreceptor cell degeneration several days

after eclosion (Harris and Stark, 1977; Stark et al., 1983). In addition, these mutant flies exhibit an abnormal light response, as recorded by the rapid deterioration of the electroretinogram (ERG), shortly after the fly's initial exposure to light. This ERG defect is manifested before any obvious physical signs of retinal degeneration (Harris and Stark, 1977), which suggests that the defect in the light response may precipitate the course of retinal degeneration.

In the photoreceptor cell, RdgB localizes to both the axon and the subrhabdomeric cisternae (SRC) (Vihtelic et al., 1993; Suzuki and Hirosawa, 1994). The SRC is an extension of the endoplasmic reticulum that functions both as an intracellular Ca²⁺ store and a compartment through which rhodopsin traffics en route to the rhabdomere (Walz, 1982; Matsumoto-Suzuki et al., 1989; Suzuki and Hirosawa, 1991). Thus, RdgB is the first identified protein required for visual transduction that is not localized in the photoreceptor rhabdomere. Genetic epistasis analyses suggest RdgB functions downstream of both rhodopsin and

S.C. Milligan and J.G. Alb, Jr. contributed equally to this work.

Address all correspondence to David R. Hyde, Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 45665. Tel.: (219) 631-8054. Fax: (219) 631-7413. E-mail: david.r.hyde.1@nd.edu

1. *Abbreviations used in this paper:* ERG, electroretinogram; PC, phosphatidylcholine; PDA, prolonged depolarizing afterpotential(s); PI, phosphatidylinositol; PITP, phosphatidylinositol transfer protein; PLC, phospholipase C; PKC, protein kinase C; *rdgB*, retinal degeneration B; RdgB, retinal degeneration B protein; RdgB-PITP, RdgB PITP domain; SM, sphingomyelin.

phospholipase C (PLC) in the visual transduction cascade as both the *ninaE* (encoding the opsin expressed in photoreceptor cells R1-6 [O'Tousa et al., 1985; Zuker et al., 1985]) and *norpA* (encoding phospholipase C [Bloomquist et al., 1988]) mutations suppress the *rdgB*-dependent, light-enhanced retinal degeneration (Harris and Stark, 1977; Stark et al., 1983). Consistent with this view, constitutive activation of the *Drosophila* G protein transducin analogue (DGq), either by application of nonhydrolyzable GTP analogues or by expression of a constitutively activated G α subunit (Dgq1), effects a rapid degeneration of *rdgB* retinas in the absence of light (Rubinstein et al., 1989; Lee et al., 1994). RdgB apparently functions downstream of the *inaC*-encoded protein kinase C (PKC) because: (a) application of phorbol ester to *rdgB* mutant retinas, which presumably activates the *inaC*-encoded PKC, stimulates retinal degeneration in the absence of light (Minke et al., 1990); and (b) the *rdgB* retinal degeneration is weakly suppressed by the *inaC* mutation (Smith et al., 1991). Thus, the available evidence identifies an execution point for RdgB downstream of PKC in the visual transduction cascade.

RdgB is a 116-kD membrane polypeptide with six potential transmembrane domains (Vihtelic et al., 1991). Additionally, the amino-terminal 281 RdgB residues share 42% amino acid identity with the rat brain phosphatidylinositol (PI) transfer protein α isoform (PITP α) (Vihtelic et al., 1993). Whereas PITPs are operationally defined by their ability to catalyze the transfer of either PI or phosphatidylcholine (PC) monomers between membrane bilayers in vitro (Bankaitis et al., 1990; Cleves et al., 1991; Wirtz, 1991), how the phospholipid transfer activity pertains to in vivo function is less clear. The yeast PITP (Sec14p) uses its PI and PC binding activities in two independent, yet complementary, ways that serve to preserve a Golgi pool of diacylglycerol that is critical for the biogenesis of Golgi-derived secretory vesicles (Kearns et al., 1997). Reconstitution studies suggest that mammalian PITPs play important roles in PLC-mediated inositol signaling, ATP-dependent, Ca²⁺-activated secretion, and constitutive secretion from the *trans*-Golgi network (Hay and Martin, 1993, 1995; Thomas et al., 1993, 1995; Ohashi et al., 1995). However, because the PITP requirement for these processes is generally satisfied by any PITP (even those lacking any primary sequence identity), the physiological relevance of these PITP involvements remains to be determined (Skinner et al., 1993; Cunningham et al., 1995; Ohashi et al., 1995; Alb et al., 1996). The recent finding that the mouse *vibrator* mutation represents a hypomorphic mutation in the *pitpn* gene, which encodes PITP α , indicates that PITP function is important to neuronal function (Hamilton et al., 1997). RdgB's PITP domain (when expressed as a soluble protein in *Escherichia coli*) is able to effect intermembrane transfer of PI in vitro (Vihtelic et al., 1993). Unlike all previously characterized PITPs, which are 32–35-kD soluble proteins (Bankaitis et al., 1989; Cleves et al., 1991; Wirtz, 1991), RdgB is a large integral membrane protein. In spite of postulated in vivo activities for PITPs, the function of RdgB in the photoreceptor cell remains unknown. Recently, vertebrate orthologues of the *rdgB* gene were identified in mice, bovines, and humans (Chang et al., 1997). Expression of the mouse

rdgB cDNA in *rdgB*² null mutant flies resulted in the elimination of the retinal degeneration and complete restoration of the wild-type ERG light response (Chang et al., 1997). Thus, the *Drosophila* RdgB protein defines a new class of functionally equivalent transmembrane PITPs.

In this work, we analyzed RdgB's involvement in the *Drosophila* phototransduction cascade and the mechanism by which it prevents the onset of retinal degeneration. This represents the first in vivo analysis of the transmembrane PITP class, and we report several novel and unanticipated aspects of RdgB function. We demonstrate that the complete repertoire of RdgB functions essential for normal phototransduction reside in the PITP domain. Expression of this domain as a soluble polypeptide fully complements the *rdgB*² null allele. Yet, other PITPs that possess PI and/or PC transfer activities in vitro cannot substitute for RdgB in the photoreceptor cell. Whereas the recessive *rdgB*² null mutation demonstrates an essential role for RdgB in proper termination of the ERG light response and dark recovery of the photoreceptor cell, one novel dominant *rdgB* mutation affects the maintenance of steady-state rhodopsin levels in photoreceptor cells. Another dominant *rdgB* mutation induces retinal degeneration and compromises the rapid regeneration of a wild-type ERG light-response amplitude subsequent to multiple or prolonged light exposure. Taken together, these data indicate an underlying complexity to the mechanism of RdgB function and its role in the photoreceptor cell that is not easily reconciled with a simple role in potentiating signal transduction via phosphoinositide-driven signaling pathways.

Materials and Methods

In Vitro Mutagenesis of the *rdgB* cDNA

The pTV vector contains a wild-type *rdgB* cDNA (consisting of 427 bp of 5' untranslated sequence, the entire *rdgB* open reading frame, and 555 bp of 3' untranslated sequence) downstream of a 317-bp *ninaE* promoter fragment, which is sufficient for gene expression in photoreceptors R1-6 (Mismer and Rubin, 1987). Single-stranded pTV DNA was prepared and in vitro mutagenized with the mutagenic primer 5'-GCTCCGAGAA-GCTTTCGTCGCTGC-3' (Kunkel et al., 1987), which introduced a HindIII site (underlined sequence) at nucleotides 827–832 (Vihtelic et al., 1991), which is the 3' boundary of the PITP domain to create pTVh1. Introduction of the HindIII site generated two conservative changes in RdgB, D276E, and V278F. An XhoI site located between the *ninaE* promoter and the *rdgB* cDNA allowed removal of the PITP domain in pTVh1 by XhoI/HindIII digestion. The mutagenized PITP fragments (see below) were subcloned into pTVh1 lacking the XhoI/HindIII fragment. The soluble rat PITP α was PCR-amplified from a rat PITP α cDNA with primers 5'-CTCGAGAAGCGACATGGTGTCTGC-3' and 5'-AAGCTTCC-TTTCACGGGGTCC-3', and clamped with XhoI and HindIII sites (underlined), respectively. The PCR product was subcloned into the pCR2.1 vector (Invitrogen, Carlsbad, CA), excised by XhoI/HindIII digestion, and ligated into pTVh1 lacking the XhoI/HindIII fragment to yield the chimeric PITP α -RdgB construct. The *rdgB-pitp* construct was made by digestion of pTVh1 with HindIII and HpaI, followed by filling in the ends with Klenow fragment and blunt end ligation. This created a stop codon immediately after residue 277 in the PITP domain. The *pitp* and *rdgB-pitp-T59E* constructs were made in the same manner as the *rdgB-pitp* construct, using *pitp*-*rdgB* and *rdgB-T59E* constructs as the starting DNAs, respectively.

To mutagenize codon 59 of *rdgB*, pTVh1 was digested with XhoI and HindIII to liberate the 1.2-kb restriction fragment carrying the entire coding sequence of RdgB-PITP domain. This fragment was subcloned into pBluescript SK (Stratagene, La Jolla, CA) to yield pSKPT. Single-stranded DNA was subsequently prepared and mutagenized using the primer 5'-GGTAATGGTCAATACGCAAAGAAAATCTATCACGT-

GGG-3' to effect the T₅₉A missense substitution (mutagenized codon underlined) (Kunkel et al., 1987). The T₅₉E substitution was constructed using a mutagenic primer of the exact same sequence with the exception that codon 59 was altered to GAA. The authenticity of each mutagenesis reaction was confirmed by sequencing each 1.2-kb restriction fragment cassette. The 1.2-kb mutagenized XhoI/HindIII fragments were subcloned into pTVh1 lacking the wild-type XhoI/HindIII region to introduce the T₅₉ mutations into the full-length RdgB. The mutagenized cassettes were also individually PCR-amplified with the forward and reverse primers 5'-GGTGAATTCATGCTGATCAAGGAGTACCG-3' (*rdgB* initiator codon in italics) and 5'-AGCGAATCAAGCTTCTTCGCTCGCT-GCCCG-3', and clamped with EcoRI and HindIII sites at the 5' and 3' ends of each product, respectively. These PCR products were subsequently digested with EcoRI and HindIII and the 0.9-kb restriction fragments, which contained the entire PITP open reading frame, were individually subcloned into the T7 RNA polymerase/promoter vector pT7-5 (Tabor, S., Harvard Medical School, Cambridge, MA). The authenticity of each construct was confirmed by nucleotide sequencing and the constructs were subsequently transformed into the *E. coli* strain BL21(DE3) (Novagen, Inc., Madison, WI) for expression of RdgB-PITP proteins.

Nucleotide Sequence Analysis

Nucleotide sequencing was accomplished by the dideoxy chain-termination method (Sanger et al., 1977) using either single- or double-stranded plasmid DNA as a template and the Sequenase version 2.0 sequencing kit (Amersham Corp., Arlington Heights, IL).

Expression of the Soluble RdgB-PITP in *E. coli*

Wild-type and mutant versions of RdgB-PITP were expressed as soluble polypeptides in *E. coli*. Briefly, 1 liter of SuperBroth (Miller, 1972), containing ampicillin (50 µg/ml) was inoculated with a 1:100 dilution of an overnight *E. coli* culture carrying the desired PITP expression plasmid and grown to an OD₆₀₀ of 0.3 at 37°C with shaking. Isopropyl B-*p*-thiogalactopyranoside (IPTG) was added (final concentration 1 mM) and the cultures were incubated for an additional 3 h. Cells were harvested by centrifugation, washed in cold lysis buffer (50 mM NaPO₄ [pH 7.1], 300 mM NaCl, 1 mM NaN₃, 0.2 mM PMSF, and 5 mM EDTA), and pellets were resuspended in 10 ml of cold lysis buffer containing 1:2 volume of 0.1 mm glass beads (BioSpec Products, Inc., Bartlesville, OK). Samples were vigorously vortexed seven times in 1-min bursts with cooling on ice between each burst. The lysates were clarified by serial centrifugation for 5 min at 2,000 g, 20 min at 20,000 g, and finally 60 min at 100,000 g to yield the final cytosolic fraction. Total protein concentration of the cytosol fraction was determined using the bicinchoninic assay (BCA) (Pierce Chemical Co., Rockford, IL).

Phospholipid Transfer Assays

E. coli cytosolic preparations (1 mg total protein per reaction) were individually assayed for PI and PC transfer in vitro as previously described (Aitken et al., 1990; Skinner et al., 1993). Sphingomyelin (SM) transfer assays were performed exactly as PC transfer assays, with the exception that [*N*-methyl-¹⁴C]SM (56 mCi/mmol; Amersham Corp.) was used as transfer substrate (0.09 µCi per assay). Quantitative ELISAs were used to normalize the RdgB content in each *E. coli* cytosolic fraction using a direct sandwich assay with polyclonal mouse anti-RdgB serum directed against RdgB-PITP. Secondary goat anti-mouse antibodies conjugated to horseradish peroxidase (Bio-Rad Laboratories, Hercules, CA) were used for development of signal in the presence of *o*-phenylenediamine. After quenching, A₄₅₀ was measured with an EL311xs automated microplate reader (Bio-Tek Instruments, Inc., Winooski, VT).

Germline Transformation

pTVh1 plasmids containing mutant and wild-type *rdgB* cDNAs were digested with XbaI and KpnI to liberate a single restriction fragment containing the *ninaE* promoter and *rdgB* cDNA. These fragments were individually subcloned into pCaSpeR-4 (Ashburner, 1989), the resulting constructs were purified on a CsCl density gradient, and coinjected with Δ2-3 helper DNA into *w¹¹¹⁸* embryos using standard techniques (Ashburner, 1989b). Multiple independent lines were isolated that expressed

each transgene. These independent lines were assayed to determine that the phenotypes described are due to expression of the transgene and not a fortuitous P element insertion into a particular genomic region. The transgenic lines (in a *rdgB²* null mutant background) were tested for the level of protein expression from the transgenes by immunoblots. The expression level of the various RdgB constructs (from one copy of a given transgene) relative to RdgB in wild-type flies is shown in Table I.

Immunoblot Analyses

Two newly eclosed (>8-h old), dark-raised flies were decapitated in room light and homogenized in 10 µl extraction buffer (2.3% SDS, 10% glycerol, 62.5 mM Tris Cl [pH 6.8], and 0.01% bromophenol blue). The homogenate was incubated at 37°C for 1 h and centrifuged briefly (Ozaki et al., 1993). The supernatant was resolved on a 12.5% polyacrylamide-SDS gel (Laemmli, 1970). Proteins were transferred to nitrocellulose using a transfer apparatus (Bio-Rad Trans-Blot SD Semi-Dry Transfer Cell; Bio-Rad Laboratories) at 15 V for 30 min, the membrane was blocked at least 2 h, and the membrane was finally washed for 20 min in TTBS (0.05% Tween-20 in TBS). The membranes were incubated overnight in either 1:1,000 dilution of anti-rhodopsin polyclonal antiserum, 1:1,000 dilution of anti-RdgC polyclonal antiserum (both provided by J.E. O'Tousa, University of Notre Dame, Notre Dame, IN), 1:500 dilution of affinity-purified anti-Dgq polyclonal antiserum (Lee et al., 1994), 1:1,000 dilution of anti-Trp polyclonal antiserum (provided by C. Montell, Johns Hopkins University, Baltimore, MD), 1:3,000 dilution of anti-Gbe ascites (provided by C. Zuker, University of California at San Diego, San Diego, CA), or anti-RdgB monoclonal supernatant. The membranes were washed three times (10 min per wash) with TTBS, and incubated an additional 2 h in the presence of goat anti-rabbit or goat anti-mouse alkaline phosphatase-conjugated secondary antibody (Sigma Chemical Co., St. Louis, MO) diluted 1:15,000 and 1:3,000, respectively, in 2% Blotto. The membranes were washed twice for 5 min with TTBS. A final 5 min wash with 0.1 M Tris (pH 9.5) preceded colorimetric detection, which was performed according to manufacturer's protocol (Bio-Rad Laboratories).

Electrophysiology

Newly eclosed, dark-reared flies were prepared for ERG analysis in dim red light. After positioning the recording electrode into the eye and the reference electrode into the back of the head, the fly was dark adapted for 1 h before recording the initial response to white light as previously described (Zars and Hyde, 1996). All *rdgB²* flies with and without transgenes were *vermillion* (*v*) eyed, whereas all *rdgB⁺* flies with and without transgenes were *w⁺*. No significant differences were observed in the ERG light-response amplitudes between the *w⁺*, *rdgB⁺*, and *v rdgB⁺* flies. Prolonged depolarizing afterpotentials (PDA) were examined from either *w rdgB⁺*, *rdgB⁺*; *cn bw, rdgB⁺*; *cn bw P[rdgB-T₅₉E], rdgB⁺*; *cn bw P[pitpa-rdgB]*; or *ninaE¹¹⁷* flies. The *cinnabar* (*cn*) and *brown* (*bw*) mutations yield a phenotypically white eye and produced the same PDA in a *rdgB⁺* background as the *white* (*w*) mutation.

Characterization and Histology of Photoreceptor Degeneration

Flies were initially analyzed for retinal degeneration by inspecting the deep pseudopupil; a virtual image produced from the rhabdomeres of ~20 adjacent ommatidia (Franceschini, 1972). Integrity of both the rhabdomeres and of the ommatidial array is required for production of the deep pseudopupil. All the flies examined were in a *w⁺* (wild-type eye color) background, which ensured that all the flies had a consistent eye color, and therefore received equivalent light exposure, regardless of the expression of the different transgenes. The flies were raised in either constant light or under a 12 h light/dark cycle. To establish a time course of degeneration, 3 or 4 replicates of 30–80 flies were examined daily for the deep pseudopupil. At least 100 flies of each genotype were examined for each time point.

Retinal degeneration was also examined by light microscopy of retinal tissue sections. Either *white* (*w*) or *vermillion* (*v*) control and experimental flies were raised in the desired light conditions for the appropriate period and then decapitated; the heads were bisected, fixed, and embedded in Polybed 812 (Polysciences Inc., Warrington, PA) as previously described (Lee et al., 1994). 1-µm sections were cut and stained with methylene blue azure II.

Results

RdgB's PITP Domain Is Sufficient to Rescue the rdgB Mutant Retinal Degeneration and Light-Response Phenotypes

The *rdgB*² null mutation causes a light-enhanced retinal degeneration characterized by the reduction and loss of rhabdomeres, photoreceptor cell death, and the appearance of holes in the retinal tissue (Fig. 1 *A*). Before any histological signs of degeneration, the *rdgB*² mutant loses its ERG light response (Fig. 1 *A*). By contrast, a wild-type retina possesses a well-ordered ommatidial arrangement containing large rhabdomeres and produces an ERG light response of some 25 mV (Fig. 1 *B*). We ectopically expressed a wild-type *rdgB* cDNA in *rdgB*² flies. After 6 d in a 12-h light/dark cycle, these transgenic flies still maintained a wild-type ERG light response and exhibited little evidence of retinal degeneration (Fig. 1 *C*). Indeed, the only detectable histological abnormality was loss of the central R7 and R8 rhabdomeres in some ommatidia; an expected result given that RdgB expression was restricted to photoreceptors R1-6 by the *ninaE* promoter. 30 d after eclosion, these transgenic flies still possessed a wild-type ERG light response and failed to exhibit any degeneration beyond the loss of R7 and R8 (data not shown).

To examine the functional importance of RdgB's PITP domain, we introduced a translational termination codon at position 277 in the wild-type *rdgB* cDNA. This resulted in RdgB's PITP domain being translated as a soluble protein (RdgB-PITP). We germline transformed this construct into flies and expressed it in *rdgB*² mutants using the *ninaE* promoter. Surprisingly, RdgB-PITP expression completely protected the R1-6 photoreceptor cells from degeneration, even at 17 d after eclosion (Fig. 1 *D*). The ERG light response of these transgenic flies was also wild-type (Fig. 1 *D*). The preservation of the R1-6 cells and the ERG light response was still apparent 30 d after eclosion (data not shown). These collective data demonstrate that RdgB-PITP, which represents only 27% of the full-length RdgB primary sequence, was sufficient for complete rescue of both the retinal degeneration and electrophysiological defects associated with *rdgB*² null alleles.

A Threonine-59 Missense Mutation Inactivates PI, but Not PC, Transfer Activity of RdgB-PITP

We previously demonstrated that RdgB-PITP exhibits PI-transfer activity (Vihtelic et al., 1993). To assess the range of phospholipid transfer capability, and to determine whether RdgB-PITP exhibited phospholipid transfer properties more similar to mammalian PITP α (which transfers PI and PC) or PITP β (which transfers PI, PC, and SM), we expressed RdgB-PITP as a soluble polypeptide in *E. coli* (Fig. 2 *A*, lane 1) and assayed its ability to mobilize PI, PC, and SM between membrane bilayers in vitro. RdgB-PITP catalyzed robust transfer of both PI and PC in vitro (Fig. 2 *B*), with $5.3 \pm 0.1\%$ and $6.4 \pm 1.2\%$ of total radiolabeled PI and PC substrate transferred, respectively. However, we did not detect SM transfer activity with RdgB-PITP (Fig. 2 *B*). Thus, RdgB-PITP exhibited biochemical properties more closely resembling those of PITP α than PITP β .

We extended this functional comparison between RdgB-




Figure 1. Wild-type RdgB rescues the *rdgB*² electrophysiological and degeneration phenotypes. (*A*) A retinal section of a 6-d-old *rdgB*² fly raised in a 12-h light/dark cycle reveals signs of massive degeneration including reduced or absent rhabdomeres and holes (*H*) in the retina. At older ages, the central R7 rhabdomere and photoreceptor (*arrowhead*) also degenerate. The ERG light response of a 1-d-old *rdgB*² mutant fly raised in a 12-h light/dark cycle has only a very small amplitude. The fly was dark adapted for 5 min before the ERG recording. (*B*) A retinal section of a 6-d-old Oregon-R fly raised in a 12-h light/dark cycle revealed no obvious signs of degeneration. The ERG light response of a 1-d-old Oregon-R (wild-type) fly exhibits a normal amplitude. (*C*) A retinal section of a 6-d-old *rdgB*²; P[*rdgB*⁺] fly raised in a 12-h light/dark cycle shows no obvious signs of *rdgB*-like retinal degeneration in R1-6. Some R7 cells are losing their rhabdomeres and becoming optically dense (*arrowheads*). The ERG of a 6-d-old *rdgB*²; P[*rdgB*⁺] fly is equivalent to wild-type. (*D*) The retinal section of a 17-d-old *rdgB*²; P[*rdgB*-*pitp*] fly raised in a 12-h light/dark cycle shows no obvious signs of *rdgB*-like retinal degeneration, except for the loss of R7 cells (*arrowheads*). The ERG of a 6-d-old sibling is wild-type. The loss of R7 cells in *C* and *D* was due to the expression of RdgB or RdgB-PITP, respectively, being limited to R1-6 by the *ninaE* promoter. All ERGs were recorded from a 2-s light stimulus, with a 5-mV scale shown in *A*. Bar, 10 μ m.




Figure 2. RdgB-PITP is functionally related to PITP α . (A) Various RdgB-PITPs (wild-type RdgB-PITP and the missense mutants T59A and T59E) were expressed in *E. coli*, partially purified and shown to be stably expressed by gel electrophoresis and immunoblot. (B) RdgB-PITP, RdgB-PITP-T59E, and RdgB-PITP-T59A were analyzed for PI and PC transfer activity in vitro. The solid bars represent the percentage of radiolabeled PI transferred relative to wild-type RdgB-PITP ($5.3 \pm 0.1\%$). The shaded bars represent the percentage of radiolabeled PC transferred relative to wild-type RdgB-PITP ($6.4 \pm 1.2\%$). SM transfer activity was assessed for mammalian PITP β , which served as positive control, and for wild-type RdgB-PITP (*striped bars*). PITP β transferred $2.3 \pm 0.1\%$ of total input radiolabeled SM and this value was set at 100%. Lysates from *E. coli* that lack a plasmid expressing PITP failed to exhibit any detectable PI and PC transfer activity (Alb et al., 1995). Each sample represents three trials and the standard deviation is shown as vertical bars.

PITP and PITP α to analyzing the biochemical effects of specific missense mutations. Previously, we demonstrated that the PI transfer activity of mammalian PITP α is sensitive to amino acid substitutions involving T59, a residue that is conserved amongst all presently known metazoan PITPs and defines a consensus PKC phosphorylation site in PITP α and RdgB (Alb et al., 1995). The T59E alteration selectively abolishes PI transfer activity in PITP α without affecting PC transfer activity, while the T59A substitution reduces the specific activity for PI transfer some twofold relative to wild-type (Alb et al., 1995). We expressed the T59E and T59A mutant forms of RdgB-PITP in *E. coli*

(Fig. 2 A) and assayed them for PI and PC transfer activities. However, the biochemical consequences associated with these individual mutations were, in each case, opposite from those anticipated from the PITP α data. The T59E RdgB-PITP transferred $5.7 \pm 0.1\%$ and $6.9 \pm 1.7\%$ of total radiolabeled PI and PC substrate, respectively. The relative specific activities for PI and PC transfer were $102 \pm 2.3\%$ and $97 \pm 6.8\%$ of wild-type RdgB-PITP, respectively (Fig. 2 B). By contrast, the T59A RdgB-PITP failed to exhibit any detectable PI transfer activity, even though PC transfer activity was essentially unaffected, with $6.6 \pm 0.8\%$ of total radiolabeled PC substrate transferred ($95 \pm 2.1\%$ of wild-type RdgB-PITP, Fig. 2 B). Thus, the T59A mutation caused the loss of PI transfer activity without affecting PC transfer capability. These data indicated that, while the PI transfer activities of both RdgB-PITP and mammalian PITP α were selectively sensitive to substitutions at T59, the spectrum of substitutions for T59 that were permissive for PI transfer clearly differed between RdgB-PITP and mammalian PITP α .

Expression of the T59E Mutant in *rdgB*² Flies Partially Rescues the ERG Defect without Suppressing the Retinal Degeneration

To dissect the functional contributions of PI and PC transfer activity to RdgB function in vivo, we introduced the T59A substitution in the full-length RdgB protein and T59E into both RdgB-PITP and RdgB. We expressed these T59 mutant proteins in *rdgB*² null flies and assessed their ability to restore a wild-type ERG light response and suppress retinal degeneration. Three independent germline transformed P[*rdgB-T59A*] mutant lines failed to exhibit detectable rescue of either mutant phenotype. A combination of immunoblotting and reverse transcriptase PCR amplification experiments (using poly[A]⁺ mRNA as PCR template) demonstrated that the T59A mutation, while not deleterious to the stability of the RdgB-PITP domain when expressed in *E. coli* (Fig. 2), nevertheless produced an unstable full-length RdgB in flies (data not shown). This destabilization precluded us from examining the effect of the T59A mutation on RdgB function in vivo.

Incorporation of the T59E mutation, however, did not affect the stable expression of either the full-length RdgB (RdgB-T59E) or RdgB-PITP (RdgB-PITP-T59E) (Table I). We compared the ERG light responses between newly

Table I. Levels of Expression of Transgene-encoded Proteins Relative to Wild-type RdgB Expression

Transgene-encoded protein	Percent of wild-type expression
RdgB	$134.0 \pm 8.0\%$
RdgB-T59E	$121.0 \pm 15.0\%$
RdgB-PITP	$24.1 \pm 3.9\%$
RdgB-PITP-T59E	$42.4 \pm 5.8\%$
PITP α -RdgB	$116.4 \pm 13.2\%$

Head protein extracts were produced from *rdgB*² flies expressing one copy of each transgene. Triplicate immunoblots of head extracts were performed using anti-RdgB antisera. Expression levels of RdgB and RdgB variants were quantitated by scanning laser densitometry and compared to the level of RdgB protein in wild-type flies. Because transgene expression was limited to the R1-6 photoreceptor cells, whereas the wild-type control expressed RdgB in the retina, antennae, and various regions in the brain (Vihtelic et al., 1993), the actual levels of transgene-encoded proteins in the retina (relative to wild type) may be higher than indicated.




Figure 3. ERG light responses of *rdgB²* null mutants expressing various transgenes. Dark-reared (<12-h old) wild-type (Oregon-R), *rdgB²*, *rdgB²*; P[*rdgB-pitp*], *rdgB²*; P[*rdgB-pitp-T59E*], *rdgB²*; P[*rdgB-T59E*], *rdgB²*; P[*pitpa*], and *rdgB²*; P[*pitpa-rdgB*] flies were mounted in dim red light for ERGs. After an additional 1-h dark adaptation, an ERG recording was made from a 2-s light stimulus. The flies were then saturated with light for 5 min and given 30 s to dark adapt, followed by another ERG recording to a 2-s light stimulus. An additional 5-min dark adaptation was given before the third ERG recording, again to a 2-s light stimulus. A 10-mV scale is shown below the initial wild-type ERG.

enclosed *rdgB²* flies that expressed or lacked T59E-containing proteins with wild-type flies. Wild-type flies maintained an ERG light-response amplitude of ~25 mV that returned to baseline within 3 s after termination of the light stimulus (Fig. 3). Whereas newly enclosed dark-raised *rdgB²* flies exhibited a wild-type light-response amplitude, the ERGs required on average 1 min to return to baseline after termination of the light stimulus (Fig. 3). Thus, *rdgB²* flies are defective in terminating the ERG light response. The *rdgB²* flies expressing either RdgB-PITP-T59E or RdgB-T59E also exhibited a wild-type light-response amplitude. While the ERG light-response termination in *rdgB²*

flies expressing T59E-containing proteins was significantly faster than *rdgB²* flies, it was still two to three times slower than wild type (Fig. 3).

We also measured the effect of light saturation on the dark recovery of these flies. After 5 min of saturating light, wild-type flies yielded a normal ERG light-response amplitude with <30 s of dark recovery (Fig. 3). By contrast, the *rdgB²* ERG amplitude was essentially eliminated by previous light treatment. Even after 5 min of dark recovery, the *rdgB²* flies lacked any significant light response to subsequent light stimuli (Fig. 3). The *rdgB²* flies required nearly 35 min of dark treatment to regenerate the wild-type ERG light-response amplitude (data not shown). This defective dark recovery was not a result of failure to return to baseline, as *rdgB²* flies displayed responses that decayed to baseline before 35 min (data not shown). Significantly, *rdgB²* flies expressing either RdgB-T59E or RdgB-PITP-T59E elicited a small ERG light response after 30 s of dark treatment and one-third of the original ERG amplitude after 5 min of dark recovery (Fig. 3). A wild-type ERG response was restored to these flies after 6–12 min of dark treatment (not shown). Thus, both the RdgB-PITP-T59E and RdgB-T59E proteins effected a substantial, but nonetheless incomplete, rescue of both the ERG light-response termination and prolonged dark recovery times characteristic of *rdgB²* flies.

The partial restoration of the ERG phenotype in *rdgB²* flies expressing either T59E mutant protein was not translated into any significant protection from retinal degeneration. Retinal sections from 6-d-old *rdgB²*, *rdgB²*; P[*rdgB-pitp-T59E*], and *rdgB²*; P[*rdgB-T59E*] flies raised in a 12 h light/dark cycle were examined by light microscopy (Fig. 4). All three genotypes displayed dark staining, photoreceptor cell bodies, small or missing rhabdomeres, and holes in the retina. Whereas the *rdgB²*; P[*rdgB-pitp-T59E*] retinas were phenotypically similar to the *rdgB²* retinas, the *rdgB²*; P[*rdgB-T59E*] flies exhibited a more severe form of degeneration, with a greater number of holes and ommatidial disorganization. Because the levels of RdgB-T59E and RdgB-PITP-T59E exceeded the level of RdgB-PITP protein needed for suppressing retinal degeneration in *rdgB²* flies (Table I), the inability to prevent degeneration was not a result of insufficient protein levels. As the T59E substitution did not measurably reduce the PI and PC transfer activities of the soluble RdgB-PITP in vitro (Fig. 2 B), the failure of the T59E-containing proteins to fully rescue *rdgB* mutant phenotypes suggests that either the phospholipid transfer activities of RdgB-PITP are more sensitive to perturbations in vivo than in vitro, or that the phospholipid transfer activity is not sufficient to completely fulfill RdgB-PITP's function in vivo.

***PITP α* Cannot Substitute for the PITP Domain of RdgB In Vivo**

To further examine the role of PI and/or PC transfer in prevention of *rdgB* mutant phenotypes, we expressed in *rdgB²* flies either the soluble rat brain PITP α or a full-length chimeric RdgB molecule containing the wild-type rat brain PITP α in place of the RdgB PITP domain (PITP α -RdgB). PITP α shares a 42% primary sequence identity with the RdgB-PITP domain (Vihtelic et al., 1993),




Figure 4. Retinal degeneration histology of *rdgB²* flies expressing various transgenes. Flies of the following genotype were raised in a 12-h light/dark cycle for 6 d: wild-type (A), *rdgB²* (B), *rdgB²*; P[*rdgB-pitp-T59E*] (C), *rdgB²*; P[*rdgB-T59E*] (D), *rdgB²*; P[*pitpα*] (E), and *rdgB²*; P[*pitpα-rdgB*] (F). The retinas were fixed and 1 μ m sections were examined by light microscopy. The *rdgB²*; P[*rdgB-pitp-T59E*] and *rdgB²*; P[*pitpα-rdgB*] flies exhibited the characteristic signs of *rdgB²*-mediated retinal degeneration, while the *rdgB²*; P[*rdgB-T59E*] and *rdgB²*; P[*pitpα*] flies exhibited retinal degeneration phenotypes that were more severe than *rdgB²*. Arrowheads indicate R7 and/or R8 cells for orientation. Bar, 10 μ m.

and exhibits the same phospholipid-transfer substrate spectrum as RdgB-PITP (Fig. 2 B). If the ability to bind and/or transfer PI and PC are the sole essential RdgB functions, then PITP α and/or PITP α -RdgB should fully rescue *rdgB²* mutant phenotypes.

Whereas immunoblot analyses confirmed the stable expression of PITP α and PITP α -RdgB in *rdgB²* flies (data not shown), neither protein restored a wild-type light response to *rdgB²* flies (Fig. 3). The *rdgB²*; P[*pitpα-rdgB*] flies exhibited an ERG response to subsequent light stimulation that was similar to *rdgB²* flies and required nearly

35 min of dark recovery to regenerate the initial light-response amplitude. However, the initial ERG light-response amplitude for *rdgB²*; P[*pitpα-rdgB*] flies was significantly smaller than *rdgB²* flies ($45 \pm 12.5\%$ of the maximal *rdgB²* ERG amplitude, Fig. 3). These data suggest that the PITP α -RdgB protein caused an additional adverse effect on the photoreceptors and did not simply fail to fulfill the requirement for RdgB. The ERG light-response amplitude and dark recovery time for *rdgB²*; P[*pitpα*] flies were nearly identical to *rdgB²* flies. Furthermore, *rdgB²*; P[*pitpα-rdgB*] flies exhibited the morphological hallmarks of *rdgB*-mediated retinal degeneration (Fig. 4), while *rdgB²*; P[*pitpα*] flies exhibited even smaller and fewer number of rhabdomeres. The failure of PITP α and PITP α -RdgB to even partially suppress either the *rdgB²* ERG defects or the retinal degeneration further separates RdgB from the classical PITPs and suggests that the RdgB-PITP domain executes a phototransduction-relevant function that mammalian PITP α cannot.

Dominant Retinal Degeneration Phenotypes in Full-Length RdgB Mutants

We expressed all the above *rdgB* transgenes in a *rdgB⁺* background to determine if they possessed a dominant mutant phenotype. We anticipated that the transgenes that previously failed to rescue the *rdgB²* mutant phenotypes would behave as inactive forms of RdgB and would, therefore, exhibit a fully recessive character. Alternatively, these RdgB variants could disrupt potential protein-protein interactions or compete with the wild-type RdgB for a particular molecule, in which case, they would exhibit a dominant phenotype. We assayed for retinal degeneration by the loss of the deep pseudopupil and further characterized all of the flies using the ERG. We found that *rdgB⁺* flies expressing any of the three soluble proteins (RdgB-PITP, RdgB-PITP-T59E, and PITP α) maintained a wild-type deep pseudopupil and ERG light response for 30 d after eclosion (data not shown). Because the RdgB-PITP-T59E and PITP α proteins lacked a dominant degeneration or ERG phenotype, the failure to completely rescue the *rdgB²* mutant phenotypes must be due to an inability to function like RdgB-PITP, rather than causing degeneration through a novel mechanism.

Surprisingly, the *rdgB-T59E* transgene, which partially restored the ERG light response in *rdgB²* flies, produced a dominant retinal degeneration. Expression of RdgB-T59E resulted in a light-enhanced and dose-dependent loss of the deep pseudopupil in *rdgB⁺* flies. Degeneration of wild-type flies expressing one copy of P[*rdgB-T59E*] was first observed 4 d after eclosion under constant light conditions and at 13 d in a 12-h light/dark cycle (Fig. 5). The deep pseudopupil loss became increasingly prevalent so that essentially all flies in the population had experienced degeneration by 17 d after eclosion in constant light and by 26 d in a 12-h light/dark cycle. Retinal degeneration was never observed, under any circumstances, in wild-type fly controls (not shown). The severity of this dominant retinal degeneration phenotype was proportional to the ratio of P[*rdgB-T59E*]/*rdgB⁺*. Both the onset and rate of degeneration were accelerated in transgenic flies raised in constant light when the P[*rdgB-T59E*]/*rdgB⁺* ratio was 2:1 rather




Figure 5. Time course of the dominant RdgB-T59E and PITP α -RdgB retinal degenerations. The time course of retinal degeneration was determined based upon the loss of the deep pseudopupil. The fraction of flies that retained a wild-type deep pseudopupil was determined daily for 30 d after eclosion. Flies raised in constant light include *rdgB*² (filled triangles), *rdgB*⁺ flies containing two copies of the P[*rdgB-T59E*] transgene (open squares), *rdgB*⁺ flies containing one copy of the P[*rdgB-T59E*] transgene (filled squares), *rdgB*⁺ flies containing two copies of the P[*pitpα-rdgB*] transgene (open circles), and *rdgB*⁺ flies containing one copy of the P[*pitpα-rdgB*] transgene (filled circles). *rdgB*⁺ flies containing one copy of the P[*rdgB-T59E*] transgene were also examined while being raised in a 12-h light/dark cycle (half-filled squares). Each point represents the average of four trials ($n > 25$ flies per trial). The standard deviation is shown as vertical lines.

than 1:1 (Fig. 5). Indeed, *rdgB*⁺ flies that possessed two copies of P[*rdgB-T59E*] lost their deep pseudopupil at essentially the same rate as did *rdgB*² mutant flies raised in identical conditions.

Expression of the PITP α -RdgB protein in *rdgB*² flies resulted in a decreased ERG amplitude relative to that observed in *rdgB*² flies alone, suggesting that the chimeric protein was further detrimental to the *rdgB*² photoreceptor (Fig. 3). We found that PITP α -RdgB expression in a wild-type background produced a dose-dependent dominant loss of the deep pseudopupil (Fig. 5). Flies with two copies of the *pitpα-rdgB* transgene first exhibited deep pseudopupil loss at 2 d after eclosion, while the onset of degeneration was not observed until 8 d after eclosion in flies expressing 1 *pitpα-rdgB* transgene (Fig. 5). Because expression of multiple *rdgB-pitp-T59E* and *pitpα* transgenes produced no dominant effects (data not shown), the RdgB-T59E- and PITP α -RdgB-associated dominant phenotypes seemingly required the carboxy-terminal region of RdgB.

Unique Histological Abnormalities Associated with Dominant-Negative RdgB-T59E and PITP α -RdgB Alleles

We compared the histology of the dominantly degenerat-




Figure 6. Histology of the dominant RdgB-T59E and PITP α -RdgB retinal degenerations. White-eyed versions (*cn bw*) of wild-type (A), *rdgB*² (B), *rdgB*⁺; P[*rdgB-T59E*] (C), and *rdgB*⁺; P[*pitpα-rdgB*] (D) flies were raised in a 12-h light/dark cycle for <6 d after eclosion. The *rdgB*² retinal sections exhibited the reduction and loss of rhabdomeres, formation of holes (long arrows), and condensation of the photoreceptor cell bodies (black arrowheads). The *rdgB*⁺; P[*rdgB-T59E*] flies revealed a reduction in the size of the R1-6 outer rhabdomeres, with few holes appearing in the retina. The R1-6 cell bodies appear to be nearly similar to the wild-type cell bodies. The *rdgB*⁺; P[*pitpα-rdgB*] flies showed signs of degeneration that more closely matched the *rdgB*² mutant flies, with holes (long arrows) appearing in the retinal sections. Additionally, the microvillar rhabdomeres are beginning to unpack (white arrowheads). Bar, 10 μ m.

ing retinas to each other and with *rdgB*² to determine if they could be undergoing the same process. We sectioned retinas from 6-d-old white-eyed (*cn bw*) *rdgB*⁺ flies that either lacked or contained P[*rdgB-T59E*] or P[*pitpα-rdgB*] (Fig. 6). The 6-d-old *rdgB*²; *cn bw* mutant flies (Fig. 6 B) exhibited the characteristic degeneration phenotype of rhabdomere loss, perforations of the retina, and the appearance of optically dense photoreceptor cell bodies. The *rdgB*⁺; P[*rdgB-T59E*] retinas exhibited significantly fewer retinal perforations and dense photoreceptor cell bodies relative to *rdgB*² flies (Fig. 6 C). Most strikingly, the *rdgB*⁺; P[*rdgB-T59E*] retinas lacked mature R1-6 rhabdomeres (Fig. 6 C). Indeed, the rhabdomeres of newly eclosed *rdgB*⁺; P[*rdgB-T59E*] flies were less well developed compared to wild-type controls, and diminished in size as the flies aged (data not shown). This rhabdomere atrophy of photoreceptors R1-6 resembled the hypomorphic *ninaE* mutant phenotype, which results from a significant reduction in rhodopsin expression in photoreceptors R1-6 (Leon-




Figure 7. The dominant RdgB-T59E ERG exhibits reduced photosensitivity. (A–D) ERGs were recorded to a 2-s light stimulus from 1-d-old wild-type (A), *rdgB*⁺; P[*rdgB-T59E*] (B), *rdgB*⁺; P[*pitpα-rdgB*] (C), and *rdgB*⁺; P[*rdgB-pitp*] (D) flies. The light response amplitude of the *rdgB*⁺; P[*rdgB-T59E*] flies is ~60% of either the wild-type or *rdgB*⁺; P[*pitpα-rdgB*] flies. A 10-mV scale is shown at the bottom. All the flies contained the *white*⁺ gene, which conferred the wild-type eye color to remove potential differences due to the expression of *w*⁺ from the P element construct. (E) The ERG amplitudes of wild-type flies (diamonds), *rdgB*⁺; P[*rdgB-T59E*] (squares), *rdgB*⁺; P[*pitpα-rdgB*] (circles), and *rdgB*⁺; P[*rdgB-pitp*] (triangles) were recorded over a range of light intensities. Neutral density units, corresponding to the filters used to modulate the light intensity, are plotted against the light response amplitude. Each point represents a minimum of four independent recordings and the standard deviation is shown as vertical lines.

ard et al., 1992; Kumar and Ready, 1995). The dominant *pitpα-rdgB* degeneration morphology was more similar to the *rdgB*² phenotype, with the most striking defects being the numerous perforations in the retina and the reduction in R1-6 rhabdomere size relative to R7 (Fig. 6 D). Additionally, the R1-6 microvillar rhabdomeres began to exhibit signs of unpacking (Fig. 6 D) that we had not previously observed in any *rdgB* mutants. Thus, while the dominant *rdgB-T59E* mutant phenotype approximated the *ninaE* hypomorphic phenotype, the dominant *pitpα-rdgB* phenotype was morphologically more like the *rdgB* mutant retina with some additional mutant characteristics.

Unique Electrophysiological Abnormalities Associated with Dominant-negative RdgB-T59E and P[*PITPα-RdgB*] Alleles

We examined whether the *rdgB*⁺; P[*rdgB-T59E*] and/or the *rdgB*⁺; P[*pitpα-rdgB*] flies exhibited an electrophysiological defect. All the flies that were tested were newly eclosed and subjected to ERG analysis after a 1-h dark adaptation period. Whereas the light-response amplitudes of *rdgB*⁺; P[*rdgB-T59E*] flies were reduced by ~40% of wild type (Fig. 7, B and A, respectively), the *rdgB*⁺; P[*pitpα-rdgB*] flies were essentially wild type (Fig. 7 C). Wild-type flies expressing multiple copies of either P[*rdgB-pitp*] (Fig. 7 D) or P[*rdgB*⁺] (data not shown), failed to mimic the reduced amplitude observed for *rdgB*⁺; P[*rdgB-T59E*] flies,




Figure 8. The dominant RdgB-T59E protein generates a *ninaE*-like ERG. White-eyed (*cn bw*) versions of wild-type (A), *ninaE*¹¹⁷ (which fails to express any of the R1-6 opsin), (B), *rdgB*⁺; P[*rdgB-T59E*] (C), and *rdgB*⁺; P[*pitpα-rdgB*] flies (D) were tested for the ERG light response using 5 s of either orange (o) or blue light (b) stimulation. Wild-type flies exhibit both a PDA and inactivation of the R1-6 light response by blue light. The *ninaE*¹¹⁷ flies possess neither the PDA nor the blue light inactivation. The *rdgB*⁺; P[*rdgB-T59E*] flies also fail to exhibit a PDA and R1-6 inactivation by blue light. The *rdgB*⁺; P[*pitpα-rdgB*] flies appear to possess a PDA, but they failed to rapidly return to baseline after the subsequent orange light stimulus. A 5-mV scale is shown at the bottom.

demonstrating the specificity of the T59E mutation. Moreover, extension of these ERG analyses to different light intensities failed to produce the wild-type light-response amplitudes in *rdgB*⁺; P[*rdgB-T59E*] flies (Fig. 7 E). These data demonstrated that expression of RdgB-T59E in a *rdgB*⁺ background resulted in a reduced photosensitivity, while P[*PITPα-RdgB*] failed to effect the light-response amplitude.

Because the reduced photosensitivity (like the small rhabdomeres) was consistent with reduced functional rhodopsin, we compared the ERG light responses between white-eyed versions (*cn bw*) of R1-6 rhodopsin-deficient (*ninaE*) flies, *rdgB*⁺; P[*rdgB-T59E*] flies, and *rdgB*⁺; P[*pitpα-rdgB*] flies. Mutations that dramatically reduce functional R1-6 rhodopsin levels exhibit aberrant ERG light responses that lack both the inactivation and the PDA (Fig. 8 B), relative to wild-type flies (Fig. 8 A; Stephenson et al., 1983). The ERGs of white-eyed *rdgB*⁺; P[*rdgB-T59E*] flies closely resembled the *ninaE* mutant phenotype (Fig. 8 C), which suggested that these flies had reduced levels of functional rhodopsin.

We confirmed that RdgB-T59E expression reduced steady-state R1-6 rhodopsin levels to ~1% of wild-type rhodopsin levels (Fig. 9). The reduced rhodopsin levels were not a result of RdgB overexpression, because overexpressing wild-type RdgB (*rdgB*⁺; P[*rdgB*⁺] flies) did not adversely affect rhodopsin levels (117 ± 34% of the wild-




Figure 9. The dominant RdgB-T59E protein preferentially affects rhodopsin protein levels. Immunoblot analyses were performed on head extracts from <1-d-old dark-raised flies of the following genotypes: *rdgB*⁺ (wild-type), *ninaE*¹¹⁷, *rdgB*², *rdgB*⁺; P[*rdgB-T59E*], *rdgB*⁺; P[*pitpα-rdgB*], and *rdgB*⁺; P[*rdgB*⁺]. Two head equivalents per sample were electrophoresed, transferred to nitrocellulose, and incubated with either anti-NinaE polyclonal or anti-Trp polyclonal antisera. Triplicate blots were used to generate the average percent of wild-type protein and standard deviation.

type levels, Fig. 9). Also, *rdgB*² null mutants failed to show reduced steady-state levels of rhodopsin (Fig. 9), which demonstrates that functional RdgB is not required for producing the wild-type rhodopsin levels. We also found that *rdgB*⁺; P[*rdgB-T59E*] flies elicited near wild-type levels of the *trp*-encoded Ca²⁺ channel (Fig. 9), the *dgq*-encoded G α subunit, the *gbe*-encoded G β_e subunit, the *rdgC*-encoded serine/threonine phosphatase, and the *ninaC*-encoded unconventional myosins, suggesting that rhodopsin is selectively sensitive to expression of RdgB-T59E (data not shown). Therefore, the dominant retinal degeneration phenotype, the reduced photoreceptor sensitivity, loss of the PDA, and the gross reduction in mature rhodopsin levels in *rdgB*⁺; P[*rdgB-T59E*] flies all resulted from expression of RdgB-T59E and its interaction with some other protein and/or organelle.

The *rdgB*⁺; P[*pitpα-rdgB*] flies appeared to have a PDA (Fig. 8 D), which was consistent with those flies expressing wild-type levels of rhodopsin (Fig. 9). However, the light response in *rdgB*⁺; P[*pitpα-rdgB*] flies remained inactivated after the conversion of metarhodopsin to rhodopsin by the orange light stimulus (Fig. 8 D). To further examine the effect of PITP α -RdgB on the dark recovery, ERGs to a 2-s light stimulus were recorded from flies, before and 30 s after a 20-min saturating light stimulus. The *rdgB*⁺; P[*pitpα-rdgB*] flies exhibited a significant difference between the initial and final ERG amplitudes (13.0 mV, Fig.




Figure 10. The dominant PITP α -RdgB protein prevents the rapid recovery of the light-response amplitude after prolonged saturating-light treatment. A 2-s ERG light response was recorded from wild-type, *rdgB*⁺; P[*rdgB-T59E*], *rdgB*⁺; P[*pitpα-rdgB*], and *rdgB*⁺; P[*rdgB*⁺] flies, followed by 20 min of saturating light, 30 s of dark recovery, and another 2-s ERG light-response recording. The difference was determined between the first and second light-response recordings. Five flies of each genotype were recorded with the average difference in the light-response amplitude and standard deviation shown. The average initial light-response amplitudes were: wild-type (28.2), *rdgB*⁺; P[*rdgB-T59E*] (19.8), *rdgB*⁺; P[*pitpα-rdgB*] (26.8), and *rdgB*⁺; P[*rdgB*⁺] (22.6 mV). The increased light-response amplitude of *rdgB*⁺; P[*rdgB-T59E*] flies in this figure, relative to Fig. 7, is due to the use of white-eyed flies in this data (*cn bw* background) and wild-type eye colored flies in Fig. 7. The other three genotypes contained some screening pigment.

10), which was 49% of the initial amplitude. Only minor amplitude differences were observed for wild-type (2.6 mV, 9% of initial amplitude), *rdgB*⁺; P[*rdgB-T59E*] (3.8 mV, 19% of initial amplitude), and *rdgB*⁺; P[*rdgB*⁺] (3.1 mV, 14% of initial amplitude) flies (Fig. 10). The *rdgB*⁺; P[*pitpα*] flies, which lacked the dominant degeneration phenotype, were similar to the wild-type controls showing a difference of only 3.1 mV (14% of the initial amplitude) under the same regimen. These data indicated that PITP α -RdgB expression negatively affected the recovery phase of the light response in an otherwise wild-type photoreceptor cell. It is unclear if there is a direct relationship between this electrophysiological defect and the retinal degeneration. However, both of these PITP α -RdgB dominant phenotypes are similar to very mild *rdgB* mutant phenotypes, which suggests that PITP α -RdgB could be interacting directly with RdgB or competing for a molecule to reduce the wild-type RdgB activity.

Discussion

In this manuscript, we detail a functional analysis of RdgB participation in the *Drosophila* phototransduction cascade and a description of the physiological function of a meta-

zoan PITP. Herein, we report four novel aspects of RdgB function in vivo. These include: (a) the demonstration that the RdgB-PITP domain houses all of the RdgB functions related to its role in phototransduction; (b) the demonstration that RdgB functions to effect a proper termination of the light response and dark recovery of the photoreceptor cell; (c) the finding that the activities of the RdgB-PITP domain required for phototransduction cannot be satisfied by other PITPs that possess in vitro PI and/or PC transfer capabilities, and (d) the generation of dominant *rdgB* mutations that suggest an underlying complexity to the mechanism of RdgB function and its role in the photoreceptor cell.

Expression of the RdgB-PITP Domain as a Soluble Polypeptide Rescues *rdgB*² Null Mutants

Expression of RdgB-PITP as a soluble domain not only restored a wild-type ERG light response to *rdgB*² flies, but also fully suppressed the hallmark retinal degeneration phenotype of these mutant flies (Fig. 1 D). These data clearly demonstrate that the RdgB activities relevant for its involvement in the *Drosophila* phototransduction cascade and photoreceptor cell maintenance are restricted to that amino-terminal PITP domain; an entirely unexpected result given that the 281-residue RdgB-PITP represents only 27% of the 1,054-residue RdgB. Moreover, because expression of RdgB-PITP as a soluble polypeptide altered its intracellular disposition from an integral membrane protein to a cytosolic species, we conclude that covalent tethering of RdgB-PITP to an integral membrane protein domain is not an obligate requirement for the efficient function of RdgB-PITP in phototransduction. This now raises the question of what the biological function of the large COOH-terminal RdgB domain is. The identification of RdgB homologues in mammalian retina that exhibit the same modular domain arrangement of *Drosophila* RdgB (i.e., NH₂-terminal PITP domain fused to a large COOH-terminal domain), share significant primary sequence identity to full-length RdgB over their entirety, and can restore the wild-type ERG light response and prevent retinal degeneration when expressed in *rdgB*² mutant flies (Chang et al., 1997), suggests that cells derive some advantage from this modular arrangement.

RdgB Is Required for Proper Termination of the Light Response and Dark Recovery of the Photoreceptor Cell

Newly eclosed, dark-adapted *rdgB*² null mutant flies displayed ERG light responses that had wild-type amplitudes, but were defective in photoreceptor cell repolarization after termination of the light stimulus (Fig. 3). This indicates that the RdgB PITP domain is not essential for establishing an initial PIP₂ pool in the rhabdomere, though its requirement for regeneration of this pool remains a formal possibility. We do not presently favor the idea that RdgB functions in regeneration of such a phosphoinositide pool because RdgB levels do not correlate closely with the gain of the light response. The amplitude size of the light response is controlled by levels of PI as observed by flies overexpressing the rate-limiting enzyme for PI biosynthesis, eye-CDP-diacylglycerol synthase (CDS), produce greater ERG amplitudes in response to light stimula-

tion (Wu et al., 1995). Correspondingly, mutants that are defective in eye-CDS activity exhibit smaller light-response amplitudes (Wu et al., 1995).

If RdgB has a critical role in replenishing rhabdomeric PI, then light saturation treatment, and therefore consumption of rhabdomeric PI by its conversion into PLC substrate, would reduce or eliminate subsequent light responses in *rdgB* mutant flies. Indeed, *rdgB*² and all *rdgB*² mutant flies containing nonrescuing transgenes produced wild-type amplitude responses 6 to 35 min after light saturation treatment, as compared to 30 s for wild-type flies (Fig. 3). Clearly, the analyses in *rdgB*² null flies are consistent with a role for RdgB in restoring the cell to a state competent for subsequent light stimulation. However, if RdgB activity is critical in replenishing rhabdomeric PI, then overexpression of RdgB, as observed for eye-CDS, should increase the amplitude of light responses in *rdgB*⁺ flies. Wild-type flies expressing multiple copies of P[*rdgB-pitp*] (Fig. 7 D) or P[*rdgB*⁺] (data not shown) failed to produce ERG light-response amplitudes that were significantly larger than wild type. Additionally, we did not detect any significant differences in the ERG light-response amplitude of *rdgB*⁺ or *rdgB*⁺; P[*rdgB*⁺] flies that were recorded over a background light (data not shown). We expected the background light would deplete the PIP₂ stores, which would result in either slower kinetics or a smaller light-response amplitude in *rdgB*⁺ flies that would be compensated for by the increased copy number of RdgB. Since overexpression of RdgB did not affect the gain of the cascade, either protein-mediated PI transfer is not limiting in phototransduction activation or RdgB does not function in this respect. Additionally, the lack of rescue by RdgB-PITP-T59E and PITP α suggests that the RdgB-PITP possesses activities separate from those characterized biochemically. Our collective data fail to support the speculation that RdgB functions in the transport of PI to specific rhabdomeric pools for phototransduction-driven consumption (Hurley, 1995; Zuker, 1996).

It remains unresolved as to how RdgB promotes termination of the light response and rapid dark recovery of the photoreceptor cell, though it is clear that all of these functions reside within the RdgB-PITP domain. This domain contains multiple consensus PKC phosphorylation sequences. One site is T59, which when mutated to T59E compromises an RdgB activity in vivo, without affecting PI transfer in vitro. As RdgB's placement in phototransduction is downstream of PKC, and since PKC is required for the wild-type rapid inactivation of the photoresponse following light cessation, the RdgB-dependent inactivation of the light response may be regulated by a phosphorylation/dephosphorylation cycle (Ranganathan et al., 1991; Smith et al., 1991; Hardie et al., 1993). If RdgB is regulated at T59 by phosphorylation in vivo, then the failure of PITP α to prevent *rdgB* phenotypes could result from PKC phosphorylation. Unlike RdgB-PITP, the T59E mutation essentially eliminated PI transfer in PITP α in vitro (Alb et al., 1995). In the fly, the phosphorylation of PITP α may lock the protein in a PI transfer incompetent state. The failure of the PI transfer competent RdgB-PITP-T59E protein to rescue may have resulted from loss of regulation at T59 pertinent to another RdgB-PITP activity. Thus, the failure of the RdgB-PITP-T59E and PITP α to rescue *rdgB*²

mutant flies does not unambiguously rule out a role for RdgB-associated PI and/or PC transfer activity in vivo. Rather, the data indicate that other activities present in RdgB-PITP are required for wild-type RdgB function.

RdgB Is Not Simply an Integral Membrane PITP

RdgB-PITP catalyzes the efficient transfer of both PI and PC between membrane bilayers in vitro, but is unable to catalyze intermembrane transfer of SM (Fig. 2 B). In this regard, RdgB-PITP is biochemically analogous to mammalian PITP α , and not to PITP β . Because RdgB-PITP represents the sole RdgB domain essential for normal function of the fly visual cycle (Fig. 1 D), it is of primary interest to determine how the PI and/or PC transfer activities of RdgB-PITP contribute to function. Although our initial efforts to selectively inactivate the PI transfer activity of RdgB (using the T59A mutation) and assess function in vivo were unsuccessful, our data with RdgB-PITP-T59E and PITP α suggest that RdgB-PITP does not simply function to transfer phospholipids in the photoreceptor cell. Neither stable expression of RdgB-PITP-T59E nor PITP α , both of which are fully active for PI and/or PC transfer in vitro, could completely rescue the *rdgB*² retinal degeneration phenotype (Fig. 4), contrary to expectations that catalysis of PI and/or PC transfer was the sole function of RdgB-PITP. It remains entirely possible that phospholipid binding serves as a molecular switch through which a second effector activity of RdgB-PITP is regulated, as has been proposed for Sec14p (McGee et al., 1994; Skinner et al., 1995; Kearns et al., 1997). It is similarly possible that the PI and/or PC bound states of RdgB-PITP may regulate an activity in the photoreceptor cell that is not responsive to mammalian PITP α presented in the RdgB context.

Although RdgB-T59E and PITP α -RdgB were both unable to phenotypically rescue *rdgB*²-associated retinal degeneration, these polypeptides nonetheless exhibited significant phenotypic differences at the level of the ERG light response. Expression of the PITP α -RdgB chimera in *rdgB*² flies failed to effect any detectable alleviation of the *rdgB*² light-response termination and dark-recovery defects, while also significantly reducing the amplitude of the light response relative to *rdgB*² (Fig. 3). This suggests that the PITP α -RdgB chimera was interacting with either a molecule and/or organelle to further reduce the photoreceptor's ability to respond to light. On the other hand, RdgB-T59E expression significantly (but not completely) improved proper termination of the light response and shortened the dark recovery time required for regeneration of a normal light response following a saturating light stimulus to *rdgB*² flies (Fig. 3). Thus, while RdgB-T59E exhibited partial function in the ERG assays, the data further suggest that either: (a) the retinal degeneration phenotype is a more sensitive indicator of RdgB-PITP perturbation than the ERG light response, or (b) that the aberrant ERGs recorded are not intimately related to the mechanism of retinal degeneration.

Dominant Mutant Forms of the RdgB-PITP Domain Reveal Unexpected and Diverse Degeneration Phenotypes

Expression of either RdgB-T59E or PITP α -RdgB in *rdgB*⁺

flies exerted powerful effects that were manifested in dominant retinal degeneration phenotypes and proved informative with regard to previously unappreciated aspects of RdgB function. In the case of RdgB-T59E, the associated dominant effects operated through the reduction of rhodopsin protein levels in photoreceptors R1-6; the primary evidence involved direct measurements of steady-state rhodopsin levels in mutant flies (Fig. 9). Additional observations include: (a) the reduced photosensitivity of *rdgB*⁺; P[*rdgB-T59E*] flies (Figs. 7, B and E); (b) the aberrant ERG responses of *rdgB*⁺; P[*rdgB-T59E*] flies that bore the signature of R1-6 opsin insufficiency (Fig. 8 C); and (c) the observation that *rdgB*⁺; P[*rdgB-T59E*] retinas exhibited degeneration morphologies closely resembling those associated with hypomorphic *ninaE* alleles (Fig. 6 C). However, this dominant mutation did not exert a general effect on the biogenesis and/or transport of rhabdomeric proteins because the levels of the *trp*-encoded Ca²⁺ channel (Fig. 9), G α q, G β e, NinaC p174, and the *rdgC*-encoded serine/threonine phosphatase were unaffected (data not shown). Because RdgB is not required for rhodopsin expression, as young *rdgB*² null mutants exhibit near wild-type steady-state levels of rhodopsin on immunoblots (Fig. 9), the specificity and mechanism of rhodopsin depletion by RdgB-T59E is unclear.

The dominant retinal degeneration phenotype associated with expression of the PITP α -RdgB chimeric molecule was distinct from that of RdgB-T59E. Retinal sections of *rdgB*⁺; P[*pitp α -rdgB*] flies bore morphological hallmarks of degeneration (Fig. 6) that more closely resembled those associated with *rdgB*² rather than *ninaE*; which is consistent with the wild-type steady-state rhodopsin levels in *rdgB*⁺; P[*pitp α -rdgB*] flies (Fig. 9). Additionally, the *rdgB*⁺; P[*pitp α -rdgB*] flies exhibited an ERG dark-recovery defect that was similar, but much more subtle, to *rdgB*² mutants (Fig. 10). However, to detect the loss of the ERG light-response amplitude in *rdgB*⁺; P[*pitp α -rdgB*] flies, the flies were exposed to a very prolonged saturating light stimulus (20 min), rather than the 5 min used in Fig. 3. Under these conditions, the *rdgB*⁺; P[*pitp α -rdgB*] flies exhibited a 49% reduction in the light-response amplitude compared to only a 9–19% reduction in three other genotypes. The *rdgB*⁺; P[*pitp α -rdgB*] flies also exhibited a very slow rate of light-response inactivation to the second orange light stimulus (Fig. 8 D). Thus, PITP α -RdgB appeared to antagonize RdgB activity in the photoreceptor cell.

Taken together, the data indicate that RdgB-T59E strongly interferes with rhodopsin biogenesis, while PITP α -RdgB potentially interferes with the activity of wild-type RdgB. Both of these dominant phenotypes suggest that RdgB physically interacts with at least one other component of the *Drosophila* phototransduction cascade. The failure to observe any dominant phenotypes associated with either RdgB-PITP-T59E or PITP α , suggests that the integral membrane nature of these proteins must be critical for these molecular interactions. Because RdgB-T59E and PITP α -RdgB have dramatically different effects in the cell, RdgB's large carboxy-terminus (which is common to both proteins) likely places the mutant protein in the proper spatial environment and the different attached PITP domains confer the phenotypes. While the identities of RdgB-interacting proteins remain unknown, the recent identification of novel

mutations that suppress *rdgB* defects may provide relevant clues (Paetkau, D., V. Elagin, and D.R. Hyde, unpublished data). In addition, these findings provide the first demonstration that dominant mutant forms of PITP molecules can be generated and that these can yield informative phenotypes when expressed in a eukaryotic cell.

We thank Joseph O'Tousa (University of Notre Dame) for providing the rabbit anti-NinaE polyclonal antiserum, Craig Montell (Johns Hopkins University) for the rabbit anti-Trp polyclonal antiserum, and Charles Zuker (University of California at San Diego) for providing the mouse anti-Gbe ascites. We gratefully acknowledge Yan Cheng (University of Notre Dame) for microinjection of embryos and Douglas McAbee (University of Notre Dame) for critical reading of the manuscript.

This work was supported by grants from the National Institutes of Health to D.R. Hyde (EY08058) and V.A. Bankaitis (GM44530), and a grant from the American Cancer Society (BE232) to V.A. Bankaitis.

Received for publication 10 March 1997 and in revised form 15 July 1997.

References

- Aitken, J.F., G.P.H. van Heusden, M. Temkin, and W. Dowhan. 1990. The gene encoding the phosphatidylinositol transfer protein is essential for cell growth. *J. Biol. Chem.* 265:4711–4717.
- Alb, J.G., A. Gedvilaite, R.T. Cartee, H.B. Skinner, and V.A. Bankaitis. 1995. Mutant rat phosphatidylinositol/phosphatidylcholine transfer proteins specifically defective in phosphatidylinositol transfer: implications for the regulation of phospholipid transfer activity. *Proc. Natl. Acad. Sci. USA.* 92:8826–8830.
- Alb, J.G., M.A. Kearns, and V.A. Bankaitis. 1996. Phospholipid metabolism and membrane dynamics. *Curr. Opin. Cell Biol.* 8:534–541.
- Ashburner, M. 1989. *Drosophila: A Laboratory Handbook*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 1331 pp.
- Bankaitis, V.A., D.E. Malehorn, S.D. Emr, and R. Greene. 1989. The *Saccharomyces cerevisiae* *SEC14* gene encodes a cytosolic factor that is required for transport of secretory proteins from the yeast Golgi complex. *J. Cell Biol.* 108:1271–1281.
- Bankaitis, V.A., J.F. Aitken, A.E. Cleves, and W. Dowhan. 1990. An essential role for a phospholipid transfer protein in yeast Golgi function. *Nature (Lond.)* 347:561–562.
- Bloomquist, B.T., R.D. Shortridge, S. Schneuwly, M. Perdew, C. Montell, H. Stellar, G. Rubin, and W.L. Pak. 1988. Isolation of a putative phospholipase C gene of *Drosophila*, *norPA*, and its role in phototransduction. *Cell* 54:723–733.
- Chang, J.T., S. Milligan, Y. Li, C.E. Chew, J. Wiggs, N.G. Copeland, N.A. Jenkins, P.A. Campochiaro, D.R. Hyde, and D.J. Zack. 1997. Mammalian homologue of *Drosophila* *retinal degeneration B* rescues the mutant fly phenotype. *J. Neurosci.* 17:5881–5890.
- Cleves, A., T. McGee, and V. Bankaitis. 1991. Phospholipid transfer proteins: a biological debut. *Trends Cell Biol.* 1:30–34.
- Cunningham, E., G.M.H. Thomas, A. Ball, I. Hiles, and S. Cockcroft. 1995. Phosphatidylinositol transfer protein dictates the rate of inositol trisphosphate production by promoting synthesis of PIP₂. *Curr. Biol.* 5:775–783.
- Franceschini, N. 1972. Pupil and pseudopupil in the compound eye of *Drosophila*. In *Information Processing in the Visual Systems of Arthropods*. R. Wehners, editor. Springer-Verlag, Berlin, Germany. 75–82.
- Hamilton, B.A., D.J. Smith, K.L. Mueller, A.W. Kerrebrock, R.T. Bronson, V. van Berkel, M.J. Daly, L. Kruglyak, M.P. Reeve, J.L. Nemhauser, et al. 1997. The *vibrator* mutation causes neurodegeneration via reduced expression of PITP: positional complementation cloning and extragenic suppression. *Neuron* 18:711–722.
- Hardie, R.C., A. Peretz, E. Susstoby, A. Romglas, S.A. Bishop, Z. Selinger, and B. Minke. 1993. Protein kinase C is required for light adaptation in *Drosophila* photoreceptors. *Nature (Lond.)* 363:634–637.
- Harris, W.A., and W.S. Stark. 1977. Hereditary retinal degeneration in *Drosophila melanogaster*: a mutant defect associated with the phototransduction process. *J. Gen. Physiol.* 69:261–291.
- Hay, J.C., and T.F.J. Martin. 1993. Phosphatidylinositol transfer protein is required for ATP-dependent priming of Ca²⁺-activated secretion. *Nature (Lond.)* 366:572–575.
- Hay, J.C., and T.F.J. Martin. 1995. ATP-dependent inositol phosphorylation required for Ca²⁺-activated secretion. *Nature (Lond.)* 374:173–177.
- Hurley, J.B. 1995. Phospholipids in action. *Nature (Lond.)* 373:194–195.
- Kearns, B.G., T.P. McGee, P. Maying, A. Gedvilaite, S.E. Phillips, S. Kagiwada, and V.A. Bankaitis. 1997. Essential role for diacylglycerol in protein transport from the yeast Golgi complex. *Nature (Lond.)* 387:101–105.
- Kumar, J.P., and D.F. Ready. 1995. Rhodopsin plays an essential structural role in *Drosophila* photoreceptor development. *Development (Camb.)* 121:4359–4370.
- Kunkel, T.A., J.D. Roberts, and R.A. Zakour. 1987. Rapid and efficient site-specific mutagenesis without phenotypic selection. *Methods Enzymol.* 154:367–382.

- Laemmli, U.K. 1970. Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature (Lond.)* 227:680–685.
- Lee, Y.J., S. Shah, E. Suzuki, T. Zars, P.M. O'Day, and D.R. Hyde. 1994. The *Drosophila* *dgg* gene encodes a G α protein that mediates phototransduction. *Neuron* 13:1143–1157.
- Leonard, D.S., V.D. Bowman, D.F. Ready, and W.L. Pak. 1992. Degeneration of photoreceptors in rhodopsin mutants of *Drosophila*. *J. Neurobiol.* 23:605–626.
- Matsumoto-Suzuki, E., K. Hirosawa, and Y. Hotta. 1989. Structure of the sub-rhabdomeric cisternae in the photoreceptor cells of *D. melanogaster*. *J. Neurocytol.* 18:87–93.
- McGee, T.P., H.B. Skinner, E.A. Whitters, S.A. Henry, and V.A. Bankaitis. 1994. A phosphatidylinositol transfer protein controls the phosphatidylcholine content of yeast Golgi membranes. *J. Cell Biol.* 124:273–287.
- Miller, J. 1972. *Experiments in Molecular Genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY. 466 pp.
- Minke, B., C.T. Rubinstein, I. Sahly, S. Bar-Nachum, R. Timberg, and Z. Selinger. 1990. Phorbol ester induces photoreceptor-specific degeneration in a *Drosophila* mutant. *Proc. Natl. Acad. Sci. USA.* 87:113–117.
- Mismer, D., and G.M. Rubin. 1987. Analysis of the promoter of the *ninaE* opsin gene in *Drosophila melanogaster*. *Genetics* 116:565–578.
- Ohashi, M., K.J. De Vries, R. Frank, G. Snoek, V. Bankaitis, K. Wirtz, and W.B. Huttner. 1995. A role for phosphatidylinositol transfer protein in secretory vesicle formation. *Nature (Lond.)* 377:544–547.
- O'Tousa, J.E., W. Baehr, R.L. Martin, J. Hirsh, W.L. Pak, and M.L. Applebury. 1985. The *Drosophila* *ninaE* gene encodes an opsin. *Cell* 40:839–850.
- Ozaki, K., H. Nagatani, M. Ozaki, and F. Tokunaga. 1993. Maturation of major *Drosophila* rhodopsin, *ninaE*, requires chromophore 3-hydroxyretinal. *Neuron* 10:1113–1119.
- Ranganathan, R., G.L. Harris, C.F. Stevens, and C.S. Zuker. 1991. A *Drosophila* mutant defective in extracellular calcium-dependent photoreceptor deactivation and rapid desensitization. *Nature (Lond.)* 354:230–232.
- Rubinstein, C.T., S. Bar-Nachum, Z. Selinger, and B. Minke. 1989. Chemically induced retinal degeneration in the *rdgB* (*retinal degeneration B*) mutant of *Drosophila*. *Vis. Neurosci.* 2:541–551.
- Sanger, F., S. Nicklen, and A.R. Coulson. 1977. DNA sequencing with chain terminating inhibitors. *Proc. Natl. Acad. Sci. USA.* 74:5463–5467.
- Skinner, H.B., J.G. Alb, Jr., E.A. Whitters, G.M. Helmkamp, and V.A. Bankaitis. 1993. Phospholipid transfer activity is relevant to but not sufficient for the essential function of the yeast *SEC14* gene product. *EMBO (Eur. Mol. Biol. Organ.) J.* 12:4775–4784.
- Skinner, H.B., T.P. McGee, C.R. McMaster, M.R. Fry, R.M. Bell, and V.A. Bankaitis. 1995. The *Saccharomyces cerevisiae* phosphatidylinositol-transfer protein effects a ligand-dependent inhibition of choline-phosphate cytidylyltransferase activity. *Proc. Natl. Acad. Sci. USA.* 92:112–116.
- Smith, D.P., R. Ranganathan, R.W. Hardy, J. Marx, T. Tsuchida, and C.S. Zuker. 1991. Photoreceptor deactivation and retinal degeneration mediated by a photoreceptor-specific protein kinase C. *Science (Wash. DC)* 254:1478–1484.
- Stark, W.S., D.-M. Chen, M.A. Johnson, and K.L. Frayer. 1983. The *rdgB* gene of *Drosophila*: retinal degeneration in different alleles and inhibition by *norPA*. *J. Insect Physiol.* 29:123–131.
- Stephenson, R.S., J. O'Tousa, N.J. Scavarda, L.L. Randall, and W.L. Pak. 1983. *Drosophila* mutants with reduced rhodopsin content. In *The Biology of Photoreception*. D. Cosens and D. Vince-Price, editors. Cambridge University Press, Cambridge, England. pp. 477–501.
- Suzuki, E., and K. Hirosawa. 1991. Immunoelectron microscopic study of the opsin distribution in the photoreceptor cell of *Drosophila melanogaster*. *JEM (J. Electron Microsc.)* 40:187–192.
- Suzuki, E., and K. Hirosawa. 1994. Immunolocalization of a *Drosophila* phosphatidylinositol transfer protein (*rdgB*) in normal and *rdgA* mutant photoreceptor cells with special reference to the subrhabdomeric cisternae. *JEM (J. Electron Microsc.)* 43:183–189.
- Thomas, G.M.H., E. Cunningham, A. Fensome, A. Ball, N.F. Totty, O. Truong, J.J. Hsuan, and S. Cockcroft. 1993. An essential role for phosphatidylinositol transfer protein in phospholipase C-mediated inositol lipid signaling. *Cell* 74:919–928.
- Vihtelic, T.S., D.R. Hyde, and J.E. O'Tousa. 1991. Isolation and characterization of the *Drosophila* *retinal degeneration B* (*rdgB*) gene. *Genetics* 127:761–768.
- Vihtelic, T.S., M. Goebel, S. Milligan, J.E. O'Tousa, and D.R. Hyde. 1993. Localization of *Drosophila* *retinal degeneration B*, a membrane-associated phosphatidylinositol transfer protein. *J. Cell Biol.* 122:1013–1022.
- Walz, B. 1982. Calcium-sequestering smooth endoplasmic reticulum in retinal cells of the blowfly. *J. Ultrastruct. Res.* 81:240–248.
- Wirtz, K.W.A. 1991. Phospholipid transfer proteins. *Annu. Rev. Biochem.* 60:73–99.
- Wu, L., B. Niemeyer, N. Colley, M. Socolich, and C.S. Zuker. 1995. Regulation of PLC-mediated signalling in vivo by CDP-diacylglycerol synthase. *Nature (Lond.)* 373:216–222.
- Zars, T., and D.R. Hyde. 1996. *rdgE*: a novel retinal degeneration mutation in *Drosophila melanogaster*. *Genetics* 144:127–138.
- Zuker, C.S. 1996. The biology of vision in *Drosophila*. *Proc. Natl. Acad. Sci. USA.* 93:571–576.
- Zuker, C.S., A.F. Cowman, and G.M. Rubin. 1985. Isolation and structure of a rhodopsin gene from *D. melanogaster*. *Cell* 40:851–858.