

## Synthesis of $\gamma$ -Aminobutyric Acid by Lactic Acid Bacteria Isolated from a Variety of Italian Cheeses<sup>∇</sup>

S. Siragusa, M. De Angelis,\* R. Di Cagno, C. G. Rizzello, R. Coda, and M. Gobbetti

*Dipartimento di Protezione delle Piante e Microbiologia Applicata, Università degli Studi di Bari, Bari 70126, Italy*

Received 12 May 2007/Accepted 10 September 2007

The concentrations of  $\gamma$ -aminobutyric acid (GABA) in 22 Italian cheese varieties that differ in several technological traits markedly varied from 0.26 to 391 mg kg<sup>-1</sup>. Presumptive lactic acid bacteria were isolated from each cheese variety (total of 440 isolates) and screened for the capacity to synthesize GABA. Only 61 isolates showed this activity and were identified by partial sequencing of the 16S rRNA gene. Twelve species were found. *Lactobacillus paracasei* PF6, *Lactobacillus delbrueckii* subsp. *bulgaricus* PR1, *Lactococcus lactis* PU1, *Lactobacillus plantarum* C48, and *Lactobacillus brevis* PM17 were the best GABA-producing strains during fermentation of reconstituted skimmed milk. Except for *L. plantarum* C48, all these strains were isolated from cheeses with the highest concentrations of GABA. A core fragment of glutamate decarboxylase (GAD) DNA was isolated from *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48 by using primers based on two highly conserved regions of GAD. A PCR product of ca. 540 bp was found for all the strains. The amino acid sequences deduced from nucleotide sequence analysis showed 98, 99, 90, and 85% identity to GadB of *L. plantarum* WCFS1 for *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48, respectively. Except for *L. lactis* PU1, the three lactobacillus strains survived and synthesized GABA under simulated gastrointestinal conditions. The findings of this study provide a potential basis for exploiting selected cheese-related lactobacilli to develop health-promoting dairy products enriched in GABA.

During the last decade, fundamental studies have opened a new field of research dealing with bioactive or biogenic substances derived from foods. Numerous definitions have been given for bioactive compounds, and one of the most appropriate could be the following: components of consumption-ready foods which may exert a regulatory activity in the human organism, irrespective of their nutritive functions (12).  $\gamma$ -Aminobutyric acid (GABA), a nonprotein amino acid, possesses well-known physiological functions such as neurotransmission, induction of hypotension, and diuretic and tranquilizer effects (24, 53). Treatments for sleeplessness, depression, and autonomic disorders (39); treatment for chronic alcohol-related symptoms (37); and stimulation of immune cells (36) have also been related to the administration of GABA. Recently, GABA has been considered as a strong secretagogue of insulin from the pancreas (1) that may prevent diabetic conditions (16). Owing to these physiological functions, several functional foods are manufactured: GABA-enriched green tea by anaerobic or cyclic treatments of tea leaves or shoots (38); GABA-enriched rice germ by soaking in water (46); GABA-enriched brown rice by high-pressure treatment and germination (26, 35); GABA-enriched germinated wheat through the activity of endogenous enzymes (30); and GABA-enriched fermented beverages such as tempeh-like beverage (3), dairy products

(18, 20, 39), and red-mold rice containing the *Monascus* fungus (44).

GABA is synthesized by glutamate decarboxylase (GAD) (EC 4.1.1.15), a pyridoxal 5'-phosphate-dependent enzyme that catalyzes the irreversible  $\alpha$ -decarboxylation of L-glutamate to GABA. GAD is widely distributed among eukaryotes and prokaryotes (51). Several reports (27, 32–34, 40, 42, 52) have shown the presence of GAD in lactic acid bacteria also. Overall, GABA may confer resistance to bacterial cells under acidic conditions (for a review see reference 5), and the GAD decarboxylation process has also been coupled with energy synthesis in *Lactobacillus* sp. strain E1 (19).

Lactic acid bacteria are largely used in a variety of fermented foods, especially for the manufacture of dairy products with functional and probiotic properties (28). The screening of lactic acid bacteria based on their capacity for synthesizing GABA may open new perspectives on production of GABA-enriched dairy products. During milk fermentation and proteolysis, a high level of L-glutamate may be theoretically liberated, since native caseins contain a high proportion of this amino acid. To our knowledge, just a few reports have considered cheeses as a potential vehicle for GABA (32–34). Overall, the protocol of manufacture, type of primary starters, and, especially, the autochthonous microbiota and ripening conditions may affect the concentration of GABA in cheese.

The aims of this study were to (i) screen 22 Italian cheese varieties, which differ for several technological traits, for their concentrations of GABA; (ii) select cheese-related GABA-producing lactic acid bacteria to be used as starters; (iii) determine the partial sequence of the GAD genes from selected GABA-producing strains; and (iv) assay the synthesis of GABA under simulated gastrointestinal (GI) conditions.

\* Corresponding author. Mailing address: Dipartimento di Protezione delle Piante e Microbiologia Applicata, Facoltà di Scienze Biologiche di Bari, Via G. Amendola 165/a, 70126 Bari, Italy. Phone: 39 080 5442948. Fax: 39 080 5442911. E-mail: m.deangelis@agr.uniba.it.

<sup>∇</sup> Published ahead of print on 21 September 2007.

## MATERIALS AND METHODS

**Cheeses.** Twenty-two Italian cheese varieties were considered in this study. Cheese varieties were manufactured by using milk from different cows (Parmigiano Reggiano, Barricato San Martino, Vento d'Estato, Ubricaco di Raboso, Caciocavallo, Gorgonzola, and Crescenza), buffaloes (Mozzarella), sheep (Pecorino Piemontese, Pecorino Marchigiano, Pecorino Umbro, Pecorino del Reatino, Pecorino Sardo, Pecorino di Filiano, Pecorino del Tarantino, and Pecorino Leccese), sheep and cows in combination (Canestrato Pugliese and Caciotta di Urbino), or goats (Caprino di Valsassina, Caprino di Cavalese, Flor di Capra, and Capritilla) which was used raw (Parmigiano Reggiano, Barricato San Martino, Vento d'Estato, Ubricaco di Raboso, Canestrato Pugliese, Casciotta di Urbino, Pecorino Piemontese, Pecorino Marchigiano, Pecorino del Reatino, Pecorino Sardo, Pecorino di Filiano, Pecorino del Tarantino, and Pecorino Leccese) or pasteurized (all other cheeses). Other main technological traits were as follows: (i) use of primary natural starters (Parmigiano Reggiano, Caciocavallo, Mozzarella, Canestrato Pugliese, Pecorino Sardo, and all goat cheeses) or commercial starters (Gorgonzola and Pecorino Umbro); (ii) use of calf powder or liquid (Parmigiano Reggiano, Gorgonzola, Crescenza, Mozzarella, Canestrato Pugliese, Casciotta di Urbino, and all goat cheeses) or lamb (Caciocavallo) or calf (all other cheeses) paste; (iii) cooking (Parmigiano Reggiano) or stretching (Caciocavallo and Mozzarella); and (iv) very long ripening (18 months; Parmigiano Reggiano), long ripening (8 months; Canestrato Pugliese), medium ripening (2 to 5 months; Barricato San Martino, Vento d'Estato, Ubricaco di Raboso, Caciotta di Urbino, Caciocavallo, and all sheep and goat cheeses), or no ripening to short ripening (0 to 45 days) (Mozzarella, Crescenza, and Gorgonzola).

Cheeses were supplied in triplicate (different batches from the same factory) by local cheese markets and were stored at 4°C for a few hours before analyses. All the analyses were carried out at least in duplicate for each batch of cheese (total of six analyses for each cheese variety).

**Compositional analysis.** Moisture and pH were determined as reported by the International Dairy Federation (22, 23). Soluble and total nitrogen (N) were determined by the micro-Kjeldahl method (21).

**Concentrations of GABA in cheeses.** Thirty grams of cheese was suspended in 90 ml of 50 mM phosphate buffer, pH 7.0, and treated for 10 min with a stomacher (PBI International, Milan, Italy). The suspension was kept at 40°C for 1 h under gentle stirring (150 rpm) and centrifuged at  $3,000 \times g$  for 30 min at 4°C. The supernatant was filtered through Whatman no. 2 paper, and the pH of the extract was adjusted to 4.6 using 1 N HCl. The suspension was centrifuged at  $10,000 \times g$  for 10 min. Finally, the supernatant was filtered through a Millex-HA 0.22- $\mu\text{m}$ -pore-size filter (Millipore Co., Bedford, MA). Total and individual free amino acids (FAAs) contained in the pH 4.6-soluble nitrogen fraction were analyzed by a Biochrom 30 series amino acid analyzer (Biochrom Ltd., Cambridge Science Park, England) with a sodium cation-exchange column (20 by 0.46 cm [inner diameter]). A mixture of amino acids at known concentrations (Sigma Chemical Co., Milan, Italy) was added with cysteic acid, methionine sulfoxide, methionine sulfone, tryptophan, ornithine, glutamic acid, and GABA and used as standard. Proteins and peptides in the samples were precipitated by addition of 5% (vol/vol) cold solid sulfosalicylic acid, holding the samples at 4°C for 1 h, and centrifuging them at  $15,000 \times g$  for 15 min. The supernatant was filtered through a 0.22- $\mu\text{m}$ -pore-size filter and diluted, when necessary, with sodium citrate (0.2 M, pH 2.2) loading buffer. Amino acids were postcolumn derivatized with ninhydrin reagent and detected by absorbance at 440 (proline and hydroxyproline) or 570 (all the other amino acids) nm.

**Enumeration and isolation of lactic acid bacteria.** Samples (20 g) of cheeses were diluted in 180 ml of sodium citrate (2%, wt/vol) solution and homogenized with a Lab-Blender 400 stomacher (PBI International Milan, Italy). Serial dilutions were made in one-quarter-strength Ringer's solution and plated on DeMan, Rogosa, Sharpe (MRS) (lactobacillus) or M17 (coccus) agar (Oxoid Ltd., Basingstoke, Hampshire, England) for viable counts. Mesophilic or thermophilic lactic acid bacteria were enumerated after incubation at 30 or 42°C for 48 to 72 h. At least 10 colonies for each medium and cheese variety, possibly with different morphologies, were isolated from the highest plate dilution. Gram-positive, catalase-negative, nonmotile rod and coccus isolates were cultivated in MRS or M17 broth (Oxoid Ltd.) at 30 or 42°C for 24 h and restreaked into MRS or M17 agar. All the isolates considered for further analyses showed the capacity of acidifying the culture medium. Microbial cultures were stored at -20°C in 10% (vol/vol) glycerol.

**Synthesis of GABA by lactic acid bacteria isolated from cheeses.** Twenty-four-hour-old cells of 440 presumptive lactic acid bacterium strains were harvested by centrifugation ( $9,000 \times g$  for 15 min at 4°C), washed twice with sterile 0.05 M potassium phosphate buffer (pH 7.0), and resuspended in sterile physiological

solution (0.85% NaCl) at an  $A_{620}$  of 2.5, which corresponded to a cell density of ca.  $8.5$  to  $9.0 \log \text{CFU ml}^{-1}$ . GAD activity was measured as described previously (33). The reaction mixture consisted of 900  $\mu\text{l}$  of 50 mM sodium acetate buffer (pH 4.7) containing 2 mM L-glutamate, 0.1 mM pyridoxal phosphate, 100  $\mu\text{l}$  of cell suspension, and 0.05% (wt/vol)  $\text{NaN}_3$  (final concentration). After 24 h at 30°C, the concentration of GABA was determined using a Biochrom 30 series amino acid analyzer (Biochrom Ltd.).

**Genotypic identification by 16S rRNA gene sequence analysis.** Genomic DNA from each strain was extracted as reported by De Los Reyes-Gavilán et al. (10) from 2-ml samples of overnight cultures grown in MRS or M17 broth. Two primer pairs (Invitrogen Life Technologies, Carlsbad, CA), LacbF/LacbR and LpCoF/LpCoR (7), were used to amplify a 16S rRNA gene fragment of presumptive lactic acid bacteria. Fifty microliters of each PCR mixture contained 200  $\mu\text{M}$  of each 2'-deoxynucleoside 5'-triphosphate (dNTP), 1  $\mu\text{M}$  of both forward and reverse primers, 2 mM  $\text{MgCl}_2$ , 2 U of *Taq* DNA polymerase (Invitrogen) in the supplied buffer, and approximately 50 ng of DNA. The expected amplicons of about 1,400 and 1,000 bp (after amplification with primer pairs LacbF/LacbR and LpCoF/LpCoR, respectively) were eluted from the gel and purified by the GFX PCR DNA and Gel Band Purification Kit (Amersham Biosciences, Piscataway, NJ). DNA sequencing reactions were performed by PRIMM srl (San Raffaele Biomedical Science Park, Milan, Italy). Taxonomic strain identification was performed by comparing the sequences of each isolate with those reported in the Basic BLAST database (2).

**Random amplified polymorphic DNA-PCR (RAPD-PCR) analysis.** Genomic DNA was extracted as reported above. Two primers (Invitrogen) with arbitrarily chosen sequences (M13, 5'-GAGGGTGGCGGTTCT-3'; P4, 5'-CCGACGCGTT-3'; and P7, 5'-AGCAGCGTGG-3') (6, 55) were used singly in two series of amplifications. The reaction mixture contained 200  $\mu\text{M}$  of each dNTP, 1 to 2  $\mu\text{M}$  primer, 1.5 to 3  $\mu\text{M}$   $\text{MgCl}_2$ , 1.25 U *Taq* DNA polymerase (Invitrogen), 2.5  $\mu\text{l}$  PCR buffer, 25 ng DNA, and sterile double-distilled water to 25  $\mu\text{l}$ . For amplifications with primer P4, the PCR program comprised 45 cycles of denaturation for 1 min at 94°C, annealing for 1 min at 35°C, and elongation for 2 min at 72°C; the cycles were preceded by denaturation at 94°C for 4 min and followed by elongation at 72°C for 5 min. For primer M13, amplification reactions were performed using one cycle at 94°C for 60 s (denaturing), 42°C for 20 s (annealing), and 72°C for 2 min (elongation). PCR products were separated by electrophoresis (2 h at 130 V) on a 1.5% (wt/vol) agarose gel (Invitrogen), and the DNA was detected by UV transillumination after staining with ethidium bromide (0.5  $\mu\text{g/ml}$ ). The molecular weight of the amplified DNA fragments was estimated by comparison with a 1 Kb Plus DNA Ladder (Invitrogen). Gels were acquired using a UNIsave gel documentation system camera, model GAS9200/1/2/3, version 11 (UVitec Limited, Cambridge, United Kingdom). Electrophoretic profiles were compared using Quantity One software (Bio-Rad, Milan, Italy).

**Acidification kinetics and synthesis of GABA during milk fermentation.** Harvested cells of lactic acid bacteria were washed in 50 mM phosphate buffer (pH 7.0), centrifuged at  $9,000 \times g$  for 15 min at 4°C, and resuspended in reconstituted skimmed milk (RSM) at a cell density of ca.  $6.8 \log \text{CFU ml}^{-1}$ . RSM was supplemented with L-monosodium glutamate (20 mM) and incubated at 30°C for 24 h, and the pH was recorded online. Acidification data were modeled according to the Gompertz equation as modified by Zwietering et al. (57), where  $y$  is  $\log(\Delta\text{pH } \Delta t^{-1})$ , units of  $\text{pH h}^{-1}$ ,  $k$  is the initial level of the dependent variable to be modeled,  $A$  ( $\Delta\text{pH}$ ) is the difference in pH (units) between the initial value ( $\text{pH}_0$ ) and the value reached after 18 h,  $\mu_{\text{max}}$  is the maximum acidification rate ( $\Delta\text{pH h}^{-1}$ ),  $\lambda$  is the length of the latency phase of acidification expressed in hours, and  $t$  is time. After incubation, the pH 4.6-soluble nitrogen fraction was prepared as described elsewhere. Total and individual FAAs contained in the pH 4.6-soluble nitrogen fraction were analyzed using a Biochrom 30 series amino acid analyzer (Biochrom Ltd.).

**Molecular characterization of the GAD gene.** Total DNAs were obtained as described above. Primers designed from highly conserved regions of GAD, CoreF/CoreR (5'-CCTCGAGAAGCCGATCGCTTAGTTCG-3' and 5'-TCATATTGACCGGTATAAGTGATGCC-3', respectively) (PRIMM), were used to amplify genes for GABA-synthesizing enzymes of selected lactic acid bacteria (34, 40). Fifty microliters of each PCR mixture contained 200  $\mu\text{M}$  of each dNTP, 1  $\mu\text{M}$  of both primers, 2  $\mu\text{M}$   $\text{MgCl}_2$ , 2 U of *Taq* DNA polymerase (Invitrogen) in the supplied buffer, and ca. 50 ng of DNA. PCR amplification was performed using the GeneAmp PCR System 9700 thermal cycler (Applied Biosystems). Amplification conditions were changed according to the primer used. PCR products were separated by electrophoresis on a 1.5% (wt/vol) agarose gel (Gibco BRL, France) and stained with ethidium bromide (0.5  $\mu\text{g ml}^{-1}$ ). The amplicons obtained were eluted from the gel and purified by the GFX PCR DNA and Gel Band Purification Kit (Amersham Biosciences). DNA sequencing reactions were performed by PRIMM srl. Sequence comparison was performed by using the

TABLE 1. Species of lactic acid bacteria producing GABA identified in 22 Italian cheese varieties

Cheese	Species
Parmigiano Reggiano	<i>L. plantarum</i> PMR1
Barricato San Martino	<i>L. mesenteroides</i> BSM10
Vento d'Estate	<i>W. cibaria</i> VE10
Ubriaco di Raboso	<i>L. paracasei</i> UB2, <i>L. brevis</i> UB5
Caciocavallo	<i>L. plantarum</i> CCV1 and CCV2, <i>L. casei</i> CCV3
Gorgonzola	<i>S. thermophilus</i> GZ1, GZ2, GZ3, and GZ4
Crescenza	<i>S. thermophilus</i> CRS1
Mozzarella	<i>S. thermophilus</i> MZZ1
Canestrato Pugliese	<i>L. plantarum</i> CNP1
Casciotta di Urbino	<i>L. lactis</i> CU8
Pecorino Piemontese	<i>L. plantarum</i> PPV1 and PPV8, <i>L. pseudomesenteroides</i> PPV6
Pecorino Marchigiano	<i>L. plantarum</i> PM8 and PM13, <i>L. paracasei</i> PM1 and PM4, <i>L. brevis</i> PM17
Pecorino Umbro	<i>L. lactis</i> PU1; <i>L. paracasei</i> PU4, PU18, PU15, and PU20; <i>Lactobacillus</i> sp. strain PU14
Pecorino del Reatino	<i>L. delbrueckii</i> subsp. <i>bulgaricus</i> PR1 and PR7; <i>L. plantarum</i> PR11, PR12, and PR20; <i>L. pseudomesenteroides</i> PR22
Pecorino Sardo	<i>L. paracasei</i> PS1 and PS11, <i>L. plantarum</i> PS15, <i>L. mesenteroides</i> PS4
Pecorino di Filiano	<i>L. paracasei</i> PF6, PF8, and PF13; <i>L. plantarum</i> PF14; <i>Lactobacillus</i> sp. strain PF7; <i>E. durans</i> PF15
Pecorino del Tarantino	<i>L. plantarum</i> PT3, <i>L. paracasei</i> PT4, <i>L. brevis</i> PT6
Pecorino Leccese	<i>L. paracasei</i> PL2, PL4, and PL13; <i>L. plantarum</i> PL7 and PL8
Caprino di Valsassina	<i>L. rhamnosus</i> CV33 and CV36, <i>L. casei</i> CV31
Caprino di Cavalese	<i>L. casei</i> CC21
Flor di Capra	<i>L. casei</i> FC13 and FC15
Capritilla	<i>L. plantarum</i> C48

Basic BLAST database. Translation of nucleotide sequences analyses was performed by using OMIGA software (Oxford Molecular, Madison, WI) or the ExPASy translation routine at the ExPASy Molecular Biology Server of the Swiss Institute of Bioinformatics (<http://ca.expasy.org/>). Similarity researches were performed with the advanced BLAST algorithm available at the National Center for Biotechnology Information site (<http://www.ncbi.nlm.nih.gov/>). Sequence alignments were conducted with the ClustalW algorithm at the ClustalW server at the European Bioinformatics Institute (<http://www.ebi.ac.uk/clustalw/index.html>).

**Resistance to simulated gastric and intestinal fluids.** Simulated gastric and intestinal fluids were created (13, 43). Stationary-phase-grown cells of lactic acid bacteria were harvested at  $9,000 \times g$  for 15 min at 4°C, washed with physiologic solution (0.85% NaCl), and resuspended in 50 ml of simulated gastric juice (ca.  $10 \log \text{CFU ml}^{-1}$ ) which contained NaCl (125 mM liter<sup>-1</sup>), KCl (7 mM liter<sup>-1</sup>), NaHCO<sub>3</sub> (45 mM liter<sup>-1</sup>), and pepsin (3 g liter<sup>-1</sup>) (Sigma) (56). The final pH was adjusted to 2.0, 3.0, and 8.0. The value pH 8.0 was used to investigate the influence of the components of the simulated gastric juice apart from the effect of low pH (13). The effect of nitrite on lactic acid bacteria was also investigated by adding N (1 mg liter<sup>-1</sup>), L-ascorbic acid (5 mg liter<sup>-1</sup>), iodide (1 mg liter<sup>-1</sup>), or thiocyanate (30 mg liter<sup>-1</sup>) (54). The suspension was incubated at 37°C with gentle agitation at 100 revolutions min<sup>-1</sup> (29, 54). Aliquots of this suspension were taken at 0, 90, and 180 min, and viable counts were determined. The effect of gastric digestion was also determined by suspending cells in RSM (25 mg ml<sup>-1</sup>) before inoculation of simulated gastric juice at pH 2.0. The final pH after the addition of RSM was ca. 3.0. This condition was assayed to simulate the effect of the food matrix during gastric transit (56). After 180 min of gastric digestion, cells were harvested and resuspended in simulated intestinal fluid which contained 0.1% (wt/vol) pancreatin and 0.15% (wt/vol) oxgall bile salt (Sigma) at pH 8.0. The suspension was incubated at 37°C under gentle agitation at 100 revolutions min<sup>-1</sup>, and aliquots were taken at 0, 90, and 180 min (13).

**Synthesis of GABA under simulated GI conditions.** RSM (25 mg/ml), containing ca.  $9 \log \text{CFU ml}^{-1}$  of lactic acid bacteria, was subjected to pepsin and pancreatin digestion as described above (43). After digestion, the suspension was further incubated for 24 h at 37°C, at 100 revolutions min<sup>-1</sup>, to mimic the synthesis of GABA by bacteria colonizing the GI tract. Digested samples were recovered after 2, 4, 16, and 24 h of incubation (8) and subjected to free fatty acid

analysis. All treatments were carried out also using RSM without bacterial inoculum as the control.

**Statistical analysis.** Experimental data were subjected to analysis of variance, and pairwise comparison of treatment means was achieved using Tukey's procedure at  $P < 0.05$  using a statistical software program, Statistica for Windows (Statistica 6.0 per Windows 1998). Multivariate statistical analysis and principal component analysis were performed using Statistica 6.0 and Statistical software for MS Excel per Windows 1998.

**Nucleotide sequence accession numbers.** The sequences in this study were deposited in GenBank, National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). The accession numbers were EF174472, EF174473, EF174474, and EF174475 for the GAD gene fragments of *Lactobacillus paracasei* PF6, *Lactobacillus delbrueckii* subsp. *bulgaricus* PR1, *Lactobacillus plantarum* FC48, and *Lactococcus lactis* PU1, respectively.

RESULTS

**Concentrations of GABA in cheeses.** Twenty-two Italian cheese varieties having features that may influence directly or indirectly the synthesis of GABA were considered in this study. The values of pH, moisture, and pH 4.6-soluble N were in the range of 4.68 to 6.70, 30 to 60%, and 7.0 to 34%, respectively. The concentrations of total FAAs and glutamate, contained in the pH 4.6-soluble N extracts of the 22 Italian cheese varieties, markedly varied from 19.17 to 231,800.0 mg kg<sup>-1</sup> and from 0.0 to 7,190.0 mg kg<sup>-1</sup>, respectively. The concentration of GABA varied from 0.260 to 391 mg kg<sup>-1</sup>. The lowest concentrations were found for Vento d'Estate (0.260 mg kg<sup>-1</sup>), Mozzarella (0.32 mg kg<sup>-1</sup>), and Crescenza (0.77 mg kg<sup>-1</sup>) cheeses. The majority of the cheeses (Parmigiano Reggiano, Barricato San Martino, Ubriaco di Raboso, Caciocavallo, Gorgonzola, Canestrato Pugliese, Casciotta di Urbino, Pecorino del Tarantino, Pecorino Piemontese, Flor di Capra, Caprino di Cavalese, Caprino di Valsassina, and Capritilla) had concen-




FIG. 1. Score plot of the first and second principal components after principal component analysis based on the concentrations of GABA, glutamic acid, and total FAAs of RSM started with lactic acid bacteria. Each fermented milk is indicated by the code of the lactic acid bacterium used for fermentation. The values are the averages of three batches of each milk fermentation.

```

C48      TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 56
PF6      TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 56
PU1      LFVRFPIWNFXAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 57
WCFS1    TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 56
PR1      TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 56
ATCC367  TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PEAESFTG 56
I11403   TFCQTYMEFEAVEIMKMDTLAKNAIDKSEYPTAEIENRCVNIIANLWHA--PDEHFTG 56
F6854    TFCQTYMEDEATKLMSETLEKNAIDKSEYPTAELENRCVNIADLWHA--PKDQKFMG 56
13       TFCQTYMDEEAVKLMMAETLEKNAIDKSEYPTAELENRCVNIADLWHA--PKDESFTG 56
DO       TFCQTYMEFEAVKLMTQTLEKNAIDKSEYPTAEIENRCVNIADLWHA--FNNEKFMG 56
100-23   TFCQTYMEFKATQMAETMOKNAIDKSEYPTAELENRCVNIIAKLWHA--QKDEEYMG 56
CFT073   TFCQTMDDDNVHKLMDLSINKNMTDKSEYPTAELENRCVNIADLWHA--PKDQKFMG 58
          * : . : . : : * : : ** * : : * : : : : : : : : : : : : : : : : *
          : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : :

C48      TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----FAGKSFVS 110
PF6      TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----VCWEKFCV 110
PU1      TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----VCWEKICG 111
WCFS1    TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----VCWEKFCV 110
PR1      TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----VCWEKFCV 110
ATCC367  TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----VCWEKFCV 110
I11403   TSTIGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQFAGKSFVCWEKFCV 116
F6854    TSTIGSSEACMLGGMAMKFAWRKRAEKGLDLIYAKKPNLVISAGYQ-----VCWEKFCV 110
13       TSTVGSSEACMLGGMAMKFAWRKRAEKGLDVTSRKPNLVISAGYQ-----VCWEKFCV 110
DO       TSTIGSSEACMLGGMAMKFAWRKRAEKGLDIQAKKPNLVISAGYQ-----VCWEKFCV 110
100-23   TSTVGSSEACMLGGLAMKFAMRKRQAAGLDLNAHRPNLVISAGYQ-----ICWEKFCV 110
CFT073   TNTIGSSEACMLGGMAMKFAWRKRAEKGLDVTSRKPNLVISAGYQ-----ICWEKFCV 109
          * : : : : * : : * : : * : : : : : : : : : : : : : : : : : : : : : : :
          : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : : :

C48      TGTLTCTFFSNGMSNTWPLTLTXSXTXXNXYTIXIVRXHGASIXP--- 153
PF6      YWDVVFVFFFSQWMSNTWPLGLNTS-TAWAENRLVFFCKQGVPTPA-- 153
PU1      YWDDDMYLAFL--MDGQH-LALDVEQIFASTWSNTAFGLNTVTTDLLFC 156
WCFS1    YWDVDMHV-VFMDEQHMALDVNHVLDYVDEYTIIGIVGIMGITYTGQYD 155
PR1      YWDVDMHV-VFMDEQHMALDVNHVLDYVDEYTIIGIVGIMGITYTGSII- 154
ATCC367  YWDIDMHV-VFMDDHMSLNVDHVLDYVDDYTIIGIVGIMGITYTGQYD 155
I11403   YWDIDMRV-VFMDKEHMSINLDKVMYDVDEYTIIGVVGILGITYTGRYD 161
F6854    YWDIDMRV-VFMDKEHMQLNLDQVLDYVDEYTIIGVVGILGITYTGRYD 155
13       YWDIDMRV-VFMDKEHMSINLDKVMYDVDEYTIIGVVGILGITYTGRYD 155
DO       YWDVRELVRE-VFMDEKHMSINLDVMDYVDEYTIIGIVGIMGITYTGRYD 155
100-23   YFDVRELVRE-VFMDEEHQSLNMNTVMDYVDEYTIIGIVGIMGITYTGRYD 155
CFT073   YWDVRELVRE-VFMRFGQLFMDPKRRIEACDENTIGVVFVTVGVYTGNYE 154
          :
          :
          :

```

FIG. 2. Alignment of the internal deduced amino acid sequences of GAD of *Lactobacillus paracasei* PF6, *Lactobacillus delbrueckii* subsp. *bulgaricus* PR1, *Lactococcus lactis* PU1, and *Lactobacillus plantarum* C48 with other similar GAD sequences from *Lactobacillus plantarum* WCFS1 (accession number NP\_786643.1), *Lactobacillus brevis* ATCC 367 (accession number YP\_795941.1), *Lactococcus lactis* subsp. *lactis* I11403 (accession number NP\_267446.1), *Listeria monocytogenes* strain F6854 (accession number ZP\_00234896.1), *Clostridium perfringens* strain 13 (accession number NP\_562974.1), *Enterococcus faecium* DO (accession number ZP\_00603789.1), *Lactobacillus reuteri* 100-23 (accession number ZP\_01274543.1), and *Escherichia coli* CFT073 (accession number NP\_753818.1). The deduced amino acid sequence was analyzed using ClustalW (1.81).

trations of GABA which ranged from 4 to 100 mg kg<sup>-1</sup>. The highest values of GABA were found in Pecorino Marchigiano (289 mg kg<sup>-1</sup>), Pecorino del Reatino (290 mg kg<sup>-1</sup>), Pecorino Leccese (290 mg kg<sup>-1</sup>), Pecorino Umbro (330 mg kg<sup>-1</sup>), and, especially, Pecorino di Filiano (391 mg kg<sup>-1</sup>). The values of pH for the cheeses containing the highest concentrations of GABA were 4.68 to 5.70. Overall, no statistical correlation was found between the concentration of GABA and the levels of L-glutamate or other free fatty acids. As determined by multivariate statistical analysis, the use of sheep milk and cheese ripening for 1.5 to 5 months seemed to be related to the highest concentration of GABA. The other technological traits or cheese characteristics did not show a statistical correlation.

**Synthesis of GABA by lactic acid bacteria isolated from cheeses.** The number of presumptive mesophilic and thermophilic lactobacilli in cheeses varied from 6.90 to 8.51 log CFU g<sup>-1</sup> and from ≤3.0 to 6.28 log CFU g<sup>-1</sup>, respectively. Lactococci and streptococci were ≤3.0 to 7.8 log CFU g<sup>-1</sup> and ≤3.0 to 5.43 log CFU g<sup>-1</sup>, respectively. As expected, the highest

numbers of thermophilic lactobacilli and streptococci were found in cheeses where primary starters were used. For each cheese, 20 gram-positive, catalase-negative, nonmotile, and acidifying isolates (total of 440) were randomly taken from the plates containing the highest dilutions and screened for GAD activity. Only 61 isolates synthesized GABA under in vitro conditions at 30°C for 24 h, pH 4.7. All these isolates were further identified by partial sequencing of the 16S rRNA gene (Table 1). Twelve species were found: *Lactobacillus plantarum* (17 isolates), *Leuconostoc mesenteroides* (two), *Weissella cibaria* (one), *Lactobacillus paracasei* (16), *Lactobacillus brevis* (three), *Lactobacillus casei* (five), *Streptococcus thermophilus* (six), *Leuconostoc pseudomesenteroides* (two), *Lactococcus lactis* (two), *Lactobacillus delbrueckii* subsp. *bulgaricus* (two), *Enterococcus durans* (one), and *Lactobacillus rhamnosus* (two). Two isolates of unidentified *Lactobacillus* species were also found. Except for Gorgonzola, Pecorino cheeses especially contained the highest numbers of GABA-producing strains (four to six). Apart from the cheese source, 17 and 16 isolates

corresponded to the species *L. plantarum* and *L. paracasei*, respectively. Overall, 53 isolates were mesophilic lactic acid bacteria (45 mesophilic lactobacilli) and only eight were thermophilic lactic acid bacteria (six *S. thermophilus* and two *L. delbrueckii* subsp. *bulgaricus* strains). To exclude clonal relatedness, three primers, M13, P4, and P7, with arbitrarily chosen sequences, were used to characterize the 61 GABA-producing isolates by RAPD-PCR analysis. The reproducibility of RAPD fingerprints was assessed by comparing the PCR products obtained from three separate cultures of the same strain. The size of the amplified fragments ranged from 100 to 1,000 bp, and the number of fragments varied from one to seven per primer per isolate. The RAPD profiles generated by the above primers were highly discriminative and reproducible with consistent fragment patterns. The polymorphic bands in total distinguished the isolates from each other with at least a 2.5% dissimilarity level (data not shown). The capacity for synthesizing GABA of the 61 isolates was further assayed in RSM. After 24 h of fermentation at 30°C, all isolates reached the cell density of 8.5 to 9.0 log CFU g<sup>-1</sup>. The kinetics of acidification was characterized by a median value of  $\Delta$ pH of 2.28. The majority of the strains caused  $\Delta$ pHs which ranged from 2.00 to 2.71. The median value for  $\mu_{\max}$  was 0.03  $\Delta$ pH min<sup>-1</sup>. The median value for the concentration of GABA was 1.71 mg kg<sup>-1</sup>, and the 25th and 75th percentiles of the data ranged from 1.0 to 3.67 mg kg<sup>-1</sup>, respectively. *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, *L. plantarum* C48, and *L. brevis* PM17, which showed the highest synthesis of GABA (99.9, 63.0, 36.0, 16.0, and 15.0 mg kg<sup>-1</sup>, respectively), were located out of the error bars of the box plot (data not shown). Overall, the concentration of total FAAs in the fermented RSM showed increases which ranged from 120 to 390.6 mg kg<sup>-1</sup>. The concentrations of GABA, glutamic acid, and total FAAs of each fermented RSM were subjected to principal component analysis using a covariance matrix (Fig. 1). The first two principal components explained ca. 83.26% of the total variance. PC1 showed the distribution of the samples according to the concentration of FAAs, while PC2 differentiated samples based on the concentration of GABA. The fermented RSMs started with *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, *L. plantarum* C48, or *L. brevis* PM17 were characterized by the highest synthesis of GABA and were located in different zones of the plane delimited by the two principal components.

**Molecular characterization of the GAD genes.** *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48, corresponding to the highest GABA-producing strains, were screened for GAD genes. Primers designed from highly conserved region of GAD genes gave a PCR product of approximately 540 bp for all the strains. The predicted amino acid sequences of *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48 showed 98, 100, 86, and 95% identity to *gadB* of *L. plantarum* WCFS1 (accession number NP\_786643.1), respectively. Alignments of the internal GAD gene fragment of *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48 with the eight most similar sequences of the GAD gene are shown in Fig. 2.

**Resistance to simulated gastric and intestinal fluids of GABA-producing strains.** *L. paracasei* PF6, *L. delbrueckii* subsp.




FIG. 3. Survival of selected *Lactobacillus paracasei* PF6 (A), *Lactobacillus delbrueckii* subsp. *bulgaricus* PR1 (B), and *Lactobacillus plantarum* C48 (C) under gastric conditions (0 to 180 min) at pH 8.0 ( $\blacktriangledown$ ), 3.0 ( $\blacksquare$ ), and 2.0 ( $\square$ ) and at 2.0 with RSM added (11%, wt/vol) ( $\nabla$ ) and with further intestinal digestion (180 to 360 min) at pH 8.0. The values were the averages of three replicates, and standard deviations are indicated by vertical bars.

*bulgaricus* PR1, *L. lactis* PU1, and *L. plantarum* C48 were incubated at 37°C in simulated gastric fluid at pH 2.0, 3.0, and 8.0. *L. lactis* PU1 was the only strain which had a very poor survival in all the conditions assayed (data not shown). Therefore, it was excluded from further characterization. After 180 min of incubation in simulated gastric juice at pH 3.0, all the other strains showed decreases lower than 1 log cycle with respect to their initial cell density (10 to 9.6 log CFU ml<sup>-1</sup>). At pH 8.0 no decrease of survival was found for all strains (Fig. 3A to C). After 180 min at pH 2.0, all strains decreased to 6.0 to 5.0 log CFU ml<sup>-1</sup>. When RSM (25 mg/ml) was added to the juice at pH 2.0, lactobacilli were resistant to the simulated gastric juice. Probably, this was due to the increased value of pH (3.0)




FIG. 4. Synthesis of GABA of *Lactobacillus paracasei* PF6 (◆), *Lactobacillus delbrueckii* subsp. *bulgaricus* PR1 (■), and *Lactobacillus plantarum* C48 (▲) in RSM after incubation under GI conditions and further incubation at 37°C for 24 h. The values were the averages of three replicates, and standard deviations are indicated by vertical bars.

caused by addition of RSM or to the direct protective effect on microbial cells by the food matrix. After 180 min of gastric digestion, cells were exposed to simulated intestinal fluid for a subsequent 180 min at pH 8.0 (Fig. 3A to C). Cell survival depended on the pH of gastric digestion. The decrease for cells previously treated at pH 8.0, 3.0, or 2.0 with the addition of RSM was always lower than 1 log cycle. When the gastric digestion was at an initial pH of 2.0, the survival of simulated intestinal fluid was approximately 3 and 4 log CFU ml<sup>-1</sup> for *L. paracasei* PF6 and *L. plantarum* C48, respectively. Nonculturable cells of *L. delbrueckii* subsp. *bulgaricus* PR1 were found in 1 ml of sample.

**Synthesis of GABA under GI conditions.** The hypothesis to be investigated concerned the capacity of selected strains to synthesize GABA during gut transit and without the addition of L-glutamate. RSM containing 9 log CFU ml<sup>-1</sup> of *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, or *L. plantarum* C48 was subjected to sequential hydrolysis by pepsin (pH 2.0) and pancreatin (pH 8.0) as described above. After digestion, the suspensions were further incubated for 24 h at 37°C under stirring conditions, to mimic the bacterial GI transit. As previously shown (Fig. 3A to C), RSM protected the cells and the numbers of the three strains did not vary with respect to the initial cell density. Nevertheless, all strains synthesized GABA at a lower concentration than that found in fermented RSM (30°C for 24 h, initial pH 6.6) (Fig. 4). Among the three strains, *L. paracasei* PF6 showed the highest synthesis of GABA, reaching ca. 20 mg kg<sup>-1</sup> after 24 h of incubation.

## DISCUSSION

Milk and dairy products provide a rich source of valuable proteins, minerals, and vitamins. The nutritional significance of proteins includes macronutrient as well as physiological and functional aspects. Besides bioactive proteins, dairy products may also provide bioactive peptides (15, 45). First, this study reports the presence of GABA in 22 Italian cheese varieties. Overall, cheeses characterized by the shortest ripening (Mozzarella and Crescenza), nonconventional ripening (Barricato San Martino and Vento d'Estato) (11), or the longest ripening (Canestrato Pugliese and Parmigiano Reggiano) (14) con-

tained the lowest concentrations of GABA. Among the technological traits differentiating the cheeses, only the type of milk and duration of ripening were statistically correlated with the concentration of GABA. Indeed, sheep cheeses ripened for 5 months such as Pecorino Marchigiano, Pecorino del Reatino, Pecorino Leccese, Pecorino Umbro, and Pecorino di Filiano contained the highest levels of GABA (289 to 391 mg kg<sup>-1</sup>). Nomura et al. (32) also analyzed seven commercial cheeses (Camembert, Gouda, blue, cream, Cheddar, Edam, and Emental) and reported concentrations of GABA of 177.0, 48.0, 7.1, and 4.2 mg kg<sup>-1</sup> for Gouda, Cheddar, blue, and Edam, respectively. All of the 22 Italian cheese varieties contained relevant numbers of mesophilic nonstarter lactic acid bacteria, as members of the autochthonous microbiota, and thermophilic lactic acid bacteria, when used as primary starters. Nevertheless, only a small part (ca. 14%) of the total 440 isolates (20 for each cheese variety) showed the capacity for synthesizing GABA under in vitro conditions and during RSM fermentation. Preliminarily, RAPD-PCR analyses showed genetic variability between the GABA-producing strains used in this study. RAPD-PCR based on at least three primers has been shown to be an effective tool in resolving intraspecific differences also for lactic acid bacteria (6, 55). Interestingly, the best GABA-producing strains, *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, and *L. brevis* PM17, were isolated from Pecorino di Filiano, Pecorino del Reatino, Pecorino Umbro, and Pecorino Marchigiano cheeses, respectively, which had the highest concentrations of GABA. The only exception was *L. plantarum* C48 isolated from Capritilla goat cheese. Other reports also showed the synthesis of GABA by primary starters such as *L. lactis*, *S. thermophilus*, and *L. delbrueckii* subsp. *bulgaricus* (17, 32–34) and nonstarter lactic acid bacteria such as *L. plantarum*, *L. paracasei*, and *L. brevis* (17, 27, 42). Strains of *L. plantarum* and *L. paracasei* corresponded to the major part of the GABA-producing isolates from the 22 Italian cheese varieties. Nevertheless, the capacity of synthesizing the highest levels of GABA was markedly strain dependent. Apart from other technological conditions, the presence of strains with particular aptitudes for synthesizing GABA during cheese ripening seemed to be the major prerequisite for the manufacture of GABA-enriched cheeses. To

the best of our knowledge, this report is the first to show the synthesis of GABA also by *L. casei*, *L. rhamnosus*, *W. cibaria*, *L. mesenteroides*, *L. pseudomesenteroides*, and *E. durans*.

The genetics of GABA have been elucidated in *Escherichia coli* (50), *Lactococcus lactis* subsp. *lactis* (34), and *L. brevis* (42). Sanders et al. (47) sequenced the *L. lactis* subsp. *lactis* *gadCB* gene and suggested that it encoded a glutamate-dependent acid resistance mechanism comprised of glutamate-GABA antiporter and GAD. Nomura et al. (34) indicated that *L. lactis* subsp. *lactis* contains a single GAD gene (*gadB*), while the gram-negative *E. coli* (50) and *Shigella* sp. (49) contain two GAD genes. The functional properties of the two *E. coli* isozymes were identical (9). The partial GAD sequences found in this study showed high identity with *gadB* sequences from *L. plantarum* WCFS1 (accession number NP\_786643.1), *L. brevis* ATCC 367 (accession number YP\_795941.1), *L. lactis* subsp. *lactis* III403 (accession number NP\_267446.1), *Listeria monocytogenes* strain F6854 (accession number ZP\_00234896.1), *Enterococcus faecium* DO (accession number ZP\_00603789.1), *Lactobacillus reuteri* 100-23 (accession number ZP\_01274543.1), *Clostridium perfringens* strain 13 (accession number NP\_562974.1), and *E. coli* CFT073 (accession number NP\_753818.1). Physiologically, the expression of GAD genes is assumed to control the acidification of the cytosolic environment by decarboxylating an acid substrate (glutamate) into a neutral compound (GABA) via incorporation of H<sup>+</sup>. GABA would then be exported into the extracellular environment, thereby contributing to alkalinization (for a review see reference 5). The partial GAD sequences found in this study could facilitate further studies of gene expression (42).

During RSM fermentation at 30°C for 24 h, *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, *L. lactis* PU1, *L. plantarum* C48, and *L. brevis* PM17 synthesized concentrations of GABA (99.9 to 15 mg kg<sup>-1</sup>) higher than those found for other cheese starters in skim milk (32, 33) and *Bifidobacterium longum* (41). A new type of GABA-enriched fermented milk was manufactured by using two starters: *L. casei* and *L. lactis*. *L. casei* hydrolyzed milk proteins into L-glutamate, and *L. lactis* subsp. *lactis* converted it into GABA (20). Potentially, the lactic acid bacteria selected in this study could be used in mixture for the manufacture of a fermented milk containing elevated levels of GABA. Foods enriched with GABA were defined as “foods for specified health use” by the Japanese government (48). Recently, some studies reported that dietary materials or products containing GABA caused a decrease of blood pressure in spontaneously hypertensive rats and hypertensive humans (20, 25, 31). A daily intake of fermented milk (10 mg of GABA) for 12 weeks decreased blood pressure by 17.4 Hg in hypertensive patients (20, 25). The amount of GABA in 100 g fermented milk synthesized by *L. paracasei* PF6 still represents the minimum effective daily dose to get positive effects. Some of the selected strains also showed tolerance and synthesis of GABA under simulated GI conditions. Acid and bile salt treatments were combined in this study, since they have both individual and combined effects (4). The time chosen for treatments in simulated gastric (180 min) and intestinal (further 180 min) fluids mimicked the in vivo times between entrance to and release from the stomach and intestine during digestive processes (4). *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, and *L. plantarum* C48 showed high resistance at pH 2.0 when RSM was added to acid and bile

salt fluids. The synthesis of GABA by *L. paracasei* PF6, *L. delbrueckii* subsp. *bulgaricus* PR1, and *L. plantarum* C48 was also found in RSM under simulated GI conditions and without addition of L-glutamate as the precursor. First, this study showed the synthesis of GABA by lactic acid bacteria under GI conditions. Overall, the screening of lactic acid bacteria based on the capacity to synthesize GABA under GI conditions may be considered another biotechnological trait to select probiotics. Fermented milk enriched in GABA by lactobacilli selected in this study would be of interest for the food industry as it could be considered a health-oriented dairy product with a potential antihypertensive effect. Further studies are in progress to show the decrease of blood pressure in spontaneously hypertensive rats and hypertensive humans.

#### REFERENCES

- Adeghate, E., and A. S. Ponery. 2002. GABA in the endocrine pancreas: cellular localization and function in normal and diabetic rats. *Tissue Cell* **34**:1–6.
- Altschul, S. F., T. L. Madden, A. A. Schäffer, J. Zhang, Z. Zhang, W. Miller, and D. J. Lipman. 1997. Grapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **25**:3389–3402.
- Aoki, H., Y. Furuya, Y. Endo, and K. Fujimoto. 2003. Effect of gamma-aminobutyric acid-enriched tempeh-like fermented soybean (GABA-tempeh) on the blood pressure of spontaneously hypertensive rats. *Biosci. Biotechnol. Biochem.* **67**:1806–1808.
- Chou, L., and B. Weimer. 1999. Isolation and characterization of acid- and bile-tolerant isolates from strains of *Lactobacillus acidophilus*. *J. Dairy Sci.* **82**:23–31.
- Cotter, P. D., and C. Hill. 2003. Surviving the acid test: responses of gram-positive bacteria to low pH. *Microbiol. Mol. Biol. Rev.* **67**:429–453.
- De Angelis, M., A. Corsetti, N. Tosti, J. Rossi, M. R. Corbo, and M. Gobbetti. 2001. Characterization of non-starter lactic acid bacteria from Italian ewe cheeses based on phenotypic, genotypic, and cell wall protein analyses. *Appl. Environ. Microbiol.* **67**:2011–2020.
- De Angelis, M., S. Siragusa, M. Berloco, L. Caputo, L. Settanni, G. Alfonsi, M. Amerio, A. Grandi, A. Ragni, and M. Gobbetti. 2006. Isolation, identification and selection of potential probiotic lactobacilli from pig faeces to be used as additives in pelleted feeding. *Res. Microbiol.* **157**:792–801.
- De Angelis, M., C. G. Rizzello, E. Scala, C. De Simone, G. A. Farris, F. Turrini, and M. Gobbetti. 2007. Probiotic preparation has the capacity to hydrolyze proteins responsible for wheat allergy. *J. Food Prot.* **70**:135–144.
- De Biase, D., A. Tramonti, R. A. John, and F. Bossa. 1996. Isolation, over-expression, and biochemical characterization of the two isoforms of glutamic acid decarboxylase from *Escherichia coli*. *Protein Expr. Purif.* **8**:430–438.
- De Los Reyes-Gavilán, C. G., G. K. Y. Limsowtin, P. Tailliez, L. Séchaud, and J. P. Acolas. 1992. A *Lactobacillus helveticus*-specific DNA probe detects restriction fragment length polymorphisms in this species. *Appl. Environ. Microbiol.* **58**:3429–3432.
- Di Cagno, R., S. Buchin, S. de Candia, M. De Angelis, P. F. Fox, and M. Gobbetti. 2007. Characterization of Italian cheeses ripened under non-conventional conditions. *J. Dairy Sci.* **90**:2689–2704.
- Diplock, A. T., P. J. Aggett, M. Ashwell, F. Bornet, E. B. Fern, and M. B. Roberfroid. 1999. Scientific concepts of functional foods in Europe—consensus document. *Br. J. Nutr.* **81**:1–27.
- Fernández, M. F., S. Boris, and C. Barbés. 2003. Probiotic properties of human lactobacilli strains to be used in the gastrointestinal tract. *J. Appl. Microbiol.* **94**:449–455.
- Gobbetti, M., M. De Angelis, R. Di Cagno, and C. G. Rizzello. 2007. The relative contributions of starter cultures and non-starter bacteria to the flavour of cheese, p. 121–156. *In* B. Weimer (ed.), *Improving the flavour of cheese*. Woodhead Publishing, Cambridge, England.
- Gobbetti, M., L. Stepaniak, M. De Angelis, A. Corsetti, and R. Di Cagno. 2002. Latent bioactive peptides in milk proteins: proteolytic activation and significance in dairy processing. *Crit. Rev. Food Sci. Nutr.* **42**:223–239.
- Hagiwara, H., T. Seki, and T. Ariga. 2004. The effect of pre-germinated brown rice intake on blood glucose and PAI-1 levels in streptozotocin-induced diabetic rats. *Biosci. Biotechnol. Biochem.* **68**:444–447.
- Hayakawa, K., Y. Ueno, S. Kawamura, R. Taniguchi, and K. Oda. 1997. Production of  $\gamma$ -aminobutyric acid by lactic acid bacteria. *Seibutsu-kogaku Kaishi* **75**:239–244.
- Hayakawa, K., M. Kimura, K. Kasaha, K. Matsumoto, H. Sansawa, and Y. Yamori. 2004. Effect of a gamma-aminobutyric acid-enriched dairy product on the blood pressure of spontaneously hypertensive and normotensive Wistar-Kyoto rats. *Br. J. Nutr.* **92**:411–417.
- Higuchi, T., H. Hayashi, and K. Abe. 1997. Exchange of glutamate and  $\gamma$ -aminobutyrate in a *Lactobacillus* strain. *J. Bacteriol.* **179**:3362–3364.

20. Inoue, K., T. Shirai, H. Ochiai, M. Kasao, K. Hayakawa, M. Kimura, and H. Sansawa. 2003. Blood-pressure-lowering effect of a novel fermented milk containing gamma-aminobutyric acid (GABA) in mild hypertensives. *Eur. J. Clin. Nutr.* **57**:490–495.
21. International Dairy Federation. 1964. Determination of the protein content of processed cheeses products. Standard 25. International Dairy Federation, Brussels, Belgium.
22. International Dairy Federation. 1970. Determination of dry matter content in whey cheese. Standard 58. International Dairy Federation, Brussels, Belgium.
23. International Dairy Federation. 1989. Determination of pH. Standard 115A. International Dairy Federation, Brussels, Belgium.
24. Jakobs, C., J. Jaeken, and K. M. Gibson. 1993. Inherited disorders of GABA metabolism. *J. Inher. Metab. Dis.* **16**:704–715.
25. Kajimoto, O., H. Hirata, S. Nakagawa, Y. Kajimoto, K. Hayakawa, and M. Kimura. 2004. Hypotensive effect of fermented milk containing  $\gamma$ -aminobutyric acid (GABA) in subjects with high normal blood pressure. *Nippon Shokuhin Kagaku Kogaku Kaishi* **51**:79–86.
26. Kinefuchi, M., M. Sekiya, A. Yamazaiki, and K. Yamamoto. 1999. Accumulation of GABA in brown rice by high pressure treatment. *Nippon Shokuhin Kagaku Kaishi* **46**:323–328.
27. Komatsuzaki, N., J. Shima, S. Kawamoto, H. Monose, and T. Kimura. 2005. Production of  $\gamma$ -aminobutyric acid (GABA) by *Lactobacillus paracasei* isolated from traditional fermented foods. *Food Microbiol.* **22**:497–504.
28. Leroy, F., and L. de Vuyst. 2004. Lactic acid bacteria as functional starter cultures for the food fermentation industry. *Trends Food Sci. Technol.* **15**:67–78.
29. Mao, Y., M. P. Doyle, and J. Chen. 2006. Role of colonic acid exopolysaccharide in the survival of enterohaemorrhagic *Escherichia coli* O157:H7 in simulated gastrointestinal fluids. *Lett. Appl. Microbiol.* **42**:642–647.
30. Nagaoka, H. 2005. Treatment of germinated wheat to increase levels of GABA and IP6 catalyzed by endogenous enzymes. *Biotechnol. Prog.* **21**:405–410.
31. Nakamura, Y., N. Yamamoto, K. Skai, and T. Takano. 1995. Antihypertensive effect of sour milk and peptides isolated from it that are inhibitors to angiotensin I-converting enzyme. *J. Dairy Sci.* **78**:1253–1257.
32. Nomura, M., H. Kimoto, Y. Someya, S. Furukawa, and I. Suzuki. 1998. Production of  $\gamma$ -aminobutyric acid by cheese starters during cheese ripening. *J. Dairy Sci.* **81**:1486–1491.
33. Nomura, M., H. Kimoto, Y. Someya, S. Furukawa, and I. Suzuki. 1999. Novel characterization for distinguishing *Lactococcus lactis* subsp. *lactis* from subsp. *cremoris*. *Int. J. Syst. Bacteriol.* **49**:163–166.
34. Nomura, M., I. Nakajima, Y. Fujita, M. Kobayashi, H. Kimono, I. Suzuki, and H. Aso. 1999b. *Lactococcus lactis* contains only glutamate decarboxylase gene. *Microbiology* **154**:1375–1380.
35. Oh, S. H. 2003. Stimulation of gamma-aminobutyric acid synthesis activity in brown rice by a chitosan/Glu germination solution and calcium/calmodulin. *J. Biochem. Mol. Biol.* **36**:319–325.
36. Oh, S. H., and C. H. Oh. 2003. Brown rice extracts with enhanced levels of GABA stimulate immune cells. *Food Sci. Biotechnol.* **12**:248–252.
37. Oh, S. H., J. R. Soh, and Y. S. Cha. 2003. Germinated brown rice extract shows a nutraceutical effect in the recovery of chronic alcohol-related symptoms. *J. Med. Food* **6**:115–121.
38. Ohmori, M., T. Yano, J. Okamoto, T. Tsushida, T. Murai, and M. Higuchi. 1987. Effect of anaerobically treated tea on blood pressure of spontaneous hypertensive rats. *Nippon Noeikagaku Kaishi* **61**:1449–1451.
39. Okada, T., T. Sugishita, T. Murakami, H. Murai, T. Saikusa, T. Hotorino, A. Onoda, O. Kajimoto, R. Takahashi, and T. Takahashi. 2000. Effect of the defatted rice germ enriched with GABA for sleeplessness, depression, autonomic disorder by oral administration. *Nippon Shokuhin Kagaku Kaishi* **47**:596–603.
40. Park, K. B., and S. H. Oh. 2004. Cloning and expression of a full-length glutamate decarboxylase gene from *Lactobacillus plantarum*. *J. Food Sci. Nutr.* **9**:324–329.
41. Park, K. B., and S. H. Oh. 2005. Production and characterization of GABA rice yogurt. *J. Food Sci. Biotechnol.* **14**:518–522.
42. Park, K. B., and S. H. Oh. 2007. Cloning, sequencing and expression of a novel glutamate decarboxylase gene from a newly isolated lactic acid bacterium, *Lactobacillus brevis* OPK-3. *Bioresour. Technol.* **98**:312–319.
43. Pasini, G., M. Simonato, M. Giannattasio, A. D. B. Peruffo, and A. Curioni. 2001. Modification of wheat flour proteins during in vivo digestion of bread dough, crumb and crust: an electrophoretic and immunological study. *J. Agric. Food Chem.* **49**:2254–2259.
44. Rhyu, M. R., E. Y. Kim, H. Y. Kim, B. H. Ahn, and C. B. Yang. 2000. Characteristics of the red rice fermented with fungus *Monascus*. *Food Sci. Biotechnol.* **9**:21–26.
45. Rizzello, C. G., I. Losito, M. Gobbetti, T. Carbonara, M. D. De Bari, and P. G. Zambonin. 2005. Antibacterial activities of peptides from the water-soluble extracts of Italian cheese varieties. *J. Dairy Sci.* **88**:2348–2360.
46. Saikusa, T., T. Horino, and Y. Mori. 1994. Accumulation of  $\gamma$ -aminobutyric acid (GABA) in the rice germ during water soaking. *Biosci. Biotechnol. Biochem.* **58**:2291–2292.
47. Sanders, J. W., K. Leenhouts, J. Burghoorn, J. R. Brands, G. Venema, and J. Kok. 1998. A chloride-inducible acid resistance mechanism in *Lactococcus lactis* and its regulation. *Mol. Microbiol.* **27**:299–310.
48. Sanders, M. E. 1998. Overview of functional foods: emphasis on probiotic bacteria. *Int. Dairy J.* **8**:341–347.
49. Small, P. L. C., and S. R. Waterman. 1998. Acid stress, anaerobiosis and *gadCB*: lessons from *Lactococcus lactis* and *Escherichia coli*. *Trends Microbiol.* **6**:214–216.
50. Smith, D. K., T. Kassam, B. Singh, and J. F. Elliott. 1992. *Escherichia coli* has two homologous glutamate decarboxylase genes that map to distinct loci. *J. Bacteriol.* **174**:5820–5826.
51. Ueno, H. 2000. Enzymatic and structural aspects on glutamate decarboxylase. *J. Mol. Catal.* **10**:67–79.
52. Ueno, Y., K. Hayakawa, S. Takahashi, and K. Oda. 1997. Purification and characterization of glutamate decarboxylase from *Lactobacillus brevis* IFO 12005. *Biosci. Biotechnol. Biochem.* **61**:1168–1171.
53. Wong, C. G., T. Bottiglieri, and O. C. Snead III. 2003. GABA,  $\gamma$ -hydroxybutyric acid, and neurological disease. *Ann. Neurol.* **54**(Suppl. 6):S3–S12.
54. Xu, J., X. Xu, and W. Verstraete. 2001. The bactericidal effect and chemical reactions of acidified nitrite under conditions simulating the stomach. *J. Appl. Microbiol.* **90**:523–529.
55. Zapparoli, G., S. Torriani, and F. Dellaglio. 1998. Differentiation of *Lactobacillus sanfranciscensis* strains by randomly amplified polymorphic DNA and pulsed-field gel electrophoresis. *FEMS Microbiol. Lett.* **166**:324–332.
56. Zárate, G. A., S. P. Chaia, and G. O. González. 2000. Viability and  $\beta$ -galactosidase activity of dairy propionibacteria subjected to digestion by artificial gastric and intestinal fluids. *J. Food Prot.* **63**:1214–1221.
57. Zwietering, M. H., I. Jongeberger, F. M. Roumbouts, and K. van 't Riet. 1990. Modeling of the bacterial growth curve. *Appl. Environ. Microbiol.* **56**:1875–1881.